

# CONTRIBUTIONS TO THE THEORY OF NONLINEAR OSCILLATIONS

---

## *VOLUME V*

---

|               |                     |
|---------------|---------------------|
| J. P. LASALLE | L. MARKUS           |
| J. JARNIK     | G. REEB             |
| J. KURZWEIL   | P. MENDELSON        |
| J. CRONIN     | J. ANDRÉ            |
| J. K. HALF    | P. SEIBERT          |
| L. CESARI     | G. COLEMAN          |
| A. STOKES     | R. P. DE FIGUEIREDO |

EDITED BY

L. Cesari, J. LaSalle, and S. Lelshetz

UNIVERSAL  
LIBRARY

**OU\_150398**

UNIVERSAL  
LIBRARY

# OSMANIA UNIVERSITY LIBRARY

Call No.

Accession No.

Author

Title

This book should be returned on or before the date last marked below.

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

# ANNALS OF MATHEMATICS STUDIES

*Edited by Robert C. Gunning, John C. Moore, and Marston Morse*

1. Algebraic Theory of Numbers, *by* HERMANN WEYL
3. Consistency of the Continuum Hypothesis, *by* KURT GÖDEL
11. Introduction to Nonlinear Mechanics, *by* N. KRYLOFF and N. BOGOLIUBOFF
16. Transcendental Numbers, *by* CARL LUDWIG SIEGEL
19. Fourier Transforms, *by* S. BOCHNER and K. CHANDRASEKHARAN
20. Contributions to the Theory of Nonlinear Oscillations, Vol. I, *edited by* S. LEFSCHETZ
21. Functional Operators, Vol. I, *by* JOHN VON NEUMANN
24. Contributions to the Theory of Games, Vol. I, *edited by* H. W. KUHN and A. W. TUCKER
25. Contributions to Fourier Analysis, *edited by* A. ZYGMUND, W. TRANSUE, M. MORSE, A. P. CALDERON, and S. BOCHNER
28. Contributions to the Theory of Games, Vol. II, *edited by* H. W. KUHN and A. W. TUCKER
30. Contributions to the Theory of Riemann Surfaces, *edited by* L. AHLFORS *et al.*
33. Contributions to the Theory of Partial Differential Equations, *edited by* L. BERS, S. BOCHNER, and F. JOHN
34. Automata Studies, *edited by* C. E. SHANNON and J. MCCARTHY
36. Contributions to the Theory of Nonlinear Oscillations, Vol. III, *edited by* S. LEFSCHETZ
38. Linear Inequalities and Related Systems, *edited by* H. W. KUHN and A. W. TUCKER
39. Contributions to the Theory of Games, Vol. III, *edited by* M. DRESHER, A. W. TUCKER and P. WOLFE
40. Contributions to the Theory of Games, Vol. IV, *edited by* R. DUNCAN LUCE and A. W. TUCKER
41. Contributions to the Theory of Nonlinear Oscillations, Vol. IV, *edited by* S. LEFSCHETZ
42. Lectures on Fourier Integrals, *by* S. BOCHNER. In preparation
43. Ramification Theoretic Methods in Algebraic Geometry, *by* S. ABHYANKAR
44. Stationary Processes and Prediction Theory, *by* H. FURSTENBERG
45. Contributions to the Theory of Nonlinear Oscillations, Vol. V, CESARI, LAsALLE, and LEFSCHETZ
46. Seminar on Transformation Groups, *by* A. BOREL *et al.*
47. Theory of Formal Systems, *by* R. SMULLYAN
48. Lectures on Modular Forms, *by* R. C. GUNNING
49. Composition Methods in Homotopy Groups of Spheres, *by* H. TODA
50. Cohomology Operations, *lectures by* N. E. STEENROD, *written and revised by* D. B. A. EPSTEIN
51. Morse Theory, *by* J. W. MILNOR
52. Advances in Game Theory, *edited by* M. DRESHER, L. SHAPLEY, and A. W. TUCKER
53. Flows on Homogeneous Spaces, *by* L. AUSLANDER, L. GREEN, F. HAHN, *et al.*
54. Elementary Differential Topology, *by* J. R. MUNKRES
55. Degrees of Unsolvability, *by* G. E. SACKS.



# CONTRIBUTIONS TO THE THEORY OF NONLINEAR OSCILLATIONS

---

## *VOLUME V*

---

|               |                     |
|---------------|---------------------|
| J. P. LaSALLE | L. MARKUS           |
| J. JARNIK     | G. REEB             |
| J. KURZWEIL   | P. MENDELSON        |
| J. CRONIN     | J. ANDRÉ            |
| J. K. HALE    | P. SEIBERT          |
| L. CESARI     | C. COLEMAN          |
| A. STOKES     | R. P. DE FIGUEIREDO |

EDITED BY

L. Cesari, J. LaSalle, and S. Lefschetz

PRINCETON, NEW JERSEY  
PRINCETON UNIVERSITY PRESS  
1960

Copyright © 1960, by Princeton University Press  
All Rights Reserved  
L. C. Card 60-12225

Second Printing 1964

Printed in the United States of America

## PREFACE

This volume is the fifth in a series of contribution to the Non-linear Oscillations published by the Annals of Mathematics Studies. As usual there are quite a variety of topics covered. The first paper by LaSalle is on a central problem on optimal control — a problem which is agitating mathematicians both in the United States and in the Soviet Union. This is followed by a contribution from Harnik and Kurzweil on a generalized differential equation depending on a parameter. This is a continuation of a series of papers published on the topic by Kurzweil. We have next an elegant paper by Jane Cronin on the perturbation theorem of Poincaré. This is followed by two papers by Hale and one by Cesari. The first paper concerns a discussion of the critical points for a linear system with periodic coefficients. The second paper of Hale deals with the important question of the stability of periodic solutions of periodic and autonomous differential systems. Simple criteria for stability are proved by the use of a convergent process of successive approximations. The paper by Cesari gives existence theorems for periodic solutions of periodic and autonomous differential systems satisfying a Lipschitz condition. These theorems are proved by application of fixed point theorems in functional spaces and from this approach an interpretation is given of the method of successive approximations used in the paper by Hale. Stokes, in his paper, applies a fixed point theorem in linear spaces to the study of certain stability and boundedness properties of the solutions of differential equations. By the use of such methods he is able to reduce the study of systems to that of related first order equations. The next paper by Markus deals with a curious application of abstract algebra to the classification of linear differential systems. The paper by Reeb deals with a property of the totality of bounded solutions of certain dynamical systems. Mendelson's contribution deals with the stable motions in a neighborhood of critical points in topological dynamics. It is connected with a topological method introduced and promoted by Ważewski. The joint paper by André and Seibert deals with piecewise continuous differential equations. It is part of the noteworthy work done by those authors in

organizing the earlier work of Mrs. Flugge-Lotz. Coleman's paper is a study of asymptotic stability of a 3-dimensional system where the leading terms are a homogeneous function of degree  $m$ . In the final paper of this volume Figueiredo studies the self-sustained oscillations of a second order system.

All but the papers by Jane Cronin, Jarnik and Kurzweil, Reeb, and Figueiredo were carried out at RIAS and supported in part by contract number AF 49(638)-382.

The work by J. Cronin and A. Stokes was supported in part by the Office of Naval Research.

Editors

L. Cesari  
J. P. LaSalle  
S. Lefschetz

# CONTENTS

|                                                                                                                             | Page |
|-----------------------------------------------------------------------------------------------------------------------------|------|
| Preface                                                                                                                     |      |
| By L. Cesari, J. P. LaSalle and S. Lefschetz                                                                                | v    |
| I. The Time Optimal Control Problem                                                                                         |      |
| By J. P. LaSalle                                                                                                            | 1    |
| II. Continuous Dependence on a Parameter                                                                                    |      |
| By J. Jarník and J. Kurzweil                                                                                                | 25   |
| III. Poincaré's Perturbation Method and Topological Degree                                                                  |      |
| By Jane Cronin                                                                                                              | 37   |
| IV. On the Behavior of the Solutions of Linear Periodic<br>Differential Systems Near Resonance Points                       |      |
| By Jack K. Hale                                                                                                             | 55   |
| V. On the Stability of Periodic Solutions of Weakly<br>Nonlinear Periodic and Autonomous Differential Systems               |      |
| By Jack K. Hale                                                                                                             | 91   |
| VI. Existence Theorems for Periodic Solutions of Nonlinear<br>Lipschitzian Differential Systems and Fixed Point<br>Theorems |      |
| By Lamberto Cesari                                                                                                          | 115  |
| VII. The Applications of a Fixed Point Theorem to a<br>Variety of Non-Linear Stability Problems                             |      |
| By Arnold Stokes                                                                                                            | 173  |
| VIII. Quadratic Differential Equations and Non-Associative<br>Algebras                                                      |      |
| By Lawrence Markus                                                                                                          | 185  |
| IX. Sur Une Propriété De L'Ensemble Des Trajectoires<br>Bornées De Certains Systèmes Dynamiques                             |      |
| By Georges Reeb                                                                                                             | 215  |
| X. On Lagrange Stable Motions in the Neighborhood of<br>Critical Points                                                     |      |
| By Pinchas Mendelson                                                                                                        | 219  |
| XI. The Local Theory of Piecewise Continuous Differential<br>Equations                                                      |      |
| By J. André and P. Seibert                                                                                                  | 225  |
| XII. Asymptotic Stability in 3-Space                                                                                        |      |
| By Courtney Coleman                                                                                                         | 257  |
| XIII. Existence and Uniqueness of the Periodic Solution<br>of an Equation for Autonomous Oscillations                       |      |
| By Rui Pacheco de Figueiredo                                                                                                | 269  |
| Appendix                                                                                                                    |      |
| By E. Pinney                                                                                                                | 285  |



CONTRIBUTIONS TO THE  
THEORY OF  
NONLINEAR OSCILLATIONS  
VOLUME V





## I. THE TIME OPTIMAL CONTROL PROBLEM

J. P. La Salle

### §1. INTRODUCTION

It has been an intuitive assumption for some time that if a control system is being operated from a limited source of power and if one wishes to have the system change from one state to another in minimum time, then this can be done by at all times utilizing properly all of the power available. This hypothesis is called "the bang-bang principle." Bushaw accepted this hypothesis and in 1952 in [2] showed for some simple systems with one degree of freedom that of all bang-bang systems (that is, systems which at all times utilize maximum power) there is one that is optimal. In 1953 I made the observation in [7] that the best of all bang-bang systems, if it exists, is then the best of all systems operating from the same power source. More recently, more general results have been obtained by Bellman, Glicksberg, and Gross in [1], and later — but seemingly independently — by Krasovskii in [6] and Gamkrelidze in [4].

We confine our attention to the time optimal problem for control systems which are linear in the sense that the elements being controlled are linear and as a function of time the steering of the system enters linearly. The differential equation for such systems is

$$(1) \quad \dot{x}(t) = A(t)x(t) + B(t)u(t) + f(t),$$

where  $x$  and  $f$  are  $n$ -dimensional vector functions ( $x(t)$  is the state of the system at time  $t$ ),  $A$  is an  $(n \times n)$  matrix function, and  $B$  is an  $(n \times r)$  matrix function. The vector equation (1) represents the system of differential equations

$$\frac{dx_i(t)}{dt} = \sum_{j=1}^n a_{ij}(t)x_j(t) + \sum_{k=1}^r b_{ik}(t)u_k(t) + f_i(t), \quad i = 1, \dots, n.$$

Our ability to control the system lies in the freedom we have to choose the "steering" function  $u$ . We assume that the admissible steering functions are piecewise continuous (or measurable) and have components less than 1 in absolute value ( $|u_k(t)| \leq 1$ ). The number 1 is selected for convenience. We could just as well assume  $-a_k \leq u_k(t) \leq b_k$  where  $a_k$  and  $b_k$  are any positive numbers. Given an initial state  $x_0$  and a moving particle  $z(t)$ , the problem of time optimal control is to hit the particle in minimum time. Let  $x(t, u)$  be the solution of (1) satisfying  $x(0) = x_0$ . An admissible steering function  $u^*$  is optimal if  $x(t^*, u^*) = z(t^*)$  for some  $t^* > 0$  and if  $x(t, u) \neq z(t)$  for  $0 < t < t^*$  and all admissible  $u$ .

In [1] Bellman, Glicksberg, and Gross consider the system

$$(2) \quad \dot{x}(t) = Ax(t) + Bu(t)$$

and restrict themselves to the problem of starting at  $x_0$  and reaching the origin in minimum time. The  $(n \times n)$  matrix  $A$  is constant and its characteristic roots are assumed to have negative real parts.  $B$  was assumed to be a constant non-singular  $(n \times n)$  matrix, and it is this restriction that is quite unrealistic. Gamkrelidze in [4] considered the same problem, removed the restriction that  $B$  be non-singular, and showed for systems which are later in this paper called "normal" the existence and uniqueness of an optimal steering function. The form of the optimal steering function is the same as that given in [1], and for normal systems one can conclude that the optimal steering is bang-bang ( $|u_k(t)| = 1$ ). Krasovskii, in [6] considers the more general control system (1) and the more general control problem of hitting a moving particle. Using results of Krein on the L-problem in abstract spaces, he proves the existence of an optimal steering function for, what are called in this paper, "proper" control systems. We prove the same existence theorem without restricting ourselves to proper systems. Krasovskii makes the further assertion that the optimal steering function is unique and simple examples show that this is not true even for proper systems.

To date, therefore, the most general bang-bang principle has been proved by Gamkrelidze for normal control systems of the form (2) and for the special problem of reaching the origin in minimum time. We shall show for the general control problem that if an admissible steering function can bring the system from one state to another in time  $t$  then there is a bang-bang steering function that can do the same thing in the same time. This extends my result in [7] and at the same time extends the bang-bang principle. This does not mean that all optimal steering functions are bang-bang. Even for proper control systems there are

examples where the objective can be reached in minimum time using a steering function which during part of the time has some zero components. As in the special case considered by Gamkrelidze, it is shown that normal systems have unique optimal steering functions and for such systems there is a true bang-bang principle: the only way to reach the objective in minimum time is by bang-bang steering. In Theorem 5 a result is established that should be of some practical importance in the synthesis problem, which is the problem of determining the optimal steering  $u^*$  as a function of the state of the system. This result shows that for some control systems optimal steering can be determined by what amounts to running the system backwards. In Section 4 we discuss the controllability properties of proper control systems. The examples have been given at the end of the paper in Section 5, and they serve to illustrate the ideas and the application of the theory.

## §2. THE GENERAL PROBLEM

The problem, described in the Introduction, for the system (1) of reaching a moving particle in minimum time will be called the general problem. For the control system (1) the state  $x(t, u)$  of the system at time  $t$  is given by

$$(3) \quad x(t, u) = X(t)x_0 + X(t) \int_0^t Y(\tau)u(\tau)d\tau + X(t) \int_0^t X^{-1}(\tau)f(\tau)d\tau$$

$X(t)$  is the principal matrix solution of  $\dot{X}(t) = A(t)X(t)$ , and  $Y(\tau) = X^{-1}(\tau)B(\tau)$ . We want at some time  $t$  to have  $x(t, u) = z(t)$ ; that is, to have

$$(4) \quad w(t) = \int_0^t Y(\tau)u(\tau)d\tau, \quad$$

where

$$w(t) = X^{-1}(t)z(t) - x_0 - \int_0^t X^{-1}(\tau)f(\tau)d\tau.$$

We assume throughout that  $A(t)$ ,  $B(t)$ ,  $f(t)$  and  $z(t)$  are continuous for  $0 \leq t < \infty$ . The proof of the following lemma was pointed out to me by L. Pukanszky.

LEMMA 1. Let  $M$  be the set of all real-valued measurable functions  $\alpha(t)$  on  $[0, 1]$  with

$|\alpha(t)| \leq 1$ . Let  $M^0$  be the subset of functions in  $M$  with  $|\alpha(t)| = 1$ . Let  $y(t)$  be any  $n$ -dimensional function in  $L^1([0, 1])$ . Define

$$K = \left\{ \int_0^1 \alpha(t)y(t)dt; \quad \alpha \in M \right\}$$

and

$$K^0 = \left\{ \int_0^1 \alpha^0(t)y(t)dt; \quad \alpha^0 \in M^0 \right\}.$$

Then  $K^0$  is closed and  $K = K^0$ .

PROOF. For each measurable set  $E$  in  $[0, 1]$  define

$$\mu(E) = \int_E y(t)dt.$$

Let  $R_\mu$  denote the range of this vector measure. Let  $c_E(t)$  be the characteristic function of  $E$ , and let  $\alpha^0(t) = 2c_E(t) - 1$ . Then  $\alpha^0(t) \in M^0$ , and each  $\alpha^0(t) \in M^0$  can be so represented. Then clearly  $K^0 = 2R_\mu - \bar{y}$ , where

$$\bar{y} = \int_0^1 y(t)dt.$$

By Liapounov's Theorem ([8], [5])  $R_\mu$  is closed and convex, and therefore  $K^0$  is closed and convex. Let

$$z = \int_0^1 \alpha(t)y(t)dt$$

be any  $n$ -vector in  $K$ . What we wish to show is that  $z$  is a limit of vectors in  $K^0$ , and hence is in  $K^0$ . It will then follow that  $K^0 = K$ . Let  $\beta(t) = \frac{1}{2}(\alpha(t) + 1)$  and  $\bar{z} = \frac{1}{2}(z + \bar{y})$ . Note that  $0 \leq \beta(t) \leq 1$  and

$$\bar{z} = \int_0^1 \beta(t)y(t)dt.$$

Define

$$\bar{z}_m = \sum_{j=1}^m \frac{1}{m} \int_{E_j} y(t)dt$$

where

$$E_j = \left\{ t; \frac{j-1}{m} < \beta(t) \leq \frac{j}{m} \right\}$$

Then

$$|\bar{z} - \bar{z}_m| = \sum_{j=1}^m \int_{E_j} \left( \frac{j}{m} - \beta(t) \right) y(t) dt \leq \frac{1}{m} \sum_{j=1}^m \int_{E_j} |y(t)| dt = \frac{1}{m} \int_0^1 |y(t)| dt$$

which shows that  $\bar{z}_m \longrightarrow \bar{z}$  as  $m \longrightarrow \infty$ . Letting

$$F_j = \bigcup_{i=j}^m E_i,$$

we obtain

$$\bar{z}_m = \frac{1}{m} \sum_{j=1}^m \int_{F_j} y(t) dt.$$

Since  $R_\mu$  is convex, it follows that  $\bar{z}_m \in R_\mu$ ; therefore  $z_m = 2\bar{z}_m - \bar{y} \in K^0$  and  $z_m \longrightarrow z$  as  $m \longrightarrow \infty$ . This completes the proof.

For our purposes here we wish to restate this lemma in the following more general fashion.

LEMMA 2. Let  $\Omega$  be the set of all  $r$ -dimensional vector functions  $u(\tau)$  measurable on  $[0, t]$  with  $|u_1(\tau)| \leq 1$ . Let  $\Omega^0$  be the subset of functions  $u^0(\tau)$  with  $|u_1^0(\tau)| = 1$ . Let  $Y(\tau)$  be any  $(n \times r)$  matrix function in  $L^1([0, t])$ . Define

$$A(t) = \left\{ \int_0^t Y(\tau) u(\tau) d\tau; \quad u \in \Omega \right\}$$

and

$$A^0(t) = \left\{ \int_0^t Y(\tau) u^0(\tau) d\tau; \quad u^0 \in \Omega^0 \right\}.$$

Then  $A^0(t)$  is closed and  $A(t) = A^0(t)$ .

PROOF. Let  $y^1(\tau), y^2(\tau), \dots, y^r(\tau)$  be the column vectors in  $Y(\tau)$ , and define

$$A_j(t) = \left\{ \int_0^t y^j(\tau) u_j(\tau) d\tau; \quad u \in \Omega \right\}$$

and

$$A_j^0(t) = \left\{ \int_0^t y^j(\tau) u_j^0(\tau) d\tau; \quad u^0 \in \Omega^0 \right\}$$

Since

$$\int_0^t Y(\tau) u(\tau) d\tau = \sum_{j=1}^r \int_0^t y^j(\tau) u_j(\tau) d\tau, \quad ,$$

we see that

$$A(t) = A_1(t) + A_2(t) + \dots + A_r(t)$$

and

$$A^0(t) = A_1^0(t) + A_2^0(t) + \dots + A_r^0(t) \quad .$$

By Lemma 1  $A_j(t) = A_j^0(t)$  for each  $j = 1, \dots, r$ , and hence  $A(t) = A^0(t)$ . Each  $A_j^0(t)$  is bounded, and it follows from Lemma 1 that each  $A_j^0(t)$  is compact. Therefore  $A^0(t)$  is closed, and this completes the proof. It is convenient to point out also two elementary properties of convex sets that we will use later.

LEMMA 3. Consider sets  $A(t)$  of  $E_n$ ,  $t \leq t^*$ , with the following properties:

- a. Each  $A(t)$  is convex.
- b. Corresponding to each  $\varepsilon > 0$  there is a  $\delta > 0$  such that  $d(p, A(t)) < \varepsilon$  for each  $p \in A(t^*)$  and all  $t^* - \delta < t < t^*$ . [ $d(p, A(t))$  is the distance of  $p$  from  $A(t)$ .]

Then, if  $q$  is in the interior of  $A(t^*)$ , there is a  $t_1 < t^*$  such that  $q$  is an interior point of  $A(t_1)$ .

PROOF. Let  $q$  be in the interior of  $A(t^*)$ , and let  $N$  be a neighborhood of  $q$  of radius  $\varepsilon > 0$  inside  $A(t^*)$ . Suppose for each  $t < t^*$  that  $q$  is not an interior point of  $A(t)$ . Then for each  $t < t^*$  there is a support plane  $P_t$  through  $q$  such that there are no points of  $A(t)$  on one side of  $P_t$  ([3]). Because of the neighborhood  $N$  about  $q$  that lies in  $A(t^*)$  we see that for each  $t < t^*$  there is a

point  $p$  in  $A(t^*)$  whose distance from  $A(t)$  is at least  $\varepsilon$ . This contradicts (b).

LEMMA 4. Let  $M$  be a convex set in  $E_n$  containing the origin with the property: given any number  $K$  and any non-zero vector  $\eta$  in  $E_n$  there is a vector  $y$  in  $M$  such that  $(\eta, y) > K$ . Then  $M = E_n$ .

PROOF. Suppose that  $M$  is not  $E_n$ . Then there is some non-zero  $\mu$  that is not in  $M$ , and hence  $a\mu$  is a boundary point of  $M$  for some  $a \geq 0$ . Then  $a\mu$  has a support plane  $P$  ([3]). Let  $\eta$  be a non-zero vector normal to  $P$  and directed toward the side of  $P$  that contains no points of  $M$ . Then  $(\eta, y)$  is bounded above for  $y$  in  $M$  — a contradiction. Therefore  $M = E_n$ .

Turning our attention back to the control problem, we see that the set  $\Omega$  in Lemma 2 is the set of admissible steering functions and  $\Omega^0$  is the set of bang-bang steering functions. The set  $A(t)$  is related by equation 4 to the set of states that can be reached in time  $t$  by the admissible steering functions, and  $A^0(t)$  is similarly related to the set of states that can be reached in time  $t$  by the bang-bang steering functions. Lemma 2 then states that anything that can be done by an admissible steering function can also be done by a bang-bang function. Suppose that there is a steering function  $u \in \Omega$  such that  $x(t_1, u) = z(t_1)$ ; the particle is then hit at time  $t_1$ , and

$$w(t_1) = \int_0^{t_1} Y(\tau)u(\tau)d\tau.$$

The lemma then states that there will be a  $u^0 \in \Omega^0$  with the property that  $x(t_1, u^0) = z(t_1)$ ; bang-bang steering can accomplish the same thing in the same time. As a direct consequence of this we obtain the following pair of theorems.

THEOREM 1. If of all bang-bang steering functions there is an optimal one relative to  $\Omega^0$ , then it is optimal (relative to  $\Omega$ ).

THEOREM 2. If there is an optimal steering function, then there is always a bang-bang steering function that is optimal.

The first of these theorems extends the result in [7], and the second is a general bang-bang principle. Although we did not state the intuitive hypothesis this way, the feeling actually is that not only should there always be bang-bang steering that is optimal but no other type of steering should be optimal. If at some time all of the available power is not being used, then it should be possible to improve the performance by using, properly, the additional power that is available. Perhaps this is true in some more general sense, but under our restriction on the amplitude of each component of the steering function this is not true without placing further restrictions on the control system. We return to this question in the next section.

We can now extend the results that have been obtained previously on the existence of and the form of optimal steering. The set  $A(t)$  is, as we have said, related by equation (4) to the set of all states of the system that can be reached in time  $t$ . The existence is a simple consequence of the fact that  $A(t)$  is a closed set, and the form (5) below for optimal steering follows from the convexity of  $A(t)$ . [For  $r$ -dimensional vectors  $a$  and  $b$ ,  $a = \text{sgn } b$  means that  $a_i = \text{sgn } b_i$ ,  $i = 1, \dots, r$ ;  $\text{sgn } b_j = 1$  if  $b_j > 0$ ,  $\text{sgn } b_j = -1$  if  $b_j < 0$ , and is considered to be undetermined if  $b_j = 0$ . In using this notation  $a$  is always a column vector and  $b$  is a row vector.]

**THEOREM 3.** If for the general problem there is a steering function  $u$  in  $\Omega$  such that  $x(t, u) = z(t)$  for some  $t > 0$ , then there is an optimal steering function in  $\Omega$ . Moreover, all optimal steering functions  $u^*$  are of the form

$$(5) \quad u^*(t) = \text{sgn } [\eta Y(t)]$$

where  $\eta$  is some non-zero  $n$ -dimensional vector.

**PROOF.** Our assumption that  $x(t, u) = z(t)$  for some  $t > 0$  and some  $u \in \Omega$  is equivalent to assuming that  $w(t) \in A(t)$  for some  $t > 0$ . Let  $t^*$  be the greatest lower bound of all positive  $t$ 's with this property. Let

$$y(t, u) = \int_0^t Y(\tau) u(\tau) d\tau.$$

Then there is a non-increasing sequence  $t_n$  converging to  $t^*$  and  $u_n \in \Omega$  such that  $w(t_n) = y(t_n, u_n) \in A(t_n)$ . Now for some number  $K$



it is easy to see that

$$|y(t_n, u_n) - y(t^*, u_n)| \leq K|t_n - t^*|.$$

Therefore  $w(t^*)$  is the limit of points in  $A(t^*)$ , and since  $A(t^*)$  is closed,  $w(t^*)$  is in  $A(t^*)$ . This proves the existence of a steering function  $u^*$  in  $\Omega$  such that  $w(t^*) = y(t^*, u^*)$ . By the definition of  $t^*$ ,  $u^*$  is optimal.

We now want to show that  $u^*$  is of the form (5). In order to do this we note first that  $w(t^*)$  cannot be an interior point of  $A(t_1)$  for  $0 < t_1 < t^*$ . Suppose this were true. Let  $N$  be a neighborhood of  $w(t^*)$  contained in  $A(t_1)$ . Then, since  $A(t_1) \subset A(t)$  for all  $t_1 < t$ ,  $N$  is contained in  $A(t)$  for all  $t > t_1$ . The trajectory  $w(t)$  is continuous and this implies  $w(t_2) \in A(t_2)$  for some  $t_2 < t^*$ . This contradicts the definition of  $t^*$ , and therefore,  $w(t^*)$  cannot be an interior point of  $A(t_1)$  for  $t_1 < t^*$ . It then follows from Lemma 3 that  $w(t^*)$  is not an interior point of  $A(t^*)$ . Therefore  $w(t^*)$  is a boundary point of the convex set  $A(t^*)$ . There is then a support plane  $P$  through  $w(t^*)$  (see [3]). Let  $\eta$  be a non-zero vector normal to  $P$  and directed to the side which contains no points of  $A(t^*)$ . Therefore for each  $y(t^*, u)$  in  $A(t^*)$

$$(\eta, y(t^*, u) - w(t^*)) \leq 0.$$

Now

$$w(t^*) = y(t^*, u^*) = \int_0^{t^*} Y(\tau) u^*(\tau) d\tau,$$

and this inequality implies

$$\int_0^{t^*} (\eta, Y(\tau) u(\tau)) d\tau \leq \int_0^{t^*} (\eta, Y(\tau) u^*(\tau)) d\tau$$

for all  $u \in \Omega$ . Let  $v(\tau) = \eta Y(\tau)$ . Then

$$\int_0^{t^*} (\eta, Y(\tau) u^*(\tau)) d\tau \leq \sum_{j=1}^r \int_0^{t^*} |v_j(\tau)| d\tau = \sum_{j=1}^r \int_0^{t^*} v_j(\tau) \operatorname{sgn} v_j(\tau) d\tau.$$

Hence we see that on any interval of positive length where  $v_j(\tau) \neq 0$  it must be that  $u_j^*(\tau) = \operatorname{sgn} v_j(\tau)$ . This completes the proof.

The form (5) for an optimal steering function enables us to state a uniqueness criterion. Let  $y^j(t)$  be the  $j^{\text{th}}$  column vector of  $Y(t)$ . The control system (1) is said to be normal if for each  $j = 1, \dots, r$  the functions  $y_1^j(t), \dots, y_n^j(t)$  are linearly (functionally) independent on each interval of positive length. This is equivalent to saying that no component of  $\eta Y(t)$ ,  $\eta \neq 0$ , is identically zero on an interval of positive length. Therefore  $u^*(t)$  is uniquely determined by (5), and for normal control systems there is at most one optimal steering function, and if there is one, it is bang-bang. Thus, for normal control systems, the only way of reaching the objective in minimum time is by bang-bang steering of the form (5). For other control systems the most that can be said is that if it is possible to reach the objective in finite time then there is a bang-bang steering function that is optimal. The optimal steering will be of the form (5) but there may be an infinity of optimal steering functions, some of which will not be bang-bang steering.

### §3. THE SPECIAL PROBLEM

The control problem for the system

$$(6) \quad \dot{x}(t) = A(t)x(t) + B(t)u(t)$$

where the objective is to start at an initial state  $x_0$  at times  $t = 0$  and to reach the origin (the equilibrium state) in minimum time will be called the special problem. The objective in the special problem is to have [see equation (4)] for some  $t > 0$

$$(7) \quad -x_0 = \int_0^t Y(\tau)u(\tau)d\tau.$$

For this special problem we can show that if there is a steering function of the form (5) that brings the system from the initial state  $x_0$  to the origin in finite time then it is an optimal steering function.

**THEOREM 5.** If for some  $t > 0$  and some  $n$ -vector  $\eta$  there is a solution  $u = \bar{u}$  of (7) of the form

$$(5) \quad \bar{u}(\tau) = \text{sgn} [\eta Y(\tau)],$$

and if  $\eta Y(\tau) \neq 0$  on an interval of positive length, then it is an optimal steering function for the special problem.

PROOF. By Theorem 3 the existence of a  $\bar{u}$  and a  $\bar{t} > 0$  satisfying

$$-x_0 = y(\bar{t}, \bar{u}) = \int_0^{\bar{t}} Y(t)\bar{u}(t)dt$$

implies the existence of an optimal  $u^*$  and a minimal time  $t^* \leq \bar{t}$ . But then  $y(t^*, u^*) = -x_0$ , and

$$\begin{aligned} (\eta, y(\bar{t}, \bar{u}) - y(t^*, u^*)) &= \int_0^{t^*} \eta Y(t)[\bar{u}(t) - u^*(t)]dt \\ &+ \int_{t^*}^{\bar{t}} \eta Y(t)\bar{u}(t)dt = 0. \end{aligned}$$

Since  $\bar{u}$  is of the form (5), we can conclude that

$$\eta Y(t) = 0 \quad \text{on} \quad [t^*, \bar{t}]$$

and that  $u_j^*(t) = \bar{u}_j(t)$  on  $[0, t^*]$  wherever the  $j^{\text{th}}$  component of  $\eta Y(t) \neq 0$ . Therefore, by the assumption that  $\eta Y(t) \neq 0$  on an interval of positive length, it follows that  $t^* = \bar{t}$ . Hence  $\bar{u}$  is an optimal steering function. This raises in a quite natural way the concept of a "proper" control system which is discussed in the next section.

Although the above result is quite elementary, it is of considerable practical significance and does offer a means of solving the synthesis problem. It has been mathematically convenient up to now to treat the steering function  $u$  as though it were a function of time. What one actually wants, is to know the optimal steering function  $u^*$  as a function of the state  $x$  of the system, and this is the "synthesis problem". If the control system is autonomous (equation 2), then we can replace  $t$  by  $-t$  in equation 2, use a steering function of the form (5), start at the origin, and look at the solution. Theorem 5 states that the particular steering function used is optimal for all states of the system through which the solution passes. The time at which the solution passes through a state of the system is then the minimal time for the system to go from that state to the origin. For normal systems the optimal steering is unique, and this procedure of reversing the system determines the optimal steering as a function of the state of the system. It is even true for some systems which are not normal that the synthesis problem can be solved in this way. This procedure leads to

the determination of the switching surfaces, which are surfaces where certain of the components change sign. This method is illustrated in Section 5, and the switching surfaces (curves in the examples considered) are easily obtained by quite elementary reasoning. In Example 3 of Section 5 the synthesis problem is solved for a system that is not normal.

#### §4. PROPER CONTROL SYSTEMS AND CONTROLLABILITY

In this section we introduce the concept of a proper control system and establish some controllability properties of such systems. We say that a control system is proper if  $\eta Y(t) = 0$  on an interval of positive length implies  $\eta = 0$ . This is equivalent to saying that the row vectors  $y_1(t), y_2(t), \dots, y_n(t)$  of  $Y(t)$  are linearly independent functions on each interval of positive length. It is clear that every normal system is proper. If the steering function has only one component ( $r = 1$ ), then the concepts of proper and normal are equivalent. However, it is not in general true that every proper system is normal, as is shown by Example 3 in Section 5. It is a direct consequence of Theorem 3 that in proper control systems optimal steering  $u^*$  has the property that at any given time some component of  $u^*$  assumes an extreme value.

Think now of removing all of the constraints on the admissible control functions, and consider any two states  $x_1$  and  $x_2$  and any two times  $t_1$  and  $t_2$ . If for each pair of states and pair of times there is a steering function such that starting at  $x_1$  at time  $t_1$  the system is brought to the state  $x_2$  at time  $t_2$ , then the system is said to be completely controllable.

THEOREM 6. Proper control systems are completely controllable.

PROOF. We may certainly assume that  $t_1 = 0$  and  $t_2 > 0$ . From (4) we see that complete controllability is equivalent to the property that for each  $t > 0$  the set  $M_t$  of all vectors

$$\int_0^t Y(\tau)u(\tau)d\tau,$$

using all possible steering functions  $u$ , is the whole phase space  $E_n$ . Let  $v(\tau) = \eta Y(\tau)$ , and define

$$y = \int_0^t Y(\tau) \operatorname{sgn} v(\tau) d\tau.$$

Then

$$(\eta, y) = \sum_{j=1}^r \int_0^t |v_j(\tau)| d\tau.$$

Since the control system is proper,

$$\sum_{j=1}^r \int_0^t |v_j(\tau)| d\tau > 0$$

if  $\eta \neq 0$ . Therefore corresponding to each direction  $\eta$  there is a vector  $y$  in  $M_t$  such that  $(\eta, y) > 0$ .  $M_t$  is obviously a linear manifold, and therefore  $M_t = E_n$ . [It is of interest to note that the theorem is also a consequence of the fact that for proper control systems the linear transformation  $F$  on  $E_n$  defined by

$$F(\eta) = \int_0^t Y(\tau) Y'(\tau) \eta' d\tau$$

is non-singular for each  $t > 0$ . The prime denotes the transpose. Thus the system is completely controllable with steering functions restricted to those of the form  $u(\tau) = Y'(\tau) \eta'$ .]

Complete controllability assumes that there is no restriction on the admissible control functions, and if unlimited power is available to a proper control system the above theorem says that it is always possible to move the system from any one state to any other state as quickly as we please. For each  $r$ -vector  $v$  define  $\|v\| = |v_1| + |v_2| + \dots + |v_r|$ . If we replace the concept of being proper by the requirement that

$$\int_0^\infty \|\eta Y(\tau)\| d\tau > 0$$

for each  $\eta \neq 0$ , then it is easily seen that with no constraints on the steering functions there is a time  $T > 0$  such that the system can be started at any state  $x_0$  at time  $t = 0$  and can be steered to any other state in time  $T$ . This fact does not appear to be of any particular importance, and for autonomous systems this condition is equivalent to

the condition that the system be proper. We wish now to return to the assumption that the system has limited power and to the special problem. We shall say that a control system is asymptotically proper if

$$\int_0^{\infty} \|\eta Y(\tau)\| d\tau = \infty$$

for each  $\eta \neq 0$ . For the special problem we are interested in systems with the property that given any initial state  $x_0$  there is a steering function in  $\Omega$  that brings the system to the origin in finite time. We shall say that such a system is controllable. Of course, by Theorem 3 we know that if a system is controllable then for each initial state  $x_0$  there is a steering function in  $\Omega$  that is optimal for the special problem.

**THEOREM 7.** Asymptotically proper control systems of the form (6) are controllable.

**PROOF.** For systems of the form (6) we know from (4) that the system is controllable if corresponding to each  $x_0$  in  $E_n$  there is a  $u$  in  $\Omega$  with the property that

$$-x_0 = y(t, u) = \int_0^t Y(\tau)u(\tau)d\tau$$

for some  $t > 0$ . Let  $A = \{y(t, u); u \in \Omega, t > 0\}$ . Clearly  $A$  is a convex set, and what we wish to show is that  $A = E_n$ .

Taking  $u(t) = \text{sgn}[\eta Y(t)]$ , we have

$$(\eta, y(t, u)) = \int_0^t \|\eta Y(\tau)\| d\tau.$$

Since the system is asymptotically proper, we know that  $(\eta, y(t, u)) \rightarrow \infty$  as  $t \rightarrow \infty$  for each  $\eta \neq 0$ . By Lemma 4,  $A = E_n$  and the system is controllable.

We now examine some special properties of autonomous control systems (equation 2). Let  $\varphi(\lambda) = \det(A + \lambda I) = \lambda^n + c_1 \lambda^{n-1} + \dots + c_n$ ;  $\varphi(-\lambda)$  is the characteristic polynomial of  $A$ , and we know that  $\varphi(-A) = 0$ . Hence  $v(t) = \eta Y(t) = \eta e^{-At} B$  satisfies the  $n^{\text{th}}$  order linear differential equation with constant coefficients

$$v^{(n)}(t) + c_1 v^{(n-1)}(t) + \dots + c_n v(t) = 0, ,$$

where  $v^{(k)}(t)$  is the  $k^{\text{th}}$  derivative. Thus  $v(t) \equiv 0$  on an interval of positive length is equivalent to  $v(0) = \eta B = 0$ ,  $v'(0) = -\eta AB = 0$ , ...,  $v^{(n-1)}(0) = (-1)^{n-1} \eta A^{n-1} B = 0$ . It then follows that an autonomous control system is proper if and only if one of the following hold ( $M = B(E_n)$  is the range of  $B$  and  $b^1, \dots, b^r$  are the column vectors of  $B$ ):

1. For each non-zero vector  $\eta$  in  $E_n$  there is an integer  $k$ ,  $0 \leq k \leq n-1$ , such that  $\eta A^k B \neq 0$ .
2.  $M + A(M) + \dots + A^{(n-1)}(M) = E_n$ .
3. The set of vectors  $b^1, \dots, b^r, Ab^1, \dots, Ab^r, \dots, A^{(n-1)}b^1, \dots, A^{(n-1)}b^r$  contains a set of  $n$  linearly independent vectors.

The requirement that a system be normal is, in general, much stronger. The system is normal if and only if no component of  $v(t) = \eta Y(t)$  is identically zero on an interval of positive length. Therefore an autonomous system is normal if and only if for each integer  $j$ ,  $0 < j \leq r$ , the vectors  $b^j, Ab^j, \dots, A^{(n-1)}b^j$  are linearly independent.

Let us now assume that an autonomous control system is both proper and stable [the characteristic roots of  $A$  have non-positive real parts]. For  $\eta \neq 0$  we know that one component of  $v(t) = \eta Y(t) = \eta e^{-At} B$  is not identically zero. We may suppose that  $v_1(t) \neq 0$ . Then

$$v_1(t) = \sum_{j=1}^m P_j(t) e^{\alpha_j t} \cos(\beta_j t + \delta_j), ,$$

$\alpha_j \geq 0$  and  $P_j(t)$  are polynomials. Hence for  $t$  sufficiently large, say  $t \geq a$ ,

$$|v_1(t)| \geq |p(t) + q(t)| ,$$

where  $p(t)$  is almost periodic, is not identically zero, and  $|q(t)| < \frac{K}{t}$ . It then follows that

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_a^T |p(t) + q(t)|^2 dt = \lim_{T \rightarrow \infty} \frac{1}{T} \int_a^T |p(t)|^2 dt = 2c > 0 .$$

Consequently, for  $T$  sufficiently large,

$$\int_a^T |v_1(t)| dt \geq \int_a^T |p(t) + q(t)| dt \geq T^{-\frac{1}{2}} \int_a^T |p(t) + q(t)|^2 dt \geq cT^{\frac{1}{2}} .$$

Therefore

$$\int_0^\infty |v_1(t)| dt = \infty ,$$

and the system is asymptotically proper. What we have shown is that:

If an autonomous system is both stable and proper,  
then it is asymptotically proper and therefore  
controllable.

Let us now see what can be said about the existence of a solution and the determination of optimal steering for the general problem of starting at  $x_0$  at time  $t = 0$  and hitting a moving particle  $z(t)$ . The control system is

$$(1) \quad \dot{x}(t) = A(t)x(t) + B(t)u(t) + f(t) .$$

We wish to know whether or not there is a  $u \in \Omega$  and a  $t_1 > 0$  such that [equation (4)]

$$w(t_1) = \int_0^{t_1} Y(\tau)u(\tau)d\tau$$

where

$$w(t_1) = X^{-1}(t_1)z(t_1) - x_0 - \int_0^{t_1} X^{-1}(\tau)f(\tau)d\tau ;$$

that is, whether or not  $w(t_1) \in A(t_1)$  for some  $t_1 > 0$ . Think for the moment that  $t_1$  is fixed and ask whether or not, for some  $t > 0$ ,  $w(t_1) \in A(t)$ . This is then the special problem for the control system

$$(6) \quad \dot{x}(t) = A(t)x(t) + B(t)u(t)$$

of starting at  $-w(t_1)$  and reaching the origin in finite time. Suppose that the synthesis problem for this special problem can be solved. Then



for each point  $-w(t_1)$  of the curve  $-w(t)$ , we would know the optimal steering function  $\bar{u}$  and the minimum time  $\bar{t}_1$  for starting at  $-w(t_1)$  and reaching the origin. If  $\bar{t}_1 \leq t_1$ , then

$$w(t_1) \in A(\bar{t}_1) \subset A(t_1),$$

and we would know by Theorem 3 that the general problem has an optimal steering function  $u^*$  and a minimum time  $t^*$ ; we would know also that  $t^* \leq t_1$ . Thus, we have the following relation between the general problem and the special problem:

The general problem  $x(0, u) = x_0$ ,  $x(t, u) = z(t)$  has a solution if and only if in the special problem it is possible, starting at some point  $-w(t_1)$  of the curve

$$-w(t) = -X^{-1}(t)z(t) + x_0 + \int_0^t X^{-1}(\tau)f(\tau)d\tau,$$

to reach the origin in time  $t_2 \leq \bar{t}_1$ .

For proper control systems we can say a bit more. Suppose that the general problem has a solution. Let  $u^*$  be the optimal steering function and let  $t^*$  be the minimum time. Let  $\bar{u}^*$  be the optimal steering for the special problem of starting at  $-w(t^*)$  and reaching the origin. Let  $\bar{t}^*$  be the minimum time. Then  $\bar{t}^* \leq t^*$ . Now, using the steering function  $\bar{u}^*$  during the time interval  $[0, \bar{t}^*]$  and coasting ( $u(t) = 0$ ) during the interval  $[\bar{t}^*, t^*]$ , we hit the particle at time  $t^*$ . But we know for proper systems that optimal steering is never identically zero on an interval of positive length, and therefore it must be that  $\bar{t}^* = t^*$ . Thus, we have shown that

For proper systems, if  $u^*$  is optimal steering for the general problem and  $t^*$  is the minimum time, then  $u^*$  is optimal for the special problem of starting at  $-w(t^*)$  and reaching the origin;  $t^*$  is the minimum time for doing this.

Thus, if the system is proper and if the synthesis problem for the special problem can be solved, one can move along the trajectory  $-w(t)$ ,  $t > 0$ , use optimal steering from each point, and locate the first point where one can go from  $-w(t)$  to the origin in time  $t$ . The steering function that does this would then be optimal steering for the

general problem. Let  $m(x)$  be the minimum time to go from the initial state  $x$  to the origin. What we wish to find is the smallest  $t > 0$  satisfying  $m(-w(t)) = t$ .

## §5. EXAMPLES

The purpose of these examples is to illustrate the concepts that have been introduced and to indicate how the theory can be applied to solve the synthesis problem. Although similar examples have been given before, the more complete theory simplifies the reasoning and the computation leading to a solution. We had pointed out previously that the restriction  $|u_1| \leq 1$  was a matter of convenience and that this can be replaced by the more general condition  $-a_1 \leq u_1 \leq b_1$ ,  $a_1 > 0$ ,  $b_1 > 0$ . This is illustrated in Example 1.

Example 2 was solved in [1], and we consider the same example to show how a more complete theory leads to a simpler solution.

In Example 3 a system which is proper but not normal is considered. The optimal steering is not unique, not necessarily bang-bang, but yet the synthesis problem, which is to define optimal steering uniquely as a function of the state of the system, can be solved.

EXAMPLE 1. The differential equation of a controlled, undamped harmonic oscillator is

$$\ddot{x} + x = u.$$

We consider the constraint  $-a \leq u \leq b$ ,  $a > 0$ ,  $b > 0$ . An equivalent system is

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= -x + u.\end{aligned}$$

$$A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, B = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, X(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}, Y(t) = X^{-1}(t)B = \begin{pmatrix} \sin t \\ \cos t \end{pmatrix},$$

$$\eta Y(t) = \eta_1 \cos t + \eta_2 \sin t = A \cos(t + \delta).$$

The system is proper, and since  $r = 1$ , is also normal. It is stable and is therefore controllable. We know therefore that for each initial point  $(x_0, y_0)$  there is a unique optimal control function of the form

$$u(t) = \operatorname{sgn}^* [A \cos(t + \delta)] = \begin{cases} b, & \text{if } A \cos(t + \delta) > 0 \\ -a, & \text{if } A \cos(t + \delta) < 0 \end{cases}$$

which brings the system to the origin in minimum time. Conversely, we know that any steering function of this form which brings the system to the origin is optimal. We replace  $t$  by  $-\tau$ , obtain

$$\frac{dx}{d\tau} = -y$$

$$\frac{dy}{d\tau} = x - u,$$

and consider steering functions of the form  $u(\tau) = \text{sgn}[A \cos(\delta - \tau)]$ . When  $u = c$  the trajectory in the phase plane is a circle with center at  $(c, 0)$  and as  $\tau$  increases ( $t$  decreases) the direction on the trajectory is counterclockwise. The arrows in the figures are in the direction of increasing  $t$ . Thus, the trajectory can leave the origin, as indicated in Figure 1, going counterclockwise on either the circle with center at  $(b, 0)$  or the circle with center at  $(-a, 0)$ . Depending on the choice of  $\eta$ , the steering function can change sign at




FIGURE 1

any time before a half-revolution and cannot go past a half-revolution without changing sign. The switching curves are therefore the chain of semicircles shown in Figure 1. Above these semicircles  $u = -a$  and below them  $u = b$ . Thus starting in the first quadrant, as illustrated, optimal steering opposes the motion and does not change sign until past the point of maximum displacement except at those exceptional points where the semicircles join. Because of the simple nature of the trajectories we know that there is a unique optimal trajectory through each point of the phase plane. This is also a consequence of the fact that these are the only possible switching curves and that the system is both controllable and normal.

EXAMPLE 2. This example was solved in [1] by reasoning, which we can now see, was more complex than is necessary. In its normalized form the system is

$$\begin{aligned}\dot{y}_1 &= -2y_1 + u_1 + u_2 \\ \dot{y}_2 &= -y_2 + u_1 + 2u_2, \quad |u_1| \leq 1, \quad |u_2| \leq 1.\end{aligned}$$

Here

$$Y(t) = \begin{pmatrix} \eta_1 e^{2t} + \eta_2 e^t \\ \eta_1 e^{2t} + 2\eta_2 e^t \end{pmatrix}$$

and it is clear that the system is normal and controllable. In order to solve the synthesis problem we consider the solutions, starting at the origin of

$$\begin{aligned}\frac{dy_1}{d\tau} &= 2y_1 - u_1 - u_2 \\ \frac{dy_2}{d\tau} &= y_2 - u_1 - 2u_2\end{aligned}$$

using steering functions of the form

$$\begin{aligned}u_1(\tau) &= \operatorname{sgn}(\eta_1 e^{-2\tau} + \eta_2 e^{-\tau}) \\ u_2(\tau) &= \operatorname{sgn}(\eta_1 e^{-2\tau} + 2\eta_2 e^{-\tau}).\end{aligned}$$

Taking  $\eta_1 + \eta_2 < 0$ ,  $\eta_1 + 2\eta_2 > 0$ ,  $\eta_2 > 0$ , we begin at the origin with  $u_1 = -1$ ,  $u_2 = -1$ , and move out along the parabola (Figure 2)




FIGURE 2

$$\alpha : y_1(\tau) = e^{2\tau} - 1, \quad y_2(\tau) = 3(e^\tau - 1) \quad .$$

Now  $u_2(\tau)$  changes sign at  $\tau_1 = \ln(-\eta_1/2\eta_2)$ , and then  $u_1(\tau)$  changes sign at  $\tau_2 = \ln(-\eta_1/\eta_2)$ . Hence  $\tau_2 - \tau_1 = \ln 2$ . Therefore, starting at the point  $(y_1(\tau_1), y_2(\tau_1))$  of  $\alpha$  with  $u_1 = -1$ ,  $u_2 = 1$ , we leave  $\alpha$  along the parabola

$$y_1(\tau) = y_1(\tau_1)e^{2\tau}$$

$$y_2(\tau) = y_2(\tau_1)e^\tau - (e^\tau - 1) \quad ,$$

and the shift in the sign of  $u_1$  occurs at

$$y_1(\ln 2) = 4y_1(\tau_1)$$

$$y_2(\ln 2) = 2y_2(\tau_1) - 1 \quad .$$

This linear transformation applied to  $\alpha$  gives the switching curve  $\beta$  where  $u_1$  changes sign. Starting at the origin with  $\eta_1 + \eta_2 < 0$  and  $\eta_1 + 2\eta_2 \geq 0$ , we have  $u_1 = -1$ ,  $u_2 = 1$ , and  $u_1$  must change sign some time not later than  $\tau = \ln 2$  and can change at any time along the line from  $(0, 0)$  to  $(-1, 0)$ . This is the switching curve  $\gamma$ . The switching curves  $\alpha'$ ,  $\beta'$ ,  $\gamma'$  are obtained by symmetry. These are the only possible switching curves, and, since the system is controllable and normal, the steering function is determined uniquely at each point of the phase plane as indicated in Figure 2.

EXAMPLE 3. Here we consider a system which is proper and controllable but not normal. The difficulty is that it is not sufficient to determine all of the possible switching curves, since the optimal steering is not unique. What one wants to do is to determine a set of switching curves for which there is a unique optimal trajectory through each point. The system is

$$\begin{aligned}\dot{x}_1 &= -x_1 + u_1 \\ \dot{x}_2 &= -2x_2 + u_1 + u_2, \quad |u_1| \leq 1, \quad |u_2| \leq 1,\end{aligned}$$

where the matrix  $A$  is in diagonal form.

$$\begin{aligned}X(t) &= \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{-2t} \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \\ Y(t) &= \begin{pmatrix} e^t & 0 \\ e^{2t} & e^{2t} \end{pmatrix}, \quad \eta Y(t) = \begin{pmatrix} \eta_1 e^t + \eta_2 e^{2t} \\ \eta_2 e^{2t} \end{pmatrix}.\end{aligned}$$

The matrix  $B$  is non-singular, and the system is proper and controllable but not normal. We shall obtain a solution in which one component of the steering is zero in two regions of the phase plane. Letting  $t = -\tau$ , we examine the solutions of

$$\frac{dx_1}{d\tau} = x_1 - u_1$$

$$\frac{dx_2}{d\tau} = 2x_2 - u_1 - u_2$$

leaving the origin with steering

$$u_1(\tau) = \text{sgn}(\eta_1 e^{-\tau} + \eta_2 e^{-2\tau}), \quad u_2(\tau) = \text{sgn}(\eta_2 e^{-2\tau}).$$

Taking  $\eta_2 = 0$ , we can choose  $u_2$  to be any value between  $-1$  and  $1$  and switch values any time we please. We restrict ourselves to values  $-1$ ,  $0$ , or  $1$ .

The curve  $\alpha$  in Figure 3 corresponds to  $\eta_1 < 0$ ,  $\eta_2 = 0$ , and  $u_1 = -1$ ,  $u_2 = 1$ . If we switch to  $u_2 = 0$ , the trajectories leaving  $\alpha$  are parabolas with vertices at  $(-1, -\frac{1}{2})$ . The curve  $\beta$  corresponds to switching at time  $\tau = 0$ , and every point between  $\alpha$  and  $\beta$  has an optimal trajectory passing through it. The curve  $\beta$  corresponds to leaving the origin with  $u_1 = -1$ ,  $u_2 = 0$ . Switching to  $u_2 = -1$ , we leave  $\beta$  at any point we please along parabolas with vertices at  $(-1, 1)$ . The curve  $\gamma$  corresponds to switching to  $u_2 = -1$  at time  $t = 0$ ; that is, to leaving the origin with  $u_1 = -1$ ,  $u_2 = -1$ . All points between  $\beta$  and  $\gamma$  can be reached in this way. Picking  $\eta_1 > 0$  and  $\eta_1 + \eta_2 < 0$ , we see that we can switch to  $u_1 = 1$  at any point along  $\gamma$  we please. The trajectories leaving  $\gamma$  are parabolas with vertices at  $(1, 0)$ , and each point between  $\gamma$  and  $\alpha$  can be reached in this way. The switching curves  $\alpha'$ ,  $\beta'$ , and  $\gamma'$  are obtained by symmetry.




FIGURE 3

## REFERENCES

- [1] BELLMAN, R., GLICKSBERG, I., and GROSS, O., On the "bang-bang" control problem, *Quart. Appl. Math.*, 14 (1956), 11-18.
- [2] BUSHAW, D. W., Ph. D. Thesis, Department of Mathematics, Princeton University, 1952; Differential equations with a discontinuous forcing term, Experimental Towing Tank, Stevens Institute of Technology, Report No. 469 (January 1953); Optimal discontinuous forcing terms, *Contributions to the Theory of Nonlinear Oscillations*, IV, Princeton, 1958.
- [3] EGGLESTON, H. G., *Convexity*, Cambridge Tracts in Math. and Math. Phys., No. 47, Cambridge, 1958.
- [4] GAMKRELIDZE, R. V., Theory of time-optimal processes for linear systems, *Izv. Akad. Nauk SSSR. Ser. Mat.*, 22 (1958), 449-474 (Russian).
- [5] HALMOS, P. R., The range of a vector measure, *Bull. Amer. Math. Soc.*, 54 (1948), 416-421.
- [6] KRASOVSKII, N. N., Concerning the theory of optimal control; *Avtomat. i Telemekh.*, 18 (1957), 960-970 (Russian).
- [7] LASALLE, J. P., Abstract 247t, *Bull. Amer. Math. Soc.*, 60 (1954), 154; Study of the basic principle underlying the "bang-bang" servo, Goodyear Aircraft Corp. Report GER-5518 (July 1953).
- [8] LIAPUNOV, A., Sur les fonctions-vecteurs completement additives, *Bull. Acad. Sci. URSS (Izv. Akad. Nauk SSSR. Ser. Mat.)* 4 (1940), 465-478.



## II. CONTINUOUS DEPENDENCE ON A PARAMETER

J. Jarník and J. Kurzweil

Gichman [1] was the first to show that the averaging principle is a consequence of a modified theorem on continuous dependence on a parameter (cf. also [2] and [3]). Let the solutions of

$$(1) \quad \frac{dx}{dt} = f(x, t)$$

be unique. In [4] it was proved that the solutions of

$$(2) \quad \frac{dx}{dt} = f_k(x, t), \quad k = 1, 2, 3, \dots$$

tend uniformly to the solutions of (1) with  $k \rightarrow \infty$  if

$$(3) \quad \int_0^t f_k(x, \tau) d\tau \rightarrow \int_0^t f(x, \tau) d\tau$$

uniformly and if the following Condition A is fulfilled. ( $x, f, f_k$  are  $n$ -vectors,  $f, f_k$  are continuous for  $x \in G, t \in \langle 0, T \rangle, G$  open.)<sup>1</sup>

Let  $\omega_1(\eta), \omega_2(\eta)$  be non-decreasing for  $0 \leq \eta \leq \sigma, \psi(\eta) = \omega_1(\eta)\omega_2(\eta), \eta^{-1}\psi(\eta)$  non-decreasing.<sup>2</sup>

CONDITION A:

$$(4) \quad \left| \int_{t_1}^{t_2} f_k(x, t) dt \right| \leq \omega_1(|t_2 - t_1|)$$

<sup>1</sup> Cf. [4], Theorems 4, 2, 1, which is formulated for generalized differential equations. In the special case of equations (1), (2) with  $f, f_k$  continuous, every solution is necessarily regular (cf. Definitions 4, 2, 1).

<sup>2</sup> It is sufficient that  $\psi(\eta)$  be non-decreasing. This will be proved in a paper by J. Kurzweil, which will be published in Czechoslovak Math. Journal.

for  $|t_2 - t_1| \leq \sigma$ ,  $k = 1, 2, 3, \dots$

$$(5) \quad \left| \int_{t_1}^{t_2} (f_k(x_2, t) - f_k(x_1, t)) dt \right| \leq |x_2 - x_1| \omega_2(|t_2 - t_1|)$$

for  $|t_2 - t_1| \leq \sigma$ ,  $|x_2 - x_1| \leq 2\omega_1(\sigma)$ ,  $k = 1, 2, 3, \dots$

$$(6) \quad \sum_{i=1}^{\infty} 2^{i\psi\left(\frac{\sigma}{2^i}\right)}$$

Consider the special sequence

$$(2') \quad \frac{dx}{dt} = xk^{1-\alpha} \cos kt + k^{1-\beta} \sin kt$$

$$0 < \alpha, \beta \leq 1, k = 1, 2, 3, \dots$$

and the equation

$$(1') \quad \frac{dx}{dt} = 0$$

$$(n = 1, G = E(x; |x| < 2)).$$

Then (3) is obviously fulfilled, (4) and (5) hold with  $\omega_1(\eta) = K_1 \eta^\gamma$ ,  $\omega_2(\eta) = K_2 \eta^\alpha$ ,  $\gamma = \min(\alpha, \beta)$  ( $K_1, K_2$  independent of  $k$ ) and (6) holds for  $\alpha + \gamma > 1$ . Consequently the solutions of (2') tend to the solutions of (1') with  $k \rightarrow \infty$  if  $\alpha + \gamma > 1$ , i.e.,

$$(7) \quad \alpha + \beta > 1, \quad \alpha > \frac{1}{2}.$$

By direct calculation it is easy to show that the solutions of (2') tend to the solutions of (1') precisely if  $\alpha + \beta > 1$  (cf. [4], p. 419).

Our aim is to prove that condition (6) cannot be weakened even if we restrict ourselves to linear equations. In the case of linear equations we shall always suppose that

$$\omega_1(\eta) \geq c\omega_2(\eta), \quad c > 0.$$

As  $\omega_1(\eta), \omega_2(\eta)$  are moduli of continuity (cf. (4) and (5)), we shall

suppose that

$$(8) \quad \omega_j(\eta_1) + \omega_j(\eta_2) \geq \omega_j(\eta_1 + \eta_2), \quad j = 1, 2.$$

THEOREM. Let  $\eta^{-\lambda}\omega_1(\eta)$ ,  $\eta^{-\lambda}\omega_2(\eta)$ , ( $0 \leq \eta \leq 1$ ) be non-decreasing for a positive  $\lambda$ ,  $\psi(\eta) = \omega_1(\eta)\omega_2(\eta)$ ,

$$(9) \quad \omega_1(\eta) \geq c\omega_2(\eta), \quad c > 0,$$

$$(10) \quad \sum_{i=1}^{\infty} 2^i \psi\left(\frac{1}{2^i}\right).$$

Then there exists a sequence of linear differential equations

$$(11) \quad \frac{dx}{dt} = a_k(t)x + b_k(t),$$

( $n = 1$ ,  $G = E(x; |x| < 2)$ ), such that (4) and (5) are fulfilled,

$$\int_0^t a_k(\tau) d\tau \longrightarrow 0, \quad \int_0^t b_k(\tau) d\tau \longrightarrow 0$$

uniformly, while the sequence of solutions  $x_k(t)$  of (11) (with  $x_k(0) = 0$ ) does not converge.

Let us explain the rôle of the inequality  $\alpha > \frac{1}{2}$  in (7). Consider the sequence

$$(2'') \quad \frac{dx}{dt} = a_k(t)x + b_k(t),$$

where  $x$ ,  $b_k$  are  $n$ -vectors,  $a_k$  are  $n \times n$ -matrices. Suppose that  $a_k(t_1)a_k(t_2) = a_k(t_2)a_k(t_1)$  for  $t_1, t_2 \in (0, T)$ ,  $k = 1, 2, 3, \dots$ . Then the solutions of (2'') tend to the solutions of

$$(1'') \quad \frac{dx}{dt} = a(t)x + b(t)$$

if

$$A_k(t) = \int_0^t a_k(\tau) d\tau \longrightarrow A(t) = \int_0^t a(\tau) d\tau$$

uniformly,

$$B_k(t) = \int_0^t b_k(\tau) d\tau \longrightarrow B(t) = \int_0^t b(\tau) d\tau$$

uniformly,

$$|B_k(t_2) - B_k(t_1)| \leq \omega_1(|t_2 - t_1|), \quad |A_k(t_2) - A_k(t_1)| \leq \omega_2(|t_2 - t_1|)^3,$$

if (6) holds (with  $\psi = \omega_1 \omega_2$ ) and if  $\psi(\eta)$  is non-decreasing. This follows from the representation

$$x_k(t) = \exp(A_k(t)) \left[ 2 + \int_0^t \exp(-A_k(\tau)) dB_k(\tau) \right],$$

where

$$\int_0^t \exp(-A_k(\tau)) dB_k(\tau)$$

tends to

$$\int_0^t \exp(-A(\tau)) dB(\tau)$$

if  $\eta^{-1}\psi(\eta)$  is non-decreasing according to [4], Theorem 3.2. (In the case that merely  $\psi(\eta)$  is non-decreasing, cf. footnote 2.) Let  $b_k(t) = b(t) = 0$ . Then the solutions of (2'') converge to the solutions of (1'') if (and only if)  $A_k(t) \longrightarrow A(t)$ . This situation specially occurs for  $n = 1$ .

In the non-commutative case it follows from the Theorem that there exists such a sequence of linear homogeneous equations that (3), (4) and (5) are fulfilled (with  $\omega_1(\eta) = \omega_2(\eta)$ ) and the sequence of solutions of these equations does not converge, if

$$\sum_{i=1}^{\infty} 2^{i\psi_2\left(\frac{1}{2^i}\right)} = \infty,$$

$\psi_2(\eta) = \omega_2^2(\eta)$ ,  $\eta^{-\lambda}\psi_2(\eta)$  non-decreasing,  $\lambda > 0$ . (It suffices to take  $n = 2$  and

<sup>3</sup> If  $Z$  is a matrix and  $Z_{ij}$  are its elements, we put, for example,  $|Z| = (\sum_{i,j} Z_{ij}^2)^{\frac{1}{2}}$ .

$$\frac{dx_1}{dt} = a_k(t)x_1 + b_k(t)x_2, \quad ,$$

$$\frac{dx_2}{dt} = 0$$

$$x_{1k}(0) = 0, \quad x_{2k}(0) = 1, \quad k = 1, 2, 3, \dots$$

with  $a_k(t)$ ,  $b_k(t)$  from the Theorem). This explains the appearance of the inequality  $\alpha > \frac{1}{2}$  in (7), as a general result on the continuous dependence on a parameter applied to the sequence (2').

PROOF. Let

$$\exp(-A_k(t)) = C_1 + \sum_{i=k}^{m_k} l_1 \omega_2\left(\frac{1}{2^i}\right) \sin 2^i t, \quad ,$$

$$B_k(t) = C_2 \sum_{i=k}^{m_k} l_1 \omega_1\left(\frac{1}{2^i}\right) \cos 2^i t, \quad ,$$

$$a_k(t) = \frac{d}{dt} A_k(t), \quad b_k(t) = \frac{d}{dt} B_k(t), \quad k = 1, 2, 3, \dots$$

where the constants  $C_1$ ,  $C_2$ ,  $l_1$ ,  $l_2$ ,  $l_3$ , ...,  $m_1$ ,  $m_2$ ,  $m_3$ , ...,  $m_k > k$ , will be chosen later.

We shall prove

$$(1) \quad A_k(t) - A_k(0) = \int_0^t a_k(\tau) d\tau \longrightarrow 0 \text{ uniformly,}$$

$$B_k(t) - B_k(0) = \int_0^t b_k(\tau) d\tau \longrightarrow 0 \text{ uniformly with } k \longrightarrow \infty.$$

$$(2) \quad \omega_2(\eta) \text{ is a modulus of continuity of } A_k(t) = \psi = \min\left(\frac{c}{4}, \frac{1}{4}\right) \\ \text{(cf. (9)) } \frac{1}{2} \omega_1(\eta) \text{ is a modulus of continuity of } B_k(t), \quad k = 1, 2, 3, \dots$$

$$(3) \quad \text{Consequently (4) and (5) are fulfilled. } (G = E(x; |x| < 2)) \\ \text{the sequence } x_k^{(t)} \text{ diverges for } t \neq 0.$$

Let us choose the sequence  $l_1$ . Let

$$L_s = E \left[ 1; 1 > s; \omega_j \left( \frac{1}{2^1} \right) \leq 2^{s-1} \omega_j \left( \frac{1}{2^s} \right) \frac{1}{C} \left( 1 + \frac{1}{C} \right)^{1-s-1} \text{ for } j = 1 \text{ or } j = 2 \right],$$

$$s = 1, 2, 3, \dots$$

$$1 = 2, 3, 4, \dots,$$

$$L = \bigcup_{s=1}^{\infty} L_s, \quad M = Q - L,$$

where  $Q$  is the set of all natural numbers and  $C > 1$  will be chosen later. We put

$$l_1 = \begin{cases} 1 & \text{for } 1 \in M, \\ 0 & \text{for } 1 \in L. \end{cases}$$

As  $\eta^{-\lambda} \omega_j(\eta)$  is non-decreasing,

$$(12) \quad \omega_j \left( \frac{1}{2^{1+1}} \right) \leq 2^{-\lambda} \omega_j \left( \frac{1}{2^1} \right), \quad j = 1, 2, \quad 1 = 1, 2, 3, \dots$$

Let us estimate

$$\begin{aligned} \sum_{1 \in L_s} 2^{1-s} \left( \frac{1}{2^1} \right) &\leq \sum_{1 > s} 2^1 \left[ \omega_1 \left( \frac{1}{2^1} \right) 2^{s-1} \omega_2 \left( \frac{1}{2^s} \right) + \right. \\ &\quad \left. + \omega_2 \left( \frac{1}{2^1} \right) 2^{s-1} \omega_1 \left( \frac{1}{2^s} \right) \right] \frac{1}{C} \left[ 1 + \frac{1}{C} \right]^{1-s-1} \leq \\ &\leq 2^s \psi \left( \frac{1}{2^s} \right) \frac{2}{C} \sum_{1 > s} \left[ 2^{-\lambda} \left( 1 + \frac{1}{C} \right) \right]^{1-s-1}. \end{aligned}$$

Let  $C$  be so large that

$$4 \sum_{1=0}^{\infty} \left[ 2^{-\lambda} \left( 1 + \frac{1}{C} \right) \right]^1 < C;$$

it follows that

$$\sum_{1 \in L_s} 2^1 \psi\left(\frac{1}{2^1}\right) < \frac{1}{2} 2^s \psi\left(\frac{1}{2^s}\right).$$

Consequently

$$\begin{aligned} \sum_{1=1}^k \ell_1 2^1 \psi\left(\frac{1}{2^1}\right) &\geq \sum_{1=1}^k 2^1 \psi\left(\frac{1}{2^1}\right) - \sum_{s=1}^k \sum_{1 \in L_s} 2^1 \psi\left(\frac{1}{2^1}\right) \geq \\ &\geq \frac{1}{2} \sum_{1=1}^k 2^1 \psi\left(\frac{1}{2^1}\right) \end{aligned}$$

and

$$(13) \quad \sum_{1=1}^{\infty} \ell_1 2^1 \psi\left(\frac{1}{2^1}\right) = \infty$$

(cf. (10)).

We shall further prove that

$$\begin{aligned} \sum_{1=1}^{r-1} \ell_1 2^{1-r} \omega_j\left(\frac{1}{2^1}\right) &\leq C(C+2) \omega_j\left(\frac{1}{2^r}\right) \\ (14) \quad j &= 1, 2, \quad r = 2, 3, 4, \dots \end{aligned}$$

If  $r \in M$ , then

$$\begin{aligned} \omega_j\left(\frac{1}{2^r}\right) &\geq 2^{1-r} \omega_j\left(\frac{1}{2^1}\right) \frac{1}{C} \left[1 + \frac{1}{C}\right]^{r-1-1}, \quad 1 = 1, 2, \dots, r-1, \\ \sum_{1=1}^{r-1} \ell_1 \omega_j\left(\frac{1}{2^1}\right) 2^{1-r} &\leq C \omega_j\left(\frac{1}{2^r}\right) \sum_{1=1}^{r-1} \left(1 + \frac{1}{C}\right)^{1+1-r} \leq C(C+1) \omega_j\left(\frac{1}{2^r}\right). \end{aligned}$$

If  $r \in L$  (obviously  $1 \in M$ ), then there exists such an  $r_1 < r$ ,  $r_1 \in M$  that  $1 \in L$  for  $r_1 < 1 \leq r$ . Then

$$\sum_{i=1}^{r_1-1} \ell_1 \omega_j \left( \frac{1}{2^i} \right) 2^{i-r} = 2^{r_1-r} \sum_{i=1}^{r_1-1} \ell_1 \omega_j \left( \frac{1}{2^i} \right) 2^{i-r_1} + 2^{r_1-r} \omega_j \left( \frac{1}{2^{r_1}} \right) \leq$$

$$\leq C(C+1) \omega_j \left( \frac{1}{2^{r_1}} \right) 2^{r_1-r} + 2^{r_1-r} \omega_j \left( \frac{1}{2^{r_1}} \right) \leq$$

$$\leq C(C+2) 2^{r_1-r} \omega_j \left( \frac{1}{2^{r_1}} \right) \leq C(C+2) \omega_j \left( \frac{1}{2^r} \right)$$

( $C > 1$ ) (cf. (8)) and (14) holds for  $r = 2, 3, 4, \dots$ .

Let us prove 1), 2), 3).

1) According to (12),

$$\sum_{i=1}^{\infty} \omega_j \left( \frac{1}{2^i} \right) < \infty.$$

If

$$C_1 > \sum_{i=1}^{\infty} \omega_2 \left( \frac{1}{2^i} \right),$$

then  $A_k(t) - A_k(0) \rightarrow 0$ ,  $B_k(t) - B_k(0) \rightarrow 0$  uniformly.

2) Let

$$S_k(t) = \sum_{i=1}^{\infty} d_1 \omega_1 \left( \frac{1}{2^i} \right) \cos 2^i t,$$

$$T_k(t) = \sum_{i=1}^{\infty} d_1 \omega_2 \left( \frac{1}{2^i} \right) \sin 2^i t,$$

$$d_1 = \ell_1 \quad \text{for } i = k, k+1, \dots, m_k,$$

$$d_1 = 0 \quad \text{for } i < k \text{ or } i > m_k,$$

$$\frac{1}{2^{r-1}} \geq |t_2 - t_1| > \frac{1}{2^r}.$$



Then

$$\begin{aligned}
 |S_k(t_2) - S_k(t_1)| &\leq \sum_{i=1}^{r-1} d_i \omega_1\left(\frac{1}{2^i}\right) \left| \sin 2^i \frac{t_2 - t_1}{2} \right| + \sum_{i=r}^{\infty} \omega_1\left(\frac{1}{2^i}\right) \leq \\
 &\leq \sum_{i=1}^{r-1} l_i \omega_1\left(\frac{1}{2^i}\right) 2^{i-r} + \omega_1\left(\frac{1}{2^r}\right) \sum_{i=0}^{\infty} 2^{-\lambda i} \leq K \omega_1\left(\frac{1}{2^r}\right) \leq \\
 &\leq K \omega_1(|t_2 - t_1|) ,
 \end{aligned}$$

$$K = C(C+2) + (1 - 2^{-\lambda})^{-1}$$

(cf. (12) and (14)). Similarly .

$$|T_k(t_2) - T_k(t_1)| \leq K \omega_2(|t_2 - t_1|) .$$

If  $C_1, C_2$  are large enough, then 2) is fulfilled.

3) Obviously

$$\begin{aligned}
 b_k(t) &= -C_2 \sum_{i=k}^{m_k} l_i 2^i \omega_1\left(\frac{1}{2^i}\right) \sin 2^i t , \\
 x_k(t) &= \exp(A_k(t)) \int_0^t \exp(-A_k(\tau)) b_k(\tau) d\tau = \\
 &= - \left[ C_1 + \sum_{i=k}^{m_k} l_i \omega_2\left(\frac{1}{2^i}\right) \sin 2^i t \right]^{-1} \times \\
 &\int_0^t \left[ C_1 + \sum_{i=k}^{m_k} l_i \omega_2\left(\frac{1}{2^i}\right) \sin 2^i \tau \right] C_2 \sum_{i=k}^{m_k} 2^i l_i \omega_1\left(\frac{1}{2^i}\right) \sin 2^i \tau d\tau .
 \end{aligned}$$

As

$$C_1 > \sum_{i=1}^{\infty} \omega_2\left(\frac{1}{2^i}\right) ,$$

it follows that

$$\begin{aligned}
|x_k(t)| &\geq C_3 \left[ \sum_{i=k}^{m_k} 2^{i\ell_1} \psi\left(\frac{1}{2^i}\right) \int_0^t \sin^2 2^i \tau \, d\tau \right. \\
&- C_1 \sum_{i=k}^{m_k} 2^{i\ell_1} \omega_1\left(\frac{1}{2^i}\right) \left| \int_0^t \sin 2^i \tau \, d\tau \right| - \\
&- \sum_{1, j=k}^{m_k} 2^{i\ell_1} \ell_j \omega_1\left(\frac{1}{2^i}\right) \omega_2\left(\frac{1}{2^j}\right) \left| \int_0^t \sin 2^i \tau \sin 2^j \tau \, d\tau \right| \Big],
\end{aligned}$$

$$C_3 > 0 \quad .$$

If  $j \neq 1$ , then

$$\begin{aligned}
2^i \left| \int_0^t \sin 2^i \tau \sin 2^j \tau \, d\tau \right| &\leq \frac{2^i}{|2^i - 2^j|} + \frac{2^i}{2^i + 2^j} \leq 3 \\
\sum_{\substack{1, j=k \\ j \neq 1}}^{m_k} 2^{i\ell_1} \ell_j \omega_1\left(\frac{1}{2^i}\right) \omega_2\left(\frac{1}{2^j}\right) \left| \int_0^t \sin 2^i \tau \sin 2^j \tau \, d\tau \right| &\leq \\
&\leq 3 \left[ \sum_{i=k}^{m_k} \omega_1\left(\frac{1}{2^i}\right) \right] \left[ \sum_{i=k}^{m_k} \omega_2\left(\frac{1}{2^i}\right) \right] \quad .
\end{aligned}$$

Consequently

$$\begin{aligned}
|x_k(t)| &\geq C_3 \left\{ \frac{t}{2} \sum_{i=k}^{m_k} 2^{i\ell_1} \psi\left(\frac{1}{2^i}\right) - \frac{1}{4} \sum_{i=k}^{m_k} \ell_1 \psi\left(\frac{1}{2^i}\right) \sin 2^{i+1} t \right. \\
&- 2C_1 \sum_{i=k}^{m_k} \omega_1\left(\frac{1}{2^i}\right) - 3 \left[ \sum_{i=k}^{m_k} \omega_1\left(\frac{1}{2^i}\right) \right] \left[ \sum_{i=k}^{m_k} \omega_2\left(\frac{1}{2^i}\right) \right] \Big\}
\end{aligned}$$

Let us choose  $m_k$  in such a manner that

$$\sum_{i=k}^{m_k} 2^{i/2} \psi\left(\frac{1}{2^i}\right) \longrightarrow \infty$$

with  $k \longrightarrow \infty$  (cf. (13)). Then  $|x_k(t)| \longrightarrow \infty$  with  $k \longrightarrow \infty$  for  $t \neq 0$  (cf. (12)).

## REFERENCES

- [1] GICHMAN, I. I., Concerning a theorem of N. N. Bogoljubov, Ukrain. Math. Journal, 4, 2, 215-219 (in Russian).
- [2] KRASNOSELSKIJ, M. A. and KREJN, S. G., Averaging principle in non-linear mechanics, Uspechi mat. nauk., 10, 3 (1955), 147-152 (in Russian).
- [3] KURZWEIL, J. and VOREL, Z., Continuous dependence on a parameter of solutions of differential equations, Czechoslovak math. journal, 7, (82), 4 (1957), 568-583 (in Russian).
- [4] KURZWEIL, J., Generalized ordinary differential equations and continuous dependence on a parameter, Czech. math. journal, 7 (82), 1957, 3, 418-449.



### III. POINCARÉ'S PERTURBATION METHOD AND TOPOLOGICAL DEGREE

Jane Cronin

#### §1. INTRODUCTION

We study the periodic solutions of a system of differential equations

$$(1.1) \quad \frac{dx}{dt} = F(x, t, \mu)$$

where  $x$ ,  $F$  are  $n$ -vectors, appropriate differentiability and periodicity conditions are imposed on the components of  $F$ , and  $\mu$  is a real parameter. Poincaré's perturbation method is used and the results extended by using topological degree. The technique developed gives a general approach to the existence problem for the degenerate case (the case for which the variational equation has periodic solutions) and new existence theorems are obtained if the degree of degeneracy exceeds one. (In problems of mechanical or electrical oscillations, the degree of degeneracy is frequently greater than one.) The technique consists in using topological degree to make an 'in the large' study of the bifurcation system (Verzweigungsgleichungen) without imposing any local uniqueness conditions on the solutions of the bifurcation system. The results obtained are generalizations and refinements of the results in a previous paper [7], (Numbers in brackets refer to the bibliography at the end of this paper.) although the present paper is independent of [7]. Existence theorems are obtained for a wider class of equations; autonomous systems are studied; and it is shown that for the totally degenerate case (degree of degeneracy equals dimension of the system), the topological degree is a kind of lower bound for the number of distinct periodic solutions.

The literature on this subject is extensive. General treatments are described briefly in [7]. Bass [2] has given a large bibliography including not only general treatments but papers on applications. The work of Friedrichs [8, 9, 10], Coddington and Levinson [3, 4] and

Lefschetz [11, 12] are most closely related to our approach. We base our treatment on the results of Coddington and Levinson concerning properties of the bifurcation system. As in Lefschetz's work, we need not impose local uniqueness conditions on the solutions of the bifurcation system, and our results may be regarded as an extension of the criterion for the existence of real periodic solutions given by Lefschetz.

In Section 2, we describe how the bifurcation system is set up. Periodic solutions of the original differential equation correspond to solutions of the bifurcation system. We study solutions of the bifurcation system by determining the topological degree of the mapping defined by the bifurcation system. By using the properties of the bifurcation system obtained by Coddington and Levinson [3, 4], we show in Section 2 that the problem of determining the topological degree can be reduced to that of determining the topological degree of a mapping in Euclidean  $q$ -space where  $q$  is the degree of degeneracy.

In Section 3, we apply the method to obtain some existence theorems for periodic solutions of non-autonomous differential equations. In particular we show that if the function  $F(x, t, \mu)$  in (1.1) is such that  $F(x, t, \mu) = Ax + \mu f(x, t, \mu)$  where  $A$  is a constant matrix and  $f(x, t, \mu) = f_1(x, \mu) + f_2(t, \mu)$  where the components of  $f_1(x, 0)$  behave like polynomials for large  $x$ , then, except for cases for which the topological degree is not defined, equation (1.1) has, for sufficiently small  $\mu$ , at least one periodic solution near the initial solution.

In Section 4, the exceptional case in which the topological degree is not defined is discussed. In Section 5, the case in which the topological degree is even is briefly considered. In Section 6, we prove that the topological degree is, for the totally degenerate case, a kind of lower bound for the number of distinct periodic solutions. Finally in Section 7, application of topological degree to an autonomous differential equation is described.

I am indebted to Professor S. Lefschetz for a number of extremely helpful discussions of this material.

Most of the work on this paper was done under the sponsorship of the Office of Naval Research contract Nonr-1858(04).

## §2. THE BIFURCATION SYSTEM

In [8], Friedrichs treated the equation

$$(2.1) \quad \dot{x} = F(x, t, \mu) \quad (\dot{\phantom{x}} = d/dt)$$

where  $x$ ,  $F$  are  $n$ -vectors; the components of  $F$  have continuous derivatives with respect to  $x$ ,  $t$ , and  $\mu$ ; these derivatives have continuous derivatives with respect to  $x$ ;  $F(x, t, \mu)$  is periodic in  $t$  with period  $T(\mu)$  where  $d^2T/d\mu^2$  is continuous; and (2.1) has for  $\mu = 0$  a solution  $x_0(t)$  of period  $T(0)$ . The problem is to determine if for sufficiently small  $\mu$ , equation (2.1) has a solution  $x(t, \mu)$  periodic of period  $T(\mu)$ . The classical result of Poincaré ([13], Vol. 1, Chapter IV) states that if the variational equation  $\dot{x} = A(t)x$ , where

$$A(t) = \left. \frac{\partial F}{\partial x} \right|_{x=x_0(t), \mu=0},$$

has no non-zero solutions of period  $T(0)$ , then for sufficiently small  $\mu$ , equation (2.1) has a unique solution  $x(t, \mu)$ , periodic of period  $T(\mu)$ , such that

$$\lim_{\mu \rightarrow 0} x(t, \mu) = x_0(t).$$

Here we treat the degenerate case in which the variational equation has  $q$  linearly independent solutions, periodic of period  $T(0)$ , where  $1 \leq q \leq n$ . The number  $q$  is called the degree of degeneracy of the problem.

Friedrichs shows that if

$$\left. \frac{\partial F}{\partial \mu} \right|_{x=x_0(t), \mu=0} = 0,$$

then the problem can be reduced to the study of the equation

$$(2.2) \quad \dot{x} = A(t)x + \mu f(x, t, \mu)$$

and then derives a bifurcation system for (2.2). Coddington and Levinson ([3], [4]), by imposing the further condition that  $A(t)$  is a constant matrix and then putting this constant matrix in a canonical form, derive a more explicit form for the bifurcation system. (Coddington and Levinson also assume that  $T(\mu)$  is a constant function. This is not a restriction in the generality because Friedrichs shows in [8] that the general case may be reduced to the case  $T(\mu)$  a constant.)

Now we describe the bifurcation system derived by Coddington and Levinson. The system is obtained by using the variation of constants formula to describe the solution and then imposing the condition that the solution be periodic of period  $2\pi$  (i.e., assume  $T(0) = 2\pi$ .) The system

is:

$$(2.3) \quad (e^{2\pi A} - E)c + \mu \int_0^{2\pi} \{e^{(2\pi-s)A} f[x(s, \mu, c), s, \mu]\} ds = 0,$$

(this is equation (1.12), p. 23, of [3] or equation (3.20), p. 360 of [4]) where  $E$  is the identity matrix,  $c$  is an  $n$ -vector such that  $x(t, \mu, c)$  is a solution of (2.2) with initial value  $c$ , i.e.,  $x(0, \mu, c) = c$ , and  $x(t, \mu, c)$  has period  $2\pi$ . Thus (2.3) is a system of  $n$  equations in the  $n$  unknowns which are the components  $c_1, \dots, c_n$  of  $c$ . The problem of finding periodic solutions for (2.2) is reduced to that of solving (2.3) for  $c_1, \dots, c_n$ .

In order to study the properties of (2.3), Coddington and Levinson assumed that matrix  $A$  already has the following real canonical form

$$A = \begin{pmatrix} A_1 & & & & & & \\ & \ddots & & & & & \\ & & A_k & & & & \\ & & & B_1 & & & \\ & & & & \ddots & & \\ & & & & & B_m & \\ & & & & & & C \end{pmatrix}$$

where the elements not shown are zeros. Each  $A_j$ ,  $j = 1, \dots, k$ , is a matrix of  $\alpha_j$  ( $\alpha_j$  even) rows and columns of the form

$$A_j = \begin{pmatrix} S_j & & & & \\ E_2 & S_j & & & \\ & \ddots & \ddots & & \\ & & E_2 & S_j & \\ & & & & S_j \end{pmatrix}$$

where all elements are zero except  $S_j$  and  $E_2$ , and

$$S_j = \begin{pmatrix} 0 & -N_j \\ N_j & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$



where  $N_j$  is a positive integer. A matrix  $A_j$  may have only two rows and columns in which case  $A_j = S_j$ . Each matrix  $B_j$  has  $\beta_j$  rows and columns,  $j = 1, \dots, m$ , and is of the form

$$B_j = \begin{pmatrix} 0 & 0 & \dots & \dots & 0 \\ 1 & 0 & & & \vdots \\ 0 & 0 & \ddots & & \vdots \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 1 & 0 \end{pmatrix}$$

where  $B_j$  may have only one row and column in which case  $B_j$  consists of the single element 0. The matrix  $C$  has  $(n - \sum_{j=1}^k \alpha_j - \sum_{j=1}^m \beta_j)$  rows and columns and has no characteristic roots of the form  $iN$  for any integer  $N$  including  $N = 0$ . Matrix  $C$  need not be in canonical form.

If  $(c_1, \dots, c_1, \dots, c_n)$  is an  $n$ -vector, the indices  $i$  corresponding to the last two rows of any  $A_j$  or to the last row of any  $B_j$  are called exceptional indices. They are the indices with the following form:

$$\begin{aligned} i &= \alpha_1 + \alpha_2 + \dots + (\alpha_j - 1) \\ i &= \alpha_1 + \alpha_2 + \dots + \alpha_j \end{aligned}$$

where  $j = 1, \dots, k$  and

$$i = \alpha_1 + \dots + \alpha_k + \beta_1 + \dots + \beta_j$$

where  $j = 1, \dots, m$ . The indices  $i$  corresponding to the first two rows of any  $A_j$  or to the first row of any  $B_j$  are called singular indices. They are the indices with the following form:

$$\begin{aligned} i &= 1, 2, \alpha_1 + 1, \alpha_1 + 2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{k-1} + 1, \\ &\quad \alpha_1 + \alpha_2 + \dots + \alpha_{k-1} + 2, \alpha_1 + \dots + \alpha_k + 1, \\ &\quad \alpha_1 + \dots + \alpha_k + \beta_1 + 1, \dots, \alpha_1 + \dots + \alpha_k + \beta_1 + \dots + \beta_{m-1} + 1 \end{aligned}$$

There are  $(2k + m)$  exceptional indices and  $(2k + m)$  singular indices. The number  $q = 2k + m$  is the degree of degeneracy of the problem. Throughout our study, we assume that  $q > 0$ , i.e., that there is at least one  $A_j$  or one  $B_j$  in the canonical form of matrix  $A$ .

Now let  $(c'_1, \dots, c'_{n-q})$  be the  $(n-q)$ -vector whose components are the components (in the same order) of  $(c_1, \dots, c_n)$  which have

non-exceptional indices and let  $(c_1'', \dots, c_q'')$  be the  $q$ -vector whose components are the components of  $(c_1, \dots, c_n)$  which have exceptional indices. Let subscript  $j$  denote a singular index and  $j'$  a non-singular index. It is shown in [3] and [4] that system (2.3) can be replaced by the system

$$(2.4) \quad N(c_1', \dots, c_{n-q}') + \mu \left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[x(s, \mu, c), s, \mu]} ds \right\}_{(j')} = 0$$

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[x(s, \mu, c), s, \mu]} ds \right\}_{(j)} = 0$$

where  $N$  is a non-singular  $(n-q) \times (n-q)$  matrix acting on vector  $(c_1', \dots, c_{n-q}')$  and

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[x(s, \mu, c), s, \mu]} ds \right\}_{(j')}$$

denotes the vector composed of the  $(n-q)$  components of

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[x(s, \mu, c), s, \mu]} ds \right\}$$

which have non-singular indices. Similarly

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[x(s, \mu, c), s, \mu]} ds \right\}_{(j)}$$

denotes the vector composed of the  $q$  components which have singular indices.

The left side of system (2.4) describes a continuous mapping (call it  $\mathcal{M}$ ) of real Euclidean  $n$ -space into itself. Let  $c''$  denote the vector  $c$  in which the  $(n-q)$  components with non-exceptional indices have been set equal to zero. Then

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[e^{SA_C''}, s, 0]} ds \right\}_{(j)}$$

defines a continuous mapping of real  $q$ -space into itself which we denote by  $\mathcal{M}_0$ . We call the system

$$\left\{ \int_0^{2\pi} e^{(2\pi-s)A_F[e^{SA_C''}, s, 0]} ds \right\}_{(j)} = 0$$

the reduced bifurcation system.

Friedrichs [8] and Coddington and Levinson, [3] and [4] assume the existence of a solution  $c''$  of the reduced bifurcation system such that the Jacobian of the reduced bifurcation system is non-zero at  $c''$ . From this, the existence of a solution of (2.3) and hence of a periodic solution of (2.2) is obtained. We use instead the notion of topological degree.

LEMMA 2.1. For given  $r$ , there is an  $\varepsilon > 0$  such that if  $|\mu| < \varepsilon$ , the topological degree of  $\mathcal{M}$  at the origin and relative to a solid  $(n-1)$ -sphere with radius  $r$  and center at the origin is equal (except possibly for sign) to the topological degree of  $\mathcal{M}_0$  at the origin (in  $q$ -space) and relative to a solid  $(q-1)$ -sphere with radius  $r$  and center at the origin.

PROOF. The proof follows from the definition of topological degree and the invariance under homotopy of the topological degree. (see [1, Deformationsatz, p. 424].)

Lemma 2.1 shows that in order to demonstrate the existence of periodic solutions  $x(t, \mu, c)$  of (2.2) it is sufficient to show that the topological degree of  $\mathcal{M}_0$  is non-zero. For then the degree of  $\mathcal{M}$  is non-zero; hence for given small  $\mu$ , system (2.3) has at least one solution  $c_0 = (c_1^0, \dots, c_n^0)$ . This implies that (2.2) has a solution  $x(t, \mu, c_0)$  of period  $2\pi$  such that  $x(0, \mu, c_0) = (c_1^0, \dots, c_n^0)$ . From the fundamental properties of topological degree, the initial values  $(c_1, \dots, c_n)$  are continuous in  $\mu$  in the following sense: if for given  $\mu_0$ ,  $(c_1^0, \dots, c_n^0)$  is an isolated solution of (2.3) which has non-zero topological index, then given  $\varepsilon > 0$ , there is a  $\delta > 0$  such that if  $|\mu_1 - \mu_0| < \delta$ , there is at least one periodic solution  $x(t, \mu, c_1)$  with initial value  $(c_1^1, \dots, c_n^1)$  such that  $|c_1^1 - c_1^0| < \varepsilon$  for  $i = 1, \dots, n$ ; if  $(c_1^0, \dots, c_n^0)$  is not an isolated solution of (2.3), i.e.,  $(c_1^0, \dots, c_n^0)$  is a limit point of a set  $S$  of solutions  $\{(c_1^v, \dots, c_n^v)\}$  of (2.3) for  $\mu = \mu_0$ , then suppose that the topological degree of an open set  $N$  containing  $S$  but no other solutions of (2.3) is different from zero. Then given  $\varepsilon > 0$ , there is a  $\delta > 0$  such that if  $|\mu - \mu_0| < \delta$ , then there is at least one solution  $x(t, \mu, c)$  with initial value  $(c_1, \dots, c_n)$  such that for some  $(c_1^v, \dots, c_n^v) \in S$ , we have  $|c_1^v - c_1| < \varepsilon$  for  $i = 1, \dots, n$ . From the usual existence theorems for differential equations, it follows that the solutions  $x(t, \mu, c)$  are themselves continuous in  $\mu$  in the same sense.

## §3. EXISTENCE THEOREMS FOR THE NON-AUTONOMOUS CASE

We obtain existence theorems for periodic solutions of the equation

$$(3.1) \quad \dot{x} = Ax + \mu f(x, t, \mu)$$

where  $A$  is a constant matrix;  $\mu$  is a real parameter; and  $f, \partial f / \partial x_1$  are continuous in  $(x, t, \mu)$  for small  $|\mu|$  and all  $(x, t)$ ; and  $f$  has period  $2\pi$  in  $t$ . The periodic solutions of the variational equation  $\dot{x} = Ax$  correspond to the characteristic roots of  $A$  that are of the form  $iN$  where  $N$  is an integer.

ASSUMPTION 1. Suppose that  $A$  has just one pair of characteristic roots of the form  $iN, -iN$  where we assume the integer  $N$  is non-zero. Then  $A$  may be put in the standard form

$$A = \left( \begin{array}{cc|c} 0 & -N & 0 \\ N & 0 & 0 \\ \hline 0 & 0 & A_0 \end{array} \right)$$

where  $A_0$  is not necessarily canonical but is real and has no characteristic roots of the form  $iN$ . We assume that  $A$  is in this standard form.

ASSUMPTION 2. The components  $f_1(x, t, \mu)$  and  $f_2(x, t, \mu)$  of  $f(x, t, \mu)$  have the form

$$f_1(x, t, \mu) = H_1(x_1, \dots, x_n, \mu) + K_1(t, \mu) \quad (i = 1, 2)$$

where  $x_1, \dots, x_n$  are the components of  $x$ , and  $H_1, K_1$  are functions of the indicated variables which have continuous second derivatives in these variables. The functions  $K_1(t, \mu)$  and  $K_2(t, \mu)$  have period  $2\pi$  in  $t$ . Also there exist polynomials  $P_1(x_1, x_2)$  and  $P_2(x_1, x_2)$  such that

$$\lim_{r \rightarrow \infty} \frac{H_1(x_1, x_2, 0, \dots, 0, 0)}{P_1(x_1, x_2)} = 1 \quad (i = 1, 2,)$$

where  $r^2 = x_1^2 + x_2^2$ .

Two real independent periodic solutions of the variational equation are:

$$\begin{aligned} x^{(1)}(t) &= (\cos Nt, \sin Nt, 0, \dots, 0) \\ x^{(2)}(t) &= (-\sin Nt, \cos Nt, 0, \dots, 0) \end{aligned}$$

Following Coddington and Levinson, we may write the reduced bifurcation system as:

$$(3.2) \quad \int_0^{2\pi} \left\{ \cos(Ns) H_1 \left[ c_1 x^{(1)}(-s) + c_2 x^{(2)}(-s), 0 \right] \right. \\ \left. - \sin(Ns) H_2 (c_1 x^{(1)} + c_2 x^{(2)}, 0) \right\} ds + \\ + \int_0^{2\pi} \left\{ \cos(Ns) K_1(-s, 0) - \sin(Ns) K_2(-s, 0) \right\} ds = 0.$$

$$(3.3) \quad \int_0^{2\pi} \left\{ \sin(Ns) H_1 \left[ c_1 x^{(1)}(-s) + c_2 x^{(2)}(-s), 0 \right] \right. \\ \left. + \cos(Ns) H_2 (c_1 x^{(1)} + c_2 x^{(2)}, 0) \right\} ds + \\ + \int_0^{2\pi} \left\{ \sin(Ns) K_1(-s, 0) + \cos(Ns) K_2(-s, 0) \right\} ds = 0.$$

(Here for convenience we have not indicated the components of  $c_1 x^{(1)} + c_2 x^{(2)}$  separately although  $H_1$  and  $H_2$  are actually functions of these components. We use this convention with  $P_1$  and  $P_2$  also.) Let  $k_1$  and  $k_2$  denote the constants which are the second integrals on the left sides of equations (3.2) and (3.3). Let  $\#_1(c_1, c_2)$  and  $\#_2(c_1, c_2)$  denote the first integrals on the left sides of (3.2) and (3.3), and let  $\mathcal{P}_1(c_1, c_2)$  and  $\mathcal{P}_2(c_1, c_2)$  denote the polynomials obtained if in  $\#_1$  and  $\#_2$ , the terms  $H_1(c_1 x^{(1)} + c_2 x^{(2)}, 0)$  and  $H_2(c_1 x^{(1)} + c_2 x^{(2)}, 0)$  are replaced by  $P_1(c_1 x^{(1)} + c_2 x^{(2)})$  and  $P_2(c_1 x^{(1)} + c_2 x^{(2)})$ .

ASSUMPTION 3. If  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are of degrees  $n_1$  and  $n_2$  respectively in  $c_1$  and  $c_2$ , let  $Q_1$  and  $Q_2$  be the homogeneous polynomials of degrees  $n_1$  and  $n_2$  in  $\mathcal{P}_1$  and  $\mathcal{P}_2$  respectively. Let  $M : (c_1, c_2) \longrightarrow (c'_1, c'_2)$  be the mapping defined by

$$Q_1(c_1, c_2) = c'_1 \\ Q_2(c_1, c_2) = c'_2.$$

We assume that the topological index at  $(0, 0)$  of  $M$  is defined.

From standard arguments, we have:

LEMMA 3.1. Let  $M_1 : (c_1, c_2) \longrightarrow (c'_1, c'_2)$  be defined by:

$$\begin{aligned}\mathcal{P}_1(c_1, c_2) + k_1 &= c'_1 \\ \mathcal{P}_2(c_1, c_2) + k_2 &= c'_2.\end{aligned}$$

The topological degree of mapping  $M_1$  at  $(0, 0)$  and relative to any sufficiently large circle with center  $(0, 0)$  is equal to the topological index at  $(0, 0)$  of  $M$ .

THEOREM 3.1. Let Assumptions 1, 2, 3 be satisfied. Then for all sufficiently small  $\mu$ , Equation (3.1) has at least one solution of period  $2\pi$ .

PROOF. Let  $\mathcal{M}_0 : (c_1, c_2) \longrightarrow (c'_1, c'_2)$  be defined by:

$$\begin{aligned}\mathcal{H}_1(c_1, c_2) + k_1 &= c'_1 \\ \mathcal{H}_2(c_1, c_2) + k_2 &= c'_2.\end{aligned}$$

It is sufficient by Lemma 2.1 to show that the topological degree of  $\mathcal{M}_0$  at  $(0, 0)$  and relative to a circle with center at  $(0, 0)$  is different from zero.

We show first that relative to any sufficiently large circle, the topological degrees of  $M_1$  and  $\mathcal{M}_0$  at  $(0, 0)$  are the same. From Assumption 3 and the fact that  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are polynomials, it follows that given  $m > 0$ , there exists  $r_1 > 0$  such that if  $c_1^2 + c_2^2 > r_1^2$ , then

$$[\mathcal{P}_1(c_1, c_2) + k_1]^2 + [\mathcal{P}_2(c_1, c_2) + k_2]^2 > m.$$

Further there exists  $\varepsilon > 0$  and  $r_2 > 0$  such that if  $|\eta_1| < \varepsilon$ ,  $|\eta_2| < \varepsilon$ , and  $c_1^2 + c_2^2 > r_2^2$ , then

$$\begin{aligned}& \left[ \int_0^{2\pi} \left\{ \cos(Ns) [\mathcal{P}_1(c_1 x^{(1)} + c_2 x^{(2)})] (1 + \eta_1) \right. \right. \\ & \left. \left. - \sin(Ns) [\mathcal{P}_2(c_1 x^{(1)} + c_2 x^{(2)})] (1 + \eta_2) \right\} ds + k_1 \right]^2 +\end{aligned}$$

$$+ \left[ \int_0^{2\pi} \left\{ \sin(Ns) [P_1(c_1 x^{(1)} + c_2 x^{(2)})] (1 + \eta_1) \right. \right. \\ \left. \left. + \cos(Ns) [P_2(c_1 x^{(1)} + c_2 x^{(2)})] (1 + \eta_2) \right\} ds + k_2 \right]^2 > \frac{1}{2} m.$$

By Assumption 2, there exists  $r_3 > 0$  such that if  $c_1^2 + c_2^2 > r_3^2$ , then

$$H_1(c_1 x^{(1)} + c_2 x^{(2)}, 0) = (1 + \eta_1) P_1(c_1 x^{(1)} + c_2 x^{(2)})$$

where  $|\eta_1| < \varepsilon$  for  $1 = 1, 2$ . Hence the homotopy  $h: (c_1, c_2, t) \longrightarrow (c_1'(t), c_2'(t))$  defined by:

$$c_1'(t) = \int_0^{2\pi} \left\{ \cos(Ns) [P_1(c_1 x^{(1)} + c_2 x^{(2)})] (1 + t\eta_1) \right. \\ \left. - \sin(Ns) [P_2(c_1 x^{(1)} + c_2 x^{(2)})] (1 + t\eta_2) \right\} ds + k_1 \\ c_2'(t) = \int_0^{2\pi} \left\{ \sin(Ns) [P_1(c_1 x^{(1)} + c_2 x^{(2)})] (1 + t\eta_1) \right. \\ \left. - \cos(Ns) [P_2(c_1 x^{(1)} + c_2 x^{(2)})] (1 + t\eta_2) \right\} ds + k_2$$

shows that the topological degrees of  $M_1$  and  $M_0$  at  $(0, 0)$  and relative to any sufficiently large circle with center  $(0, 0)$  are the same.

A trivial computation based on the fact that

$$\int_0^{2\pi} \sin^m(x) \cos^n(x) dx \neq 0$$

only if  $m$  and  $n$  are even shows that  $\mathcal{P}_1$  and  $\mathcal{P}_2$  are polynomials of odd degree in  $c_1$  and  $c_2$ . Then the topological index at  $(0, 0)$  of  $M$  is odd [5] and hence the topological degrees of  $M_1$  and  $M_0$  are different from zero. This completes the proof of Theorem 3.1.

As a second example, we consider Equation (3.1) under the following assumptions.

ASSUMPTION 4. Matrix  $A$  has just one pair of characteristic roots  $iN, -iN$  where  $N$  is a non-zero integer and  $A$  has 0 as a characteristic root. Also  $A$  has the form

$$A = \left( \begin{array}{ccc|c} 0 & -N & 0 & 0 \\ N & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline & 0 & & A_0 \end{array} \right)$$

where  $A_0$  is not necessarily canonical but is real and has no characteristic roots of the form  $iN$ .

ASSUMPTION 5. The components  $f_1(x, t, \mu)$ ,  $f_2(x, t, \mu)$  and  $f_3(x, t, \mu)$  of  $f(x, t, \mu)$  have the form:

$$f_1(x, t, \mu) = H_1(x_1, \dots, x_n, \mu) + K_1(t, \mu) \quad (i = 1, 2, 3)$$

where  $x_1, \dots, x_n$  are the components of  $x$ ; and  $H_1$  and  $K_1$  are functions of the indicated variables and have continuous second derivatives in these variables; and  $K_1, K_2, K_3$  have period  $2\pi$  in  $t$ . There exist polynomials  $P_1(x_1, x_2, x_3)$ ,  $P_2(x_1, x_2, x_3)$ ,  $P_3(x_1, x_2, x_3)$  such that

$$\lim_{r \rightarrow \infty} \frac{H_1(x_1, x_2, x_3, 0, \dots, 0)}{P_1(x_1, x_2, x_3)} = 1 \quad (i = 1, 2, 3).$$

Also the degree of  $P_3$  is  $k$ , odd, and there is a term  $ax_3^k$  (with  $a \neq 0$ ) in  $P_3$ .

Three real independent periodic solutions of the variational equation are:

$$\begin{aligned} x^{(1)} &= (\cos(Nt), \sin(Nt), 0, \dots, 0) \\ x^{(2)} &= (-\sin(Nt), \cos(Nt), 0, \dots, 0) \\ x^{(3)} &= (0, 0, 1, \dots, 0). \end{aligned}$$

The reduced bifurcation system is:

$$\begin{aligned} & \int_0^{2\pi} [\cos(Ns)H_1(c_1x^{(1)} + c_2x^{(2)} + c_3x^{(3)}, 0) \\ & - \sin(Ns)H_2(c_1x^{(1)} + c_2x^{(2)} + c_3x^{(3)}, 0)] ds + \\ & + \int_0^{2\pi} [\cos(Ns)K_1(-s, 0) - \sin(Ns)K_2(-s, 0)] ds = 0 \end{aligned}$$



$$\int_0^{2\pi} [\sin(Ns)H_1 + \cos(Ns)H_2] ds + \int_0^{2\pi} [\sin(Ns)K_1 + \cos(Ns)K_2] ds = 0$$

$$\int_0^{2\pi} H_3(c_1x^{(1)} + c_2x^{(2)} + c_3x^{(3)}, 0) ds = 0.$$

Let  $k_1$ ,  $H_1$ ,  $P_1$ , and  $Q_1$  ( $i = 1, 2, 3$ ) be defined analogously to their counterparts in the preceding example.

ASSUMPTION 6. Let  $M : (c_1, c_2, c_3) \longrightarrow c_1', c_2', c_3'$  be the mapping defined by  $c_i' = Q_i(c_1, c_2, c_3)$  for  $i = 1, 2, 3$ . We assume that the topological index at  $(0, 0)$  of  $M$  is defined.

By the same type of argument as for the preceding example, we obtain:

THEOREM 3.2. Let Assumptions 4, 5, 6 be satisfied.

Then for all sufficiently small  $\mu$ , Equation (3.1)

has at least one solution of period  $2\pi$ .

More complicated examples may be treated in a similar manner.

#### §4. EXCEPTIONAL CASES

By Assumption 3, the topological index at  $(0, 0)$  of mapping  $M$  is defined. Now suppose that this assumption is not satisfied, i.e., suppose the topological index is not defined. This means that  $Q_1, Q_2$  have a common real factor. Hence by varying one coefficient in  $Q_1$  (i.e., varying  $H_1$ ) arbitrarily slightly, we obtain a mapping for which the topological index is defined. When such a change is made, two different cases arise. First suppose that after  $Q_1$  is changed slightly in some definite manner, the topological index  $j$  of the resulting mapping is defined and suppose that  $j$  is odd or that  $|j| > 2$ . Then by using the analysis of [5, Section 3] it can be seen that regardless of how  $Q_1$  or  $Q_2$  is changed (provided the magnitude of the change is sufficiently small) the topological index will be non-zero and there will be at least one periodic solution.

Now suppose that after  $Q_1$  is changed slightly in some definite way, the index  $j$  is zero. Then no conclusion can be drawn about whether there is a periodic solution. Further suppose that  $j = 2$  or  $-2$ . Then, again by using the analysis of [5, Section 3] it can be shown that

a different small change in  $Q_1$  will yield a mapping of index zero. Again no conclusion about the existence of periodic solutions can be drawn.

Another exceptional case which may occur is this. In the first example of Section 3, functions  $H_1(c_1, c_2)$  or  $H_2(c_1, c_2)$  may be identically zero. If  $H_1(c_1, c_2) \equiv 0$  and  $k_1 \neq 0$ , the bifurcation system is inconsistent and there are no periodic solutions with initial values continuous in  $\mu$  for small  $\mu$ . If  $H_1(c_1, c_2) \equiv 0$ , and  $k_1 = 0$ , no immediate conclusion can be drawn. It is worth noting that if we consider a problem in which

$$\left. \frac{dT}{d\mu} \right|_{\mu=0} \neq 0,$$

then as is shown in [8], neither  $H_1$  or  $H_2$  is identically zero.

#### §5. IF THE TOPOLOGICAL DEGREE IS EVEN

The condition (in Assumptions 2 and 5) that the  $f_1(x, t, \mu)$  can be written as the sum of a term that depends only on  $x$  and  $\mu$  and a term that depends only on  $t$  and  $\mu$  implies that the topological degree is odd and therefore non-zero. If we do not impose this condition, the topological degree may be odd or even. By the results of [5] and Lemma 3.1, if there are two equations in the reduced bifurcation system, the topological degree can be computed in all cases and the computation consists simply in approximating the real roots of certain polynomials. (See [7], Theorems 1 and 2.) If there are three or more equations in the reduced bifurcation system, the degree can be computed in many special cases.

If the topological degree is zero, Equation (3.1) may or may not have periodic solutions as the example in [7, Section 5] shows.

#### §6. THE NUMBER OF DISTINCT PERIODIC SOLUTIONS

For the totally degenerate case ( $q = n$ ), the topological degree also yields an estimate of the number of distinct periodic solutions. We illustrate this by considering the first example of Section 3 with the additional hypothesis that  $n = q$ . As before, the reduced bifurcation system is:

$$H_1(c_1, c_2) + k_1 = 0$$

$$H_2(c_1, c_2) + k_2 = 0.$$

LEMMA 6.1. Let  $M$  be a continuous mapping defined on the closure  $\bar{O}$  of an open set  $O \subset \mathbb{R}^n$ , real Euclidean  $n$ -space, and suppose  $M$  is differentiable in  $O$ . Suppose the topological degree of  $M$  at point  $p_0$  and relative to  $O$  is  $d \neq 0$ . Then there is a neighborhood  $U$  of  $p_0$  and a set  $E$  of  $n$ -dimensional measure zero,  $E \subset U$ , such that if  $p \in U - E$ , then  $M^{-1}(p)$  is a finite set consisting of at least  $|d|$  points.

This lemma is proved in [6, p. 213].

Now suppose that the topological degree at  $(0, 0)$  and relative to a solid circle  $S$  with center  $(0, 0)$  and radius  $r$  of the mapping defined by the reduced bifurcation system is  $d$  for all sufficiently large  $r$ . Let mapping  $M_3$  be defined by  $c_1' = \mathcal{H}_1(c_1, c_2)$ ,  $c_2' = \mathcal{H}_2(c_1, c_2)$ . Then if  $r$  is sufficiently large, the topological degree of  $M_3$  at  $(-k_1, -k_2)$  and relative to  $S$  is also equal to  $d$ . Applying this fact and Lemma 6.1 to the bifurcation system, we see that if  $k_1, k_2$  are changed arbitrarily slightly, there will be at least  $|d|$  distinct solutions of the bifurcation system. Since

$$k_1 = \int_0^{2\pi} \left\{ \cos(Ns)K_1(-s, 0) - \sin(Ns)K_2(s, 0) \right\} ds$$

and similarly for  $k_2$ , we obtain:

THEOREM 6.1. Suppose  $K_1$  and  $K_2$  are independent of  $\mu$  and suppose the topological degree at  $(0, 0)$  of the mapping defined by the left side of the reduced bifurcation system is equal to  $d$  if the topological degree is taken relative to any sufficiently large solid circle with center at  $(0, 0)$ . Let  $\mu$  have a fixed value, say  $\mu_0$ . Then given  $\eta > 0$ , there exist functions  $K_{11}(t)$ ,  $K_{12}(t)$ , both of period  $2\pi$ , with continuous second derivatives, such that

$$\max_{0 \leq t \leq 2\pi} |K_{1i}(t) - K_i(t)| < \eta \quad (i = 1, 2)$$

and such that if  $\mu = \mu_0$  in Equation (3.1) and  $K_1(t), K_2(t)$  are replaced by  $K_{11}(t), K_{12}(t)$ ,

then the resulting equation has at least  $|d|$  distinct solutions of period  $2\pi$ .

### §7. THE AUTONOMOUS CASE

Topological degree can also be used to study autonomous systems, but because the bifurcation system is quite different in the autonomous case, a different approach must be used. We describe the application to an example.

Consider the equation:

$$(7.1) \quad \dot{x} = Ax + \mu f(x, \mu) \quad .$$

Assume  $A$  has just one pair of characteristic roots of the form  $iN$ ,  $-iN$ , where  $N$  is a non-zero integer, that  $A$  has zero as a characteristic root, and that  $A$  has the standard form described in Assumption 4. Assume further that  $n = q$ . The bifurcation system may be written:

$$\begin{aligned} & -\nu c_2 + \int_0^{2\pi} \left\{ \cos(Ns) f_1(c_1 x^{(1)} + c_2 x^{(2)} + c_3 x^{(3)}, 0) \right. \\ & \quad \left. - \sin(Ns) f_2(c_1 x^{(1)} + c_2 x^{(2)} + c_3 x^{(3)}, 0) \right\} ds = 0 \\ & \nu c_1 + \int_0^{2\pi} \left\{ -\sin(Ns) f_1 + \cos(Ns) f_2 \right\} ds = 0 \\ & \nu c_3 + \int_0^{2\pi} \left\{ f_3(c_1 x^{(1)} + c_2 x^{(2)} + c_3 x^{(3)}, 0) \right\} ds = 0 \quad , \end{aligned}$$

where  $f_1(x, \mu)$ ,  $f_2$ ,  $f_3$  are the components of  $f$ , and  $x^{(1)}$ ,  $x^{(2)}$ , and  $x^{(3)}$  have the same meaning as before and

$$\nu = \lim_{\mu \rightarrow 0} \frac{\tau(\mu)}{\mu}$$

where  $\tau(\mu)$  is the period of the solutions of (7.1). The physical problem represented by the autonomous case is such that the corresponding mathematical problem can be posed as: set one  $c_1 = 0$  and solve for the remaining  $c_1$ 's and  $\nu$ . (See [3, pp. 30-31] and [4, pp. 364-366].)

Let us assume that  $f_1(x, 0)$ ,  $f_2(x, 0)$ ,  $f_3(x, 0)$  are polynomials

in the components of  $x$ . Then the bifurcation system may be written:

$$(7.2) \quad -vc_2 + P_1(c_1, c_2, c_3) = 0$$

$$(7.3) \quad vc_1 + P_2(c_1, c_2, c_3) = 0$$

$$(7.4) \quad vc_3 + P_3(c_1, c_2, c_3) = 0$$

where  $P_1, P_2, P_3$  are polynomials in  $c_1, c_2, c_3$ . We set  $c_3 = 0$  and solve (7.2), (7.3), (7.4) for  $c_1, c_2$ , and  $v$ . Multiplying (7.2) by  $c_1$  and (7.3) by  $c_2$ , and adding, we obtain the system:

$$(7.5) \quad c_1 P_1(c_1, c_2, 0) + c_2 P_2(c_1, c_2, 0) = 0$$

$$(7.6) \quad P_3(c_1, c_2, 0) = 0.$$

If  $\mu$  is varied slightly, Equations (7.5) and (7.6) become

$$(7.5)' \quad c_1 R_1(c_1, c_2, \mu) + c_2 R_2(c_1, c_2, \mu) = 0$$

$$(7.6)' \quad R_3(c_1, c_2, \mu) = 0$$

where  $R_1, R_2, R_3$  are continuous functions of  $c_1, c_2$ , and  $\mu$ . Suppose the topological degree of the mapping defined by the left sides of (7.5) and (7.6) is non-zero. Then for a sufficiently small  $\mu$ , (7.5)' and (7.6)' have a solution  $(c_1^0, c_2^0)$ . When  $\mu$  is varied slightly in (7.2), (7.3), and (7.4) (but keeping  $c_3 = 0$ ) we obtain:

$$(7.2)' \quad -vc_2 + R_1(c_1, c_2, \mu) = 0$$

$$(7.3)' \quad vc_1 + R_2(c_1, c_2, \mu) = 0$$

$$(7.4)' \quad R_3(c_1, c_2, \mu) = 0.$$

Now suppose one of the  $c_1^0, c_2^0$ , say  $c_1^0$ , is different from zero. (This will occur if, for example,  $P_3(c_1, c_2, 0)$  has a constant term.) We set

$$v_0 = \frac{R_2(c_1^0, c_2^0, \mu)}{c_1^0}.$$

Then  $c_1, c_2, v_0$  are a solution of (7.2)', (7.3)', and (7.4)'.

## REFERENCES

- [1] ALEXANDROFF, P., and HOPF, H., Topologie 1, Berlin, 1935.
- [2] BASS, R. W., Extension of frequency method of analyzing relay-operated servomechanisms, Section III, Final report, Contract DA-36-034-ORD-1273 RD, Johns Hopkins Institute for Cooperative Research (1955).
- [3] CODDINGTON, E. A., and LEVINSON, N., "Perturbations of linear systems with constant coefficients possessing periodic solutions", Contributions to the theory of nonlinear oscillations, Vol. II, Annals of Mathematics Studies No. 29, Princeton, 1952.
- [4] CODDINGTON, E. A., and LEVINSON, N., Theory of ordinary differential equations, New York, 1953.
- [5] CRONIN, J., "Topological degree of some mappings", Proc. Amer. Math. Soc., 5 (1954), pp. 175-178.
- [6] CRONIN, J., "Branch points of solutions of equations of Banach Space, II, Trans. Amer. Math. Soc., 76 (1954), pp. 207-222.
- [7] CRONIN, J., "Note to Poincaré's perturbation method", Duke Mathematical Journal, 26 (1959), pp. 251-262.
- [8] FRIEDRICHS, K. O., Advanced ordinary differential equations (mimeographed) New York University, 1949.
- [9] FRIEDRICHS, K. O., "Fundamentals of Poincaré's theory", Proceedings of the symposium on nonlinear circuit analysis, New York, 1953, pp. 56-67.
- [10] FRIEDRICHS, K. O., Special topics in analysis (mimeographed), New York University, 1954.
- [11] LEFSCHETZ, S., "Complete families of periodic solutions of differential equations", Comment. Math. Helv., 28 (1954), pp. 341-345.
- [12] LEFSCHETZ, S., Differential equations: geometric theory, New York, 1957.
- [13] POINCARÉ, H., Les méthodes nouvelles de la mécanique céleste, Vol. I, II, III, (1892-1899). Reprinted by Dover Publications, New York, 1957.

#### IV. ON THE BEHAVIOR OF THE SOLUTIONS OF LINEAR PERIODIC DIFFERENTIAL SYSTEMS NEAR RESONANCE POINTS

Jack K. Hale

##### §1. INTRODUCTION

In the last few years, systems of differential equations of the form

$$(1.1) \quad y_j'' + \sigma_j^2 y_j = \varepsilon \sum_{k=1}^n \varphi_{jk} y_k, \quad j = 1, \dots, n,$$

where  $\varepsilon$  is a real small parameter, the constants  $\sigma_j$  are positive, and the real functions  $\varphi_{jk}(t)$  are periodic of period  $T = 2\pi/\omega$  and  $L$ -integrable in  $[0, T]$ , have been investigated by many authors [1, 3a, 4a, 5, 7, 8, 9, 10b, 10e, 12, 13]. (See also the book [3b]). For  $n = 1$ , classical results of O. Haupt [9] assure that all solutions of (1.1) are bounded for  $|\varepsilon|$  sufficiently small and an arbitrary periodic function  $\varphi_{11}$  provided  $2\sigma_1 \neq 0 \pmod{\omega}$ .

For the study of the system (1.1) for  $|\varepsilon|$  small and  $n \geq 1$ , L. Cesari [3a] in 1940 considered a method of successive approximations, which was successively developed by L. Cesari, J. K. Hale and R. A. Gambill [6, 4b, 10c, 10e] and applied to questions of existence and stability of periodic solutions of weakly nonlinear differential systems (cf. the book [3b] and [3c]). By using this method it was first proved [3a, 10b] that all solutions of (1.1) are bounded provided  $2\sigma_j = 0$ ,  $\sigma_j \pm \sigma_k \neq 0 \pmod{\omega}$ ,  $j \neq k$ ,  $j, k = 1, 2, \dots, n$ , and the matrix  $\Phi(t) = [\varphi_{jk}(t)]$  is either symmetric or even in  $t$ . Under the same restrictions on the numbers  $\sigma_j$  more general boundedness theorems have then been proved [5a, 10e] by the same method, but they all involve some type of "symmetry" conditions on the matrix  $\Phi(t)$ . Again using the same method it has been proved [3a, 5b] that some "symmetry" condition is necessary to assure that all solutions of (1.1) are bounded for  $\varepsilon$

sufficiently small. For systems more general than (1.1), see [1, 10e].

Analogous boundedness theorems involving "symmetry" conditions have been proved by M. Golomb [7] by a different method. Finally, J. Moser [13], J. M. Gel'fand and V. B. Lidskii [14] have recently shown that it is sufficient to require  $2\sigma_j \neq 0$ ,  $\sigma_j + \sigma_k \neq 0 \pmod{\omega}$ ,  $j \neq k$ ,  $j, k = 1, 2, \dots, n$ , when  $\Phi$  is a symmetric matrix.

In the present paper a procedure based on the same method developed by Cesari, Hale and Gambill is given for the study of the behavior of the solutions of (1.1) (and more general systems) near the "resonance points"  $2\sigma_j = s\omega$ ,  $\sigma_j \pm \sigma_k = s\omega$ , for some integer  $s$ . The ultimate goal, of course, is to try to classify systems of type (1.1) according to the "resonance points" for which some solutions are unbounded for every  $\varepsilon \neq 0$ . This general problem is not solved in this paper. However, we do give some general theorems concerning the behavior of the solutions of a class of differential equations (1.1) (and even a more general type) around the resonance points (A)  $\sigma_1 - \sigma_2 = s\omega$ ,  $s$  an integer or zero,  $2\sigma_j \neq 0$ ,  $\sigma_j \pm \sigma_k \neq 0 \pmod{\omega}$ ,  $j \neq k$ ,  $j, k = 2, 3, \dots, n$  [Theorems (5.2) and (5.3)] and (B)  $\sigma_1^2 = \varepsilon^2$ ,  $\sigma_j \neq 0$ ,  $2\sigma_j \neq 0$ ,  $\sigma_j \pm \sigma_k \neq 0 \pmod{\omega}$ ,  $j \neq k$ ,  $j, k = 2, 3, \dots, n$  [Theorem (5.4)].

More specifically, if the  $\sigma_j$  satisfy (A) and Theorem (5.2) is applied to system (1.1) with (C)  $\Phi = (\Phi_{jk})$ , where each  $\Phi_{jk}(-t) = (-1)^{k+j} \Phi_{jk}(t)$  is a matrix, then the AC (absolutely continuous) solutions of (1.1) are bounded for  $\varepsilon$  sufficiently small if a certain function  $G(s, \sigma)$  of the Fourier coefficients of the functions  $\varphi_{jk}(t)$  is positive and some AC solutions are unbounded for every  $\varepsilon \neq 0$  if this function is negative. If  $\Phi$  is symmetric and also satisfies (C), then  $G(s, \sigma) > 0$  (Remark 5.3), which agrees with the conclusions of J. Moser [13]. Some examples are given in §5 to show that  $G(s, \sigma)$  may be  $< 0$  for some matrices  $\Phi$  if  $\Phi$  satisfies (C) and is not symmetric. If Theorem (5.4) is applied to system (1.1) satisfying (B), (C), then all of the AC solutions of (1.1) are bounded for  $\varepsilon$  sufficiently small.

In §6, the above method is applied to systems of Mathieu type equations of the form (1.1) where each  $\varphi_{jk}(t) = d_{jk} \cos 2t$ , each  $d_{jk}$  is a constant and  $2\sigma_j \neq 0$ ,  $\sigma_j \pm \sigma_k \neq 0 \pmod{1}$ ,  $j \neq k$ ,  $j, k = 2, 3, \dots, n$ ,  $\sigma_1(0) = m$ , where  $m$  is a positive integer. Sufficient conditions are given (Theorem 6.1) to insure that there are unbounded absolutely continuous solutions of (1.1) in every neighborhood of the point  $(m, 0)$  in the  $(\sigma_1, \varepsilon)$ -plane, where  $m$  is a positive integer. A corollary of Theorem (6.1) is the well known fact [11] that there are unbounded solutions of the Mathieu equation  $x'' + \sigma^2 x + \varepsilon(\cos 2t)x = 0$ , in every neighborhood of the point  $(m, 0)$  of the  $(\sigma, \varepsilon)$ -plane for every positive integer  $m$ .



Using determinants, the behavior of the solutions of (1.1),  $n > 1$ , near the "resonance points" has also been discussed by E. Mettler [12] and W. Haacke [3] for the case where  $\phi(t)$  is even in  $t$ . General theorems of the above type seem to be easier to obtain using the method of this paper since it does not involve determinants.

Another application of the method discussed here concerns the stability of periodic solutions of weakly nonlinear differential systems. The linear variational equations associated with such a periodic solution is a linear differential equation with periodic coefficients of the form (1.1) and, in many cases, some of the basic frequencies  $\sigma_j$  are "in resonance" with  $\omega$ . The asymptotic stability of periodic solutions of weakly nonlinear periodic differential systems has been discussed recently by H. R. Bailey and R. A. Gambill [2] using essentially the same method. For a more complete discussion of the application of this method, in particular, Theorems (2.1) and (3.1), to the stability of periodic solutions of both weakly nonlinear autonomous and weakly nonlinear periodic differential systems, see [10g].

## §2. DESCRIPTION OF THE METHOD

Let  $C_\omega$  denote the family of all functions which are finite sums of functions of the form  $f(t) = e^{\alpha t} \varphi(t)$ ,  $-\infty < t < +\infty$ , where  $\alpha$  is any complex number and  $\varphi(t)$  is any complex-valued function of the real variable  $t$ , periodic of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ . Following L. Cesari [3a], if  $f(t) = e^{\alpha t} \varphi(t) \in C_\omega$  and  $\varphi(t)$  has the Fourier series

$$\varphi(t) \sim \sum_{k=-\infty}^{+\infty} c_n e^{in\omega t},$$

$i = \sqrt{-1}$ , then we denote by mean value,  $M[f]$ , the number  $M[f] = 0$  if  $in\omega + \alpha \neq 0$  for all  $n$ ;  $M[f] = c_n$  if  $in\omega + \alpha = 0$  for some  $n$ . It is clear that  $M[f]$  is uniquely defined. Finally define  $M[f]$  in the class  $C_\omega$  as an additive functional. Obviously  $M[f]$  reduces to the usual mean value for periodic functions  $f(t)$  of period  $T$ . The following lemma is needed in the sequel.

LEMMA (2.1) If  $f(t) \in C_\omega$  and  $M[f] = 0$ , then there is a unique primitive of  $f(t)$ , say  $F(t) = \int f(t)dt$ , which belongs to  $C_\omega$  and  $M[f] = 0$ . (L. Cesari [3a], J. K. Hale [10a]).

Consider a linear system of differential equations of the form

$$(2.1) \quad z' = Az + \varepsilon C(t)z$$

where  $\varepsilon$  is a parameter,  $A, C$  are  $N \times N$  matrices,  $C = (c_{jk}(t))$ , each  $c_{jk}(t)$  is periodic in  $t$  of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ ,  $A = \text{diag}(\rho_1, \dots, \rho_N)$ , where the  $\rho_j$  are complex numbers such that

$$(2.2) \quad \rho_j - \rho_k = m_{jk}\omega i, \quad m_{jk} \text{ an integer or zero, } j, k = 1, 2, \dots, \nu,$$

$$\rho_j \not\equiv \rho_k \pmod{\omega i}, \quad j = 1, 2, \dots, \nu; \quad k = \nu + 1, \dots, N.$$

Notice that all of the  $\rho_j$ ,  $j = 1, 2, \dots, \nu$ , could be equal and also the remaining  $N - \nu$  of the  $\rho_k$ ,  $k = \nu + 1, \dots, N$ , can be two by two congruent mod  $\omega i$ . This becomes important in discussing the stability of periodic solutions of nonlinear differential systems [10g].

In this section, a method of successive approximations is given for the determination of the characteristic exponents as well as the solutions of (2.1). This method coincides with the one given by L. Cesari [3a] for the case where  $\nu = 1$  in (2.2). More specifically, we try to determine solutions of (2.1) of the form

$$(2.3) \quad z(t) = e^{\tau t} p(t), \quad p(t + T) = p(t),$$

where  $\tau$  is a complex number to be determined. The equation for  $p(t)$  is then

$$(2.4) \quad p' = (A - \tau I)p + \varepsilon C(t)p.$$

In the following, the characteristic exponent  $\tau$  which is "close to"  $\rho_1$  is to be determined, i.e.,  $\tau$  as a function of  $\varepsilon$  is such that  $\tau(0) = \rho_1$ . From (2.2) it follows that  $\rho_k - \tau$  will be "close to" a multiple of  $\omega i$  for  $k = 1, 2, \dots, \nu$ . Rather than solve (2.4) directly, consider an auxiliary equation

$$(2.5) \quad p' = Bp + \varepsilon C(t)p,$$

where

$$B = \text{diag}(\rho_1^*, \dots, \rho_N^*),$$

(2.6)

$$\rho_1^* = 0, \quad \rho_k^* = m_{k1}\omega i, \quad k = 2, 3, \dots, \nu, \quad \rho_k^* = \rho_k - \tau, \quad k = \nu + 1, \dots, N.$$

and the  $m_{jk}$  are defined in (2.2). Suppose that  $S_1$  is a circle in the complex plane with the center at  $\rho_1$  such that

$$(2.7) \quad \rho_j^* \neq 0 \pmod{\omega 1}, \quad j = v + 1, \dots, N,$$

for all  $\tau \in S_1$ .

If the  $m^{\text{th}}$  approximation to  $p(t)$  is denoted by

$$(2.8) \quad p^{(m)} = x^{(0)} + \varepsilon x^{(1)} + \dots + \varepsilon^m x^{(m)},$$

where each  $x^{(r)}$  is periodic in  $t$  of period  $T = 2\pi/\omega$ , let

$$(2.9) \quad \begin{aligned} s^{(r)}(t) &= e^{-Bt} C(t) x^{(r-1)}(t), \\ s^{(r)} &= (s_1^{(r)}, \dots, s_N^{(r)}) \end{aligned}$$

and define

$$(2.10) \quad \begin{aligned} S^{(r)} &= \text{diag}(S_1^{(r)}, \dots, S_v^{(r)}, 0, \dots, 0) \\ S_j^{(r)} &= 0 \quad \text{if } M[s_j^{(r)}] = 0, \\ d_j S_j^{(r)} &= M[s_j^{(r)}] \quad \text{if } M[s_j^{(r)}] \neq 0 \quad j = 1, 2, \dots, v, \end{aligned}$$

where  $M[s_j^{(r)}]$  denotes the mean value for functions of the class  $C_\omega$  and  $d_1, \dots, d_v$  are complex numbers. If  $M[s_j^{(r)}] \neq 0$  for any  $r$ , it is understood in definition (2.10) that the corresponding  $d_j$  is chosen  $\neq 0$  so that  $S_j^{(r)}$  will be well defined in all cases. In fact, if  $d_j$  is a function of  $\varepsilon$ , then  $d_j$  is  $\neq 0$  for  $\varepsilon = 0$ .

With these notations, the successive approximations are defined by

$$(2.11) \quad \begin{aligned} x^{(0)} &= (d_1 e^{\rho_1^* t}, \dots, d_v e^{\rho_v^* t}, 0, \dots, 0) \\ x^{(r)} &= e^{Bt} \int e^{-Bt} [C(t) x^{(r-1)} - (S^{(r)} x^{(0)} + \dots \\ &\quad S^{(1)} x^{(r-1)})] dt, \quad r = 1, 2, 3, \dots \end{aligned}$$

It is clear that condition (2.7) assures that each of the integrands in (2.11) belongs to the class  $C_\omega$  and has mean value zero provided that the integrations are always performed so as to obtain the unique primitive of mean value zero. It is assumed that the integrations are performed in

this manner. Furthermore, the  $x^{(r)}(t)$  defined by (2.11) are periodic of period  $T$ .

The method of successive approximations (2.11) is precisely the one given in [6] for obtaining periodic solutions of nonlinear systems of differential equations. Exactly as in [6], it follows that  $p^{(m)}$  defined by (2.8), (2.11) converges uniformly in  $t$  for  $|\varepsilon|$  sufficiently small to a solution of the equation

$$(2.12) \quad p' = [B - \varepsilon H(\tau, d, \rho, \varepsilon)]p + \varepsilon C(t)p,$$

where  $d = (d_1, \dots, d_v)$ ,  $\rho = (\rho_1, \dots, \rho_N)$  and  $H = \text{diag}(h_1, \dots, h_v, 0, \dots, 0)$  and each  $h_j(\tau, d, \rho, \varepsilon)$  is an analytic function of  $\varepsilon$  at  $\varepsilon = 0$  and its power series expansion is given by

$$(2.13) \quad h_j = S_j^{(1)} + \varepsilon S_j^{(2)} + \varepsilon^2 S_j^{(3)} + \dots, \quad j = 1, 2, \dots, v.$$

Therefore, if the numbers  $\tau, d_1, \dots, d_v, \rho_1, \dots, \rho_v$  are chosen so that the matrix equation,

$$(2.14) \quad B - \varepsilon H(\tau, d, \rho, \varepsilon) = A - \tau I$$

or, equivalently, the equations,

$$m_{k1}\omega_1 - \varepsilon h_k(\tau, d, \rho, \varepsilon) = \rho_k - \tau,$$

$$(2.14')$$

$$k = 1, 2, \dots, v, m_{11} = 0,$$

are satisfied, then the periodic solution  $p(t)$  of (2.12) will be a periodic solution of (2.4) and  $z(t)$  defined by (2.3) will be a solution of (2.1). We shall refer to the equations (2.14), (2.14') as the determining equations for  $\tau, d_1, \dots, d_v, \rho_1, \dots, \rho_v$ . These represent  $v$  equations for the  $2v + 1$  parameters  $\tau, d_1, \dots, d_v, \rho_1, \dots, \rho_v$ , and the particular problem of interest will, in general, determine which of these parameters are to be determined and which are fixed.

REMARK 2.1. In the above discussion, it was assumed that the numbers  $\rho_j, j = 1, 2, \dots, N$ , and the matrix  $C(t)$  were independent of  $\varepsilon$ . This is not necessary and systems of the form

$$(2.15) \quad z' = A(\varepsilon)z + \varepsilon C(t, \varepsilon)z$$

could have been considered, where  $A(\varepsilon) = \text{diag}(\rho_1(\varepsilon), \dots, \rho_N(\varepsilon))$ , each  $\rho_j(\varepsilon)$  is an analytic function of  $\varepsilon$  at  $\varepsilon = 0$  with

$$\rho_j(0) - \rho_k(0) = m_{jk}\omega_1, m_{jk} \text{ an integer or zero, } j, k = 1, 2, \dots, v, \quad (2.16)$$

$$\rho_j(0) \neq \rho_k(0), \pmod{\omega_1}, j = 1, 2, \dots, v; k = v+1, \dots, N;$$

$C(t, \varepsilon) = (c_{jk}(t, \varepsilon))$ ,  $j, k = 1, 2, \dots, N$ , where each  $c_{jk}(t, \varepsilon)$  is analytic in  $\varepsilon$  for  $0 \leq |\varepsilon| \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ ,  $|c_{jk}(t, \varepsilon)| < \eta(t)$ ,  $0 \leq |\varepsilon| \leq \varepsilon_0$ ,  $j, k = 1, 2, \dots, N$ , and  $\eta(t)$  is  $L$ -integrable in  $[0, T]$ . The matrix  $B$  in (2.6) is exactly the same as before and the method of successive approximations is applied in the same manner by replacing everywhere  $\rho_j$  by  $\rho_j(\varepsilon)$ ,  $j = 1, 2, \dots, N$ , and the matrix  $C(t)$  by  $C(t, \varepsilon)$ . The determining equations are

$$(2.17) \quad m_{k1}\omega_1 - \varepsilon h_k(\tau, d, \rho(\varepsilon), \varepsilon) = \rho_k(\varepsilon) - \tau, k = 1, 2, \dots, v,$$

where  $m_{11} = 0$  and the functions  $h_k$  are given by (2.13) with the  $S_j^{(m)}$  analytic functions of  $\varepsilon$  for  $0 \leq |\varepsilon| \leq \varepsilon_0$ . The remainder of the discussion will deal with this more general system. It is also true that the above analyticity at  $\varepsilon = 0$  can be replaced by continuity in  $\varepsilon$  at  $\varepsilon = 0$  if  $\rho_j(\varepsilon) - \rho_j(0) = o(\varepsilon)$ , but to make the presentation as simple as possible we will not consider this generalization.

A solution  $\tau, d_1, \dots, d_v, \rho_1, \dots, \rho_v$  of (2.17) must be such that  $d_j \neq 0$  for  $\varepsilon = 0$  if  $M[s_j^{(r)}] \neq 0$  for any  $r = 1, 2, \dots$ . In case some  $d_j$  is  $o(\varepsilon)$ , then the method of successive approximations must be modified slightly taking the zero<sup>th</sup> approximation to have this  $d_j = o(\varepsilon)$ . We will not give the details of the method for such a case, since it is clear how one would proceed.

In case the numbers  $\rho_1, \dots, \rho_v$  are fixed, then the determining equations represent  $v$  equations for the  $v+1$  parameters  $\tau, d_1, \dots, d_v$ . Since (2.1) is a linear system, it is clear that one of the numbers  $d_j$  may be chosen equal to one, but, in general, this decision will depend upon the form of equations (2.17). Two solutions  $(\tau, d_1, \dots, d_v), (\tau^*, d_1^*, \dots, d_v^*)$  of (2.17) will be called distinct if they correspond to two linearly independent solutions of (2.1) of the form (2.3). A similar definition applies to  $k$  solutions of (2.17). If  $\rho_j = \rho_k$ ,  $j, k = 1, 2, \dots, v$ , one would hope that there would be  $v$  distinct solutions of (2.17). However, this is not always the case since some of the characteristic exponents  $\tau$  may be equal.

From (2.10) the functions  $h_j(\tau, d, \rho(\varepsilon), \varepsilon)$  in (2.17) satisfy the property

$$(2.18) \quad d_j h_j(\tau, d, \rho(0), 0) = \frac{1}{T} \int_0^T \sum_{k=1}^v c_{jk}(t, 0) d_k e^{1(m_{k1} - m_{j1})\omega t} dt,$$

$$j = 1, 2, \dots, v,$$

which is a linear function of  $d_1, \dots, d_v$  and is independent of  $\tau$ . If  $\varepsilon \beta = \tau - \rho_1(0)$ , then for  $\varepsilon \neq 0$  equations (2.17) are equivalent to the system of equations

$$(2.19) \quad (G - \beta I)d + \varepsilon F(d, \tau, \rho(\varepsilon), \varepsilon) = 0$$

where  $G = (g_{jk})$ ,  $j, k = 1, 2, \dots, v$ , is independent of  $\varepsilon$  and

$$g_{jj} = \frac{1}{T} \int_0^T c_{jj}(t, 0) dt + \lim_{\varepsilon \rightarrow 0} \frac{\rho_j(\varepsilon) - \rho_j(0)}{\varepsilon}$$

$$(2.20) \quad g_{jk} = \frac{1}{T} \int_0^T e^{1(m_{k1} - m_{j1})\omega t} c_{jk}(t, 0) dt, \quad m_{11} = 0,$$

$$j \neq k, \quad j, k = 1, 2, \dots, v,$$

and  $F(d, \tau, \rho(\varepsilon), \varepsilon)$  is an analytic vector function of  $\varepsilon$  for  $0 \leq |\varepsilon| \leq \varepsilon_0$ . These results are summarized in the following theorem.

**THEOREM (2.1).** Consider the system of differential equations (2.15), (2.16), and suppose the matrix  $G$  is defined by (2.20). If  $\beta_0$  is a simple characteristic root of the matrix  $G$  and the corresponding eigenvector,  $d_0$ , has no zero components, then there is a characteristic exponent  $\tau(\varepsilon)$  of (2.15) which is analytic in  $\varepsilon$  for  $0 \leq |\varepsilon| \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ , and

$$(2.21) \quad \tau(\varepsilon) = \rho_1(\varepsilon) + \varepsilon \beta_0 + o(\varepsilon^2).$$

The proof of this theorem follows immediately by applying the implicit function theorem to (2.19). If the matrix  $d_0$  has some components which are zero, then the above method of successive approximations must be modified slightly taking the first approximation (2.11) to have some  $d_j = 0(\varepsilon)$ . In case the matrix  $G$  in Theorem (2.1) is the zero matrix, then the theorem says nothing about the characteristic exponents. However, by obtaining more of the functions  $S_j^{(m)}$  in (2.17), one can

determine the characteristic exponents. This theorem and the following corollary are not used in this paper but are important for the study of stability of periodic solutions of weakly nonlinear systems of differential equations (see [2], [10g]).

COROLLARY (2.1). Consider the system of differential equations (2.15) with

$$(2.22) \quad \rho_j(0) \not\equiv \rho_k(0) \pmod{\omega_1}, k \neq j, k = 1, 2, \dots, N,$$

where  $j$  is a fixed integer,  $1 \leq j \leq N$ . Then there is a characteristic exponent of (2.15),  $\tau_j = \tau_j(\varepsilon)$ , analytic in  $\varepsilon$  for  $0 \leq |\varepsilon| \leq \varepsilon_0$ , and

$$(2.23) \quad \tau_j(\varepsilon) = \rho_j(\varepsilon) + \frac{\varepsilon}{T} \int_0^T c_{jj}(t, 0) dt + o(\varepsilon^2),$$

where the terms of higher order in  $\varepsilon$  may be calculated by using the above method of successive approximations.

REMARK 2.2. L. Cesari [3a] had previously observed that one could calculate the characteristic exponents one at a time to obtain relation (2.23) but under the more restrictive requirement that  $\rho_j \not\equiv \rho_k \pmod{\omega_1}$   $j \neq k$ ,  $j, k = 1, 2, \dots, N$ . It is also clear from his work that the above corollary holds for a particular characteristic exponent  $\tau_j$ , with the weaker restriction (2.22).

For specific examples, the method of successive approximations of this section is probably as efficient as any, but to obtain general boundedness theorems and the form of the functions  $h_j$  in (2.13) for classes of differential equations, the following formulation is somewhat more convenient when the  $\rho_k$  satisfy certain additional properties.

### §3. AN ALTERNATIVE FORMULATION OF THE METHOD

Suppose the numbers  $\rho_k$  in (2-1) satisfy the property\*

$$\rho_{2j-1} = \bar{\rho}_{2j}, j = 1, 2, \dots, v,$$

$$(3.1) \quad \rho_{2j-1} - \rho_{2k-1} = m_{jk}\omega_1, m_{jk} \text{ an integer or zero, } j, k = 1, 2, \dots, v,$$

$$\rho_j \not\equiv \rho_k \pmod{\omega_1}, j = 1, 2, \dots, 2v, k = 2v + 1, \dots, N.$$

\* Throughout this paper,  $\bar{a}$  will always denote the complex conjugate of  $a$ .

Notice that  $\rho_{2j-1}$  may or may not be congruent to  $\rho_{2j}$  modulo  $\omega_1$ ,  $j = 1, 2, \dots, v$ . Both situations are important in the applications. Rather than make the substitution (2.3), consider the auxiliary equation

$$(3.2) \quad z' = Bz + \varepsilon C(t)z$$

where  $B = \text{diag}(\rho_1^*, \dots, \rho_N^*)$ ,

$$(3.3) \quad \rho_1^* = \tau = \bar{\rho}_2^*, \quad \rho_{2k-1}^* = \tau - m_{k1}\omega_1 = \bar{\rho}_{2k}^*, \quad k = 1, 2, \dots, v,$$

$$\rho_k^* = \rho_k, \quad k = 2v+1, \dots, N,$$

where the numbers  $m_{k1}$  are defined by (3.1), and  $\tau$  is an undetermined complex number such that

$$(3.4) \quad \rho_j^* \neq \rho_k^*, \quad j = 1, 2, \dots, 2v; \quad k = 2v+1, \dots, N.$$

If the  $m^{\text{th}}$  approximation is defined by

$$(3.5) \quad z^{(m)} = x^{(0)} + \varepsilon x^{(1)} + \dots + \varepsilon^m x^{(m)},$$

$s^{(r)}$  is defined as in (2.9) with  $B$  as in (3.3) and

$$(3.6) \quad \begin{aligned} S^{(r)} &= \text{diag}(S_1^{(r)}, \dots, S_{2v}^{(r)}, 0, \dots, 0) \\ S_j^{(r)} &= 0 \quad \text{if } M[s_j^{(r)}] = 0 \\ d_j S_j^{(r)} &= M[s_j^{(r)}] \quad \text{if } M[s_j^{(r)}] \neq 0, \quad j = 1, 2, \dots, 2v, \end{aligned}$$

where  $M[s_j^{(r)}]$  again denotes the mean value for functions of the class  $C_\omega$  and  $d_1, \dots, d_{2v}$  are complex numbers. If  $M[s_j^{(r)}] \neq 0$  for any  $r$ , it is understood in definition (3.6) that the corresponding  $d_j$  is chosen  $\neq 0$  so that  $S_j^{(r)}$  will be well defined in all cases. In fact, if  $d_j$  is a function of  $\varepsilon$ , then  $d_j$  is  $\neq 0$  for  $\varepsilon = 0$ . The successive approximations are now defined by

$$(3.7) \quad \begin{aligned} x^{(0)} &= (d_1 e^{\rho_1^* t}, \dots, d_{2v} e^{\rho_{2v}^* t}, 0, \dots, 0), \\ x^{(r)} &= e^{Bt} \int e^{-Bt} [C(t)x^{(r-1)} - (S^{(r)}x^{(0)} + \dots + S^{(1)}x^{(r-1)})] dt, \\ r &= 1, 2, 3, \dots, \end{aligned}$$



where the integrations are performed so as to obtain the unique primitive of mean value zero. The proof of convergence of the functions  $z^{(m)}$  defined by (3.5), (3.7) to a solution of the equation

$$(3.8) \quad z' = [B - \varepsilon V(\tau, d, \varepsilon)]z + \varepsilon C(t)z,$$

where  $V = \text{diag}(V_1, V_2, \dots, V_{2\nu}, 0, \dots, 0)$ ,  $(d = (d_1, \dots, d_{2\nu}))$ ,

$$(3.9) \quad V_j = S_j^{(1)} + \varepsilon S_j^{(2)} + \dots$$

is exactly the same as before. The determining equations for  $\tau, d_1, \dots, d_{2\nu}, \rho_1, \dots, \rho_{2\nu}$ , are

$$(3.10) \quad \tau - \varepsilon V_j(\tau, d, \varepsilon) = \rho_j, \quad j = 1, 2, \dots, 2\nu.$$

Remark (2.1) at the end of §2 also applies to the method defined in this manner. In the following, our choice between these two methods will depend upon the application and the convenience which it affords.

In case system (2.1) arises from a real system of first and second order differential equations, some of the determining equations (3.10) are redundant as we show in the next few lines. Consider the system of real equations

$$y_j'' + \alpha_j y_j' + \sigma_j^2 y_j = \varepsilon \sum_{k=1}^n \varphi_{jk} y_k + \varepsilon \sum_{k=1}^{\mu} \psi_{jk} y_k', \quad j = 1, 2, \dots, \mu,$$

(3.11)

$$y_j' + \beta_j y_j = \varepsilon \sum_{k=1}^n \varphi_{jk} y_k + \varepsilon \sum_{k=1}^{\mu} \psi_{jk} y_k', \quad j = \mu + 1, \dots, n,$$

where  $\alpha_j, \beta_j, \sigma_j, \varepsilon$  are real numbers,  $\gamma_j \equiv (4\sigma_j^2 - \alpha_j^2)^{\frac{1}{2}} > 0$ ,  $j = 1, 2, \dots, \mu$ , and the functions  $\varphi_{jk}(t), \psi_{jk}(t)$  are real, periodic in  $t$  of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ . Suppose the roots of the equation  $z^2 + \alpha_j z + \sigma_j^2 = 0$  are  $\lambda_{2j-1}, \lambda_{2j} = \bar{\lambda}_{2j-1}$ ,  $j = 1, 2, \dots, \mu$ , and let  $\lambda_{\mu+j} = -\beta_j$ ,  $j = \mu + 1, \dots, n$ . The transformation

$$(3.12) \quad y_j = (2i\gamma_j)^{-1} (z_{2j-1} + z_{2j}), \quad y_j' = (2i\gamma_j)^{-1} (\lambda_{2j-1} z_{2j-1} + \lambda_{2j} z_{2j}),$$

$$y_j = z_{\mu+j}, \quad j = \mu + 1, \dots, n,$$

yields the equivalent system of first order equations,

$$(3.13) \quad z' = Az + \varepsilon C(t)z$$

where  $A = \text{diag}(\lambda_1, \dots, \lambda_{n+\mu})$  and  $C$  is an  $(n + \mu) \times (n + \mu)$  matrix whose elements  $c_{jk}(t)$  are given by

$$c_{2j-1, 2k-1} = (2i\gamma_k)^{-1}(\varphi_{jk} + \rho_{2k-1}\psi_{jk}); \quad c_{2j-1, 2k} = (2i\gamma_k)^{-1}(\varphi_{jk} + \rho_{2k}\psi_{jk}),$$

$$k = 1, 2, \dots, \mu,$$

$$c_{2j-1, \mu+h} = \varphi_{jh}, \quad h = \mu + 1, \dots, n; \quad c_{2j-1, \ell} = -c_{2j, \ell}, \quad \ell = 1, 2, \dots, n+\mu;$$

$$(3.14) \quad j = 1, 2, \dots, \mu;$$

$$c_{\mu+h, 2k-1} = (2i\gamma_k)^{-1}(\varphi_{hk} + \rho_{2k-1}\psi_{hk}); \quad c_{\mu+h, 2k} = (2i\gamma_k)^{-1}(\varphi_{hk} + \rho_{2k}\psi_{hk}),$$

$$k = 1, 2, \dots, \mu; \quad c_{\mu+h, \mu+\ell} = \varphi_{h\ell}, \quad \ell = \mu + 1, \dots, n; \quad h = \mu + 1, \dots, n.$$

We also suppose that the equations in (3.13) are reordered so that

$$(3.15) \quad A = \text{diag}(\rho_1, \dots, \rho_{n+\mu})$$

where each  $\rho_j$  is one of the numbers  $\lambda$  and the  $\rho_j$  satisfy condition (3.1). It is very easy to prove the following lemma.

LEMMA (3.1). If the alternative method of successive approximations is applied to the auxiliary equation of (3.13), (3.15) with the  $d_j$  in (3.7) such that  $d_{2j-1} = -\bar{d}_{2j}$ ,  $j = 1, 2, \dots, \nu$ , then

$$(3.16) \quad V_{2j-1}(\tau, d, \varepsilon) = \bar{V}_{2j}(\tau, d, \varepsilon), \quad j = 1, 2, \dots, \nu,$$

for every  $\tau, d$  where  $V_j$  is defined by (3.9). Therefore, the determining equations (3.10) become

$$(3.17) \quad \begin{aligned} \tau - \varepsilon V_{2j-1}(\tau, d, \varepsilon) &= \rho_1, & j &= 1, 2, \dots, \nu, \\ d &= (d_1, \dots, d_{2\nu}), \quad d_{2j} = -\bar{d}_{2j-1}, & j &= 1, 2, \dots, \nu. \end{aligned}$$

For applications to the stability of periodic solutions of non-linear differential equations (see [10g]), it is convenient to have the

following theorem.

THEOREM (3.1). Consider the system of differential equations (3.11) with

$$(3.18) \quad \begin{aligned} \lambda_{2j-1} &= \bar{\lambda}_{2j}, \quad \lambda_{2j-1} \neq \lambda_{2j} \pmod{\omega_1}, \\ \lambda_{2j-1} - \lambda_{2k-1} &= m_{jk}\omega_1, \quad m_{jk} \text{ an integer or zero, } j, k = 1, 2, \dots, \nu, \\ \lambda_j &\neq \lambda_k \pmod{\omega_1}, \quad j = 1, 2, \dots, 2\nu; \quad k = 2\nu + 1, \dots, N, \end{aligned}$$

where  $\lambda_{2j-1} = \frac{1}{2}(-\alpha_j + i\gamma_j)$ ,  $\gamma_j = (4\sigma_j^2 - \alpha_j^2)^{\frac{1}{2}} > 0$ ,  
 $j = 1, 2, \dots, \mu$ ,  $\lambda_{\mu+j} = -\beta_j$ ,  $j = \mu + 1, \dots, n$ ,  
 and let  $H = (h_{jk})$ ,  $j, k = 1, 2, \dots, \nu$ , be the  
 $\nu \times \nu$  matrix defined by

$$(3.19) \quad h_{jk} = \frac{1}{T} \int_0^T c_{2j-1, 2k-1}(t) e^{i(m_{k1} - m_{j1})\omega t} dt,$$

$$j, k = 1, 2, \dots, \nu,$$

where the functions  $c_{2j-1, 2k-1}$  are defined by (3.14). If  $\beta_0$  is a simple root of the equation  $|H - \beta I| = 0$  and the corresponding eigenvector has no zero components, then there are two characteristic exponents  $\tau, \bar{\tau}$  of (3.11) given by

$$(3.20) \quad \tau = \rho_1 + \varepsilon \beta_0 + o(\varepsilon^2)$$

for  $\varepsilon$  sufficiently small.

PROOF. For  $\varepsilon \neq 0$  the determining equations (3.17) can be written in the form

$$(H - \beta I)d = o(\varepsilon),$$

where  $\varepsilon \beta = \tau - \rho_1$ ,  $d = (d_1, d_3, \dots, d_{2\nu-1})$  and  $H = (h_{jk})$  is the  $\nu \times \nu$  matrix given by (3.19). It then follows immediately from the implicit function theorem that Theorem (3.1) is true for  $\varepsilon$  sufficiently small.

## §4. SOME BASIC LEMMAS

Consider the system of linear differential equations,

$$y_j'' + \sigma_j^2 y_j = \varepsilon \sum_{k=1}^{\mu} (\phi_{jk} y_k + \psi_{jk} y_k') + \varepsilon \sum_{k=\mu+1}^n \psi_{jk} y_k, \quad j = 1, 2, \dots, \mu,$$

(4.1)

$$y_j' = \varepsilon \sum_{k=1}^{\mu} (\phi_{jk} y_k + \psi_{jk} y_k') + \varepsilon \sum_{k=\mu+1}^n \psi_{jk} y_k, \quad j = \mu+1, \dots, n,$$

where  $\varepsilon$  is a real parameter, each  $\sigma_j$  is a positive real number (and if  $\sigma_j = \sigma_j(\varepsilon)$  is an analytic function of  $\varepsilon$  at  $\varepsilon = 0$ , then  $\sigma_j > 0$  for  $\varepsilon \neq 0$ ), each  $\phi_{jk}(t)$ ,  $\psi_{jk}(t)$  is a real function, periodic in  $t$  of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ . Suppose the matrices  $\Phi = (\phi_{jk})$ ,  $j = 1, 2, \dots, n$ ,  $k = 1, 2, \dots, \mu$ ;  $\Psi = (\psi_{jk})$ ,  $j, k = 1, 2, \dots, n$ , are partitioned so that  $\Phi = (\phi_{jk})$ ,  $j = 1, 2, 3$ ,  $k = 1, 2$ ,  $\Psi = (\psi_{jk})$ ,  $j, k = 1, 2, 3$ , where  $\Phi_{11}$ ,  $\Psi_{11}$  are  $p \times p$  matrices,  $\Phi_{22}$ ,  $\Psi_{22}$  are  $(\mu - p) \times (\mu - p)$  matrices and  $\Psi_{33}$  is an  $(n - \mu) \times (n - \mu)$  matrix. Throughout the present section, we always assume that

$$(4.2) \quad \phi_{jk}(t) = (-1)^{k+j} \phi_{jk}(-t), \quad \psi_{jk}(t) = (-1)^{k+j+1} \psi_{jk}(-t)$$

for all  $j, k$ .

LEMMA (4.1). If  $\sigma_j = \sigma_j(\varepsilon)$  is an analytic function of  $\varepsilon$  at  $\varepsilon = 0$ ,  $\sigma_j(0) \neq 0 \pmod{\omega_1}$ ,  $j = 1, 2, \dots, \mu$ , and the matrices  $\Phi, \Psi$  satisfy (4.2), then there exists an  $\varepsilon_0 > 0$  such that there are  $n - \mu$  linearly independent periodic solutions of (4.1) for  $|\varepsilon| < \varepsilon_0$ .

PROOF. The transformation  $y_{\mu+j} = z_j$ ,  $j = 1, 2, \dots, n - \mu$ ,  $y_j = (2i\sigma_j)^{-1}(z_{n-\mu+j} + z_{n+j})$ ,  $y_j' = 2^{-1}(z_{n-\mu+j} - z_{n+j})$ ,  $j = 1, 2, \dots, \mu$ , leads to an equivalent system of first order equations

$$(4.3) \quad z' = Az + \varepsilon C(t)z,$$

where  $A = \text{diag}(0, \dots, 0, i\sigma_1, \dots, i\sigma_\mu, -i\sigma_1, \dots, -i\sigma_\mu)$ ,  $C = (C_{jk})$ ,  $j = 1, 2, \dots, 5$ ;

$$C_{jk} = -C_{j+2,k}, \quad j = 2, 3, \quad k = 1, 2, \dots, 5; \quad C_{j,k} = -\bar{C}_{j,k+2}, \quad k = 2, 3, \\ j = 1, 2, \dots, 5;$$

$$C_{11} = \Psi_{33}; \quad C_{21} = \Psi_{13}; \quad C_{31} = \Psi_{23}; \quad 2C_{1,k+1} = \Phi_{3k} A_k^{-1} + \Psi_{3k}, \quad k = 1, 2; \quad 2C_{j+1,k+1} = \\ = \Phi_{j,k} A_k^{-1} + \Psi_{jk}, \quad j, k = 1, 2; \quad A_1 = \text{diag}(1\sigma_1, \dots, 1\sigma_p); \quad A_2 = \text{diag}(1\sigma_{p+1}, \dots, 1\sigma_\mu).$$

With this partitioning of the matrix  $C$  it is convenient to also partition the vector  $z = (z_{(1)}, \dots, z_{(5)})$ , where  $z_{(j)}$  has a dimension compatible to the matrix multiplication. We now apply the above method of successive approximations of §2 directly to (4.3) with  $x^{(0)} = (d, 0, 0, 0, 0)$ ,  $d = (d_1, \dots, d_{n-\mu})$ , where  $d_1, \dots, d_{n-\mu}$  are arbitrary real numbers. The solution obtained in this manner will obviously be periodic. If the successive approximations  $x^{(r)}$  are partitioned the same as  $z$ , i.e.,  $x^{(r)} = (x_{(1)}^{(r)}, \dots, x_{(5)}^{(r)})$ , then we first show by induction that

$$H^{(r)} = \text{diag}(S_1^{(r)}, \dots, S_{n-\mu}^{(r)}) = 0, \\ (4.4) \quad x_{(1)}^{(r)}(-t) = x_{(1)}^{(r)}(t), \quad x_{(2)}^{(r)}(-t) = -x_{(2)}^{(r)}(t), \quad x_{(3)}^{(r)}(-t) = x_{(3)}^{(r)}(t), \\ r = 0, 1, 2, \dots$$

The induction is on the functions  $x_{(j)}^{(r)}$ . The assertion is clearly true for  $r = 0$ . Assume the assertion true for  $r = 0, 1, 2, \dots, v-1$ . Then the matrix  $C(t)x^{(r)}(t)$  satisfies the property

$$C(-t)x^{(r)}(-t) = E C(t)x^{(r)}(t), \quad E = \text{diag}(-1, -1, 1, -1, 1) \\ r = 0, 1, 2, \dots, v-1.$$

Since the first vector component of this matrix product is an odd function of  $t$  it follows immediately from (2.10) that  $H^{(r)} = 0$ ,  $r = 1, 2, \dots, v$ , for every  $d = (d_1, \dots, d_{n-\mu})$ . To complete the induction, we have from (2.11), since  $B = A$ ,

$$x^{(v)}(-t) = e^{-At} \int e^{-A\alpha} C(\alpha) x^{(v-1)}(\alpha) d\alpha = \\ = -e^{-At} \int e^{A\alpha} C(-\alpha) x^{(v-1)}(-\alpha) d\alpha = \\ = -e^{-At} \int e^{A\alpha} E C(\alpha) x^{(v-1)}(\alpha) d\alpha.$$

This is clearly assertion (4.4) for  $r = v$ , since  $C_{j,k} = -C_{j+2,k}$ ,  $j = 2, 3, k = 1, 2, \dots, 5$ ; therefore, (4.4) is true for all  $r$ . Since the  $d_j$ ,  $j = 1, 2, \dots, n - \mu$ , are arbitrary, one can find  $n - \mu$  such linearly independent solutions and the lemma is proved.

Suppose the  $\sigma_j = \sigma_j(\varepsilon)$  of (4.1) are analytic functions of  $\varepsilon$  at  $\varepsilon = 0$ ,  $\sigma_j(0) > 0$ ,  $j = 1, 2, \dots, \mu$ ,  $\sigma_j(0) - \sigma_k(0) = m_{jk}\omega$ ,  $m_{jk}$  an integer or zero,  $j, k = 1, 2, \dots, q$ , and the transformation (3.12) is applied to (4.1), to obtain the equivalent first order system

$$(4.5) \quad z' = Az + \varepsilon C(t)z$$

where  $A = \text{diag}(i\sigma_1, -i\sigma_1, \dots, i\sigma_\mu, -i\sigma_\mu, 0, \dots, 0)$  and  $C = (\frac{1}{2} C_{jk})$ ,  $j, k = 1, 2, \dots, n + \mu$ , where

$$C_{2j-1,2k-1} = \frac{\varphi_{jk}}{i\sigma_k} + \psi_{jk}, \quad C_{2j-1,2k} = \frac{\varphi_{jk}}{i\sigma_k} - \psi_{jk}, \quad k = 1, 2, \dots, \mu;$$

$$C_{2j-1,h} = \psi_{j,h}, \quad h = \mu + 1, \dots, n; \quad C_{2j,\ell} = -C_{2j-1,\ell},$$

$$(4.6) \quad \ell = 1, 2, \dots, n; \quad j = 1, 2, \dots, \mu;$$

$$C_{h,2k-1} = \frac{\varphi_{hk}}{i\sigma_k} + \psi_{hk}, \quad C_{h,2k} = \frac{\varphi_{hk}}{i\sigma_k} - \psi_{hk}, \quad k = 1, 2, \dots, \mu;$$

$$C_{h,m} = \psi_{hm}, \quad m = \mu + 1, \dots, n; \quad h = \mu + 1, \dots, n.$$

The auxiliary equation of (4.5) is chosen as

$$(4.7) \quad z' = Bz + \varepsilon C(t)z$$

where  $B = \text{diag}(i\tau_1, i\tau_2, \dots, i\tau_{2q}, i\tau_{q+1}, -i\tau_{q+1}, \dots, i\tau_\mu, -i\tau_\mu, 0, \dots, 0)$ , and the  $\tau$ 's are real numbers such that

$$\tau_{2j-1} = -\tau_{2j}, \quad \tau_{2j-1} - \tau_{2k-1} = m_{jk}\omega, \quad j, k = 1, 2, \dots, q,$$

$$(4.8)$$

$$\tau_{2j-1} \neq 0, \quad \tau_{2j-1} \pm \sigma_k(0) \neq 0 \pmod{\omega}, \quad j = 1, 2, \dots, q,$$

$$k = q + 1, \dots, n;$$

and the  $m_{jk}$  are defined above by the numbers  $\sigma_j$  at  $\varepsilon = 0$ . Notice that  $2\tau_j$  may or may not be a multiple of  $\omega$ .

LEMMA (4.2). If the matrices  $\Phi, \Psi$  associated with (4.1) satisfy (4.2) and if the alternative method of successive approximations is applied to (4.7), (4.8) with the zero<sup>th</sup> approximation given by

$$x^{(0)} = (d_1 e^{i\tau_1 t}, \dots, d_{2q} e^{i\tau_{2q} t}, 0, \dots, 0),$$

with  $d_j = b_j$ ,  $j = 1, 2, \dots, 2 \min(p, q)$ ;  $d_j = i b_j$ ,  $j = 2p + 1, \dots, 2q$ ; where each  $b_j$  is a non zero real number, then the numbers  $V_j$ ,  $j = 1, 2, \dots, 2q$ , defined by (3.9) are purely imaginary.

PROOF. For simplicity in notation we only prove the lemma for  $p \geq q$ , but it will be clear how to discuss the other situation. Applying the method (3.7), it is clear that  $x^{(r)}(t)$  has the form

$$(4.9) \quad x^{(r)}(t) = \sum_{j=1}^{2q} e^{i\tau_j t} \sum_{k=-\infty}^{+\infty} B_{jk}^{(r)} e^{ik\omega t},$$

where  $B_{jk}^{(r)}$  is an  $n + \mu$  dimensional column vector. We show by induction that

$$(4.10) \quad B_{jk}^{(r)} = \text{col}(B_{jk1}^{(r)}, 1 B_{jk2}^{(r)}, B_{jk3}^{(r)}), \quad r = 0, 1, 2, \dots,$$

and all  $j, k$ , where  $B_{jk\ell}^{(r)}$ ,  $\ell = 1, 2, 3$ , are real column vectors of dimension  $2p$ ,  $2\mu - 2p$ ,  $n - \mu$ , respectively. The assertion is clearly true for  $r = 0$ . Assume that (4.10) is true for  $r = 0, 1, \dots, v - 1$  and observe that the Fourier series for

$$C(t) \sim \sum_{k=-\infty}^{+\infty} C^{(k)} e^{ik\omega t}$$

in (4.5) is such that  $C^{(k)} = (\alpha_{h+j} C_{h,j}^{(k)})$ ,  $h, j = 1, 2, 3$ , where  $\alpha_{h+j} = 1$  if  $h + j$  is odd;  $= i$ , if  $h + j$  is even, and each  $C_{h,j}^{(k)}$  is a real matrix with  $C_{jj}^{(k)}$ ,  $j = 1, 2, 3$ , of dimensions  $2p \times 2p$ ,  $(2\mu - 2p) \times (2\mu - 2p)$  and  $(n - \mu) \times (n - \mu)$ , respectively. Therefore,

$$C(t)x^{(r)}(t) = \sum_{j=1}^{2q} e^{i\tau_j t} \sum_{k=-\infty}^{+\infty} D_{jk}^{(r)} e^{ik\omega t}, \quad r = 1, 2, \dots, v - 1,$$

where  $D_{jk}^{(r)} = \text{col}(iD_{jk1}^{(r)}, D_{jk2}^{(r)}, D_{jk3}^{(r)})$  where each of the real vectors  $D_{jkl}^{(r)}, l = 1, 2, 3$ , has the same dimension as  $B_{jkl}^{(r)}$ . From (3.6) and the assumption on the numbers  $d_j$  in the statement of the lemma,

$$(4.11) \quad S_j^{(r)} \text{ is purely imaginary, } j = 1, 2, \dots, 2q; r = 1, 2, \dots, v.$$

It then follows immediately from (3.7) and the above results that (4.10) is true for  $r = v$  and, therefore, (4.10) is true for all  $r$ . Finally, (4.11) also holds for all  $r$  and the statement of the lemma is true.

REMARK (4.1). Lemma (4.2) holds also in a slightly more general situation, namely when  $q$  of the  $\sigma$ 's differ by a multiple of  $\omega$  at  $\varepsilon = 0$ , but the ones which satisfy this property are not necessarily the first  $q$ . The notation for the statement of this result as in Lemma (4.2) is extremely complicated, but it should be clear that the same conclusion is true, the only thing necessary is the proper choice of the numbers  $d_j$ .

REMARK (4.2). When  $\mu = n$  in (4.1), it is obvious that the condition  $\tau_{2j-1} \neq 0 \pmod{\omega}$  in (4.8) may be eliminated.

REMARK (4.3). The determining equations for the  $2q + 1$  real numbers  $b_1, \dots, b_{2q}$  and  $\tau_1$  [only  $\tau_1$ , since the  $\tau_j$  are related by (4.8)] in Lemma (4.2) are the  $2q$  real equations

$$(4.12) \quad \tau_j - \varepsilon I[V_j(\tau, b, \varepsilon)] = (-1)^{j+1} \sigma_j, \quad j = 1, 2, \dots, 2q,$$

$b = (b_1, \dots, b_{2q})$  and  $I[w]$  is the imaginary part of a complex number  $w$ . If, in addition to condition (4.8), we assume that  $2\tau_j \neq 0 \pmod{\omega}$ ,  $j = 1, 2, \dots, 2q$ ; and choose  $b_{2j-1} = -b_{2j}$ ,  $j = 1, 2, \dots, p$ ,  $b_{2j-1} = b_{2j}$ ,  $j = p + 1, \dots, q$  in Lemma (4.2), then we know from Lemma (3.1) that  $V_{2j-1} = \bar{V}_{2j}$ ,  $j = 1, 2, \dots, q$ , and the determining equations (4.12) are equivalent to the equations

$$(4.13) \quad \tau_{2j-1} - \varepsilon I[V_{2j-1}(\tau, b, \varepsilon)] = \sigma_j, \quad j = 1, 2, \dots, q,$$

for the  $q + 1$  real numbers  $b_1, b_3, \dots, b_{2q-1}$  and  $\tau_1$ . This remark will be important for the applications.

REMARK (4.4). In the previous remark, equations (4.12), (4.13) were considered as determining equations for  $b$  and  $\tau_1$ , but we could just as well have considered  $\tau_1$  as fixed and determined the numbers  $\sigma, b$  so as to obtain a solution of (4.5). This will be important in determining the stable and unstable regions around "resonance points".



## §5. SOME BOUNDEDNESS THEOREMS

As an application of the lemmas in §4, we state a few theorems concerning the behavior of the characteristic exponents of (4.1) and also some theorems concerning the boundedness or unboundedness of the AC solutions of (4.1) in  $(-\infty, +\infty)$ .

THEOREM (5.1). If the matrices  $\Phi, \Psi$  associated with system (4.1) satisfy (4.2) and the numbers  $\sigma_j = \sigma_j(\varepsilon)$  are real analytic functions of  $\varepsilon$  at  $\varepsilon = 0$ , with

$$(5.1) \quad \sigma_j(0) \not\equiv 0 \pmod{\omega}, \quad j = 1, 2, \dots, \mu,$$

then there exists an  $\varepsilon_0 > 0$  such that there are always  $n - \mu$  linearly independent periodic solutions of (4.1) for  $|\varepsilon| < \varepsilon_0$ . If one of the numbers  $\sigma_j$ , say  $\sigma_1$ , is such that

$$(5.2) \quad 2\sigma_1(0) \not\equiv 0, \quad \sigma_1(0) \not\equiv \pm \sigma_k(0) \pmod{\omega}, \quad k = 2, 3, \dots, \mu,$$

then there are always two characteristic exponents  $i\tau_1, i\tau_2$  of (4.1) which are purely imaginary for  $|\varepsilon| < \varepsilon_0$ ,  $\varepsilon_0 > 0$ , are analytic in  $\varepsilon$  at  $\varepsilon = 0$  and  $\tau_1 = -\tau_2$ ,  $\tau_1(0) = \sigma_1$ .

PROOF. The first part of the theorem is a restatement of Lemma (4.1). The second part of the theorem is an immediate consequence of Lemma (4.2) and Remark (4.3). For since  $q = 1$ , the implicit function theorem assures us that Equation (4.13) has a real solution  $\tau_1$  analytic in  $\varepsilon$  at  $\varepsilon = 0$  with  $\tau_1(0) = \sigma_1$ .

COROLLARY (5.1). If the conditions of Theorem (5.1) are satisfied and

$$(5.3) \quad 2\sigma_j(0) \not\equiv 0, \quad \sigma_j(0) \not\equiv \pm \sigma_k(0) \pmod{\omega}, \quad j \neq k, \quad j, k = 1, 2, \dots, \mu;$$

then all of the AC solutions of (4.1) are bounded for  $|\varepsilon| < \varepsilon_0$ ,  $\varepsilon_0 > 0$ .

PROOF. This result follows immediately by applying Theorem (5.1) to each of the numbers,  $\sigma_j$ ,  $j = 1, 2, \dots, \mu$ .

Corollary (5.1) coincides with a previous result obtained by the author [10e]. For  $\mu = n - 1$ , this corollary has also been obtained by the author [10f] without using successive approximations. For  $\mu = n$ , this result has also been obtained by M. Golomb [7].

Consider the system of differential equations (4.1) satisfying (4.2) and suppose that  $\sigma_1(0) - \sigma_2(0) = s\omega$  where  $s$  is an integer or zero and each  $\sigma_j$  is a real positive analytic function of  $\varepsilon$  at  $\varepsilon = 0$  with

$$(5.4) \quad 2\sigma_j(0) \neq 0, \quad \sigma_j(0) \pm \sigma_k(0) \neq 0 \pmod{\omega} \quad j \neq k, \quad j, k = 2, 3, \dots, \mu.$$

If  $j, k$  in (5.4) are allowed to take on all integer values from 1 to  $\mu$ , it follows from Corollary (5.1) that all of the AC solutions of (4.1) are bounded for  $|\varepsilon|$  sufficiently small. We now wish to determine additional conditions on the functions  $\varphi_{jk}, \psi_{jk}$  in (4.1) so that all of the AC solutions are still bounded for  $|\varepsilon|$  sufficiently small.

It follows from Theorem (5.1) that there are  $n + \mu - 4$  linearly independent bounded AC solutions of (4.1) for  $|\varepsilon|$  sufficiently small. Furthermore, the characteristic exponents corresponding to these solutions are either zero or close to one of the numbers  $i\sigma_j, -i\sigma_j, j = 3, 4, \dots, \mu$ . It remains only to determine the four characteristic exponents close to the numbers  $i\sigma_j, -i\sigma_j, j = 1, 2$ . Consider the equivalent first order system (4.5), (4.6) and apply the alternative method of successive approximations (3.7) to the auxiliary equation

$$(5.5) \quad z' = Bz + \varepsilon C(t)z, \quad$$

where

$$(5.6) \quad B = \text{diag}(i\tau, -i\bar{\tau}, i(\tau - s\omega), -i(\bar{\tau} - s\omega), i\sigma_3, -i\sigma_3, \dots, i\sigma_\mu, -i\sigma_\mu, 0, \dots, 0),$$

and the zero<sup>th</sup> approximation is

$$(5.7) \quad x^{(0)} = (i b e^{i\tau t}, i \bar{b} e^{-i\bar{\tau} t}, i a e^{i(\tau - s\omega)t}, i \bar{a} e^{-i(\bar{\tau} - s\omega)t}, 0, \dots, 0),$$

where  $\tau, a, b$  are undetermined complex numbers, and if  $\tau$  is a function of  $\varepsilon$ , then  $\tau$  is close enough to  $\sigma_1$  so that (5.4) is satisfied with  $\sigma_2(0)$  replaced by  $\tau(0) - s\omega$ .

From Lemma (3.1),  $S_1^{(r)} = \bar{S}_2^{(r)}, S_3^{(r)} = \bar{S}_4^{(r)}$  for every  $r = 1, 2, \dots$ , and all complex numbers  $a, b$ . Therefore, the determining equations

$$(5.8) \quad \begin{aligned} 1\tau - \varepsilon V_1(\tau, a, b, \varepsilon) &= 1\sigma_1 \\ 1\tau - \varepsilon V_3(\tau, a, b, \varepsilon) &= 1(\sigma_2 + s\omega) \end{aligned}$$

represent two equations for the undetermined parameters  $\tau, a$ . We now calculate the first terms of  $V_1$  and  $V_3$  to obtain the characteristic exponent  $\tau$ .

Suppose the Fourier series of  $\varphi_{jk}(t), \psi_{jk}(t)$  are given by

$$(5.9) \quad \varphi_{jk}(t) \sim \sum_{\ell=-\infty}^{+\infty} b_{jk\ell} e^{i\ell\omega t}, \quad \psi_{jk}(t) \sim \sum_{\ell=-\infty}^{+\infty} d_{jk\ell} e^{i\ell\omega t}, \quad j, k = 1, 2, \dots, n.$$

From (4.2) it follows that each  $b_{jk\ell}, d_{jk\ell}$  must be either purely imaginary or real and  $b_{jk,\ell} = \bar{b}_{jk,-\ell}$ . If

$$(5.10) \quad b_{jjo} = d_{jjo} = b_{jks} = d_{jks} = 0, \quad j \neq k, \quad j, k = 1, 2, \dots,$$

then  $S_1^{(1)} = S_3^{(1)} = 0$ . Calculating the functions  $x_j^{(1)}, j = 1, 2, \dots, n + \mu$ , from (3.7) and (4.6), it is then very easy to show that

$$(5.11) \quad S_1^{(2)} = -\frac{1}{4i} [b\alpha_{11}^{(s)}(\tau) + a\alpha_{12}^{(s)}(\tau)],$$

$$S_3^{(r)} = -\frac{1}{4ia} [b\alpha_{21}^{(s)}(\tau) + a\alpha_{22}^{(s)}(\tau)],$$

where

$$(5.12) \quad \begin{aligned} \alpha_{11}^{(s)}(\tau) &= \beta_{11}(0, 0), \quad \alpha_{12}^{(s)}(\tau) = \beta_{21}(s, 0), \quad \alpha_{21}^{(s)}(\tau) = \beta_{12}(0, s), \\ \alpha_{22}^{(s)}(\tau) &= \beta_{22}(s, s) \end{aligned}$$

$$\begin{aligned} \beta_{uv}(\ell_1, \ell_2) &= \sum_{k=1}^{\mu} \sum_{\ell=-\infty}^{+\infty} \gamma_{ku, \ell+\ell_1}^{(+)} [\gamma_{vk, -(\ell+\ell_2)}^{(+)} \delta_{\ell}(\nu_k) - \gamma_{vk, -(\ell+\ell_2)}^{(-)} \delta_{\ell}(-\bar{\nu}_k)] + \\ &+ \sum_{k=\mu+1}^n \sum_{\ell=-\infty}^{+\infty} \gamma_{ku, \ell+\ell_1}^{(+)} \cdot b_{vk_1, -(\ell+\ell_2)} \cdot \delta_{\ell}(0) \end{aligned}$$

$$\gamma_{kj\ell}^{(+)} = \left( \frac{b_{kj\ell}}{\sigma_j} + id_{kj\ell} \right), \quad \gamma_{kj\ell}^{(-)} = \left( \frac{b_{kj\ell}}{\sigma_j} - id_{kj\ell} \right), \quad \delta_\ell(v_k) = \frac{1}{\ell\omega + \tau - v_k},$$

$$v_1 = \tau, \quad v_2 = \tau - s\omega, \quad v_j = \sigma_j, \quad j = 3, 4, \dots, \mu,$$

and the index  $\ell + \ell_j$  in (5.12) does not take on the values in (5.10).

In order to make the statement of the following theorem meaningful, first observe that the conditions (4.2) and Lemma (4.2) imply that each of the numbers  $\alpha_{jk}^{(s)}$  is real if  $\tau$  is real and either  $p \geq 2$  or  $p = 0$ , and  $\alpha_{11}^{(s)}, \alpha_{22}^{(s)}$  are real,  $\alpha_{12}^{(s)}, \alpha_{21}^{(s)}$  are purely imaginary if  $\tau$  is real and  $p = 1$ , since the  $\alpha_{jk}^{(s)}$  are independent of the numbers  $d_j$  in Lemma (4.2).

**THEOREM (5.2).** Consider the system of differential equations (4.1) with the functions  $\varphi_{jk}(t), \psi_{jk}(t)$  satisfying the condition (4.2) and having the Fourier series

$$\varphi_{jk}(t) \sim \sum_{\ell=-\infty}^{+\infty} b_{jk\ell} e^{i\ell\omega t}, \quad \psi_{jk}(t) \sim \sum_{\ell=-\infty}^{+\infty} d_{jk\ell} e^{i\ell\omega t},$$

$b_{jj0} = d_{jj0} = b_{jks} = d_{jks} = 0, \quad j \neq k, \quad j, k = 1, 2$ .  
If  $\sigma_1(0) - \sigma_2(0) = s\omega$ , where  $s$  is an integer or zero and each  $\sigma_j$  is a real positive analytic function of  $\varepsilon$  at  $\varepsilon = 0$  with  $2\sigma_j(0) \neq 0, \sigma_j(0) \pm \sigma_k(0) \neq 0 \pmod{\omega}, j \neq k, j, k = 2, 3, \dots, \mu$ ; if the numbers  $\alpha_{uv}^{(s)}(\tau)$  are defined by (5.12) and  $G(s, \sigma_1, \sigma) = [\alpha_{22}^{(s)}(\sigma_1) - \alpha_{11}^{(s)}(\sigma_1) + 4s], H(s, \sigma_1) = 4\alpha_{21}^{(s)}(\sigma_1)\alpha_{12}^{(s)}(\sigma_1)$  where  $\sigma = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-2} [\sigma_2(\varepsilon) - \sigma_1(\varepsilon) + s\omega]$  is finite, then the following conclusions hold:

- (i) If  $H(s, \sigma_1) \neq 0, [G(s, \sigma_1, \sigma)]^2 + H(s, \sigma_1) > 0$  for  $\varepsilon = 0$ , then there exists an  $\varepsilon_0 > 0$  such that all of the AC solutions of (4.1) with the  $\sigma_j$  as above are bounded for  $|\varepsilon| < \varepsilon_0$ ;  
(ii) If  $H(s, \sigma_1) \neq 0, [G(s, \sigma_1, \sigma)]^2 + H(s, \sigma_1) < 0$  for  $\varepsilon = 0$ , then some of the AC solutions of (4.1) with the  $\sigma_j$  as above are unbounded for every  $\varepsilon \neq 0$ .

**PROOF.** From the remarks preceding the statement of the theorem

it is sufficient to discuss the four characteristic exponents which are close to the numbers  $i\sigma_j$ ,  $-i\sigma_j$ . For  $\epsilon \neq 0$ , the determining equations for these characteristic exponents are given by

$$1F_1(\tau, a, b, \epsilon) = 1\tau + \frac{\epsilon^2}{4i} [\alpha_{11}^{(s)}(\tau) + \frac{a}{b} \alpha_{12}^{(s)}(\tau)] - i\sigma_1 + o(\epsilon^3) = 0 \quad (5.13)$$

$$F_2(\tau, a, b, \epsilon) = \frac{a}{b} \alpha_{12}^{(s)}(\tau) - G(s, \sigma_1, \sigma) - \frac{b}{a} \alpha_{21}^{(s)}(\tau) + o(\epsilon) = 0$$

where the  $\alpha_{jk}^{(s)}$  are given by (5.12). Any solution  $\tau, a, b$  of Equation (5.13) yields two characteristic exponents  $i\tau, -i\tau$ .

Suppose that the conditions of (1) are satisfied. If the integer  $p$  associated with system (4.1), (4.2) is  $\geq 2$ , then, from the remark preceding the statement of the theorem, all of the numbers  $\alpha_{jk}^{(s)}(\sigma_1)$  are real. Furthermore, from Lemma (4.2), the functions  $F_1, F_2$  in (5.13) are real if  $\tau, a$  are real and  $b = 1$ . The numbers  $\tau_0 = \sigma_1(0)$ ,  $a_0 = [G(s, \sigma_1, \sigma) \pm [G^2(s, \sigma_1, \sigma) + H(s, \sigma_1)]^{\frac{1}{2}}] \cdot [2\alpha_{12}^{(s)}(\sigma_1)]^{-1}$  are such that  $F_1(\tau, a_0, 0) = F_2(\tau_0, a_0, 0) = 0$  and the Jacobian of these two functions with respect to  $\tau, a$  at  $\tau = \tau_0, a = a_0, \epsilon = 0$  is different from zero. Consequently, Equations (5.13) have two real distinct solutions  $\tau = \tau(\epsilon), a = a(\epsilon)$  analytic in  $\epsilon$  at  $\epsilon = 0$  and

$$\tau = \sigma_1 + \frac{\epsilon^2}{4} [\alpha_{11}^{(s)}(\sigma_1) + a_0 \alpha_{12}^{(s)}(\sigma_1)] + o(\epsilon^3), a = a_0 + o(\epsilon).$$

For  $p = 1$ , take  $a$  purely imaginary,  $b = 1$  and for  $p = 0$ , take  $a$  purely imaginary,  $b = -1$ , and apply the same reasoning.

The reasoning for the proof of (ii) of the theorem is exactly the same as above except that we do not have to apply Lemma (4.2) but apply the implicit function theorem directly to the functions  $F_1, F_2$  in Equations (5.13). The two  $\tau$ 's obtained are obviously such that one  $i\tau$  has a real part positive and the other has a real part negative. Therefore, the theorem is proved.

REMARK (5.1). In the statement of Theorem (5.2) some cases have been excluded, in particular, the case where  $H(s, \sigma_1) \neq 0$  and the other expression in Theorem (5.2) is equal to zero. Reasoning as in the proof of (i) Of Theorem (5.2), one sees that two of the remaining characteristic roots are still purely imaginary, but one cannot decide using the above method whether the solutions are bounded or unbounded. The other cases may be treated by going to higher approximations in the method.

Theorem (5.2) gives some insight into the behavior of the solutions of (4.1) near the resonance point  $\sigma_1(0) = \sigma_2(0) = s_0$ . More

specifically, if  $\sigma_1, \sigma_3, \dots, \sigma_n$  are fixed and say independent of  $\varepsilon$ , if the conditions of Theorem (5.2) are satisfied and  $H(s, \sigma_1) \neq 0$ , then all of the AC solutions of (4.1) are bounded for  $\varepsilon$  sufficiently small along the curve  $\sigma_2 = \sigma_1 - s\omega + \varepsilon^2\sigma$  in the  $(\sigma_2, \varepsilon)$  plane if the discriminant  $D(\sigma) = [G(s, \sigma_1, \sigma)]^2 + H(s, \sigma_1) > 0$  and some solutions are unbounded along this curve for every  $\varepsilon \neq 0$  if  $D(\sigma) < 0$ . Furthermore, if  $H(s, \sigma_1) > 0$ , then  $D(\sigma) > 0$  for every  $\sigma$  and the AC solutions of (4.1) are bounded along every curve  $\sigma_2 = \sigma_1 - s\omega + \varepsilon^2\sigma$  in the  $(\sigma_2, \varepsilon)$  plane for  $\varepsilon = \varepsilon(\sigma)$  sufficiently small. On the other hand if  $H(s, \sigma_1) < 0$ , then there exists two  $\sigma$ 's such that  $G(\sigma) = 0$ . If these two  $\sigma$ 's are distinct then there is a region in the neighborhood of the point  $(\sigma_1 - s\omega, 0)$  of the  $(\sigma_2, \varepsilon)$  plane where some solutions are unbounded no matter how small  $\varepsilon \neq 0$ . These results are summarized in the following theorem.

THEOREM (5.3). If the conditions of Theorem (5.2) are satisfied and  $\sigma_1, \sigma_j, j = 3, 4, \dots, n$ , are fixed numbers independent of  $\varepsilon$ , then

(i)  $H(s, \sigma_1) > 0$  for  $\varepsilon = 0$  implies that the AC solutions of (4.1) are bounded in a sufficiently small neighborhood of the point  $(\sigma_1 - s\omega, 0)$  in the  $(\sigma_2, \varepsilon)$  plane:

(ii)  $H(s, \sigma_1) < 0$  for  $\varepsilon = 0$  implies that some of the AC solutions of (4.1) are unbounded in every neighborhood of the point  $(\sigma_1 - s\omega, 0)$  of the  $(\sigma_2, \varepsilon)$  plane.

REMARK (5.2). Theorems (5.2) and (5.3) deal only with the case  $\sigma_1(0) - \sigma_2(0) = s\omega$ . The same type of analysis could be used to discuss the resonance points  $\sigma_1(0) + \sigma_2(0) = s\omega$ . The essential element of the argument was (5.11) and Lemma (4.2). It should be clear that a statement similar to Lemma (4.2) is true for the more general situation where  $\sigma_j(0) + \sigma_k(0) = n_{jk}\omega$ ,  $n_{jk}$  an integer or zero,  $j, k \in T_1$ ,  $\sigma_j(0) \mp \sigma_k(0) = m_{jk}\omega$ ,  $m_{jk}$  an integer or zero,  $j, k \in T_2$ , where  $T_1, T_2$  are subsets of the set of integers  $(1, 2, \dots, \mu)$ , if the matrix  $B$  in (4.7) is chosen properly and the numbers  $\tau$  satisfy a condition similar to (4.7). That the functions  $S_1, S_3$  have the form (5.11) when  $\sigma_1(0) + \sigma_2(0) = s\omega$ ,  $2\sigma_j(0) \neq 0$ ,  $\sigma_j(0) \pm \sigma_k(0) \neq 0 \pmod{\omega}$ ,  $j \neq k$ ,  $j, k = 2, 3, \dots, \mu$ , follows immediately from the method of successive approximations. With the generalization of Lemma (4.2) mentioned above, one can discuss the region of stability and instability of systems (4.1) when any of the numbers  $\sigma_j$  are "in resonance" with  $\omega$ . However, the results in general cannot be stated explicitly as in Theorems (5.2) and (5.3) since the determining

equations are extremely complicated.

REMARK (5.3). As remarked in the Introduction, if  $\mu = n$  in (4.1),  $\Psi = 0$ , and  $\Phi$  is symmetric, J. M. Gel'fand and V. B. Lidskii [14] and J. Moser [13] have shown that only the points  $\sigma_j + \sigma_k = s\omega$ ,  $\sigma_j > 0$  may lead to unbounded solutions. On the other hand, for systems (4.1), (4.2), Theorem (5.2) implies the behavior of the solutions at the resonance points  $\sigma_1 - \sigma_2 = s\omega$  depends upon the value of a certain discriminant. Therefore if  $\mu = n$ ,  $\Psi = 0$ ,  $\Phi$  is symmetric and also satisfies (4.2), it must be true that the function  $H(s, \sigma_1)$  in Theorem (5.3) is  $> 0$  for  $\varepsilon = 0$ . This is indeed the case; for the functions  $\alpha_{jk}^{(s)}$  in (5.12) are given by

$$\alpha_{11}^{(s)}(\tau) = \sum_{k=1}^{\mu} \sum_{\ell=-\infty}^{+\infty} \frac{|b_{k1,\ell}|^2}{\sigma_1 \sigma_k} A_{\ell k}; \quad \alpha_{12}^{(s)}(\tau) = \sum_{k=1}^{\mu} \sum_{\ell=-\infty}^{+\infty} \frac{b_{k2,\ell+s} \bar{b}_{k1,\ell}}{\sigma_2 \sigma_k} A_{\ell k};$$

(5.14)

$$\alpha_{21}^{(s)}(\tau) = \sum_{k=1}^{\mu} \sum_{\ell=-\infty}^{+\infty} \frac{b_{k1,\ell} \bar{b}_{k2,\ell+s}}{\sigma_1 \sigma_k} A_{\ell k}; \quad \alpha_{22}^{(s)}(\tau) = \sum_{k=1}^{\mu} \sum_{\ell=-\infty}^{+\infty} \frac{|b_{k2,\ell+s}|^2}{\sigma_2 \sigma_k} A_{\ell k}$$

where  $A_{\ell k} = \delta_{\ell}(v_k) - \delta_{\ell}(-\bar{v}_k)$ ,  $v_1 = \tau$ ,  $v_2 = \tau - s\omega$ ,  $v_k = \sigma_k$ ,  $k = 3, 4, \dots, n$ . Consequently,

$$\alpha_{12}^{(s)}(\sigma_1) = \frac{\sigma_1}{\sigma_2} \bar{\alpha}_{21}^{(s)}(\sigma_1)$$

and  $H(s, \sigma_1) > 0$  for all  $s$  if  $\alpha_{12}^{(s)}(\sigma_1) \neq 0$ . Actually, all that we have required to obtain boundedness at  $\sigma_1 - \sigma_2 = s\omega$  is  $\varphi_{1k} = \varphi_{k1}$ ,  $\varphi_{2k} = \varphi_{k2}$ ,  $k = 1, 2, \dots, n$ . To obtain boundedness at all of the points  $\sigma_j - \sigma_k = s\omega$ ,  $j \neq k$ ,  $j, k = 1, 2, \dots, n$ , the matrix  $\Phi$  must be symmetric.

In case  $\Phi$  is not symmetric, then some solution may not be bounded for an  $\varepsilon \neq 0$ . For example, consider the system (4.1), (4.2) with  $\mu = n = 2$ ,  $\Psi = 0$ ,  $\sigma_1 = \sigma_2(0)$ ,  $\sigma_1$  independent of  $\varepsilon$ ,

$$\Phi = \begin{bmatrix} 2p \cos t & 2 \cos t \\ 2q \cos t & 2r \cos t \end{bmatrix}.$$

From (5.12),  $\alpha_{12}^{(0)}(\sigma_1) = \beta(p+r)$ ,  $\alpha_{21}^{(0)}(\sigma_1) = \beta q(p+r)$ ,  $\beta = \sigma_1^{-1}(4\sigma_1^2 - 1)^{-1}$  for  $\varepsilon = 0$ . Therefore, from Theorem 5.3 the AC solutions are bounded in

a neighborhood of the point  $(\sigma_1, 0)$  in the  $(\sigma_2, \varepsilon)$  plane for  $\varepsilon$  sufficiently small if  $p + r \neq 0$ ,  $q > 0$ , and some solutions are unbounded in every neighborhood of this point if  $p + r \neq 0$ ,  $q < 0$ .

As another example, consider system (4.1), (4.2) with  $\mu = n = 2$ ,  $\Psi = 0$ ,  $\sigma_1 - \sigma_2(0) = 1$ ,  $\sigma_1$  independent of  $\varepsilon$ ,

$$\Phi = \begin{bmatrix} 2p \cos t & 2 \cos 2t \\ 2q \cos 2t & 2r \cos 2t \end{bmatrix}.$$

From (5.12),  $\alpha_{12}^{(1)}(\sigma_1) = \beta(\sigma_1 - 1)^{-1}$ ,  $\alpha_{21}^{(1)}(\sigma_1) = \beta q \sigma_1^{-1}$ ,  $\beta = 2p(1 + 2\sigma_1)^{-1}$  for  $\varepsilon = 0$ . Therefore, from Theorem (5.3) all of the AC solutions are bounded in a neighborhood of the point  $(\sigma_1 - 1, 0)$  of the  $(\sigma_2, \varepsilon)$  plane for  $\varepsilon$  sufficiently small if  $p \neq 0$ ,  $\sigma_1 > 1$ ,  $q > 0$ , and some solutions are unbounded in every neighborhood of this point if  $p \neq 0$ ,  $\sigma_1 > 1$ ,  $q < 0$ .

The following theorem concerns the behavior of the solutions of (4.1), (4.2) with  $\mu = n$  when one of the  $\sigma$ 's approaches zero as  $\varepsilon$  approaches zero.

THEOREM (5.4). Consider the system of differential equations of order  $2n$ ,

$$(5.15) \quad y'' = Dy = \varepsilon \Phi(t)y + \varepsilon \Psi(t)y'$$

where  $\varepsilon > 0$ ,  $D = \text{diag}(\varepsilon \sigma_1^2, \sigma_2^2, \dots, \sigma_n^2)$ ,  $\Phi$  and  $\Psi$  are  $n \times n$  real matrices whose elements  $\phi_{ij}(t)$ ,  $\psi_{ij}(t)$  are periodic in  $t$  of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$  and have mean value zero;  $\Phi = (\phi_{jk})$ ,  $\Psi = (\psi_{jk})$ ,  $j, k = 1, 2, \dots, n$ , where  $\phi_{jk}$ ,  $\psi_{jk}$  are matrices of the same dimension and

$$(5.16) \quad \phi_{jk}(-t) = (-1)^{k+j} \phi_{jk}(t), \quad \psi_{jk}(-t) = (-1)^{k+j+1} \psi_{jk}(t), \\ j, k = 1, 2, \dots, n.$$

If  $\sigma_j > 0$ ,  $j = 1, 2, \dots, n$ ,

$$(5.17) \quad 2\sigma_k \neq 0, \sigma_j \pm \sigma_k \neq 0, j \neq k \pmod{\omega}, j = k, j, k = 2, 3, \dots, n,$$

[if the  $\sigma_j$  are analytic functions of  $\varepsilon$  at  $\varepsilon = 0$ , then  $\sigma_k$  in (5.17) is replaced by  $\sigma_k(0)$ ], then there exists an  $\varepsilon_0 > 0$  such that all of the absolutely



continuous (AC) solutions of (5.15) are bounded for  $0 < \varepsilon < \varepsilon_0$  (notice this is not true for  $\varepsilon = 0$ ). Furthermore, the characteristic exponents of (5.15) are analytic functions of  $\sqrt{\varepsilon}$  at  $\varepsilon = 0$ .

PROOF. By using an elementary argument, we show that there are  $2n - 2$  linearly independent bounded AC solutions of (5.15). This same result could also be proved by the method of successive approximations. The characteristic exponents  $\tau_k$  are only determined up to a multiple of  $\omega_1$ , but without loss of generality we may assume that the  $\tau_k(\varepsilon)$ ,  $k = 3, 4, \dots, 2n$  are such that  $\tau_{2j-1}(0) = i\sigma_j$ ,  $\tau_{2j}(0) = -i\sigma_j$ ,  $j = 2, 3, \dots, n$ . From the Floquet theory condition (5.17) assures us that these functions  $\tau_k(\varepsilon)$ ,  $k = 3, 4, \dots, 2n$ , are analytic in  $\varepsilon$  at  $\varepsilon = 0$  and

$$\begin{aligned} \tau_{2j-1}(\varepsilon) &= \bar{\tau}_{2j}(\varepsilon), \quad j = 2, 3, \dots, n; \\ (5.19) \quad \tau_j &\neq 0, \quad \tau_j \not\equiv \tau_k \pmod{\omega_1}, \quad j \neq k, \quad j, k = 3, 4, \dots, n; \\ 0 &\leq \varepsilon \leq \varepsilon_0, \quad \varepsilon_0 > 0, \end{aligned}$$

provided only that  $\varepsilon_0$  is sufficiently small. If  $y(t)$  is a solution of (5.15), there exist, from (5.16), a nonsingular matrix  $P$  such that  $Py(-t)$  is also a solution of (5.15). This, together with (5.18) implies that each  $\tau_k$ ,  $k = 3, 4, \dots, 2n$ , is purely imaginary and this obviously implies the existence of  $2n - 2$  linearly independent solutions of (5.15).

It does not seem possible to obtain the other two characteristic exponents by such an elementary argument so we apply the above method of successive approximation.

The transformation of variables  $y_j = (2i\sigma_j)^{-1}(z_{2j-1} + z_{2j})$ ,  $y_j' = 2^{-1}(z_{2j-1} - z_{2j})$ ,  $j = 2, 3, \dots, n$ ,  $y_1 = z_1$ ,  $y_1' = \lambda z_2$ ,  $\lambda = \sqrt{\varepsilon}$ , leads to the equivalent system of first order equations

$$(5.19) \quad z' = Az + \lambda C(t, \lambda)z$$

where  $A = \text{diag}(0, 0, i\sigma_2, -i\sigma_2, \dots, i\sigma_n, -i\sigma_n)$ ,  $C = (c_{jk})$ ,  $j, k = 1, 2, \dots, 2n$ ,

$$c_{11} = 0, c_{12} = 1, c_{1j} = 0, j = 2, 3, \dots, 2n, c_{21} = \varphi_{11} - \sigma_1^2, \\ c_{22} = \psi_{11}$$

$$c_{2,2j-1} = (\frac{\varphi_{1j}}{1\sigma_j} + \psi_{1j}), c_{2,2j} = (\frac{\varphi_{1j}}{1\sigma_j} - \psi_{1j}), j = 2, 3, \dots, n,$$

$$(5.20) \quad c_{2k,j} = -c_{2k-1,j}, k = 2, 3, \dots, n; j = 1, 2, \dots, 2n,$$

$$c_{2k-1,1} = \lambda \varphi_{k,1}, c_{2k-1,2} = \lambda^2 \psi_{k,1}, k = 2, 3, \dots, n,$$

$$c_{2j-1,2k-1} = \lambda (\frac{\varphi_{jk}}{1\sigma_k} + \psi_{jk}), c_{2j-1,2k} = \lambda (\frac{\varphi_{jk}}{1\sigma_k} - \psi_{jk}), \\ j, k = 2, 3, \dots, n.$$

To obtain the two characteristic exponents close to zero, we apply the first method of successive approximations to obtain a solution of the form  $z = e^{1\tau t} p(t)$ ,  $p(t+T) = p(t)$ , where  $\tau$  is a real number. The equation for  $p(t)$  is

$$(5.21) \quad p' = (A - 1\tau I)p + \lambda C(t, \lambda)p.$$

If  $\tau$  is close enough to zero so that

$$(5.22) \quad \sigma_k \pm \tau \neq m\omega, k = 2, 3, \dots, n; m = 0, 1, 2, \dots,$$

then the method of successive approximations may be applied to the auxiliary equation

$$(5.23) \quad p' = Bp + \lambda C(t, \lambda)p$$

where  $B = \text{diag}(0, 0, 1(\sigma_2 - \tau), -1(\sigma_2 + \tau), \dots, 1(\sigma_n - \tau), -1(\sigma_n + \tau))$  and the zero<sup>th</sup> approximation is taken to be

$$(5.24) \quad p^{(0)} = x^{(0)} = (1, 1b, 0, \dots, 0)$$

where  $b$  is a nonzero real number. The functions  $x^{(r)}$  obtained in this manner are periodic of period  $T = 2\pi/\omega$ . If  $x^{(r)}(t)$  has the Fourier series

$$(5.25) \quad x^{(r)}(t) \sim \sum B_k^{(r)} e^{ik\omega t}$$

where  $B_k^{(r)}$  is a  $2n$  dimensional column vector, then we show by induction that

$$B_k^{(r)} = \text{col} (B_{k1}^{(r)}, iB_{k2}^{(r)}, iB_{k3}^{(r)}, B_{k4}^{(r)}), \quad k = 0, \pm 1, \pm 2, \dots; \quad (5.26)$$

$$r = 0, 1, 2, \dots$$

where  $B_{k1}^{(r)}, B_{k2}^{(r)}$  are real scalars,  $B_{k3}^{(r)}$  is a  $2\mu - 2$  dimensional real vector and  $B_{k4}^{(r)}$  is a  $2(n - \mu)$  dimensional real vector. The assertion is clearly true for  $r = 0$ . Assume that it is true for  $r = 0, 1, 2, \dots, v - 1$ . Then from the definition of the  $c_{jk}$  in (5.20) and the assumptions on the  $\varphi_{jk}, \psi_{jk}$  of the theorem, it is clear that  $C(t)x^{(r)}(t) \sim \sum D_k^{(r)} e^{ik\omega t}$  where

$$D_k^{(r)} = \text{col} (iD_{k1}^{(r)}, D_{k2}^{(r)}, D_{k3}^{(r)}, iD_{k4}^{(r)}) \quad (5.27)$$

and the real vectors  $D_{kj}^{(r)}$  have the same dimension as the  $B_{kj}^{(r)}$ . From (3.6) and (5.24)

$$S_1^{(r)}, S_2^{(r)} \text{ are purely imaginary, } r = 1, 2, \dots, v. \quad (5.28)$$

It is now very easy to show from (3.7), (5.27), (5.28) that (5.26) is true for  $r = v$  and the induction is completed. Consequently (5.28) is true for all  $r$  and the determining equations

$$\begin{aligned} -i\tau &= -\lambda S_1^{(1)} - \lambda^2 S_1^{(2)} \dots \\ -i\tau &= -\lambda S_2^{(1)} - \lambda^2 S_2^{(2)} \dots \end{aligned} \quad (5.29)$$

are purely imaginary. Furthermore, it is immediate from (3.6) and (5.24) that

$$S_1^{(1)} = ib, \quad S_2^{(1)} = -\frac{\sigma_1^2}{ib}$$

and for  $\lambda \neq 0$ , the determining equations (5.29) become

$$\begin{aligned} \tau &= i\lambda b + o(\lambda^2) \\ b - \frac{\sigma_1^2}{b} &= o(\lambda^2). \end{aligned} \quad (5.30)$$

For  $\lambda = 0$ , these equations have two solutions  $\tau = 0, b = \pm \sigma_1$  and since the Jacobian of the functions  $\tau, b - \sigma_1^2 b^{-1}$ , with respect to  $b, \tau$  is  $\neq 0$  at each of these solutions, it follows from the implicit function theorem that there are two solutions of (5.30) analytic in  $\lambda$  at  $\lambda = 0$  and the solutions are given by

$$\tau = \pm i\lambda\sigma_1 + o(\lambda^2)$$

$$b = \pm \sigma_1 + o(\lambda) ,$$

where  $\tau$  is purely imaginary and  $b$  is real. These two solutions lead to two linearly independent solutions of (5.15). These solutions together with the  $2n - 2$  mentioned previously will form a fundamental system of bounded AC solutions of (5.15) and Theorem (5.4) is proved.

#### §6. ZONES OF INSTABILITY FOR SYSTEMS OF MATHIEU TYPE EQUATIONS

Consider the equation

$$(6.1) \quad x_j'' + \sigma_j^2 x_j + 4\varepsilon(\cos 2t) \sum_{k=1}^n \sigma_k d_{jk} x_k = 0, \quad j = 1, 2, \dots, n ,$$

where  $\varepsilon$  is a real parameter and  $\sigma_j, d_{jk}, j, k = 1, 2, \dots, n$ , are real constants. If  $\tau_1, \dots, \tau_{2n}$  are the characteristic exponents of (6.1) and  $\rho_j = e^{\pi \tau_j}$ ,  $j = 1, 2, \dots, 2n$ , are the characteristic multipliers, then the  $\rho_j$  are the solutions of a polynomial equation,

$$(6.2) \quad F(\rho) = \prod_{j=1}^n (\rho^2 - 2A_j \rho + B_j) ,$$

where the  $A_j, B_j$  are real numbers and  $\prod_{j=1}^n B_j = 1$ . If  $\sigma_1, \dots, \sigma_n$  are analytic functions of  $\varepsilon$  at  $\varepsilon = 0$ ,

$$(6.3) \quad \sigma_1(0) = m, \quad m \text{ a positive integer} ,$$

$$2\sigma_j(0) \neq 0, \quad \sigma_j \pm \sigma_k \neq 0 \pmod{2}, \quad j \neq k, \quad j, k = 2, 3, \dots, n ,$$

then it follows from Theorem (5.1) that  $2n - 2$  of the characteristic exponents are purely imaginary, say  $\tau_3, \dots, \tau_{2n}$ , and  $\tau_{2j-1}(\varepsilon) = \bar{\tau}_{2j}(\varepsilon)$ ,  $\tau_{2j-1}(0) = i\sigma_j(0)$ ,  $j = 2, 3, \dots, n$ , for  $|\varepsilon|$  sufficiently small. Therefore,  $|\rho_j| = 1$ ,  $j = 3, 4, \dots, 2n$ , and  $B_j = 1$ ,  $|A_j| < 1$ ,  $j = 2, 3, \dots, n$ . Since  $\prod_{j=1}^n B_j = 1$ ,  $B_1 = 1$  and

$$(6.4) \quad F(\rho) = \prod_{j=1}^n (\rho^2 - 2A_j \rho + 1) ,$$

The remaining characteristic multipliers  $\rho_1, \rho_2$  will have absolute value less than one if and only if  $|A_1| < 1$  and have absolute value greater than one if and only if  $|A_1| > 1$ . Consequently, the transition curves in the  $(\sigma_1, \varepsilon)$ -plane from stability to instability will occur when  $A_1 = +1$  or  $A_1 = -1$ . But this implies that  $\rho_1 + \rho_2 = +2$  or  $\rho_1 + \rho_2 = -2$ , i.e., there is a periodic solution of (6.1) of period  $\pi$  or  $2\pi$ . These comments are generalizations of the well known facts for the Mathieu equations [11].

The purpose of the present section is to determine the nature of these transition curves for system (6.1) satisfying condition (6.3).

To obtain these curves transform system (6.1) into an equivalent system [see (3.12)].

$$(6.5) \quad z' = Az + i\varepsilon(e^{2it} + e^{-2it})Dz$$

where  $A = \text{diag}(i\sigma_1, -i\sigma_1, \dots, i\sigma_n, -i\sigma_n)$ , and

$$(6.6) \quad D = \begin{bmatrix} d_{jk} & d_{jk} \\ -d_{jk} & -d_{jk} \end{bmatrix}, j, k = 1, 2, \dots, n.$$

The alternative method of successive approximations is now applied to the auxiliary system

$$(6.7) \quad z' = Bz + i\varepsilon(e^{2it} + e^{-2it})Dz$$

where  $B = \text{diag}(im, -im, i\sigma_2 - i\sigma_2, \dots, i\sigma_n, -i\sigma_n)$ ,  $m$  a positive integer, and the zero<sup>th</sup> approximation is given by

$$(6.8) \quad z^{(0)} = x^{(0)} = (e^{imt}, be^{-imt}, 0, \dots, 0)$$

and  $b$  is a real number. From Lemma (4.2), we know that the functions  $V_j$  obtained from the successive approximations are purely imaginary. Consequently, the determining equations for  $\varepsilon \neq 0$  are equivalent to the equations

$$(6.9) \quad \begin{aligned} m - \varepsilon I [V_1(m, \sigma_1, b, \varepsilon)] &= \sigma_1 \\ I [V_1(m, \sigma_1, b, \varepsilon)] + I [V_2(m, \sigma_1, b, \varepsilon)] &= 0. \end{aligned}$$

Since  $m$  is fixed, equations (6.9) may be used to determine  $\sigma_1$  and  $b$  as functions of  $\varepsilon$  for  $|\varepsilon|$  sufficiently small and the corresponding

function  $\sigma_1 = \sigma_1(\varepsilon)$  will be a transition curve in the  $(\sigma_1, \varepsilon)$ -plane near the point  $\varepsilon = 0$ ,  $\sigma_1 = m$ . Therefore, the only thing that remains to be done is to find the first nonzero terms of  $V_1$  and  $V_2$  containing  $b$  and solve equations (6.9).

First, it is clear that  $x_j^{(\ell)}(t)$ ,  $j = 1, 2, \dots, 2n$ , for all  $\ell$ , contains only terms of the form  $e^{1(m+2p)t}$ ,  $e^{-1(m+2p)t}$ ,  $p = 0, \pm 1, \dots, \pm \ell$ . Let

$$x_j^{(\ell)}(t) = \sum_{p=-\ell}^{+\ell} [a_{p,j}^{(\ell)} e^{1(m+2p)t} + c_{p,j}^{(\ell)} e^{-1(m+2p)t}] , \quad (6.10)$$

$$j = 1, 2, \dots, 2n ,$$

for all  $\ell$ , where  $a_{p,j}^{(\ell)}$ ,  $c_{p,j}^{(\ell)}$  are constants.

We now show by induction that

$$c_{p,2j}^{(\ell)} = ba_{p,2j-1}^{(\ell)}, \quad c_{p,2j-1}^{(\ell)} = ba_{p,2j}^{(\ell)}, \quad j = 1, 2, \dots, n; \quad (6.11)$$

$$p = 0, \pm 1, \dots, \pm \ell; \quad \ell = 0, 1, 2, \dots, m-1 ,$$

where the constants  $a_{p,2j-1}^{(\ell)}$ ,  $a_{p,2j}^{(\ell)}$  are independent of  $b$ . For  $k = 0$ , the constants  $a_{p,j}^{(0)}$ ,  $c_{p,j}^{(0)}$  satisfy (6.11) for all  $p, j$ . If the constants  $a_{p,j}^{(\ell)}$ ,  $c_{p,j}^{(\ell)}$  satisfy (6.11) for all  $p, j$  and  $\ell = 0, 1, 2, \dots, v-1$ , then the terms  $S_1^{(\ell)}$ ,  $S_2^{(\ell)}$  in (3.6) are given by

$$S_1^{(\ell)} = 1 \sum_{k=1}^n d_{1k} \sum_{p=-1,1} (a_{p,2k-1}^{(\ell-1)} + a_{p,2k}^{(\ell-1)}) , \quad (6.12)$$

$$S_2^{(\ell)} = -ib^{-1} \sum_{k=1}^n d_{1k} \sum_{p=-1,1} (c_{p,2k-1}^{(\ell-1)} + c_{p,2k}^{(\ell-1)}) = -S_1^{(\ell)} ,$$

$$\ell = 1, 2, \dots, v ,$$

if  $v < m$  and  $S_1^{(\ell)}$ ,  $S_2^{(\ell)}$  are independent of  $b$ . From the definition of  $x_j^{(v)}$  in (3.7), it now follows immediately that the coefficients  $a_{p,j}^{(v)}$ ,  $c_{p,j}^{(v)}$  satisfy (6.11) for all  $v < m$ . Therefore, (6.12) is also satisfied for all  $v < m$ .

From (3.6) and (6.11),

$$S_1^{(m)} = 1 \sum_{k=1}^n d_{1k} \left[ \sum_{p=-1,1} (a_{p,2k-1}^{(m-1)} + a_{p,2k}^{(m-1)}) + c_{-(m-1),2k-1}^{(m-1)} + c_{-(m-1),2k}^{(m-1)} \right]$$

(6.13)

$$S_1^{(m)} + S_2^{(m)} = 1(b - b^{-1}) \sum_{k=1}^n d_{1k} \left[ a_{-(m-1),2k-1}^{(m-1)} + a_{-(m-1),2k}^{(m-1)} \right]$$

where this last expression is independent of  $b$ . If we define  $\gamma_m$  by the relation  $S_1^{(m)} + S_2^{(m)} = 1(b - b^{-1})\gamma_m$ , then it is a simple calculation to show that

$$(6.14) \quad \gamma_m = \sum_{\substack{k_1, \dots, k_{m-1}=1 \\ k_0=1}}^n \prod_{j=0}^{m-1} \frac{2d_{k_j k_{j+1}} \sigma_{k_{j+1}}}{(m-2j-2)^2 - \sigma_{k_{j+1}}} \cdot d_{k_{m-1}, 1}$$

where  $\sigma_1 = m$  in this expression.

Consequently, for  $\varepsilon \neq 0$ , the determining equations (6.9) are

$$\sigma_1 = m - \varepsilon I [V_1(m, \sigma_1, b, \varepsilon)]$$

(6.15)

$$\gamma_m(b - b^{-1}) = o(\varepsilon) .$$

If  $\gamma_m \neq 0$  for  $\varepsilon = 0$ , then these two equations have two distinct solutions  $\sigma_1 = m + o(\varepsilon)$ ,  $b = \pm 1 + o(\varepsilon)$ , where the two curves  $\sigma_1 = \sigma_1(\varepsilon)$  coincide up to terms of order  $\varepsilon^{m-1}$ . We have proved the following theorem.

**THEOREM (6.1).** Consider the system of equations (6.1) satisfying (6.3). If  $\gamma_m$  defined by (6.14) is different from zero, then there are unbounded AC solutions of (6.1) in every neighborhood of the point  $(m, 0)$  in the  $(\sigma_1, \varepsilon)$ -plane, where  $m$  is a positive integer. Furthermore, the transition curves in the  $(\sigma_1, \varepsilon)$ -plane from a region of stability to instability coincide up to terms of order  $\varepsilon^{m-1}$ .

COROLLARY (6.1). There are unbounded solutions of the Mathieu equation  $x'' + \sigma^2 x + 4\epsilon(\cos 2t)x = 0$  in every neighborhood of the point  $(m, 0)$  in the  $(\sigma, \epsilon)$ -plane for every positive integer  $m$ .

PROOF. In this case (6.14) implies  $\gamma_m(0) = [-2^{m-1}m!(m-1)!]^{-1} \neq 0$  for every positive integer  $m$ .

COROLLARY (6.2). Consider the system of equations (6.1) satisfying condition (6.3) and suppose that  $d_{jkd_{rs}} \geq 0$  for every  $j, k, r, s$ , at least one  $d_{jk}$  is  $\neq 0$ , and  $\sigma_k > M$ ,  $k = 2, 3, \dots, n$ ,  $M$  a positive integer. There are unbounded solutions of (6.1) in every neighborhood of the point  $(m, 0)$  in the  $(\sigma_1, \epsilon)$ -plane for every positive integer  $m \leq M + 2$ .

PROOF. In this case (6.14) implies  $\gamma_m \neq 0$  if at least one  $d_{jk}$  is  $\neq 0$ .

In case  $\gamma_m = 0$ , then there still may be two distinct solutions to the determining equations (6.9). It should be clear from the proof of relations (6.11) and (6.12) that if the first  $S_j^{(\ell)}$  which depends on  $b$  occurs in the  $m_1^{\text{th}}$  approximation, then relations (6.12) are valid for  $v = m_1 - 1$ , and (6.13), (6.14) will hold for  $m = m_1$  and the determining equations (6.15) for  $m = m_1$  will have two different solutions. Therefore, Theorem (6.1) will hold for  $m = m_1$ . In case  $S_j^{(\ell)}$  is independent of  $b$  for every  $\ell$ , then the determining equations (6.9) will always have only one solution  $\sigma_1 = \sigma_1(\epsilon)$  which is independent of  $b$ . Therefore, there are two linearly independent periodic solutions along this curve and the AC solutions are bounded in a neighborhood of the point  $(m, 0)$  in the  $(\sigma_1, \epsilon)$ -plane.

#### BIBLIOGRAPHY

- [1] BAILEY, H. R. and CESARI, L., Boundedness of solutions of linear differential systems with periodic coefficients, *Archive Rat. Mech. Ana.* 3(1958), 246-271.
- [2] BAILEY, H. R. and GAMBILL, R. A., On stability of periodic solutions of weakly nonlinear differential systems, *Journ. Math. Mech.* 6(1957), 655-668.
- [3] CESARI, L., (a) Sulla stabilita delle soluzioni dei sistemi di equazioni differenziali lineari a coefficienti periodici, *Atti. Accad. Italia, Mem. Cl. Fis. Mat. Nat.* (6) 11(1940), 633-692; (b) Asymptotic behavior and stability problems in ordinary differential equations, *Ergbn. d. Math. N.F.* Heft 16, Springer 1959;



(c) Existence theorems for nonlinear Lipschitzian differential systems and fixed point theorems, this Study.

- [4] CESARI, L. and HALE, J. K., (a) Second order linear differential systems with periodic L-integrable coefficients, Riv. Mat. Univ. Parma 5(1954), 55-61; (b) A new sufficient condition for periodic solutions of weakly nonlinear differential systems, Proc. Amer. Math. Soc. 8(1957), 757-764.
- [5] GAMBILL, R. A., (a) Stability criteria for linear differential systems with periodic coefficients, Riv. Mat. Univ. Parma 5(1954), 169-181; (b) Criteria for parametric instability for linear differential systems with periodic coefficients, Riv. Mat. Univ. Parma. 6(1955), 37-43.
- [6] GAMBILL, R. A. and HALE, J. K., Subharmonic and ultraharmonic solutions for weakly nonlinear differential systems, Journ. Rat. Mech. Ana. 5(1956), 353-394.
- [7] GOLOMB, M., Expansion and boundedness theorems for solutions of linear differential systems with periodic or almost periodic coefficients, Archive Rat. Mech. Anal. 2(1958), 284-308.
- [8] HAACKKE, W., Über die Stabilität eines Systems von gewöhnlichen linearen Differentialgleichungen zweiter Ordnung mit periodischen Koeffizienten, die von Parametern abhängen (I und II) Math. Z. 56(1952), 65-79, (1952), 34-45.
- [9] HAUPT, O., Über lineare homogene Differentialgleichungen zweiter Ordnung mit periodischen Koeffizienten, Math. Ann. 79(1919), 278.
- [10] HALE, J. K., (a) Evaluations concerning products of exponential and periodic functions, Riv. Mat. Univ. Parma, 5(1954), 63-81; (b) On boundedness of the solutions of linear differential systems with periodic coefficients, Riv. Mat. Univ. Parma, 5 (1954), 137-167; (c) Periodic solutions of nonlinear systems of differential equations, Riv. Mat. Univ. Parma, 5(1954), 281-311; (d) Sufficient conditions for the existence of periodic solutions of systems of weakly nonlinear first and second order differential equations, Journ. Math. Mech. 2(1958), 163-172; (e) Linear systems of first and second order differential equations with periodic coefficients, Ill. Journ. Math. 2, 586-591, (1958); (f) A short proof of a boundedness theorem for linear differential systems with periodic coefficients, Archive Rat. Mech. Ana. 2 (1959), 429-434; (g) On the stability of periodic solutions of weakly nonlinear periodic and autonomous differential systems. This Study.
- [11] McLACHLAN, N. W., Theory and application of Mathieu functions, Clarendon Press, Oxford, 1947.
- [12] MEYTLER, E., Allgemeine Theorie der Stabilität erzwungener Schwingungen elastischer Körper, Ing. Arch. 17(1949), 418-449.
- [13] MOSER, J., New aspects in the theory of stability of Hamiltonian systems, Comm. Pure Appl. Math. 9(1958), 81-114.
- [14] GEL'FAND, J. M. and LIDSKII, V. B., On the structure of the regions of stability of linear canonical systems of differential equations with periodic coefficients, Uspehi Mat. Nauk (N.S.), 10(1955), 3-40; Am. Math. Soc. Trans. (2) 8(1958), 143-182.



V. ON THE STABILITY OF PERIODIC SOLUTIONS OF WEAKLY  
NONLINEAR PERIODIC AND AUTONOMOUS  
DIFFERENTIAL SYSTEMS

J. K. Hale

§1. INTRODUCTION

Consider the system of differential equations

$$(1.1) \quad x'' + Ax = \varepsilon f(x, x', \varepsilon, t)$$

where  $\varepsilon$  is a small real parameter,  $x = (x_1, \dots, x_n)$ ,  $A = \text{diag}(\sigma_1^2, \dots, \sigma_n^2)$ , and the vector  $f = (f_1, \dots, f_n)$  is periodic in  $t$  of period  $T = 2\pi/\omega$ . For  $\varepsilon$  small, and certain conditions of smoothness on  $f$ , various methods have been given for the determination of periodic solutions,  $x_0(\varepsilon, t)$  of (1.1) of the form

$$(1.2) \quad \begin{aligned} x_{j0}(0, t) &= a_j \cos(r_j \omega t + \varphi_j), \quad j = 1, 2, \dots, \mu, \\ x_{j0}(0, t) &= 0, \quad j = \mu + 1, \dots, n, \end{aligned}$$

where each  $a_j, \varphi_j$  is a real number and each  $r_j = k_j/m_j$  is a rational number,  $j = 1, 2, \dots, \mu$  (for example, see [4a, Section (8.5)] and [4b]. If the partial derivatives of  $f_j$  with respect to  $x_k, x'_k$  exist, the standard procedure for determining whether or not such a solution is stable is to discuss the solutions of the linear variational equation

$$(1.3) \quad y'' + Ay = \varepsilon \frac{\partial f(x_0, x'_0, \varepsilon, t)}{\partial x} y + \varepsilon \frac{\partial f(x_0, x'_0, \varepsilon, t)}{\partial x'} y',$$

where  $\partial f/\partial x, \partial f/\partial x'$  are matrices defined by  $\partial f/\partial x = (\partial f_j/\partial x_k)$ ,  $\partial f/\partial x' = (\partial f_j/\partial x'_k)$ ,  $j, k = 1, 2, \dots, n$ . The coefficients in (1.3) are periodic of period  $mT$ ,  $m = m_1 \dots m_\nu$ ,  $T = 2\pi/\omega$ . If all of the characteristic

exponents of system (1.3) have negative real parts, then the solution (1.2) of (1.1) is asymptotically stable for  $\varepsilon$  sufficiently small ([Liapunov, 9]).

From the Floquet theory [5, p. 78], the characteristic exponents of system (1.3) are obtained from the eigenvalues of a  $2n \times 2n$  matrix, which depends upon a fundamental system of solutions of (1.3). Suppose that the coefficients in (1.3) are such that the characteristic exponents are continuous functions of  $\varepsilon$  at  $\varepsilon = 0$ . Some of the properties of the characteristic exponents of (1.3) can be obtained by using the fact that they are continuous functions of  $\varepsilon$  at  $\varepsilon = 0$  and are only determined up to a multiple of  $\omega_1/m$ . In fact, the characteristic exponents can be grouped into sets  $S_1$ , which are disjoint for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , such that each  $S_1$  satisfies one of the following properties: (i)  $S_1$  contains more than one characteristic exponent all of which are equal to zero for  $\varepsilon = 0$ ; (ii)  $S_1$  contains more than one characteristic exponent all of which are equal to a complex or real number  $\neq 0$  for  $\varepsilon = 0$ ; (iii)  $S_1$  contains only one characteristic exponent. Furthermore, if there exists an  $S_1$  which satisfies (ii) there is a set  $S_j$  (which may coincide with  $S_1$ ) whose characteristic exponents are the complex conjugates of those characteristic exponents of  $S_1$ . Even with this more detailed information about the characteristic exponents, it does not seem possible to use the Floquet theory directly to reduce the order of the  $2n \times 2n$  matrix from which the exponents are obtained.

For  $\varepsilon$  sufficiently small and using the method of successive approximations developed by L. Cesari, J. K. Hale and R. A. Gambill in a series of papers (see [4], Section 4.5 and 8.5), it was shown in [8] that the characteristic exponents of (1.3) which belong to any one of the sets  $S_j$  above, say  $S$ , are determined by the eigenvalues of a  $p \times p$  matrix where  $p$  is the number of elements in  $S$ , provided that all of the eigenvalues of this matrix are distinct for  $|\varepsilon|$  sufficiently small.<sup>1</sup> Furthermore, this matrix of order  $p$  can be written explicitly in terms of the coefficients in (1.3). In particular, for  $p = 1$ , this characteristic exponent can be obtained directly from the method. The purpose of the present paper is to illustrate the application of some of the results of [8] (see Lemmas 2.1 and 2.2) to the stability problem. We state some explicit stability theorems in §3, §4 and give a few examples.

In §3, under the assumption that each  $f_j$  in (1.1) is analytic

<sup>1</sup> Added in proofs: Since writing this paper, the author has become aware of some other recent literature on this same problem in which similar results have been obtained (see [13], [14]).

in  $x, x', \varepsilon$  and satisfies (2.2) these results are applied to systems (1.1) (and even more general systems) to determine a number of sufficient conditions for asymptotic stability (see Theorems 3.1, 3.2, 3.3). Using the same method, Theorem 3.1 had already been proved by H. R. Bailey and R. A. Gambill [3] for the case where  $\mu = n$  in (1.1), (1.2). As a corollary of Theorem 3.2 one obtains a result proved by L. Mandelstam and N. Papalexli [10] using another method for the case where  $\mu = n = 1$  in (1.1), (1.2).

In §4, these same methods are applied to the linear variational equations for a periodic solution of an autonomous nonlinear differential system to obtain a number of sufficient conditions for asymptotic orbital stability (see Theorems 4.1, 4.2, 4.3). In his thesis at Purdue University, E. W. Thompson [12] has obtained Theorem 4.2 by first eliminating the zero characteristic root of the linear variational system and applying the method of successive approximations mentioned above; in particular, the results of H. R. Bailey and L. Cesari [3]. Theorem (4.2) is obtained in this paper by the above method in a more straightforward way. A special case of Theorem (4.2) is the following result which seems particularly significant. Consider the system of second order equations

$$(1.4) \quad x_j'' + \sigma_j^2 x_j = \varepsilon f_j(x_1, \dots, x_n, x_1', \dots, x_n'), \quad j = 1, 2, \dots, n,$$

where  $2\sigma_j \neq m\sigma_1$ ,  $\sigma_j \pm \sigma_k \neq m\sigma_1$ ,  $j \neq k$ ,  $j, k = 2, 3, \dots, n$ ;  $m = 0, \pm 1, \pm 2, \dots$  and each  $f_j$  is analytic in  $x_j, x_j'$  for  $|x_j| \leq b, |x_j'| \leq b$ ,  $b > 0$ . and suppose that  $x_{j0}(\varepsilon, t) = x_{j0}(\varepsilon, t + 2\pi/\omega)$ ,  $j = 1, 2, \dots, n$ ;  $\omega = \sigma_1 + o(\varepsilon)$ , is a periodic solution of (1.4) and is analytic in  $\varepsilon$  at  $\varepsilon = 0$  with  $x_{10}(0, t) = a \cos(\sigma_1 t + \varphi)$ ,  $x_{k0}(0, t) = 0$ ,  $k = 2, 3, \dots, n$ , where  $a, \varphi$  are real numbers. If

$$(1.5) \quad \int_0^{2\pi} f_{jx_j'}(a \cos t, 0, \dots, 0, -a \sigma_1 \sin t, 0, \dots, 0) dt < 0,$$

$$j = 1, 2, \dots, n,$$

then this periodic solution of (1.4) is asymptotically orbitally stable in  $[t_0 + \infty)$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ . This result was obtained by A. Andronov and A. Witt [1] for  $n = 2$ , by evaluating the characteristic polynomial of a matrix of order four. The result above for arbitrary  $n$  is proved

by the above method and the properties of the characteristic exponents without evaluating any determinants.

In §3 and §4, the nonlinear functions in the differential equations are assumed to satisfy a certain condition of analyticity. It is shown in §5 that it is sufficient to require that these functions possess continuous second derivatives.

Throughout the present paper, the following notations are used:  $i = \sqrt{-1}$ ;  $\bar{a}$  is the complex conjugate of  $a$ ;  $R(a)$  is the real part of  $a$ ;  $I(a)$  is the imaginary part of  $a$ . Also  $\varepsilon_0$  is used to denote a sufficiently small positive number.

## §2. DEFINITIONS OF STABILITY

Consider a system of differential equations

$$(2.1) \quad x' = q(x, t)$$

where  $q = (q_1, \dots, q_n)$  is a continuous vector function of the real vector  $x = (x_1, \dots, x_n)$  and the time  $t$  and suppose that (2.1) satisfies a uniqueness condition in a region  $R$  of the  $(n+1)$ -dimensional  $(x, t)$  space. A solution  $x^*(t) = x(t, x_0, t_0)$ ,  $t_0 \leq t < \infty$ , of (2.1) with  $x^*(t_0) = x_0$  is said to belong to  $R$  if  $[x^*(t), t]$  is an interior point of  $R$  for every  $t \geq t_0$ . A solution  $x^*(t) = x(t; x_0, t_0)$  is said to be asymptotically stable (to the right) if there exists a  $\delta > 0$  such that (i) every solution  $x(t; x_1, t_0)$  exists for all  $t \geq t_0$  and belongs to  $R$  if  $\|x_1 - x_0\| < \delta$ ; (ii)  $\|x(t; x_1, t_0) - x(t; x_0, t_0)\| \rightarrow 0$  as  $t \rightarrow \infty$  if  $\|x_1 - x_0\| < \delta$ , where  $\|x\| = \sum_{j=1}^n |x_j|$ .

Suppose now that  $q(x, t)$  in (2.1) is periodic in  $t$  and the solution  $x^*(t)$  is also periodic and the second partial derivatives of the functions  $q_j$ ,  $j = 1, 2, \dots, n$ , with respect to  $x_k$ ,  $k = 1, 2, \dots, n$ , in the region  $R$  of  $(x, t)$  space. Then the linear variational equation for the solution  $x^*$  is given by

$$(2.2) \quad y' = Q(t)y, \quad Q(t) = (\partial q_j / \partial x_k)_{x=x^*}, \quad j, k = 1, 2, \dots, n,$$

where  $Q(t)$  is periodic in  $t$ . We need the following known theorem.

**THEOREM 2.1.** If  $q(x, t)$  satisfies the above conditions and the characteristic exponents of (2.2) have negative real parts, then the periodic solution  $x^*(t)$  of (2.1) is asymptotically stable to the right ([Liapunov, 9], see also [5, p. 314]).

If system (2.1) is autonomous, i.e.,  $x' = q(x)$ , then any periodic solution  $x^*(t)$  defines a closed curve  $C$  in the  $n$ -dimensional  $x$ -space,  $E_n$ . If  $d(x, c)$  denotes the distance of a point  $x$  in  $E_n$  from the curve  $C$  in  $E_n$ , then the solution  $x^*(t)$  is said to be asymptotically orbitally stable (to the right) provided, given  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that every solution  $x(t)$  with  $d(x(t_0), C) < \delta$  for some  $t_0$  implies  $d(x(t), C) < \varepsilon$  for  $t \geq t_0$  and  $d(x(t), C) \rightarrow 0$  as  $t \rightarrow +\infty$ .

THEOREM 2.2. If  $x^*(t)$  is a periodic solution of the autonomous system  $x' = q(x)$  where each component of  $q_j$  has continuous second derivatives in a region  $R$  of  $E_n$ , and  $n-1$  of the characteristic exponents of the linear variational equation for  $x^*(t)$  have negative real parts, then the solution  $x^*(t)$  is asymptotically stable to the right. [5, p. 323].

### §3. PERIODIC DIFFERENTIAL SYSTEMS

Consider the periodic system of equations

$$(3.1) \quad \begin{aligned} x_j'' + \alpha_j x_j' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', \varepsilon, t), \quad j = 1, 2, \dots, \mu, \\ x_j' + \beta_j x_j &= \varepsilon f_j(x, x', \varepsilon, t), \quad j = \mu + 1, \dots, n; \quad t_0 \leq t < +\infty, \end{aligned}$$

where  $\varepsilon > 0$  is a real parameter,  $x = (x_1, \dots, x_n)$ ,  $x' = (x_1', \dots, x_n')$ , each  $f_j$  is real and analytic in a neighborhood  $U$  of the origin of the  $(x, x', \varepsilon)$  space; the power series expansion of each  $f_j$  in  $U$  has coefficients periodic in  $t$  of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ , and there exists a function  $\eta(t)$ ,  $L$ -integrable in  $[0, T]$ , such that

$$(3.2) \quad |f_j(x, x', \varepsilon, t)| < \eta(t), \quad j = 1, 2, \dots, n,$$

for all  $(x, x', \varepsilon)$  in  $U$ . The parameters  $\alpha_j, \beta_j, \sigma_j$  are real, analytic functions of  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ ;  $\gamma_j(\varepsilon) = (4\sigma_j^2 - \alpha_j^2)^{\frac{1}{2}} > 0$ ,  $j = 1, 2, \dots, \mu$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ . Furthermore, we define numbers  $\rho_j$ ,  $j = 1, 2, \dots, n + \mu$ , by the relations

$$(3.3) \quad \rho_{2j-1} = \frac{1}{2}(-\alpha_j + i\gamma_j) = \bar{\rho}_{2j}, \quad j = 1, 2, \dots, \mu, \quad \rho_{\mu+j} = -\beta_j,$$

$$j = \mu + 1, \dots, n.$$

The system of periodic differential equations (3.1) satisfying all of the above conditions shall be referred to as system (3.1). Notice the vectors  $x, x'$  used above do not have the same dimension.

Suppose there exists a real periodic solution,  $X(\epsilon, t) = (X_1, \dots, X_n)$ , of system (3.1), analytic in  $\epsilon$  for  $0 \leq \epsilon \leq \epsilon_0$ ,  $\epsilon_0 > 0$  with

$$X_{j0} \equiv X_j(0, t) = a_j \cos(r_j \omega t + \varphi_j), \quad j = 1, 2, \dots, \nu; \quad \nu \leq \mu, \quad (3.4)$$

$$X_{j0} \equiv X_j(0, t) = 0, \quad j = \nu + 1, \dots, n,$$

where  $a_j, \varphi_j$  are real numbers,  $r_j = k_j/m_j$ ,  $k_j, m_j$  relatively prime positive integers,  $j = 1, 2, \dots, \nu$ , and  $X(\epsilon, t + mT) = X(\epsilon, t)$ ,  $T = 2\pi/\omega$ ,  $m = m_1 \dots m_\nu$ . Many methods have been given for obtaining such periodic solutions (for example, see [4a, Section 8.5] and [4b]).  
(3.1),

$$(3.5) \quad x_j = X_j + y_j, \quad j = 1, 2, \dots, n,$$

then the linear variational equation for  $y_j$  is

$$y_j'' + \alpha_j y_j + \sigma_j^2 y_j = \epsilon \sum_{k=1}^n f_{jx_k}^* y_k + \epsilon \sum_{k=1}^{\mu} f_{jx_k^1}^* y_k^1, \quad j = 1, 2, \dots, \mu, \quad (3.6)$$

$$y_j'' + \beta_j y_j = \epsilon \sum_{k=1}^n f_{jx_k}^* y_k + \epsilon \sum_{k=1}^{\mu} f_{jx_k^1}^* y_k^1, \quad j = \mu + 1, \dots, n,$$

where

$$(3.7) \quad f_{jx_k}^* = f_{jx_k}(X, X', \epsilon, t), \quad f_{jx_k^1}^* = f_{jx_k^1}(X, X', \epsilon, t),$$

and  $X = (X_1, \dots, X_n)$ ,  $X' = (X_1', \dots, X_n')$ . Each of the coefficients of  $y_k, y_k^1$  in (3.7) is periodic in  $t$  of period  $mT = 2\pi m/\omega$ .

The transformation of variables

$$(3.8) \quad y_j = (2i\gamma_j)^{-1}(z_{2j-1} + z_{2j}), \quad y_j^1 = (2i\gamma_j)^{-1}(\rho_{2j-1}z_{2j-1} + \rho_{2j}z_{2j}),$$

$$j = 1, 2, \dots, \mu, \quad y_k = z_{\mu+k}, \quad k = \mu + 1, \dots, n,$$



leads to the equivalent system of first order equations

$$(3.9) \quad z' = A(\varepsilon)z + \varepsilon C(t, \varepsilon)z$$

where  $A = \text{diag}(\rho_1, \dots, \rho_{n+\mu})$ , the  $\rho_j$  are defined by (3.3), and

$$C = (c_{jk}), \quad j, k = 1, 2, \dots, n + \mu,$$

$$c_{2j-1, 2k-1} = (2i\gamma_k)^{-1} (f_{jx_k}^* + \rho_{2k-1} f_{jx_k}^*); \quad c_{2j-1, 2k} = -\bar{c}_{2j-1, 2k-1},$$

$$k = 1, 2, \dots, \mu;$$

$$c_{2j-1, \mu+h} = f_{jx_k}^*, \quad h = \mu + 1, \dots, n; \quad c_{2j-1, l} = -c_{2j, l}, \quad l = 1, 2, \dots, n + \mu,$$

$$(3.10) \quad j = 1, 2, \dots, \mu;$$

$$c_{\mu+h, 2k-1} = (2i\gamma_k)^{-1} (f_{hx_k}^* + \rho_{2k-1} f_{hx_k}^*); \quad c_{\mu+h, 2k} = -\bar{c}_{\mu+h, 2k-1},$$

$$k = 1, 2, \dots, \mu;$$

$$c_{\mu+h, \mu+l} = f_{hx_l}^*, \quad l = \mu + 1, \dots, n; \quad h = \mu + 1, \dots, n,$$

and the functions  $f_{jx_k}^*$ ,  $f_{jx_k}^*$  are defined by (3.7).

In the sequel, we need the following known results.

LEMMA 3.1. (J. K. Hale [8, Theorem 2.1]). Consider the system of differential equations (3.9) with  $C(t + mT, \varepsilon) = C(t, \varepsilon)$ ,  $T = 2\pi/\omega$ ,  $m = m_1 \dots m_p$ . Suppose that  $\omega' = \omega/m$ ,

$$\rho_j(0) - \rho_k(0) = m_{jk}\omega' i, \quad m_{jk} \text{ an integer or zero,}$$

$$(3.11) \quad j, k = 1, 2, \dots, p;$$

$$\rho_j(0) \neq \rho_k(0) \pmod{\omega' i}, \quad j = 1, 2, \dots, p; \quad k = p + 1, \dots, n + \mu,$$

and define the  $p \times p$  matrix  $G = H - D$ ,  $D = \text{diag}(d_1, \dots, d_p)$ ,  $H = (h_{jk})$ ,  $j, k = 1, 2, \dots, p$ , where

$$d_j = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\rho_j(\varepsilon) - \rho_j(0)]$$

$$(3.12)$$

$$h_{jk} = (mT)^{-1} \int_0^{mT} e^{i(m_{k1} - m_{j1})\omega' t} c_{jk}(t, 0) dt, \quad j, k = 1, 2, \dots, p.$$

If  $\lambda_0$  is a simple characteristic root of the matrix  $G$  and the corresponding eigenvector has no zero components, there is a characteristic exponent  $\tau(\varepsilon)$  of (3.9) which is analytic in  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ , and

$$(3.13) \quad \tau(\varepsilon) = \rho_1(0) + \varepsilon \lambda_0 + o(\varepsilon^2) .$$

COROLLARY 3.1. Consider the system of differential equations (3.9) with  $C(t + mT, \varepsilon) = C(t, \varepsilon)$ ,  $T = 2\pi/\omega$ ,  $m = m_1 \dots m_\nu$ . If  $\omega' = \omega/m$  and

$$(3.14) \quad \rho_j(0) \not\equiv \rho_k(0) \pmod{\omega'1}, \quad k \neq j, \quad k = 1, 2, \dots, n + \mu ,$$

where  $j$  is a fixed integer,  $1 \leq j \leq n + \mu$ , there is a characteristic exponent of (3.9),  $\tau_j = \tau_j(\varepsilon)$ , analytic in  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ , with

$$(3.15) \quad \tau_j(\varepsilon) = \rho_j(\varepsilon) + \varepsilon T^{-1} \int_0^T c_{jj}(t, 0) dt + o(\varepsilon^2) .$$

LEMMA (3.2) (J. K. Hale, [8, Theorem 3.1]). Consider the system of differential equations (3.9) with  $C(t + mT, \varepsilon) = C(t, \varepsilon)$ ,  $T = 2\pi/\omega$ ,  $m = m_1 \dots m_\nu$ . Suppose  $\omega' = \omega/m$ ,

$$\rho_{2j-1} = \bar{\rho}_{2j}, \quad \rho_{2j-1}(0) \not\equiv \rho_{2j}(0) \pmod{\omega'1},$$

$$\rho_{2j-1}(0) = \rho_{2k-1}(0) = m_{jk}\omega'1, \quad m_{jk} \text{ and integer or zero,}$$

$$(3.16) \quad j, k = 1, 2, \dots, p ;$$

$$\rho_j(0) \not\equiv \rho_k(0) \pmod{\omega'1}, \quad j = 1, 2, \dots, 2p; \quad k = 2p + 1, \dots, n + \mu,$$

and define the  $p \times p$  matrix  $G = H - D$ ,

$D = \text{diag}(d_1, \dots, d_p)$ ,  $H = (h_{jk})$ ,  $j, k = 1, 2, \dots, p$ , where

$$d_j = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\rho_{2j-1}(\varepsilon) - \rho_{2j-1}(0)], \quad j = 1, 2, \dots, p ,$$

$$(3.17)$$

$$h_{jk} = (mT)^{-1} \int_0^{mT} c_{2j-1, 2k-1}(t, 0) e^{i(m_{k1} - m_{j1})\omega't} dt ,$$

$$j, k = 1, 2, \dots, p .$$

If  $\lambda_0$  is a simple characteristic root of the matrix  $G$  and the corresponding eigenvector has no zero components, then there are two characteristic exponents  $\tau, \bar{\tau}$  of (3.9) analytic in  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$  and

$$(3.18) \quad \tau(\varepsilon) = \rho_1(0) + \varepsilon \lambda_0 + o(\varepsilon^2).$$

From the Floquet theory [5, p. 78] the characteristic exponents of system (3.9) are obtained by finding the eigenvalues of a matrix of order  $2n$ . We shall illustrate by means of a few theorems how Lemmas 3.1 and 3.2 can be of assistance in reducing the order of this determinant. More specifically, if there are  $p$  of the characteristic exponents close to the origin in the complex plane, then they will be obtained from the eigenvalues of a determinant of order  $p$ . If the remaining characteristic exponents are isolated for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , then they will be obtainable immediately from Corollary 3.1. If there are say  $2p$  characteristic exponents which are two by two complex conjugate none of which are close to the origin in the complex plane, and  $p$  are equal for  $\varepsilon = 0$ , then they will be obtained from a determinant of order  $p$ .

**THEOREM (3.1).** Consider the system (3.1) and suppose that (3.4) is a periodic solution of (3.1). Suppose  $\varepsilon > 0$ ,  $\alpha_1 = \alpha_2 = \dots = \alpha_q = 0$ ,  $\alpha_j > 0$ ,  $j = q+1, \dots, \mu$ ;  $\beta_j > 0$ ,  $j = \mu+1, \dots, n$ ; for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , and

$$\rho_{2j-1}(0) = i k_j \omega/m_j = \bar{\rho}_{2j}(0), \quad j = 1, 2, \dots, \nu; \quad \nu \leq \mu,$$

$$(3.19) \quad \rho_j(0) \not\equiv \rho_k(0) \pmod{\omega^1 i}, \quad j = 1, 2, \dots, 2\nu; \quad k = 2\nu+1, \dots, n+\mu;$$

$$\rho_j(0) \not\equiv \rho_k(0) \pmod{\omega^1 i}, \quad j \neq k, \quad j, k = 2\nu+1, \dots, n+\mu,$$

where  $\omega^1 = \omega/m$ ,  $m = m_1 \dots m_\nu$ , and the  $\rho_j$  are defined by (3.3). The periodic solution (3.4) of (3.1) is asymptotically stable to the right for  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ , if the following conditions are satisfied:

$$(1) \quad \int_0^{mT} f_{jx_j}(X_0, X_0^1, 0, t) dt < 0, \quad j = \nu+1, \dots, q,$$

where  $X_0 = (X_{10}, \dots, X_{n0})$ ,  $X_0^1 = (X_{10}^1, \dots, X_{\mu 0}^1)$  are defined by (3.4);

$$(11) \quad \text{the eigenvalues } \lambda_1, \dots, \lambda_{2\nu} \text{ of the } 2\nu \times 2\nu \text{ matrix}$$

$G = H - D$ ,  $D = \text{diag}(d_1, \dots, d_{2v})$ ,  $H = (h_{jk})$ ,  
 $j, k = 1, 2, \dots, 2v$ ,

$$d_{2j-1} = i \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\sigma_j(\varepsilon) - \sigma_j(0)], \quad d_{2j} = -d_{2j-1}, \\ j = 1, 2, \dots, v,$$

$$h_{2j-1, 2k-1} = \frac{1}{2\pi T} \int_0^{mT} [(1\sigma_k)^{-1} f_{jx_k} + f_{jx_k'}] e^{i(m_{k1} - m_{j1})\omega' t} dt \\ (3.20)$$

$$h_{2j-1, 2k} = \frac{1}{2\pi T} \int_0^{mT} [(1\sigma_k)^{-1} f_{jx_k} - f_{jx_k'}] e^{-i(m_{k1} + m_{j1})\omega' t} dt$$

$$h_{2j, 2k-1} = \bar{h}_{2j-1, 2k}, \quad h_{2j, 2k} = \bar{h}_{2j-1, 2k-1}, \quad j, k = 1, 2, \dots, v,$$

where the functions under the integrals are evaluated at  $(X_0, X_0', 0, t)$ ,  $\rho_{2j-1}(0) - \rho_{2k-1}(0) = m_{jk}\omega' i$ ,  $m_{jk}$  an integer or zero,  $j, k = 1, 2, \dots, v$ , are distinct, have negative real parts and the corresponding eigenvectors have no zero components.

PROOF. From Corollary 3.1, it follows immediately that the characteristic exponents  $\tau_j(\varepsilon)$ ,  $j = 1, 2, \dots, n + \mu$ ,  $\tau_{2j-1}(0) = \bar{\tau}_{2j}(0) = \frac{1}{2} [-\alpha_j(0) + i\gamma_j(0)]$ ,  $j = q + 1, \dots, \mu$ ;  $\tau_{\mu+j}(0) = -\beta_j(0)$   $j = \mu + 1, \dots, n$ , have negative real parts for  $0 \leq \varepsilon \leq \varepsilon_0$ . Furthermore, from the same corollary and (3.10),

$$R(\tau_{2j-1}) = R(\tau_{2j}) = \varepsilon(2\pi T)^{-1} \int_0^{mT} f_{jx_j}(X_0, X_0', 0, t) dt + o(\varepsilon^2),$$

for  $j = v + 1, \dots, q$ . Therefore, from (i) of the theorem, the real parts of these characteristic exponents are negative for  $0 < \varepsilon \leq \varepsilon_0$ . The other characteristic exponents  $\tau_1, \dots, \tau_{2v}$  are determined from Lemma 3.1 for  $p = 2v$ . The matrix  $G$  of the theorem is precisely the matrix  $G$  of Lemma 3.1. Therefore, from (ii) and (3.13), it follows that these characteristic exponents also have negative real parts for  $0 < \varepsilon \leq \varepsilon_0$ . Theorem 2.1 now implies that Theorem 3.1 is true.

REMARK 3.1. Theorem 3.1 has been previously obtained by H. R. Bailey and R. A. Gambill [3] for the case where  $v = \mu = q = n$ ; i.e., for systems of second order equations,  $x_j'' + \sigma_j^2 x_j = \varepsilon f_j(x_1, \dots, x_n, x_1', \dots, x_n', \varepsilon, t)$

$j = 1, 2, \dots, n$ , where each  $\sigma_j = \sigma_j(\varepsilon)$  satisfies the relation  $\sigma_j(0) = k_j \omega / m_j$ ,  $j = 1, 2, \dots, n$ . An interesting example is also given in [3] for  $v = \mu = q = n = 2$  illustrating the application of this theorem.

REMARK 3.2. If any one of the roots of  $|G - \beta I| = 0$  in (11) have a positive real part or if any of the functions in (1) are positive, then the periodic solution (3.4) of (3.1) is unstable in  $[t_0 + \infty]$  for every  $\varepsilon \neq 0$ .

THEOREM 3.2. Consider the system (3.1) and suppose that (3.4) with  $v = 1$  is a periodic solution of (3.1). Suppose  $\varepsilon > 0$  and the  $\alpha_j, \beta_j, \sigma_j$  satisfy the conditions of Theorem 3.1 with  $v = 1$ ; i.e.,

$$\begin{aligned} \rho_1(0) &= 1 \quad k_1 \omega / m_1 = \bar{\rho}_2(0), \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega' 1} \quad j = 1, 2; \quad k = 3, 4, \dots, n + \mu, \\ (3.21) \quad \rho_j(0) &\neq \rho_k(0) \pmod{\omega' 1} \quad j \neq k, \quad j, k = 3, 4, \dots, n + \mu; \end{aligned}$$

where  $\omega' = \omega / m_1$ . The periodic solution (3.4) of (3.1) is asymptotically stable to the right for  $0 < \varepsilon \leq \varepsilon_0$  if the following conditions are satisfied:

$$(1) \quad \int_0^{m_1 T} f_{jx_j}(X_0, X'_0, 0, t) dt < 0, \quad j = 1, 2, \dots, q;$$

where  $X_0, X'_0$  are defined by (3.4) for  $v = 1$ ;

(11) The numbers  $A, B$  defined by

$$\begin{aligned} (2m_1 T)A &= \int_0^{m_1 T} f_{1x_1} dt, \\ (3.22) \quad (2m_1 T)^2 B &= \left( \int_0^{m_1 T} f_{1x_1} dt \right)^2 + \left[ 2m_1 T d_1 - \sigma_1^{-1}(0) \int_0^{m_1 T} f_{1x_1} dt \right]^2 \\ &\quad - \left[ \sigma_1^{-1}(0) \int_0^{m_1 T} f_{1x_1} \cos 2\sigma_1(0)t dt - \int_0^{m_1 T} f_{1x_1} \sin 2\sigma_1(0)t dt \right]^2 \\ &\quad - \left[ \sigma_1^{-1}(0) \int_0^{m_1 T} f_{1x_1} \sin 2\sigma_1(0)t dt + \int_0^{m_1 T} f_{1x_1} \cos 2\sigma_1(0)t dt \right]^2 \\ d_1 &= \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\sigma_1(\varepsilon) - \sigma_1(0)], \end{aligned}$$

where all of the functions are evaluated at  $(\bar{X}_0, \bar{X}'_0, 0, t)$ , satisfy the property  $A \neq B$ ,  $A^2 \neq B$ ,  $B > 0$ .

REMARK 3.3. This theorem has been proved by H. R. Bailey and R. A. Gambill [3] and L. Mandelstam and N. Papalex [10] for the case where  $q = v = \mu = n = 1$ ; i.e., a second order equation  $x'' + \sigma^2 x = \varepsilon f(x, x', \varepsilon)$ , where  $x, f$  are scalars and  $\sigma = \sigma(\varepsilon)$ ,  $\sigma(0) = km/m$ .

PROOF. The system under consideration in Theorem 3.2 is a special case of the system in Theorem 1 for  $v = 1$ . Condition (1) of Theorem 3.2, for  $j = 2, 3, \dots, q$  implies condition (1) of Theorem 1 is satisfied for  $v = 1$ . For  $v = 1$ , the characteristic equation of the matrix  $G$  in Theorem 3.1 is given by  $\lambda^2 - 2A\lambda + B = 0$ , where  $A, B$  are given in (3.22). The roots of this equation have negative real parts if and only if  $A > 0$ ,  $A^2 \neq B$ ,  $B > 0$  which is assured by condition (1) and (11) of Theorem 3.2. The condition  $A \neq B$  assures that the corresponding eigenvectors of  $G$  have no zero components and Theorem 3.2 is proved.

EXAMPLE. Consider the system of equations

$$x''_1 + x_1 = \varepsilon(1 - x_1^2 - x_2^2)x'_1 + \varepsilon p \cos t \equiv \varepsilon f_1(x_1, x_2, x'_1, x'_2, \varepsilon, t), \quad (3.23)$$

$$x''_2 + \sigma_2^2 x_2 = \varepsilon(1 - x_1^2 - x_2^2)x'_2 \equiv \varepsilon f_2(x_1, x_2, x'_1, x'_2, \varepsilon, t),$$

where  $\varepsilon > 0$ ,  $p < 0$  and  $\sigma_2 > 0$ ,  $\sigma_2 \neq m$ ,  $m$  an integer. Applying the method of successive approximations [6], it is easy to see that (3.23) has a periodic solution  $X_j(\varepsilon, t) = X_j(\varepsilon, t + 2\pi)$ ,  $j = 1, 2$ , analytic in  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ ,

$$(3.24) \quad X_1(0, t) = a \sin t, \quad X_2(0, t) = 0,$$

where  $a > 2$  is the positive solution of the equation  $1 - \frac{(a^2)^2}{2} + \frac{p}{a} = 0$ . Furthermore,

$$\int_0^{2\pi} f_{jX'_j}(a \sin t, 0, a \cos t, 0, 0, t) dt = \pi(2 - a^2) < 0,$$

$j = 1, 2$ , and the numbers  $A, B$  of Theorem 3.2 are given by  $4a = (2 - a^2)$ ,  $16B = (2 - a^2)^2 + 16a^2 - 9a^2/4$  and  $A \neq B$ ,  $A^2 \neq B$ ,  $B > 0$  for every real  $a \neq 0$ . Therefore, from Theorem 3.2, the solution (3.24) of (3.23) is asymptotically stable to the right for  $0 < \varepsilon \leq \varepsilon_0$ .

**THEOREM 3.3.** Consider the system (3.1) and suppose that (3.4) is a periodic solution of (3.1). Suppose  $\varepsilon > 0$ ,  $\alpha_1 = \alpha_2 = \dots = \alpha_q = 0$ ,  $\alpha_j > 0$ ,  $j = q + 1, \dots, \mu$ ,  $\beta_j > 0$ ,  $j = \mu + 1, \dots, n$ , for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , and

$$\rho_{2j-1}(0) = 1 \cdot k_j \omega / m_j = \bar{\rho}_{2j}(0), \quad j = 1, 2, \dots, v; \quad v \leq q;$$

$$\rho_j(0) \neq \rho_k(0) \pmod{\omega'1}, \quad j = 1, 2, \dots, 2v;$$

$$k = 2v + 1, \dots, n + \mu;$$

$$(3.25) \quad \rho_{2j-1}(0) \neq \rho_{2j}(0) \pmod{\omega'1}, \quad \rho_{2j-1}(0) - \rho_{2k-1}(0) = m_{jk} \omega'1,$$

$$m_{jk} \text{ an integer or zero, } j, k = v + 1, \dots, r; \quad v \leq r \leq q;$$

$$\rho_j(0) \neq \rho_k(0) \pmod{\omega'1}, \quad j = 2v + 1, \dots, 2r;$$

$$k = 2r + 1, \dots, n + \mu;$$

$$\rho_j(0) \neq \rho_k(0) \pmod{\omega'1}, \quad j \neq k, \quad j, k = 2r + 1, \dots, n + \mu,$$

where  $\omega' = \omega/m$ ,  $m = m_1 \dots m_v$  and the  $\rho_j$  are defined by (3.3). The periodic solution (3.4) of (3.1) is asymptotically stable to the right,  $0 \leq \varepsilon \leq \varepsilon_0$ , if condition (ii) of Theorem 3.1 is satisfied and, in addition,

$$(3.26) \quad \int_0^{mT} f_{jx'_j}(X_0, X'_0, 0, t) dt < 0, \quad j = r + 1, \dots, q,$$

where  $X_0 = (X_{10}, \dots, X_{n0})$ ,  $X'_0 = (X'_{10}, \dots, X'_{\mu 0})$  are defined by (3.4); and the eigenvalues  $\lambda_1, \dots, \lambda_{r-v}$  of the  $(r-v) \times (r-v)$  matrix  $M = N - P$ ,  $P = \text{diag}(p_1, \dots, p_{r-v})$ ,  $N = (n_{jk})$ ,  $j, k = 1, 2, \dots, r - v$ ,

$$p_j = 1 \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\sigma_{v+j}(\varepsilon) - \sigma_{v+j}(0)], \quad j = 1, 2, \dots, r - v;$$

(3.27)

$$n_{j-v, k-v} = \frac{1}{mT} \int_0^{mT} c_{2j-1, 2k-1}(t, 0) e^{1(m_{k1} - m_{j1})\omega' t} dt,$$

$$j, k = v + 1, \dots, r;$$

where the  $c_{jk}$  are defined by (3.10), are distinct, have negative real parts and the corresponding eigenvectors have no zero components.

PROOF. The proof that the characteristic exponents  $\tau_1, \dots, \tau_{2v}, \tau_{2r+1}, \dots, \tau_{n+\mu}$  have negative real parts is exactly the same as the proof in Theorem 3.1. The remaining characteristic exponents  $\tau_{2v+1}, \dots, \tau_{2r}$  are obtained from Lemma 3.2. The matrix  $M$  of the theorem is precisely the matrix  $G$  of Lemma 3.2 if the equations (3.9) are reordered so that  $A = \text{diag}(\rho_{2v+1}, \dots, \rho_{2r}, \rho_1, \dots, \rho_{2v}, \rho_{2r+1}, \dots, \rho_{n+\mu})$ . Therefore, each of the eigenvalues  $\lambda_{j-v}$  of the matrix  $M$  determines two characteristic exponents  $\tau_{2j-1}, \tau_{2j} = \bar{\tau}_{2j-1}$ , with  $\tau_{2j-1} = 1\rho_j(0) + \varepsilon \lambda_{j-v} + o(\varepsilon)$ ,  $j = v+1, \dots, r$ . The real parts of these characteristic exponents are also negative for  $0 < \varepsilon \leq \varepsilon_0$ , and the theorem follows from Theorem 2.1.

REMARK 3.3. For  $v = 1$  in Theorem 3.3, the condition (ii) referred to can be replaced by condition (ii) of Theorem 3.2.

This theorem should clarify the remark preceding Theorem 3.1. In fact, this theorem is mentioned only to illustrate the procedure for calculating the characteristic exponents of (3.9) which are two by two complex conjugate,  $r - v$  which are equal for  $\varepsilon = 0$  and none of which are close to the origin in the complex plane.

#### §4. AUTONOMOUS DIFFERENTIAL SYSTEMS

Consider the autonomous system of equations

$$(4.1) \quad \begin{aligned} x_j'' + \alpha_j x_j' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', \varepsilon), \quad j = 1, 2, \dots, \mu, \\ x_j' + \beta_j x_j &= \varepsilon f_j(x, x', \varepsilon), \quad j = \mu + 1, \dots, n; \quad t_0 \leq t < +\infty, \end{aligned}$$

where  $\varepsilon > 0$  is a real parameter  $x = (x_1, \dots, x_n)$ ,  $x' = (x_1', \dots, x_\mu')$ , each  $f_j$  is real and analytic in a neighborhood  $U$  of the origin of the  $(x, x', \varepsilon)$  space. The parameters  $\alpha_j, \beta_j, \sigma_j$  are real analytic functions of  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ ;  $\gamma_j(\varepsilon) = (4\sigma_j^2 - \alpha_j^2)^{\frac{1}{2}} > 0$ ,  $j = 1, 2, \dots, \mu$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ . Furthermore, we define numbers  $\rho_j$ ,  $j = 1, 2, \dots, n + \mu$ , by the relations

$$(4.2) \quad \begin{aligned} \rho_{2j-1} &= \frac{1}{2}(-\alpha_j + i\gamma_j) = \bar{\rho}_{2j}, \quad j = 1, 2, \dots, \mu, \quad \rho_{\mu+j} = -\beta_j, \\ j &= \mu + 1, \dots, n. \end{aligned}$$



The system of autonomous differential equations (4.1) satisfying all of the above conditions shall be referred to as system (4.1). Notice that the vectors  $x, x'$  used above do not have the same dimension.

Suppose there exists a real periodic solution,  $X(\varepsilon, t) = (X_1, \dots, X_n)$ , of system (4.1), analytic in  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$  with

$$X_{j0} = X_j(0, t) = a_j \cos(r_j \omega t + \varphi_j), \quad j = 1, 2, \dots, v; \quad v \leq \mu, \quad (4.3)$$

$$X_{j0} = X_j(0, t) = 0, \quad j = v + 1, \dots, n,$$

where  $a_j, \varphi_j$  are real numbers,  $r_j = k_j/m_j$ ,  $k_j, m_j$  relatively prime positive integers  $j = 1, 2, \dots, v$ , and  $X(\varepsilon, t + mT) = X(\varepsilon, t)$ ,  $T = 2\pi/\omega$ ,  $m = m_1 \dots m_v$ . Many methods have been given for obtaining such periodic solutions (for example, see [4a, Section 8.5] and [4b]).

As in the previous section, the linear variational equation associated with the solution  $X(\varepsilon, t)$  of (4.1) is given by

$$(4.4) \quad z' = A(\varepsilon)z + \varepsilon C(t, \varepsilon)z$$

where  $A = \text{diag}(\rho_1, \dots, \rho_{n+\mu})$ , the  $\rho_j$  are defined by (4.2) and the matrix  $C = (c_{jk})$ ,  $j, k = 1, 2, \dots, n + \mu$ , is given by (3.10) with

$$(4.5) \quad f_{jx_k}^* = f_{jx_k}(X, X', \varepsilon), \quad f_{jx_k'}^* = f_{jx_k'}(X, X', \varepsilon)$$

and  $X = (X_1, \dots, X_n)$ ,  $X' = (X_1', \dots, X_n')$ . The matrix  $C(t, \varepsilon)$ , therefore, satisfies the relation  $C(t + mT, \varepsilon) = C(t, \varepsilon)$ .

We wish to apply Lemmas 3.1 and 3.2 to obtain sufficient conditions for the asymptotic orbital stability of the periodic solution  $X(\varepsilon, t)$  of (4.1). Before proceeding to the formal discussion, observe that one expects each of the given numbers  $\sigma_j$  to satisfy a relation of the form  $\sigma_j = r_j \sigma_j + o(\varepsilon)$ ,  $j = 2, 3, \dots, v$ , to obtain a periodic solution of (4.1) of the form (4.3). The number  $\omega$  (also the numbers  $r_j, \varphi_j$ ,  $j = 1, 2, \dots, v$ ) is then determined as a function of  $\varepsilon$  and the numbers  $\sigma_j$  so that (4.3) satisfies (4.1). In particular  $\omega = \sigma_1 + o(\varepsilon)$ . As we know, one could just as well obtain  $\sigma_1$  as a function of  $\omega$  and  $\varepsilon$  simply by making a convenient change of scale in the variable  $t$ . In the present context, it is convenient to assume that the latter alternative is chosen, namely,  $\omega$  is independent of  $\varepsilon$  and the numbers  $\sigma_j$ ,  $j = 1, 2, \dots, v$ , are convenient functions of  $\varepsilon$  so that (4.3) satisfies (4.1).

Another observation to be made is that the linear variational equation (4.4) always has a periodic solution of period  $mT$  [5, p. 322] and, therefore, one of the characteristic exponents (since they are only determined up to a multiple of  $\omega_1/m$ ) must be  $k\omega_1/m$ , where  $k$  is an integer or zero.

**THEOREM 4.1.** Consider the system (4.1) and suppose that (4.3) is a periodic solution of (4.1). Suppose  $\varepsilon > 0$ ,  $\alpha_1 = \alpha_2 = \dots = \alpha_q = 0$ ,  $\alpha_j > 0$ ,  $j = q+1, \dots, \mu$ ;  $\beta_j > 0$ ,  $j = \mu+1, \dots, n$ ; for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ , and

$$(4.6) \quad \begin{aligned} \rho_{2j-1}(0) &= 1 \quad k_j \omega / m_j = \bar{\rho}_{2j}(0), \quad j = 1, 2, \dots, \nu, \quad \nu \leq \mu, \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega'1}, \quad j = 1, 2, \dots, 2\nu; \quad k = 2\nu+1, \dots, n+\mu; \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega'1}, \quad j \neq k, \quad j, k = 2\nu+1, \dots, n+\mu, \end{aligned}$$

where  $\omega' = \omega/m$ ,  $m = m_1 \dots m_\nu$ , and the  $\rho_j$  are defined by (4.2). The periodic solution (4.3) of (4.1) is asymptotically orbitally stable to the right,  $0 < \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ , if the following conditions are satisfied:

$$(1) \quad \int_0^{mT} f_{jx_j}(X_0, X'_0, 0) dt < 0, \quad j = \nu+1, \dots, q,$$

where  $X_0 = (X_{10}, \dots, X_{n0})$ ,  $X'_0 = (X'_{10}, \dots, X'_{\mu 0})$  are defined by (4.3);

$$(11) \quad \begin{aligned} &2\nu - 1 \text{ of the eigenvalues of the } 2\nu \times 2\nu \text{ matrix} \\ &G = H - D, \quad D = \text{diag}(d_1 \dots d_{2\nu}), \quad H = (h_{jk}), \quad j, k = \\ &1, 2, \dots, 2\nu, \end{aligned}$$

$$d_{2j-1} = 1 \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [\sigma_j(\varepsilon) - \sigma_j(0)], \quad d_{2j-1} = -d_{2j}, \quad j = 1, 2, \dots, \nu;$$

$$(4.7) \quad \begin{aligned} h_{2j-1, 2k-1} &= \frac{1}{2mT} \int_0^{mT} [(i\sigma_k)^{-1} f_{jx_k} + f_{jx'_k}] e^{i(m_{k1} - m_{j1})\omega' t} dt, \\ h_{2j-1, 2k} &= \frac{1}{2mT} \int_0^{mT} [(i\sigma_k)^{-1} f_{jx_k} - f_{jx'_k}] e^{i(m_{k1} + m_{j1})\omega' t} dt, \end{aligned}$$

$$h_{2j, 2k-1} = \bar{h}_{2j-1, 2k}, \quad h_{2j, 2k} = \bar{h}_{2j-1, 2k-1}; \quad j, k = 1, 2, \dots, \nu;$$

where the functions under the integrals are evaluated at  $(X_0, X_0', 0)$ ,  $\rho_{2j-1}(0) - \rho_{2k-1}(0) = m_{jk}\omega'1$ ,  $m_{jk}$  an integer or zero,  $j, k = 1, 2, \dots, v$ , are distinct, have negative real parts and the corresponding eigenvectors have no zero components.

PROOF. The proof of this theorem is exactly the same as the proof of Theorem 3.1 if we observe that one of the characteristic exponents of (4.4) being a multiple of  $\omega'1/m$  is equivalent to one of the roots  $\beta$  of the equation  $|G - \beta I| = 0$ , being zero. One then applies Theorem 2.2 rather than Theorem 2.1 to complete the proof.

THEOREM 4.2. Consider the system (4.1) and suppose that (4.3) with  $v = 1$  is a periodic solution of (4.1). If  $\varepsilon > 0$  and the  $\alpha_j, \beta_j, \sigma_j$  satisfy the conditions of Theorem 4.1 with  $v = 1$ , i.e.,

$$(4.8) \quad \begin{aligned} \rho_1(0) &= i\omega = \bar{\rho}_2(0), \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega'1}, \quad j = 1, 2; \quad k = 3, 4, \dots, n + \mu; \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega'1}, \quad j \neq k, \quad j, k = 3, 4, \dots, n + \mu; \end{aligned}$$

the periodic solution (4.3) with  $v = 1$  of (4.1) is asymptotically orbitally stable to the right,  $0 < \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ , if

$$(4.9) \quad \int_0^T f_{jx_j'}(X_0, X_0', 0) dt < 0, \quad j = 1, 2, \dots, q;$$

where  $X_0 = (X_{10}, \dots, X_{n0})$ ,  $X_0' = (X_{10}', \dots, X_{n0}')$ ,  $X_{10} = a \cos(\omega t + \varphi)$ ,  $X_{k0} = 0$ ,  $k = 2, 3, \dots, n$ ;  $a, \varphi$  real numbers.

REMARK 4.1. In his thesis at Purdue University, E. W. Thompson [12] has obtained this same result by another method. The method used by E. Thompson was to first eliminate the zero root of the variational equations and then apply some known results of H. R. Bailey and L. Cesari [2]. Also, A. Andronov and A. Witt [1], (see also [11, p. 153]) have obtained this result for the case  $v = 1$ ,  $q = \mu = n = 2$ ; i.e., for a system of two second order equations with no "large" damping terms. The method used by A. Andronov and A. Witt was to evaluate the characteristic polynomial of a matrix of order four. As we shall see in the proof of this

theorem, no determinants are required.

PROOF. Exactly as in the proof of Theorem 4.1, we apply Corollary 3.1 to obtain the real parts of the characteristic exponents,  $\tau_{2j-1} = \bar{\tau}_{2j}$ ,  $j = 2, 3, \dots, \mu$ ;  $\tau_{\mu+j}$ ,  $j = \mu + 1, \dots, n$  as

$$R(\tau_{2j-1}) = -\frac{\alpha_j}{2} + \frac{\varepsilon}{2T} \int_0^T f_{jx_j^1}(X_0, X_0^1, 0) dt + o(\varepsilon^2), \quad j = 1, 2, \dots, \mu, \quad (4.10)$$

$$R(\tau_{\mu+j}) = -\beta_j + \frac{\varepsilon}{T} \int_0^T f_{jx_j}(X_0, X_0^1, 0) dt + o(\varepsilon^2), \quad j = \mu + 1, \dots, n.$$

We may assume that  $\tau_1 = 0$ ,  $\tau_2 = o(\varepsilon)$  and we know [5, p. 81] that

$$T \sum_{j=1}^{n+\mu} \tau_j = \int_0^T \text{tr} (A(\varepsilon) + \varepsilon C(t, \varepsilon)) dt = - \sum_{j=1}^{\mu} \alpha_j - \sum_{j=\mu+1}^n \beta_j$$

$$+ \varepsilon \int_0^T \left( \sum_{j=1}^{\mu} f_{jx_j^1} + \sum_{j=\mu+1}^n f_{jx_j} \right) dt.$$

From (4.10) and the fact that  $\tau_1 = 0$ , we have

$$R(\tau_2)T = \varepsilon \int_0^T f_{1x_1^1}(X_0, X_0^1, 0) dt + o(\varepsilon^2),$$

and Theorem 4.2 now follows immediately. One could prove this theorem in another way by evaluating directly the eigenvalues of the  $2 \times 2$  matrix  $G$  in Theorem 4.1.

EXAMPLE. Consider the system of equations

$$x_1'' + \sigma_1^2 x_1 - \varepsilon(1 - x_1^2 - x_2^2)x_1^1 = \varepsilon f(x_1, x_2, x_2^1, \varepsilon) \quad (4.11)$$

$$x_2'' + \sigma_2^2 x_2 - \varepsilon(1 - x_1^2 - x_2^2)x_2^1 = \varepsilon g(x_1, x_1^1, x_2, \varepsilon), \quad \varepsilon > 0,$$

$\sigma_1, \sigma_2$  are analytic functions of  $\varepsilon$  at  $\varepsilon = 0$  with  $\sigma_2(0) = \sqrt{2}\sigma_1(0)$ ,  $f(-x_1, x_2, x_2^1, \varepsilon) = -f(x_1, x_2, x_2^1, \varepsilon)$ ,  $g(x_1, x_1^1, -x_2, \varepsilon) = -g(x_1, x_1^1, x_2, \varepsilon)$  are any analytic functions in a neighborhood  $U$  of the origin in

$(x_1, x_2, x_1', x_2', \varepsilon)$  space. It is known [7, p. 300] that there are two periodic solutions of this equation of the form

$$(i) \quad x_1 = a_1 \sin(t + \varphi) + o(\varepsilon), \quad x_2 = 0, \quad a_1 = 2 + o(\varepsilon), \quad \sigma_1 = 1 + o(\varepsilon)$$

$$(ii) \quad x_1 = 0, \quad x_2 = a_2 \sin(\sqrt{2}t + \varphi) + o(\varepsilon), \quad a_2 = 2 + o(\varepsilon), \quad \sigma_2 = \sqrt{2} + o(\varepsilon)$$

of periods  $2\pi$  and  $\sqrt{2}\pi$ , respectively (we are choosing  $\sigma_1, \sigma_2$  as functions of  $\varepsilon$  rather than choose the period). For (i) conditions (4.9) for  $j = 1, 2$ , are equivalent to

$$\int_0^{2\pi} [1 - 4 \sin^2 t] dt = -2 \int_0^{2\pi} \sin^2 t dt < 0.$$

It follows from Theorem 4.2 that (i) is asymptotically orbitally stable to the right,  $0 < \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ . Similarly, one shows that (ii) is also asymptotically orbitally stable to the right for  $\varepsilon$  sufficiently small.

**THEOREM 4.3.** Consider the system (4.1) and suppose that (4.3) is a periodic solution of (4.1). Suppose  $\varepsilon > 0$ ,  $\alpha_1 = \alpha_2 = \dots = \alpha_q = 0$ ,  $\alpha_j > 0$ ,  $j = q+1, \dots, \mu$ ;  $\beta_j > 0$ ,  $j = \mu+1, \dots, n$ , for all  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ ,

$$(4.12) \quad \begin{aligned} \rho_{2j-1}(0) &= 1 \cdot k_j \omega / m_j = \bar{\rho}_{2j}(0), \quad j = 1, 2, \dots, v; \quad v \leq q; \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega^1}, \quad j = 1, 2, \dots, 2v; \\ &\quad k = 2v+1, \dots, n+\mu; \\ \rho_{2j-1}(0) &\neq \rho_{2j}(0) \pmod{\omega^1}, \quad \rho_{2j-1}(0) - \rho_{2k-1}(0) = m_{jk} \omega^1, \\ &\quad m_{jk} \text{ an integer or zero, } j, k = v+1, \dots, r; \quad v \leq r \leq q; \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega^1}, \quad j = 2v+1, \dots, 2r; \\ &\quad k = 2r+1, \dots, n+\mu; \\ \rho_j(0) &\neq \rho_k(0) \pmod{\omega^1}, \quad j \neq k, \quad j, k = 2r+1, \dots, n+\mu, \end{aligned}$$

where  $\omega^1 = \omega/m$ ,  $m = m_1 \dots m_v$  and the  $\rho_j$  are defined by (4.2). The periodic solution (4.3) of (4.1) is asymptotically orbitally stable to the right,  $0 < \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$  if condition (ii) of Theorem 4.1 is satisfied, and, in addition,

$$(4.13) \quad \int_0^{mT} f_{jx_j} (X_0, X_0^1, 0) dt < 0, \quad j = r+1, \dots, q,$$

where  $X_0 = (X_{10}, \dots, X_{n0})$ ,  $X_0^1 = (X_{10}^1, \dots, X_{\mu 0}^1)$  are defined by (4.3); and the eigenvalues  $\lambda_1, \dots, \lambda_{r-v}$  of the  $(r-v) \times (r-v)$  matrix  $M = N - P$ ,  $P = \text{diag}(p_1, \dots, p_{r-v})$ ,  $N = (n_{jk})$ ,  $j, k = 1, 2, \dots, r-v$ ;

$$(4.14) \quad p_j = 1 \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} \sigma_{v+j}(\varepsilon) - \sigma_{v+j}(0), \quad j = 1, 2, \dots, r-v,$$

$$n_{j-v, k-v} = \frac{1}{mT} \int_0^{mT} c_{2j-1, 2k-1}(t, 0) e^{i(m_{k1} - m_{j1})\omega^1 t} dt,$$

$$j, k = v+1, \dots, r,$$

where the functions  $c_{jk}$  are defined in (3.10) with the  $f^*$  given by (4.5) are distinct, have negative real parts and the corresponding eigenvectors have no zero components.

PROOF. The proof is exactly the same as the proof of Theorem 3.3, replacing everywhere the words Theorem 3.1, Theorem 2.1 by Theorem 4.1 and Theorem 2.2, respectively.

REMARK 4.2. For  $v = 1$  in Theorem 4.3, the condition (ii) referred to can be replaced by

$$\int_0^{mT} f_{1x_1^1} (X_0, X_0^1, 0) dt < 0.$$

EXAMPLE. Consider the system of equations

$$(4.15) \quad \begin{aligned} x_1'' + \sigma_1^2 x_1 &= \varepsilon(1 - x_1^2) x_1^1 + \varepsilon g_1(x_1, x_2, x_2^1, x_3, x_3^1) \equiv \varepsilon f_1(x, x^1) \\ x_2'' + 2x_2 &= \varepsilon(1 - x_1^2 - x_2^2 - x_3^2) x_2^1 + \varepsilon x_1^2 x_3 + \varepsilon g_2(x_1, x_1^1) \equiv \varepsilon f_2(x, x^1) \\ x_3'' + 2x_3 &= \varepsilon(1 - x_1^2 - x_2^2 - x_3^2) x_3^1 + \varepsilon x_1^2 x_2 + \varepsilon g_3(x_1, x_1^1) \equiv \varepsilon f_3(x, x^1) \end{aligned}$$

where  $x = (x_1, x_2, x_3)$ ,  $\varepsilon > 0$ ,  $\sigma_1 = \sigma_1(\varepsilon)$ ,  $\sigma_1(0) = 1$ ,  $g_1(-x_1, x_2, x_2^1, x_3, x_3^1) = g_1(x_1, x_2, x_2^1, x_3, x_3^1)$  and  $g_1, g_2, g_3$  are analytic functions of their arguments for  $|x_j|, |x_j^1| \leq A$ ,  $A > 0$ . Using the method in [7], it is easy to see that there exists  $\sigma_1 = \sigma_1(\varepsilon)$ ,  $a = a(\varepsilon)$  analytic in  $\varepsilon$  for

$0 \leq \varepsilon \leq \varepsilon_0$ ,  $\sigma_1(0) = 1$ ,  $a(0) = 2$  such that there is a periodic solution  $X(\varepsilon, t) = X(\varepsilon, t + 2\pi)$  analytic in  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $X = (X_1, X_2, X_3)$  where

$$(4.14) \quad X_1(0, t) = 2 \sin(t + \varphi), \quad X_2(0, t) = X_3(0, t) = 0,$$

and  $\varphi$  is an arbitrary constant. This solution will be asymptotically orbitally stable if we can show that five of the six characteristic exponents of the linear variational equation associated with  $X(\varepsilon, t)$  have negative real parts. Since this system is a special case of Theorem 4.3 for  $v = 1$ ,  $r = q = \mu = n = 3$ , it follows from Remark 4.2 it is sufficient to show that

$$\int_0^{2\pi} f_{1X_1'} [X(0, t), X'(0, t)] dt < 0$$

and the eigenvalues of the matrix  $M$ ,

$$2\pi M = (m_{jk}), \quad m_{jk} = \int_0^{2\pi} [-12^{-\frac{1}{2}} f_{jX_k}(X(0, t), X'(0, t)) + f_{jX_k'}(X(0, t), X'(0, t))] dt,$$

$j, k = 2, 3$ , have negative real parts. One finds immediately that

$$\int_0^{2\pi} f_{jX_j} dt = -2\pi, \quad j = 1, 2, 3, \quad \int_0^{2\pi} f_{jX_j} dt = 0 = \int_0^{2\pi} f_{jX_k'} dt, \quad \int_0^{2\pi} f_{jX_k} dt = 4\pi,$$

$j \neq k$ ,  $j, k = 2, 3$ , where the functions under the integral sign are evaluated at  $X(0, t)$ ,  $X'(0, t)$ . The characteristic equation for  $M$  is  $\beta^2 + 2\beta + 3 = 0$  and both of the roots of this equation have negative real parts. Also the corresponding eigenvectors have no zero components and the solution (4.14) of (4.13) is asymptotically orbitally stable to the right,  $0 < \varepsilon \leq \varepsilon_0$ ,  $\varepsilon_0 > 0$ .

## §5. GENERALIZATIONS

In the preceding sections, we have assumed that the  $f_j$ ,  $\alpha_j$ ,  $\beta_j$  were analytic functions. This is not necessary but was assumed only to make the presentation simple. The basic Lemmas 3.1 and 3.2 hold under very general conditions on these functions [8], the most severe restriction arising from applying the implicit function theorem to show that the

eigenvalues  $\lambda$  of the matrices  $G$  in these lemmas will yield a characteristic exponent of the form (3.13) or (3.18). One can show that the Lemmas are still valid if we assume only that the functions  $f_j$  have continuous second derivatives with respect to  $x_k, x_k^1, \varepsilon, k = 1, 2, \dots, n$ , and the functions  $\alpha_j, \beta_j, \sigma_j^2$  have continuous second derivatives with respect to  $\varepsilon$  [8]. Therefore, the stability criteria in Sections 3 and 4 hold for this more general situation. For the existence of periodic solutions of systems (3.1), (4.1) under these and even weaker hypotheses, see L. Cesari [4b].

Another assumption that has been made in all of the theorems is  $\beta_j(\varepsilon) > 0, j = \mu + 1, \dots, n, 0 \leq \varepsilon \leq \varepsilon_0$ . This restriction was introduced only to simplify the notation. In fact, if one of the  $\beta_j = 0$ , say  $\beta_{\mu+1} = 0$ , then the periodic solution (3.4) or (4.3) may have  $X_{\mu+1}(0, t) \neq 0$ . One then has one additional characteristic exponent which is close to the origin in the complex plane. There is a generalization of Lemma 3.1 [8] which will yield sufficient conditions for stability in these cases.

#### BIBLIOGRAPHY

- [1] ANDRONOV, A. and WITT, A., On the mathematical theory of self-oscillatory systems with two degrees of freedom, J. Technical Physics (USSR), (Russian) 4(1933), 249-271.
- [2] BAILEY, H. R. and CESARI, L., Boundedness of solutions for linear differential systems with periodic coefficients, Archive Rat. Mech. Ana. 3(1958), 246-271.
- [3] BAILEY, H. R. and GAMBILL, R. A., On stability of periodic solutions of weakly nonlinear differential systems, Jour. Math. Mech. 6(1957), 655-668.
- [4] CESARI, L., a) Asymptotic behavior and stability problems in ordinary differential equations, Ergbn. d. Math. N.F. Heft 16, Springer, 1959; b) Existence theorems for nonlinear Lipschitzian differential systems and fixed point theorems. This study.
- [5] CODDINGTON, E. A. and LEVINSON, N., Theory of ordinary differential equations, McGraw-Hill, 1955.
- [6] GAMBILL, R. A. and HALE, J. K., Subharmonic and ultraharmonic solutions for weakly nonlinear differential systems, Jour. Rat. Mech. Ana. 5(1956), 353-394.
- [7] HALE, J. K., Periodic solutions of nonlinear systems of differential equations, Riv. Mat. Univ. Parma 5(1954), 281-311.
- [8] HALE, J. K., On the behavior of the solutions of linear periodic differential systems near resonance points. This study.
- [9] LIAPOUNOV, A., Problème général de la stabilité du mouvement, Annals. Math. Studies No. 17, Princeton, 1947.



- [10] MANDELSTAM, L. and PAPALEXI, N., Über Resonanzerscheinungen bei Frequenzteilung, Zeit. Physik 73(1932), 223.
- [11] MINORSKY, N., Introduction to nonlinear mechanics, J. W. Edwards, 1947.
- [12] THOMPSON, E. W., On stability of periodic solutions of autonomous differential systems. To appear.
- [13] BLEHMAN, I. I., On the stability of periodic solutions of quasi-linear systems with many degrees of freedom (Russian). Dokl. Akad. Nauk. SSSR 104(1955), 809-812; 112(1957), 183-186.
- [14] NOHEL, J. A., Stability of perturbed motions. J. Reine Angew. Math. 1960.



# VI. EXISTENCE THEOREMS FOR PERIODIC SOLUTIONS OF NONLINEAR LIPSCHITZIAN DIFFERENTIAL SYSTEMS AND FIXED POINT THEOREMS

Lamberto Cesari

## INTRODUCTION

In the present paper we consider nonlinear systems of ordinary differential equations in the complex field containing a small parameter and satisfying merely conditions of continuity, or a Lipschitz condition. We prove existence theorems for periodic solutions, and families of periodic solutions. This is done by considering a convenient transformation  $\psi = \mathfrak{z}\phi$  from a space  $\Omega$  of periodic vector functions into itself, and by proving that  $\mathfrak{z}$  transforms a compact convex set  $\Omega_0^* \subset \Omega$  into itself under mere hypotheses of continuity. Thus  $\mathfrak{z}$  has at least a fixed element  $y \in \Omega_0^*$ ,  $y = \mathfrak{z}y$ , and  $y$  is a periodic solution of the given differential system provided a (finite) equation is satisfied (determining equation). The analysis of  $\mathfrak{z}$  and of the determining equation in the case of complex or real systems leads to final and simple existence theorems for periodic solutions (and families of periodic solutions). Under a Lipschitz condition  $\mathfrak{z}$  is a contraction in a closed sphere  $\Omega_0 \subset \Omega$  and thus  $\mathfrak{z}$  has a unique fixed element  $y \in \Omega$ . Under a Lipschitz condition, the method of successive approximations  $\phi^{(k+1)} = \mathfrak{z}\phi^{(k)}$ , as usually associated with  $\mathfrak{z}$ , is uniformly convergent toward the periodic solution  $y$ .

This method of successive approximations, or variants of it, has been already studied and its convergence proved directly by the author, by J. K. Hale, and R. A. Gambill [4b, 8, 9c] in the analytic and in the linear case. In the present paper, therefore, a functional interpretation of it is given, and this assures the convergence of the method under a mere Lipschitz condition. Also, known existence theorems for periodic solutions

---

<sup>†</sup> This research was partially supported by AF Contract No. 49 (638)-382 at RIAS, Baltimore, Maryland.

and families of them are extended, particularly the ones recently proved by J. K. Hale [9 cf ] in the analytic case using the same method. Also, other theorems are extended in various ways. The method just mentioned, or variants of it, has a bearing on two papers of J. K. Hale [9 hi, this volume of the Contributions], one dedicated to the study of the critical frequencies of linear systems with periodic coefficients, and the other to the proof of new and simple criteria for stability of periodic solutions of nonlinear systems. Finally, let it be mentioned that the same method of successive approximations was studied and applied by the author, J. K. Hale, R. A. Gambill, H. R. Bailey, W. R. Fuller in a series of papers listed in the references.<sup>(2)</sup>

All applications and results under consideration belong to the class of the problems of perturbation of linear differential systems. In successive papers I shall deal with straightforward nonlinear differential systems as well as with perturbation problems of nonlinear differential systems.

#### §1. THE CONCEPT OF MEAN VALUE AND OTHER REMARKS

(a). The concept of mean value. Consider the family  $C_\omega$  of all functions  $f(t)$ ,  $-\infty < t < +\infty$ , which are finite sums of functions of the form  $e^{\sigma t}\varphi(t)$ , where  $\sigma$  is any complex number, and  $\varphi(t)$  is any complex-valued function, periodic of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ . Obviously  $C_\omega$  is an additive class. If  $\varphi(t)$  has the Fourier series

$$\varphi(t) \sim \sum_{n=-\infty}^{+\infty} c_n e^{in\omega t},$$

then the series

$$(1.1) \quad f(t) = e^{\sigma t}\varphi(t) \approx \sum_{m=-\infty}^{+\infty} c_m e^{(im\omega + \sigma)t}$$

is said to be the series associated with  $f(t) = e^{\sigma t}\varphi(t) \in C_\omega$ . Note that the decomposition  $e^{\sigma t}\varphi(t)$  is not unique since also  $e^{(\sigma+in\omega)t} \cdot e^{-in\omega t}\varphi(t)$ ,

<sup>2</sup> The present results contain some of those proved by W. R. Fuller [10] in his thesis at Purdue University. For instance, W. R. Fuller proved the convergence of the method of successive approximations in the autonomous case under a Lipschitz condition. Other results of this thesis, (e.g., the applications of the same method to systems containing differences) will appear independently.

$n$  any integer, has the same properties. Nevertheless, the series (1.1) is uniquely determined. The mean value  $\mathcal{M}[f]$  of  $f(t) = e^{\sigma t} \varphi(t)$  is defined by  $\mathcal{M}[f] = c_m$  if  $im\omega + \sigma = 0$  for some  $m$ , by  $\mathcal{M}[f] = 0$  otherwise. For any function  $f(t) \in C_\omega$ , then  $\mathcal{M}[f]$  is defined as an additive linear functional. For periodic functions  $f$ ,  $\mathcal{M}[f]$  is the usual mean value of Fourier series theory.

For vector functions  $f = (f_1, \dots, f_n)$ , all  $f_j \in C_\omega$ , we shall denote by  $\mathcal{M}[f]$  the vector  $(m_1, \dots, m_n)$  with  $m_j = \mathcal{M}[f_j]$ ,  $j = 1, \dots, n$ .

This concept of mean value has been proposed by L. Cesari [4b] and further studied by J. K. Hale [9a]. It was proved first that  $\mathcal{M}[f]$  is uniquely defined in  $C_\omega$ , and that, if  $f = f_1 + \dots + f_N$  is any finite decomposition of  $f$  in  $C_\omega$ , then  $\mathcal{M}[f] = \mathcal{M}[f_1] + \dots + \mathcal{M}[f_N]$ . Also the following simple statement holds:

(1.1). If  $f(t) \in C_\omega$  then any primitive  $F(t)$  of  $f(t)$  belongs to  $C_\omega$  if and only if  $\mathcal{M}[f] = 0$ . In addition, if  $\mathcal{M}[f] = 0$ , there exists one and only one primitive  $F(t) \in C_\omega$  with  $\mathcal{M}[F] = 0$ . We shall denote this primitive by

$$\int f(t) dt$$

[see 4b and 9a].

If  $f(t) = e^{(\alpha+i\beta)t} \varphi(t) \in C_\omega$ ,  $\alpha, \beta$  real,  $\mathcal{M}[f] = 0$ ,

$$\varphi(t) \sim \sum c_m e^{im\omega t},$$

the unique primitive  $F(t)$  of  $f(t)$  of mean value zero is given by

$$(1.2) \quad F(t) = e^{(\alpha+i\beta)t} \Phi(t), \quad \Phi(t) = \sum c_m (\alpha + i\beta + im\omega)^{-1} e^{im\omega t},$$

where in the last series the term with  $\alpha + i\beta + im\omega = 0$  (if any) is omitted, the series is convergent for all  $t$ , and  $\Phi(t)$  is absolutely continuous in each finite interval [see 9a]. The following simple examples may be considered, all with  $\omega = 1$ ,  $\mathcal{M}[f] = 0$ :

$$\int \cos t \, dt = \sin t, \quad \int \sin t \, dt = -\cos t$$

$$\int e^{(\alpha+i\beta)t} dt = (\alpha + i\beta)^{-1} e^{(\alpha+i\beta)t}, \quad \alpha + i\beta \neq 0.$$

REMARK. Though not needed in the sequel the following considerations may be of some interest. Given a function  $f(t) \in C_\omega$  with  $\mathcal{M}[f] = 0$ , it is of interest to know whether its unique primitive  $F(t) \in C_\omega$  with  $\mathcal{M}[F] = 0$  is a definite integral of the form

$$\int_{\xi}^t f(u) du$$

with  $0 \leq \xi < T$ . For functions  $f(t) = e^{(\alpha+i\beta)t} \varphi(t) \in C_\omega$ ,  $\mathcal{M}[f] = \mathcal{M}[\varphi] = 0$ , this is true, for instance, if the function  $\varphi(t)$  above happens to be real, and then we may take for  $\xi$  any zero of  $\varphi(t)$ . In particular this is true for  $\alpha + i\beta = 0$ ,  $f = \varphi$  real. For  $\alpha + i\beta = 0$ ,  $f = \varphi = \varphi_1 + i\varphi_2$ ,  $\varphi_1, \varphi_2$  real,  $\mathcal{M}[f] = \mathcal{M}[\varphi_1] = \mathcal{M}[\varphi_2] = 0$ , there are two points  $0 \leq \xi_1, \xi_2 < T$ , such that

$$F(t) = \int_{\xi_1}^t \varphi_1 du + \int_{\xi_2}^t \varphi_2 du ,$$

and the points  $\xi_1, \xi_2$  may be distinct, as for  $f(t) = e^{it} = \cos t + i \sin t$ ,  $F(t) = -ie^{it} = \sin t - i \cos t$ ,  $\xi_1 = 0$ ,  $\xi_2 = \pi/2$ ,  $\xi_1 \neq \xi_2 \pmod{\pi}$ . If  $f(t) = e^{(\alpha+i\beta)t} \varphi(t) \in C_\omega$ ,  $\mathcal{M}[f] = \mathcal{M}[\varphi] = 0$ , then there is always a decomposition  $f = f_1 + f_2$  and two points  $0 \leq \xi_1, \xi_2 < T$  such that  $f_j \in C_\omega$ ,  $\mathcal{M}[f_j] = 0$ ,

$$\int f_j(t) dt = \int_{\xi_j}^t f_j du ,$$

$j = 1, 2$  [J. K. Hale, 9a, proved this by the use of faltung integrals]. If  $\mathcal{M}[\varphi] \neq 0$  then not even this is true as the example shows:  $f(t) = e^{\sigma t}$ ,  $\mathcal{M}(f) = 0$ ,  $\mathcal{M}(\sigma) \neq 0$ ,

$$F(t) = \sigma^{-1} e^{\sigma t} = \sigma^{-1} \int_{-\infty}^t e^{\sigma u} du ,$$

$\pm$  according as  $\mathcal{M}(\sigma) < 0$ , or  $\mathcal{M}(\sigma) > 0$ . Note that for  $f(t) = e^{\sigma t} \varphi(t) \in C_\omega$ ,  $\sigma \neq 0 \pmod{\omega 1}$ , we have

$$F(t) = (e^{\sigma T} - 1)^{-1} \int_t^{t+T} f(u) du .$$

For  $\sigma = 0 \pmod{\omega 1}$  we have

$$F(t) = T^{-1} \int_t^{t+T} u f(u) du$$

[J. Moser].

(1.11). If  $f(t) = e^{\sigma t} \varphi(t) \in C_\omega$ ,  $\mathcal{M}[f] = 0$ , and  $F(t) \in C_\omega$  is its unique primitive with  $\mathcal{M}[F] = 0$ , if  $V \geq T$  is any constant, then there is a constant  $N$ , depending only on  $\sigma, T, V$ , such that

$$(1.3) \quad |F(t)| \leq N \int_0^T |\varphi(u)| du, \quad 0 \leq t \leq V.$$

If  $\sigma \equiv 0 \pmod{\omega 1}$  we may take  $N = 2$ , if  $|\sigma + im\omega| \geq \delta$  for all  $m = 0, \pm 1, \pm 2, \dots$ , and some  $\delta > 0$ , we may take  $N$  depending only on  $\delta, T, V$ . If  $F(t)$  is periodic then (1.3) holds for all  $t$  [J. K. Hale, 9a; see also H. R. Bailey and L. Cesari, 2].

Given  $\sigma = \alpha + i\beta$ ,  $\alpha, \beta$  real, with  $\sigma + im\omega \neq 0$ ,  $m = 0, \pm 1, \pm 2, \dots$ , let  $\delta = \min [|\sigma + im\omega|, m = 0, \pm 1, \dots]$ . For every  $\sigma' = \alpha' + i\beta'$ ,  $\alpha', \beta'$  real, with  $|\sigma' - \sigma| \leq \delta/2$  we have  $|\sigma' + im\omega| \geq \delta/2$  for the same  $m$ .

(1.111). Let  $\sigma = \alpha + i\beta$ ,  $\alpha, \beta$  real, be any complex number with  $\sigma + im\omega \neq 0$ ,  $m = 0, \pm 1, \dots$ , and let  $0 < \delta \leq \min [|\sigma + im\omega|, m = 0, \pm 1, \dots]$ ,  $0 < \delta \leq \omega$ , let  $\sigma'$  be any other number with  $|\sigma' - \sigma| \leq \delta/2$ , let  $\varphi(t)$  be any complex valued function periodic of period  $T = 2\pi/\omega$ ,  $L$ -integrable in  $[0, T]$ , let  $f(t) = e^{\sigma t} \varphi(t)$ ,  $f_1(t) = e^{\sigma' t} \varphi(t)$ , hence  $\mathcal{M}[f] = \mathcal{M}[f_1] = 0$ , and let  $V \geq T$  be any given constant. Then the unique primitives  $F, F_1$  of  $f, f_1$  of class  $C_\omega$  and mean values zero verify the relation

$$(1.5) \quad |F(t) - F_1(t)| \leq |\sigma - \sigma'| N' \int_0^T |\varphi(u)| du, \quad 0 \leq t \leq V,$$

where  $N'$  is a constant depending only on  $\delta, T, V$  ( $\sigma$  bounded).

PROOF. A simple proof can be given by making use of the last lines of the remark above. By taking

$$\alpha = (e^{\sigma T} - 1)^{-1} (e^{\sigma' T} - 1)^{-1},$$

we have

$$\begin{aligned}
 F(t) - F_1(t) &= \int_t^{t+T} [(e^{\sigma T} - 1)^{-1} e^{\sigma u} - (e^{\sigma' T} - 1)^{-1} e^{\sigma' u}] \varphi(u) du = \\
 &= \alpha \int_t^{t+T} [e^{\sigma' T} (1 - e^{(\sigma' - \sigma)(u-T)}) + (e^{(\sigma' - \sigma)u} - 1)] e^{\sigma u} \varphi(u) du.
 \end{aligned}$$

For  $0 \leq t \leq V$ ,  $T \leq V$ , we have also

$$|F(t) - F_1(t)| \leq |\sigma' - \sigma| N \int_0^T |\varphi(u)| du.$$

This proves (1.iii).

(1.iv). If  $A$  is a constant  $n \times n$  matrix with characteristic roots  $\rho_1, \dots, \rho_n$ , if  $f(t) = (f_1, \dots, f_n)$  is a vector function whose components  $f_j$  are periodic of period  $T = 2\pi/\omega$  and  $L$ -integrable in  $[0, T]$ , if  $|\rho_j + im\omega| \geq \delta > 0$  for some  $\delta > 0$ , all  $j = 1, \dots, n$ , and  $m = 0, \pm 1, \pm 2, \dots$ , then the differential system  $y' = Ay + f(t)$ ,  $y = (y_1, \dots, y_n)$ , has exactly one solution  $y(t) = (y_1, \dots, y_n)$  whose components are periodic of period  $T$ . Also, there is a constant  $N$  depending only on  $A$  and  $T$  and not on  $f(t)$  such that

$$(1.6) \quad |y_j(t)| \leq N \sum_{h=1}^n \int_0^T |f_h(t)| dt, \quad j = 1, \dots, n, \quad -\infty < t < +\infty.$$

PROOF. It is not restrictive to suppose that  $A = [a_{jh}]$  is given in triangular form, i.e.,  $a_{jh} = 0$ ,  $j > h$ ,  $a_{jj} = \rho_j$ ,  $j = 1, \dots, n$ . Let  $\rho_j = \alpha_j + i\beta_j$ ,  $\alpha_j, \beta_j$  real,  $j = 1, \dots, n$ ,  $\gamma = \max |\alpha_j|$ ,  $M = e^{\gamma T}$ ,  $Q$  the sum in (1.6),  $a = \max |a_{jh}|$ . The last equation of the system,

$$y_n' = \rho_n y_n + f_n(t),$$

has a unique periodic solution of period  $T = 2\pi/\omega$ , which, by using the notation of this Section, can be written as

$$y_n(t) = e^{\rho_n t} \int e^{-\rho_n u} f_n(u) du,$$

where the integrand is of class  $C_\omega$  and has mean value zero, the integral is the unique primitive of class  $C_\omega$  and mean value zero, has the form



$e^{-\rho_n t} \phi(t)$  considered in (1.2), and hence  $y_n = \phi(t)$  is periodic of period  $T$ . Let  $N$  be the constant of (1.11) depending only on  $\delta$ ,  $T$ , and  $V = T$ . Then by (1.11) for all  $0 \leq t \leq T$ , we have

$$|y_n(t)| \leq MN \int_0^T |f_n(t)| dt \leq MNQ.$$

The last but one equation of the system is

$$y'_{n-1} = \rho_{n-1} y_{n-1} + [a_{n-1,n} y(t) + f_{n-1}(t)],$$

where the expression in brackets is periodic of period  $T$ . Hence this equation has a unique periodic solution given by

$$y_{n-1}(t) = e^{\rho_{n-1} t} \int e^{-\rho_{n-1} u} [a_{n-1,n} y_n(u) + f_{n-1}(u)] du,$$

and we have, for  $0 \leq t \leq T$ ,

$$\begin{aligned} |y_{n-1}(t)| &\leq MN \int_0^T [|a_{n-1,n}| |y_n(u)| + |f_{n-1}(u)|] du \\ &\leq MNQ(a_{MN}T + 1). \end{aligned}$$

By repeating this procedure  $n$  times we find, for  $0 \leq t \leq T$ ,

$$|y_j(t)| \leq MNQ(a_{MN}T + 1)^{n-j}, \quad j = 1, \dots, n,$$

and these relations hold for all  $t$  since  $y_j(t)$  is periodic.

(b). A form of Brouwer's fixed point theorem. For the use in §5 we mention here the following statement which has been shown to be equivalent to the Brouwer fixed point theorem for a cell in the Euclidean space  $E_n$  [C. Miranda, Un'osservazione su un teorema di Brouwer, Boll. Unione Mat. Ital. 3, 1941, 5-7].

(1.v). If  $K \subset E_n$  is a cube whose  $2n$  opposite (closed) faces are  $K_j^I, K_j^{II}$ ,  $j = 1, \dots, n$ , if  $f(x) = (f_1, \dots, f_n)$ ,  $x \in K$ ,  $x = (x_1, \dots, x_n)$ , is a real vector function, continuous in  $K$ , if  $f_j(x)$  has opposite constant signs on the two faces  $K_j^I, K_j^{II}$ , ( $j = 1, \dots, n$ ), then there is at least one point  $x_0 \in K^0$ , such that  $f(x_0) = 0$ , i.e.,  $f_j(x_0) = 0$ ,  $j = 1, \dots, n$ .

An immediate corollary of (1.v) is

(1.vi). If  $K \subset E_n$  is a cube whose  $2n$  opposite (closed) faces are  $K_j^I, K_j^{II}, j = 1, \dots, n$ , if  $M$  is a compact topological space, and  $f(x, m) = (f_1, \dots, f_n), x \in K, m \in M$ , is a real vector function, continuous in  $K \times M$ , if for some  $m_0 \in M, f_j(x, m_0)$  has opposite signs on the two faces  $K_j^I, K_j^{II}, (j = 1, \dots, n)$ , then there is a neighborhood  $N$  of  $m_0$  in  $M$  and, for each  $m \in N$ , at least one  $x_0 = x_0(m), m \in N, x_0 \in K^0$ , such that  $f(x_0(m), m) = 0$  for every  $m \in N$ .

Indeed, by continuity, there is a neighborhood  $N$  of  $m_0$  such that, for every  $m \in N, f_j(x, m)$  has the same constant sign on  $K_j^I, K_j^{II}$  as  $f_j(x, m_0)$ , and thus  $f(x, m)$  has opposite constant signs on  $K_j^I, K_j^{II}, (j = 1, \dots, n)$ .

A slightly different form of (1.vi) is the following one:

(1.vii). If  $K \subset E_n$  is a cube whose  $2n$  opposite (closed) faces are  $K_j^I, K_j^{II}, j = 1, \dots, n$ , if  $M_1, M_2$  are compact topological spaces and  $f(x, m_1, m_2) = (f_1, \dots, f_n), x \in K, m_1 \in M_1, m_2 \in M_2$ , is a real vector function, continuous in  $K \times M_1 \times M_2$ , if for some  $m_{20} \in M_2, f_j(x, m_1, m_{20})$  has opposite constant signs on the two faces  $K_j^I, K_j^{II}, (j = 1, \dots, n)$ , independently from  $m_1 \in M_1$ , then there is a neighborhood  $N_2$  of  $m_{20}$  in  $M_2$  and for each  $m_1 \in M_1$ , and  $m_2 \in N_2$ , at least one  $x_0 = x_0(m_1, m_2), m_1 \in M_1, m_2 \in N_2, x_0 \in K^0$ , such that  $f(x_0(m_1, m_2), m_1, m_2) = 0$  for every  $m_1 \in M_1, m_2 \in N_2$ .

(c). Schauder's fixed point theorem. We shall need in §3 the following form of Schauder's fixed point theorem:

(1.viii). Any continuous mapping  $f : K \rightarrow K$ , from a convex, closed, compact subset  $K$  of a linear space  $M$  has at least one fixed point  $y \in K, fy = y$ .

[See J. Schauder, Der Fixpunkt in Funktionalräumen, *Studia Math.* 2, 1930, 171-180. Also, S. Lefschetz, *Topics in Topology*, *Annals of Math. Studies*, No. 10, 1942.] We do not repeat all definitions, but only the following ones:  $K$  convex means that  $x, y \in K$  implies  $tx + (1-t)y \in K$  for all  $0 \leq t \leq 1$ ;  $K$  closed means that  $M - K$  is open, and hence  $x_n \rightarrow x, x_n \in K, x \in M$ , implies  $x \in K$ ;  $K$  compact means that every sequence  $\{x_n\}, x_n \in K$ , possesses a convergent subsequence.

We shall need in §3 also the following statement:

(1.ix). Any continuous mapping  $f : K \rightarrow K$ , from a complete subspace  $K$  of a metric space  $M$ , which is a contraction in  $K$ , has exactly one fixed point in  $K$  (Banach's fixed point theorem).

If  $d(x, y)$  is the metric in  $M$ , then  $f$  a contraction means that  $d(fx, fy) \leq md(x, y)$  for all  $x, y \in K$ , and a constant  $m < 1$ . That  $K$  is complete means that every Cauchy sequence in  $K$  has its limit

in  $K$ . Thus  $K$  is closed [H. Hahn, *Reelle Funktionen I*, Chelsea 1948, p. 118]. The uniqueness follows by the obvious remark that  $fx = x$ ,  $fy = y$ ,  $x, y \in K$ ,  $x \neq y$ , implies  $0 < d(x, y) = d(fx, fy) \leq md(x, y)$  with  $m < 1$ , a contradiction. The existence follows by proving that any sequence  $\{x_n\}$  with  $x_{n+1} = f(x_n)$ ,  $n = 0, 1, \dots$ ,  $x_0 \in K$  arbitrary, is a Cauchy sequence [Cf. E. A. Coddington and N. Levinson, *Ordinary differential equations*, McGraw-Hill, 1955, p. 41].

## §2. A TRANSFORMATION IN FUNCTIONAL SPACES

We shall consider first, for the sake of simplicity, a differential system of the form

$$(2.1) \quad dy/dt = Ay + \varepsilon q(y, t, \varepsilon), \quad -\infty < t < +\infty,$$

where  $y = (y_1, \dots, y_n)$ ,  $A = A(\varepsilon)$  is a constant real or complex  $n \times n$  matrix,  $\varepsilon \geq 0$  is a real parameter which will be supposed to be sufficiently small,  $q = (q_1, \dots, q_n)$ , and each  $q_j(y, t, \varepsilon)$  is a real or complex function of  $t, y, \varepsilon$ . We shall make on  $A$  and  $q$  the following assumptions:

( $\alpha$ ). There are numbers  $\omega > 0$ ,  $\delta > 0$ ,  $\varepsilon_0 > 0$ , and integers  $0 \leq \nu \leq n$ ,  $a_j \geq 0$ ,  $b_j > 0$ ,  $j = 1, \dots, \nu$ , such that the  $n$  characteristic roots  $\rho_j(\varepsilon)$ ,  $j = 1, \dots, n$ , of  $A$  are continuous functions of  $\varepsilon$  in  $0 \leq \varepsilon \leq \varepsilon_0$ , and verify the relations

$$\rho_j(0) = i a_j \omega / b_j, \quad j = 1, \dots, \nu,$$

(2.2)

$$|\rho_j(0) - i m \omega / b_0| > \delta > 0, \quad j = \nu + 1, \dots, n, \quad b_0 = b_1 \dots b_\nu,$$

$$m = 0, \pm 1, \pm 2, \dots$$

In addition we assume  $A = \text{diag}(A_1, A_2)$ , where  $A_1, A_2$  are  $\nu \times \nu$  and  $(n-\nu) \times (n-\nu)$  matrices,  $A = \text{diag}[\rho_1(\varepsilon), \dots, \rho_\nu(\varepsilon)]$ , and (\*)  $A_2 = \text{diag}[\rho_{\nu+1}(\varepsilon), \dots, \rho_n(\varepsilon)]$ .

Actually, the last requirement (\*) is unnecessarily restrictive, and all of the present results are valid under a much weaker assumption replacing (\*); namely (\*\*)  $B_2$  is any matrix whose coefficients are continuous functions of  $\varepsilon$  in  $0 \leq \varepsilon \leq \varepsilon_0$ . It is only for the sake of simplicity that we use condition (\*) [See remarks at the end of §2]. Also,  $b_0$  in (2.2) need only be any common multiple of  $b_1, \dots, b_\nu$ .

Finally,  $A(\varepsilon)$  could actually be any  $n \times n$  matrix whose elements are continuous functions of  $\varepsilon$  in  $0 \leq \varepsilon \leq \varepsilon_0$ , having a matrix  $A_0$

as above for its canonical form, i.e., provided there exists a matrix  $P(\varepsilon)$  whose coefficients are continuous functions of  $\varepsilon$  in  $0 \leq \varepsilon \leq \varepsilon_0$ , with  $\det P(\varepsilon) \neq 0$  and  $PAP^{-1} = A_0$ .

(K) There exists a number  $R > 0$  and a function  $\psi(t) > 0$ ,  $-\infty < t < \infty$ ,  $L$ -integrable in every finite interval such that  $|y_\ell| \leq R$ ,  $\ell = 1, \dots, n$ ,  $-\infty < t < \infty$ , implies  $|q_j(y, t, \varepsilon)| \leq \psi(t)$ ,  $j = 1, \dots, n$ . Given  $\xi > 0$  there exists  $\varepsilon > 0$  such that  $|y_\ell^1|, |y_\ell^2| \leq R$ ,  $0 \leq \varepsilon^1, \varepsilon^2 \leq \varepsilon_0$ ,  $|y_\ell^1 - y_\ell^2| \leq \xi$ ,  $\ell = 1, \dots, n$ ,  $|\varepsilon^1 - \varepsilon^2| \leq \xi$ , implies

$$|q_j(y^1, t, \varepsilon^1) - q_j(y^2, t, \varepsilon^2)| \leq \xi \psi(t), \quad j = 1, \dots, n, \quad -\infty < t < \infty.$$

Finally, (p) the functions  $q_j(y, t, \varepsilon)$  are periodic in  $t$  of period  $2\pi/\omega$ , or are independent of  $t$  and then  $\psi(t)$  is replaced by a constant  $K > 0$ .

Sometimes we shall replace (K) by the stronger assumption:

(L). Condition (K) holds, and, in addition,  $y_\ell^1, y_\ell^2 \leq R$ ,  $\ell = 1, \dots, n$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , implies

$$|q_j(y^1, t, \varepsilon) - q_j(y^2, t, \varepsilon)| \leq \psi(t) \sum_{\ell=1}^n |y_\ell^1 - y_\ell^2|, \quad j = 1, \dots, n, \quad -\infty < t < \infty.$$

Condition (p) will be replaced by a condition of quasi periodicity in some theorems. Also, it may well occur that we can take  $R = \infty$ , or that we may take  $R$  any arbitrary constant and  $\psi(t)$  depends on  $R$ , all with obvious simplifications in the statements of the present paper.

REMARK 1. Let us give here in a few words the actual meaning of condition ( $\alpha$ ). In most cases (2.1) will be the canonical form of a real system containing at least one equation of the form  $x_1'' + \sigma^2 x_1 = \varepsilon f_1$ , and then  $\rho_1 = i\sigma$ ,  $\rho_2 = -i\sigma$ . Then the first condition (2.2) is satisfied by simply taking  $\nu = 2$ ,  $\omega = \sigma$ ,  $a_1 = 1$ ,  $a_2 = -1$ ,  $b_1 = b_2 = 1$ , while the second condition (2.2) requires that the remaining characteristic roots are not "close" to any one of the numbers  $im\sigma$ ,  $m = 0, \pm 1, \pm 2, \dots$ . If the real system contains, besides the equation above, a first order equation of the form  $x_3' = \varepsilon f_3$ , then we have  $\rho_3 = 0$ , and the first condition (2.2) is satisfied by simply taking  $\nu = 3$ ,  $\omega = \sigma$ ,  $a_1 = 1$ ,  $a_2 = -1$ ,  $a_3 = 0$ ,  $b_1 = b_2 = b_3 = 1$ , while the second condition (2.2) requires again that the remaining characteristic roots are not "close" to any one of the same numbers above. In §5 we shall actually consider these and more general situations. Integers  $a_j, b_j \neq \pm 1$  certainly occur in connection with subharmonic and ultraharmonic solutions [cf. 8]. As a further informal

comment on the general process discussed rigorously below we wish to mention that periodic solutions of system (2.1) can be expected to be of the form  $y_j = c_j e^{i\tau_j t} + o(\varepsilon)$ ,  $j = 1, \dots, v$ ,  $y_j = 0(\varepsilon)$ ,  $j = v+1, \dots, n$ , where  $i\tau_j = \rho_j(0) + o(\varepsilon)$  in the autonomous case, and where the numbers  $c_j = c_j(\varepsilon)$  do not necessarily approach zero as  $\varepsilon \rightarrow 0$ . This is actually the case in innumerable known examples [see, e.g., 8, 9c for references]. Thus it is natural to look for periodic solutions of (2.1) in the "neighborhood" of solutions of (2.1) for  $\varepsilon = 0$  of the form  $y_j = c_j e^{\rho_j(0)t}$ ,  $j = 1, \dots, v$ ,  $y_j = 0$ ,  $j = v+1, \dots, n$ .

Note that, by force of the continuity requirement in condition ( $\alpha$ ), we may assume  $\varepsilon_0 > 0$  and  $\delta > 0$  sufficiently small so as

$$|\rho_j(\varepsilon) - ia_j\omega/b_j| < \delta, \quad j = 1, \dots, v, \quad 0 \leq \varepsilon \leq \varepsilon_0, \quad (2.3)$$

$$|\rho_j(\varepsilon) - im\omega/b_0| > \delta, \quad j = v+1, \dots, n, \quad 0 \leq \varepsilon \leq \varepsilon_0,$$

$$m = 0, \pm 1, \dots$$

Note that  $\omega$  appears in both conditions ( $\alpha$ ) and (K). In case the functions  $q_j$  do not depend on  $t$  (autonomous case), then we may assume that relations (2.3) hold for all  $\omega$  of a sufficiently small neighborhood  $U$  of the number, say  $\omega_0$ , for which (2.2) holds.

Note that the points  $z = \rho_j(0)$ ,  $j = 1, \dots, v$ , are apart from all points  $z = \rho_j(0)$ ,  $j = v+1, \dots, n$ , and  $z = im\omega/b_0$ ,  $m/b_0 \neq a_j/b_j$ ,  $m = 0, \pm 1, \dots$ , in the complex plane. Then we can take  $\delta$  in (2.3) sufficiently small in such a way that (2.3) still holds, and the  $\delta$ -neighborhoods of the points  $z = \rho_j(0)$ ,  $j = 1, \dots, v$ , are  $\delta$  apart from all remaining points.

Let us consider the auxiliary matrices  $B = \text{diag}(B_1, B_2)$ ,  $B_1 = A_1(0) = \text{diag}(i\tau_1, \dots, i\tau_v)$ ,  $i\tau_j = \rho_j(0) = ia_j\omega/b_j$ ,  $j = 1, \dots, v$ , and  $B_2 = A_2(\varepsilon) = (\rho_j(\varepsilon), j = v+1, \dots, n)$ . Thus  $e^{Bt} = \text{diag}(e^{B_1 t}, e^{B_2 t})$ ,  $e^{B_1 t} = \text{diag}(e^{i\tau_1 t}, \dots, e^{i\tau_v t})$ ,  $e^{-B_1 t} = \text{diag}(e^{-i\tau_1 t}, \dots, e^{-i\tau_v t})$ ,  $e^{B_2 t} = \text{diag}(e^{\rho_j(\varepsilon)t}, j = v+1, \dots, n)$ . Let us denote by  $z(t)$  any particular solution of system (2.1) for  $\varepsilon = 0$  of the form

$$(2.4) \quad z(t) = (c_1 e^{i\tau_1 t}, \dots, c_v e^{i\tau_v t}, 0, \dots, 0) = e^{Bt} C,$$

where  $c_j \neq 0$ ,  $j = 1, \dots, v$ , are constants,  $C = \text{col}(c_1, \dots, c_v, 0, \dots, 0)$ . Let  $\Omega = \Omega(c_1, \dots, c_v, b_0, \omega)$  be the class of all continuous periodic vector functions  $\varphi(t) = (\varphi_1, \dots, \varphi_n)$  of period  $T = 2\pi b_0/\omega$ ,

such that

$$(2.5) \quad \mathbb{M} \left[ e^{-1\tau_j t} \varphi_j(t) \right] = c_j, \quad j = 1, \dots, v,$$

that is, the first  $v$  component  $\varphi_j$  of  $\varphi$  have the form

$$(2.6) \quad \varphi_j(t) = e^{1\tau_j t} \left[ c_j + \varphi_j^*(t) \right], \quad j = 1, \dots, v,$$

with  $\mathbb{M}[\varphi_j^*] = 0$ . Let us denote by  $\Omega_R$  the subclass of  $\Omega$  of all  $\varphi = (\varphi_1, \dots, \varphi_n)$  with  $|\varphi_j(t)| \leq R$ ,  $j = 1, \dots, n$ . Let  $\tau$  be the transformation of  $\Omega_R$  into  $\Omega$  defined by  $\psi(t) = (\tau\varphi)(t)$  with

$$(2.7) \quad \psi(t) = z(t) + \varepsilon e^{Bt} \int e^{-Bu} \{q[\varphi(u), u, \varepsilon] - D[\varphi]\varphi(u)\} du,$$

$$(2.8) \quad D[\varphi]C = \mathbb{M} \{e^{-Bu} q[\varphi(u), u, \varepsilon]\},$$

where  $D = D[\varphi] = \text{diag}(d_1, \dots, d_v, 0, \dots, 0)$ . The following remarks are essential:

- (a) As we shall prove, each component of the integrand is of class  $C_\omega$ , with  $\omega' = \omega/b_0$  and mean value zero;
- (b) The integral in (2.7) denote the unique primitive of class  $C_\omega$ , and mean value zero;
- (c) For  $\varphi \in \Omega_R$  we have  $\psi \in \Omega$ , i.e.,  $\tau$  transforms  $\Omega_R$  into  $\Omega$  [see proof below].

In component form relations (2.7), (2.8) become

$$(2.9) \quad \psi_j(t) = c_j e^{1\tau_j t} + \varepsilon e^{1\tau_j t} \int e^{-1\tau_j u} \{q_j[\varphi(u), u, \varepsilon] - d_j \varphi_j(u)\} du, \quad j = 1, \dots, v,$$

$$(2.10) \quad \psi_j(t) = \varepsilon e^{\rho_j t} \int e^{-\rho_j u} q_j[\varphi(u), u, \varepsilon] du, \quad j = v+1, \dots, n,$$

$$(2.11) \quad c_j d_j = \mathbb{M} \{e^{-1\tau_j u} q_j[\varphi(u), u, \varepsilon]\}, \quad j = 1, \dots, v,$$

$$(2.12) \quad 0 = \mathbb{M} \{e^{-\rho_j u} q_j[\varphi(u), u, \varepsilon]\}, \quad j = v+1, \dots, n,$$

and we shall analyze these relations below.

Obviously the products  $e^{-i\tau_j u} q_j(\varphi(u), u, \varepsilon)$ ,  $j = 1, \dots, \nu$ , are periodic of period  $T = 2\pi b_0/\omega$ , and are  $L$ -integrable in  $[0, T]$ . Thus  $\mathfrak{M}$  in (2.11) is the usual mean value, and since  $c_j \neq 0$ ,  $d_j$  is defined,  $j = 1, \dots, \nu$ . By (2.6) we have

$$\begin{aligned} (2.13) \quad & e^{-i\tau_j u} (q_j(\varphi(u), u, \varepsilon) - d_j \varphi_j(u)) = \\ & = (e^{-i\tau_j u} q_j(\varphi(u), u, \varepsilon) - c_j d_j) - d_j \varphi_j^*(u), \end{aligned}$$

and, by (2.11), these functions are periodic of period  $T = 2\pi b_0/\omega$ , of class  $C_\omega$ , and mean value zero. Hence, by §1, there is one and only one primitive, say  $\psi_j^*(t)$ , of (2.13), also periodic and of mean value zero, and finally, by (2.9), we have

$$(2.14) \quad \psi_j(t) = [c_j + \varepsilon \psi_j^*(t)] e^{i\tau_j t}, \quad j = 1, \dots, \nu,$$

that is, the functions  $\psi_j(t)$  satisfy (2.5). Also, the same functions  $\psi_j(t)$  are absolutely continuous. The functions  $e^{-i\rho_j u} q_j(\varphi(u), u, \varepsilon)$ ,  $j = \nu + 1, \dots, n$ , are obviously of class  $C_\omega$ , with  $|- \rho_j + i m \omega / b_0| > \delta > 0$  for all  $j = \nu + 1, \dots, n$ ,  $m = 0, \pm 1, \pm 2, \dots$ , and hence have mean value zero according to §1. Thus (2.12) is justified. Also, there is one and only one primitive of class  $C_\omega$ , and of mean value zero, and of the form  $e^{-i\rho_j t} \psi_{j0}(t)$ ,  $\psi_{j0}(t)$  periodic of period  $T$  ( $\psi_{j0}$  of mean value not necessarily zero), and then  $\psi_j(t) = \varepsilon \psi_{j0}(t)$ ,  $j = \nu + 1, \dots, n$ . Also, the same functions  $\psi_{j0}(t)$  are periodic and absolutely continuous. Thus (a), (b), (c) are completely proved, and  $\mathfrak{z} \cap \mathfrak{R} \subset \mathfrak{n}$ .

We shall take in  $\mathfrak{n}$  the uniform topology (§3) and we shall prove that, for  $\varepsilon$  sufficiently small,  $\mathfrak{z}$  transforms a closed sphere  $\mathfrak{n}_0$  around  $z(t)$ ,  $\mathfrak{n}_0 \subset \mathfrak{R}$ , into itself under the hypotheses of continuity (K). Also, (§3), there is a compact convex set  $\mathfrak{n}_0^* \subset \mathfrak{n}_0 \subset \mathfrak{R}$  which is transformed into itself by  $\mathfrak{z}$ . This implies (§3) that  $\mathfrak{z}$  has in  $\mathfrak{n}_0^*$  at least one fixed element  $y(t)$ ,  $y = \mathfrak{z} y$ , by Schauder's fixed point theorem. Under the Lipschitz condition (L),  $\mathfrak{z}|_{\mathfrak{n}_0}$  is a contraction (§3) and thus the fixed element  $y = \mathfrak{z} y$  is unique in  $\mathfrak{n}_0$ . In any case,  $y(t)$  satisfies the integral equation

$$(2.15) \quad y(t) = z(t) + \varepsilon e^{Bt} \int e^{-Bu} \{q[y(u), u, \varepsilon] - D[y]y(u)\} du,$$

$$D[y]C = \mathfrak{M} \{e^{-Bu} q[y(u), u, \varepsilon]\}.$$

Since both  $y(t)$  and  $D$  depend upon the integers  $a_j \geq 0$ ,  $b_j > 0$ , and the complex constants  $c_j \neq 0$ ,  $j = 1, \dots, v$ , besides  $\omega$  and  $\varepsilon$ , we shall write  $y(t, a, b, c, \omega, \varepsilon)$ ,  $D(a, b, c, \omega, \varepsilon)$ , where  $a = (a_1, \dots, a_v)$ ,  $b = (b_1, \dots, b_v)$ ,  $c = (c_1, \dots, c_v)$ . By (2.15),  $y(t)$  is absolutely continuous, has first derivative  $y'(t)$  a.e., and, by differentiation, we obtain

$$(2.16) \quad y'(t) = (B - \varepsilon D)y(t) + \varepsilon q[y(t), t, \varepsilon] ;$$

in other words,  $y(t)$  satisfies a differential equation analogous to (2.1). The vector function  $y(t)$  will be a solution of (2.1) provided the equation  $B - \varepsilon D = A$  is satisfied, or, in component form,

$$(2.17) \quad ia_j\omega/b_j - \varepsilon d_j(a, b, c, \omega, \varepsilon) = \rho_j(\varepsilon), \quad j = 1, \dots, v .$$

These equations are called the determining equations of system (2.1).

Under the Lipschitz condition (L),  $\tau | \Omega_0$  is a contraction, and the actual determination of the unique fixed element  $y(t) \in \Omega_0$  and of the numbers  $d_1, \dots, d_v$  can be obtained by a method of successive approximations, namely

$$y^{(0)}(t) = z(t) = (c_1 e^{i\tau_1 t}, \dots, e^{i\tau_v t}, 0, \dots, 0) ,$$

$$y^{(m)}(t) = z(t) + \varepsilon e^{Bt} \int e^{-Bu} \{ q[y^{(m-1)}(u), u, \varepsilon] - D^{(m-1)} y^{(m-1)} \} du ,$$

$$D^{(m-1)} C = D[y^{(m-1)}] C = \mathcal{M} \{ e^{-Bu} q[y^{(m-1)}(u), u, \varepsilon] \}, \quad m = 1, 2, \dots,$$

$$C = \text{col}(c_1, \dots, c_v, 0, \dots, 0), \quad D^{(m-1)} = \text{diag}(d_1^{(m-1)}, \dots, d_v^{(m-1)}, 0, \dots, 0),$$

where each integrand is of class  $C_\omega$  and mean value zero, and each integral is the unique primitive of class  $C_\omega$  and mean value zero. The uniform convergence of the process, i.e.,

$$y(t) = \lim y^{(m)}(t) , \quad D[y] = \lim D[y^{(m)}] ,$$

as  $m \rightarrow \infty$ , is a consequence of the fact that  $\tau$  is a contraction in  $\Omega_0$ . This is the method of successive approximations mentioned in the Introduction and used in most papers listed in the references.

REMARK 2. A few informal words on the definition (2.7) of the transformation  $\tau$  may be of interest. First, the term  $D[\varphi](u)$ , which is subtracted under  $\{ \}$  in (2.7) has the effect to make the integrand



of mean value zero in the class  $C_0$ , so as the integral is also in  $C_0$ . Otherwise, there could be constant terms  $c \neq 0$  in the integrand, and terms  $ct$  in the integral (the so called secular terms of classical analysis). Secondly, we actually subtract more than secular terms, but, by this device [cf. Cesari 4b] it is assured that the fixed element  $y(t)$  of  $\mathfrak{z}$ , or the limit  $y(t)$  in the method of successive approximations, satisfies the differential equation (2.16) which is similar to (2.1) and in fact has exactly the same terms  $\varepsilon q$  as (2.1).

### §3. PROOF THAT $\mathfrak{z}$ HAS A FIXED POINT. EXTENSIONS

(a). Proof that  $\mathfrak{z}$  has a fixed point under condition (K). We shall introduce a uniform topology in  $\Omega$  by defining as norm  $\mathfrak{N}[\varphi]$  of an element  $\varphi(t) = (\varphi_1, \dots, \varphi_n)$  of  $\Omega$  ( $\varphi_j$  periodic of period  $T = 2\pi b_0/\omega$  and continuous,  $j = 1, \dots, n$ ), the number  $\mathfrak{N}[\varphi] = \max |\varphi_j(t)|$ , where the maximum is taken with respect to all  $-\infty < t < +\infty$ , and  $j = 1, \dots, n$ . Thus  $\Omega_R$  is the closed set of all  $\varphi \in \Omega$  with  $\mathfrak{N}[\varphi] \leq R$ .

(3.1). Under hypotheses  $(\alpha)$  and  $(K)$ , given  $a = (a_1, \dots, a_v)$ ,  $b = (b_1, \dots, b_v)$ , and any two numbers  $r_1, r_2$ ,  $0 < r_1 < r_2 < R$ , there is an  $\varepsilon_1$ ,  $0 < \varepsilon_1 \leq \varepsilon_0$ , such that, for every  $c = (c_1, \dots, c_v)$  and  $\varepsilon$  with  $r_1 \leq |c_j| \leq r_2$ ,  $j = 1, \dots, v$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ , the transformation  $\mathfrak{z}$  is continuous, maps the closed sphere  $\Omega_0$  into itself,  $\Omega_0 = \{\mathfrak{N}[\varphi(t) - z(t)] \leq r\}$ ,  $r = R - r_2$ , and has at least a fixed point in  $\Omega_0$ .

PROOF. Obviously  $z(t) = (c_1 e^{i\tau_1 t}, \dots, c_v e^{i\tau_v t}, 0, \dots, 0) \in \Omega$ , and  $\mathfrak{N}[z(t)] = \max |c_j| \leq r_2 < R$ . Hence,  $z(t) \in \Omega_R$ . Also, for every  $\varphi(t) \in \Omega_0$  we have  $\mathfrak{N}[\varphi] \leq \mathfrak{N}[z] + r \leq r_2 + r = R$ , and  $\Omega_0 \subset \Omega_R \subset \Omega$ . Let  $K$  be any number such that

$$(3.1) \quad \int_0^T \psi(t) dt \leq KT.$$

For every  $\varphi \in \Omega_0$  and  $j = 1, \dots, v$ , we have, by (2.11),

$$c_j d_j = \mathfrak{N} \left\{ e^{-i\tau_j u} q_j(\varphi(u), u, \varepsilon) \right\} = T^{-1} \int_0^T e^{-i\tau_j u} q_j(\varphi(u), u, \varepsilon) du,$$

(3.2)

$$|d_j| \leq |c_j|^{-1} T^{-1} \int_0^T |q_j(\varphi(u), u, \varepsilon)| du \leq |c_j|^{-1} T^{-1} \int_0^T \psi(u) du \leq r_1^{-1} K.$$

For  $j = 1, \dots, v$ , we have, by (2.4), (2.9),

$$\psi_j(t) - z_j(t) = \varepsilon e^{1\tau_j t} \int e^{-1\tau_j u} \{q_j[\varphi(u), u, \varepsilon] - d_j \varphi_j(u)\} du,$$

and, by (1.11) with  $V = T$ ,  $(K)$ , and (3.2), also

$$\begin{aligned} |\psi_j(t) - z_j(t)| &\leq 2\varepsilon \int_0^T [ |q_j[\varphi(u), u, \varepsilon]| + |d_j| |\varphi_j(u)| ] du \\ (3.3) \quad &\leq 2\varepsilon [KT + Tr_1^{-1}KR] = 2\varepsilon (1 + r_1^{-1}R)KT, \end{aligned}$$

for all  $-\infty < t < +\infty$ , since  $\psi_j, z_j$  are periodic of period  $T$ ,  $j = 1, \dots, v$ . For  $j = v+1, \dots, n$ , we have, by (2.11), (2.10),

$$\psi_j(t) - z_j(t) = \psi_j(t) = e^{\rho_j t} \int e^{-\rho_j u} q_j[\varphi(u), u, \varepsilon] du,$$

with  $|\rho_j + i m \omega / b_0| \geq \delta > 0$ ,  $j = v+1, \dots, n$ ,  $m = 0, \pm 1, \dots$ . Let  $\gamma = \max |\Re \rho_j|$ ,  $j = v+1, \dots, n$ , and  $H = e^{\gamma T}$ . By (1.11) there is a constant  $N = N(\delta, \omega, b)$  such that

$$(3.4) \quad |\psi_j(t) - z_j(t)| \leq \varepsilon H N \int_0^T |q_j[\varphi(u), u, \varepsilon]| du \leq \varepsilon H N K T, \quad j = v+1, \dots, n.$$

By (3.3) and (3.4), for  $M' = \max[2(1 + r_1^{-1}R)KT, HNK T]$ ,  $\varepsilon' = \min[\varepsilon_0, r/M]$ ,  $0 \leq \varepsilon \leq \varepsilon'$ , we have

$$\|\varphi(t) - z(t)\| \leq \varepsilon M' \leq \varepsilon' M' \leq r.$$

Thus  $\varphi \in \Omega_0$  for every  $\varphi \in \Omega_0$ , i.e.,  $\Omega_0 \subset \Omega_0$ .

Let  $L > 0$  denote any number with  $\rho_j(\varepsilon) \leq L$ ,  $j = 1, \dots, n$ . Let us consider the subset  $\Omega_0^* \subset \Omega_0$  of all  $\varphi(t) \in \Omega_0$  whose components  $\varphi_j(t)$  are absolutely continuous, with first derivatives (a.e.) satisfying the relations

$$|\varphi_j'(t)| \leq 2LR + \varepsilon_0 \psi(t), \quad j = 1, \dots, n, \quad -\infty < t < \infty.$$

Obviously  $\Omega_0^*$  is convex. Note that the functions  $\varphi_j(t)$ ,  $j = 1, \dots, n$ , with  $\varphi(t) \in \Omega_0^*$  are equibounded, equiabsolutely continuous, and equicontinuous. Thus, by Ascoli's theorem,  $\Omega_0^*$  is a compact family.

Finally, if  $y^n = (y_{jn}, j = 1, \dots, n) \in \Omega_0^*$ ,  $n = 1, \dots$ , and

$y_{jn}(t) \rightrightarrows y_j(t)$  as  $n \rightarrow \infty$ , the equiabsolute continuity of the functions  $y_{jn}$  implies that  $y_j(t)$  is absolute continuous,  $j = 1, \dots, n$ . If  $v_j, v_{jn}$  denote the total variations of  $y_j, y_{jn}$  in any interval  $[t', t'']$ , by the lower semicontinuity of the total variations we have

$$\int_{t'}^{t''} |y_j'(t)| dt = v_j \leq \liminf_{n \rightarrow \infty} v_{jn} = \liminf_{n \rightarrow \infty} \int_{t'}^{t''} |y_{jn}'(t)| dt.$$

Hence

$$\int_{t'}^{t''} |y_j'(t)| dt \leq \int_{t'}^{t''} [2LR + \varepsilon_0 \psi(t)] dt,$$

and, by a limit, also  $|y_j'(t)| \leq 2LR + \varepsilon_0 \psi(t)$  for almost all  $t$ . Thus  $\Omega_0^*$  is a closed set.

Let us prove that  $\Omega_0^* \subset \Omega_0^*$ . By (2.9), (2.10), by differentiation, we have

$$\psi_j'(t) = \rho_j(0)\psi_j(t) - \varepsilon d_j \varphi_j(t) + \varepsilon q_j[\varphi(t), t, \varepsilon], \quad j = 1, \dots, v,$$

$$\psi_j'(t) = \rho_j(\varepsilon)\psi_j(t) + \varepsilon q_j[\varphi(t), t, \varepsilon], \quad j = v+1, \dots, n,$$

and hence

$$|\psi_j'(t)| \leq LR + \varepsilon r_1^{-1}KR + \varepsilon \psi(t), \quad j = 1, \dots, n,$$

Thus, for  $\varepsilon'' = \min[\varepsilon', r_1 LK^{-1}]$ ,  $0 \leq \varepsilon \leq \varepsilon''$ , we have

$$|\psi_j'(t)| \leq 2LR + \varepsilon_0 \psi(t), \quad j = 1, \dots, n.$$

This proves that  $\psi = \tau \varphi \in \Omega_0^*$  for every  $\varphi \in \Omega_0^*$ , i.e.,  $\tau \Omega_0^* \subset \Omega_0^*$ .

We must now prove that, for every  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon''$ ,  $\tau$  is a continuous transformation in  $\Omega_0$ . Given  $\xi > 0$ , let  $\xi > 0$  be the number defined in (K) relative to  $\xi$ . For every two elements  $\varphi^s = (\varphi_{s1}, \dots, \varphi_{s2}) \in \Omega_0$ ,  $s = 1, 2$ , let  $\psi^s = \tau \varphi^s$ ,  $\psi^s = (\psi_{s1}, \dots, \psi_{sn})$ , and  $D_s = D[\varphi^s] = (d_{s1}, \dots, d_{sv}, 0, \dots, 0)$ . By (3.2) we have  $|d_{sj}| \leq r_1^{-1}K$ ,  $j = 1, \dots, v$ ,  $s = 1, 2$ . We have also, by (K) and (3.1), for  $\|\varphi^1 - \varphi^2\| < \xi$ , and by taking  $\xi \leq \xi$ ,

$$c_j d_{1j} - c_j d_{2j} = e^{-1\tau_j u} \left\{ q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon] \right\},$$

$$(3.5) \quad |d_{1j} - d_{2j}| \leq |c_j|^{-1} T^{-1} \int_0^T |q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon]| du$$

$$\leq |c_j|^{-1} T^{-1} \xi \int_0^T \psi(u) du \leq r_1^{-1} K \xi.$$

For  $j = 1, \dots, v$ , we have now, by (2.4), (2.9),

$$\psi_{1j}(t) - \psi_{2j}(t) = \varepsilon e^{1\tau_j t} \int e^{-1\tau_j u} \left\{ q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon] - d_{1j} \varphi_{1j} + d_{2j} \varphi_{2j} \right\} du,$$

and, by (1.11), (K), (3.1), (3.2), (3.5), also

$$|\psi_{1j}(t) - \psi_{2j}(t)| \leq 2\varepsilon \int_0^T \left\{ |q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon]| + |d_{1j}| |\varphi_{1j}(u) - \varphi_{2j}(u)| + |d_{1j} - d_{2j}| |\varphi_{2j}(u)| \right\} du$$

$$(3.6) \quad \leq 2\varepsilon [KT + r_1^{-1}KT + r_1^{-1}KTR] \xi \leq 2\varepsilon KT(1 + r_1^{-1} + r_1^{-1}R) \xi.$$

For  $j = v + 1, \dots, n$ , we have, by (2.10),

$$\psi_{1j}(t) - \psi_{2j}(t) = \varepsilon e^{\rho_j t} \int e^{-\rho_j u} \left\{ q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon] \right\} du$$

and by (1.11) for the same constants  $H = e^{\gamma T}$ ,  $N = N(\delta, \omega, b)$ , we have

$$|\psi_{1j}(t) - \psi_{2j}(t)| \leq \varepsilon HN \int_0^T |q_j[\varphi^1(u), u, \varepsilon] - q_j[\varphi^2(u), u, \varepsilon]| du$$

$$(3.7) \quad \leq \varepsilon HNK T \xi, \quad j = v + 1, \dots, n.$$

By (3.6), (3.7), for  $M'' = \max [2KT(1 + r_1^{-1} + r_1^{-1}R), HNK T]$  we have

$$\mathfrak{N}[\psi^1 - \psi^2] \leq \varepsilon M'' \xi \leq (\varepsilon_0 M'') \xi,$$

and this assures that  $\tau$  is a continuous transformation in  $\Omega_0$ .

Since  $\tau$  is a continuous transformation and  $\tau$  maps the convex closed compact set  $\Omega_0^*$  into itself, where  $\Omega_0^* \subset \Omega_0 \subset \Omega_R \subset \Omega$ , and  $\Omega$  is linear, by Schauder's fixed point theorem we conclude that  $\tau$  has a fixed element in  $\Omega_0^*$ .

(b). Proof that  $\tau$  is a contraction in  $\Omega_0$  under condition (L).

(3.11). Under hypotheses ( $\alpha$ ) and (L), we can take  $\varepsilon_1$  sufficiently small in (3.1) so that  $\tau|_{\Omega_0}$  is a contraction, namely  $\mathfrak{N}[\tau\varphi^1 - \tau\varphi^2] \leq 1/2 \mathfrak{N}[\varphi^1 - \varphi^2]$  for every  $\varphi^1, \varphi^2 \in \Omega_0$ . Thus, the fixed element  $y(t) \in \Omega_0^* \subset \Omega_0$  is unique in  $\Omega_0$ .

PROOF. We have already proved that  $\tau\Omega_0 \subset \Omega_0$ . We need not consider  $\Omega_0^*$ . Relations (3.1), (3.2), (3.3), (3.4) will be needed. Relation (3.5) can be replaced by

$$\begin{aligned} |d_{1j} - d_{2j}| &\leq |c_j|^{-1} T^{-1} \sum_{\ell=1}^n \int_0^T \psi(u) |\varphi_{1\ell}(u) - \varphi_{2\ell}(u)| du \\ (3.5L) \quad &\leq r_1^{-1} nK \mathfrak{N}[\varphi^1 - \varphi^2], \quad j = 1, \dots, v. \end{aligned}$$

Relation (3.6) can be replaced by

$$\begin{aligned} |\psi_{1j}(t) - \psi_{2j}(t)| &\leq 2\varepsilon [nKT + r_1^{-1}KT + r_1^{-1}nKTR] \mathfrak{N}[\varphi^1 - \varphi^2] \\ (3.6L) \quad &\leq 2\varepsilon nKT[1 + r_1^{-1}(1+R)] \mathfrak{N}[\varphi^1 - \varphi^2], \\ &j = 1, \dots, v. \end{aligned}$$

Relation (3.7) is replaced by

$$(3.7L) \quad |\psi_{1j}(t) - \psi_{2j}(t)| \leq \varepsilon H n n K T \mathfrak{N}[\varphi^1 - \varphi^2], \quad j = v+1, \dots, n.$$

By (3.6L) and (3.7L), for  $M''' = \max [2nKT[1 + r_1^{-1}(1+R)], nHnKT]$ ,  $\varepsilon_1 = \min [\varepsilon'', 1/2M''']$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ , we have

$$\begin{aligned} \mathfrak{N}[\psi^1 - \psi^2] &\leq \varepsilon M''' \mathfrak{N}[\varphi^1 - \varphi^2] \leq \varepsilon_1 M''' \mathfrak{N}[\varphi^1 - \varphi^2] \leq \\ (3.8L) \quad &\leq (1/2) \mathfrak{N}[\varphi^1 - \varphi^2]. \end{aligned}$$

Thus  $z|_{\Omega_0}$  is a contraction and (3.11) is proved.

REMARK 1. If we denote by  $\|D\|$ , or norm of  $D = (d_1, \dots, d_v, 0, \dots, 0)$  the number  $\|D\| = \max |d_j|$ ,  $j = 1, \dots, v$ , and by  $M_1$  the constant  $M = nr_1^{-1}K$ , then, under the same conditions of (3.11), by (3.5L) and (3.8L) and for  $0 \leq \varepsilon \leq \varepsilon_1$  we have

$$(3.9) \quad \begin{aligned} \mathfrak{N}[z\varphi^1 - z\varphi^2] &\leq (1/2) \mathfrak{N}[\varphi^1 - \varphi^2] , \\ \|D[\varphi^1] - D[\varphi^2]\| &\leq M_1 \mathfrak{N}[\varphi^1 - \varphi^2] . \end{aligned}$$

Note that under the present hypotheses, i.e., (L) instead of (K), the existence of a fixed element in  $\Omega_0$  could be deduced by the fact that  $z|_{\Omega_0}$  is a contraction and  $\Omega_0$  is a complete set.

REMARK 2. The fixed element  $y(t) = (y_1, \dots, y_n)$ ,  $y = z y$ , whose existence has been proved above (3.1) satisfies the integral equation

$$(3.10) \quad y(t) = z(t) + \varepsilon e^{Bt} \int e^{-Bu} \{q[y(u), u, \varepsilon] - D[y]y(u)\} du .$$

As noted in §2, the components  $y_j(t)$  are absolutely continuous and thus, by differentiation of (3.10), we conclude that the differential equation (2.16) is satisfied for almost all  $t$ . By (2.6), (2.9), (2.10), relation (3.10) in component form is

$$(3.11) \quad \begin{aligned} y_j(t) &= c_j e^{1\tau_j t} + \varepsilon e^{1\tau_j t} \int e^{-1\tau_j u} \{q_j[y(u), u, \varepsilon] - d_j y_j(u)\} du, \\ &\quad j = 1, \dots, v , \\ y_j(t) &= \varepsilon e^{\rho_j(\varepsilon)t} \int e^{-\rho_j(\varepsilon)u} q_j[y(u), u, \varepsilon] du, \quad j = v+1, \dots, n . \end{aligned}$$

By (3.3), (3.4), (2.4) we know that, for all  $t$ , we have

$$(3.12) \quad \begin{aligned} |y_j(t) - c_j e^{1\tau_j t}| &\leq 2\varepsilon K T (1 + r_1^{-1}R), \quad j = 1, \dots, v , \\ |y_j(t)| &\leq \varepsilon H N K T, \quad j = v+1, \dots, n . \end{aligned}$$

Having already denoted by  $L$  any number such that  $|\rho_j(\varepsilon)| \leq L$ ,  $j = 1, \dots, n$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , by (2.16) and (3.2) we have

$$(3.13) \quad |y_j'(t)| \leq LR + \varepsilon r_1^{-1}KR + \varepsilon \psi(t) .$$

(c). Case of large forcing terms. We shall consider the system analogous to (2.1):

$$dy/dt = Ay + F(t) + \varepsilon q(y, t, \varepsilon) ,$$

where  $F(t) = (F_1, \dots, F_n)$  is a periodic vector function of period  $2\pi/\omega$ , whose components are  $L$ -integrable in  $[0, 2\pi/\omega]$ . For what all previous considerations are concerned we may actually replace (3.14) by the system

$$(3.14) \quad dy/dt = By + F(t) + \varepsilon q(y, t, \varepsilon) .$$

We must suppose that

$$\forall j [e^{-i\tau_j t} F_j(t)] = T^{-1} \int_0^T e^{-i\tau_j t} F_j(t) dt = 0, \quad j = 1, \dots, v .$$

Under this condition, system

$$dy/dt = By + F(t)$$

has a particular solution given by

$$Y(t) = e^{Bt} \int e^{-Bu} F(u) du$$

with the usual conventions. Let  $y = Y(t) + z$ ,  $z = (z_1, \dots, z_n)$ . Then (3.14) is transformed into the system

$$dz/dt = Bz + \varepsilon q[Y(t) + z, t, \varepsilon]$$

which is of the type (2.1).

(d). Extensions.

(d<sub>1</sub>). The considerations above can be repeated for systems much more general than (2.1). For instance, we may consider systems of the form

$$(3.15) \quad dy/dt = Ay + q(y, t, \varepsilon), \quad -\infty < t < +\infty ,$$

$y = (y_1, \dots, y_n)$ ,  $q = (q_1, \dots, q_n)$ , under conditions ( $\alpha$ ) and (K) provided  $q_j(y, t, 0) = 0$ ,  $j = 1, \dots, n$ . Given  $\xi > 0$  there is a  $\varepsilon > 0$  such that

$$(3.16) \quad |q_j(y, t, \varepsilon)| < \xi \psi(t), \quad j = 1, \dots, n ,$$

for all  $|y_j| \leq R$ ,  $-\infty < t < +\infty$ ,  $0 \leq \varepsilon \leq \xi$ .

(d<sub>2</sub>). The considerations above can be repeated for systems of the form

$$(3.17) \quad dy/dt = Ay + q(y, t, \varepsilon) + g(y, t, \varepsilon), \quad -\infty < t < +\infty,$$

under conditions ( $\alpha$ ) and (K) (both  $q$  and  $g$ ) provided  $q$  is as above in (d<sub>1</sub>) and the following holds: given  $\xi > 0$  there is a  $\xi_0 > 0$  such that, for all  $0 \leq \varepsilon \leq \varepsilon_0$ ,  $-\infty < t < +\infty$ ,  $|y_j| \leq R$ , we have

$$(3.18) \quad |g_j(y, t, \varepsilon)| \leq \xi(|y_1| + \dots + |y_n|)\psi(t).$$

In this case we must take  $\Omega_R$  with  $R$  sufficiently small. The last condition is certainly satisfied if  $|g_j(y, t, \varepsilon)| \leq M(|y_1|^2 + \dots + |y_n|^2)$  for some constant  $M$ . Finally the case

$$(3.19) \quad dy/dt = Ay + F(t) + q(y, t, \varepsilon) + g(y, t, \varepsilon),$$

where  $q, g$  are as above and  $F(t)$  is a given periodic function of period  $2\pi/\omega$ , can be reduced to the previous ones by the remark (c) above.

(d<sub>3</sub>). Note that systems of the form

$$dy/dt = A(t)y + \varepsilon q(y, t, \varepsilon),$$

or the similar ones, where  $A(t)$  is a matrix whose elements are continuous periodic functions of  $t$  of period  $2\pi/\omega$ , can be reduced to systems (2.1) by a change of coordinates  $z = Py$  where  $P$  is a matrix whose elements are continuous periodic functions of  $t$ ,  $\det P \neq 0$  for all  $t$ , such that  $PAF^{-1}$  is a constant matrix.

(d<sub>4</sub>). Another important extension concerns the case where  $B_2 = B_2(\varepsilon)$  is an arbitrary  $(n-\nu) \times (n-\nu)$  matrix (§2, condition ( $\alpha$ )) whose elements are continuous functions of  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ , and whose characteristic roots  $\rho_j(\varepsilon)$ ,  $j = \nu + 1, \dots, n$ , verify the relations  $\rho_j(0) \neq im\omega/b_0$ ,  $b_0 = b_1 \dots b_\nu$ ,  $m = 0, \pm 1, \pm 2, \dots$ . Then the coefficients of the characteristic polynomials of  $B_2(\varepsilon)$  are also continuous functions of  $\varepsilon$ , and finally the characteristic roots  $\rho_j(\varepsilon)$  are continuous functions of  $\varepsilon$  (the latter statement can be proved in terms of the theory of functions of one complex variable and the use of the logarithmic indicator. See, e.g., [4a]). Also, it is known that there exists a  $(n-\nu) \times (n-\nu)$  matrix  $P = P(\varepsilon)$  whose elements are continuous functions of  $\varepsilon$  in  $0 \leq \varepsilon \leq \varepsilon_0$  for some  $\varepsilon_0 > 0$ , with  $\det P(\varepsilon) \neq 0$ , and  $PB_2P^{-1}$



is a triangular matrix. See, e.g., for analogous considerations [4a]. See also, S. P. Diliberto, On systems of ordinary differential equations. These Contributions, Vol. 1, 1950, 1-38. Thus we can suppose  $B_2 = B_2(\varepsilon)$  a triangular matrix, say  $B_2 = [b_{jh}]$ ,  $j, h = v+1, \dots, n$ , such that  $b_{jh} = 0$  for  $j > h$ ,  $b_{jj} = \rho_j(\varepsilon)$ . Note that the least  $n - v$  equations (2.1), written in inverse order, are

$$y_n' = \rho_n(\varepsilon)y_n + \varepsilon q_n, \quad ,$$

$$y_{n-1}' = \rho_{n-1}(\varepsilon)y_{n-1} + b_{n-1,n}y_n + \varepsilon q_{n-1}, \quad ,$$

$$y_{n-2}' = \rho_{n-2}(\varepsilon)y_{n-2} + b_{n-2,n-1}y_{n-1} + b_{n-2,n}y_n + \varepsilon q_{n-2}, \quad ,$$

.....

$$y_{v+1}' = \rho_{v+1}(\varepsilon)y_{v+1} + \sum_{l=v+2}^n b_{v+1,l}y_l + \varepsilon q_{v+1}. \quad .$$

The transformation  $\psi = \mathcal{I}\phi$  can now be defined by the same relations (2.9), (2.11) for  $j = 1, \dots, v$ , and, for  $j = v+1, \dots, n$ , by

$$\psi_n(t) = \varepsilon e^{\rho_n t} \int e^{-\rho_n u} q_n[\phi(u), u, \varepsilon] du, \quad ,$$

$$\psi_{n-1}(t) = e^{\rho_{n-1} t} \int e^{-\rho_{n-1} u} \left\{ \varepsilon q_{n-1}[\phi(u), u, \varepsilon] + b_{n-1,n} \psi_n(u) \right\} du, \quad ,$$

.

$$\psi_{n-2}(t) = e^{\rho_{n-2} t} \int e^{-\rho_{n-2} u} \left\{ \varepsilon q_{n-2}[\phi(u), u, \varepsilon] + b_{n-2,n-1} \psi_{n-1}(u) + b_{n-2,n} \psi_n(u) \right\} du, \quad ,$$

.....

$$\psi_{v+1}(t) = e^{\rho_{v+1} t} \int e^{-\rho_{v+1} u} \left\{ \varepsilon q_{v+1}[\phi(u), u, \varepsilon] + \sum_{l=v+2}^n b_{v+1,l} \psi_l(u) \right\} du. \quad .$$

All considerations of the present paper extend to this case without difficulties. Note that for  $v = 0$  no condition is imposed on  $A(\varepsilon)$  but  $\rho_j(0) \neq im\omega$ ,  $j = 1, \dots, n$ ,  $m = 0, \pm 1, \pm 2, \dots$ . For the sake of simplicity we shall refer in the following mainly to systems of the form (2.1). All considerations of §3 extend with obvious changes to the more general situations mentioned above.

(e). The elementary case  $v = 0$ .

## (3.iii). System

$$(3.20) \quad y' = A(\varepsilon)y + \varepsilon q(y, t, \varepsilon),$$

$y = (y_1, \dots, y_n)$ ,  $q = (q_1, \dots, q_n)$ , satisfying  $(\alpha)$  and  $(K)$  with  $v = 0$ , has always a periodic solution for all  $0 \leq \varepsilon \leq \varepsilon_1$  and some  $\varepsilon_1 > 0$  sufficiently small.

Indeed, for  $v = 0$ , there are no  $a_j$ ,  $b_j$ ,  $c_j$ , and  $(\alpha)$  requires only that  $\rho_j(0) \neq im\omega$ ,  $j = 1, \dots, n$ ,  $m = 0, \pm 1, \dots$ , where  $2\pi/\omega$  is a period of  $q$ . Then, for  $0 \leq \varepsilon \leq \varepsilon_0$  and some  $\varepsilon_0 > 0$ ,  $\delta > 0$ , we have also  $|\rho_j(\varepsilon) - im\omega| > \delta > 0$  for the same  $j$ 's and  $m$ 's. The transformation  $z$ , or  $\psi = z\varphi$ , of §2, is now the obvious one

$$\psi(t) = e^{iAt} \int e^{-iAu} q(\varphi(u), u, \varepsilon) du,$$

where the integrands are of class  $C_\omega$  and mean value zero and the integrals are the unique primitives of class  $C_\omega$  and mean value zero. Actually  $\psi(t)$  is the unique periodic solution of the linear system  $y' = Ay + \varepsilon q(\varphi(t), t, \varepsilon)$ . Note that we have here  $z(t) = (0, \dots, 0)$ , and  $A = B$ . Also  $\Omega_0$  is a sphere about  $\varphi = 0$ ,  $z\Omega_0 \subset \Omega_0$ , and  $z\Omega_0 \subset \Omega_0$  for  $0 \leq \varepsilon \leq \varepsilon_1$  and some  $\varepsilon_1 > 0$ . Finally  $z$  has a fixed element  $y(t) \in \Omega_0$ . The determining equations (2.16) are nonexistent (thus certainly satisfied) and  $y(t)$  is a periodic solution of (3.20). Under condition (L) the corresponding method of successive approximations is the usual one  $y^{(0)} = 0$ ,  $y^{(m)} = z y^{(m-1)}$ ,  $m = 1, 2, \dots$  [See A. Lyapunov, *Problème général de la stabilité du mouvement*. Ann. of Math. Studies, 17, 1949].

## (3.iv). System

$$(3.21) \quad y' = A(\varepsilon)y + F(t) + \varepsilon q(y, t, \varepsilon),$$

where  $y$ ,  $q$ ,  $F$  are  $n$ -vectors, where  $A$  and  $q$  satisfy  $(\alpha)$  and  $(K)$  with  $v = 0$ , and  $F(t)$  is periodic of period  $2\pi/\omega$ ,  $L$ -integrable in  $[0, 2\pi/\omega]$ ,  $|F_j(t)| \leq r < R$ ,  $j = 1, \dots, n$ , has always a periodic solution for all  $0 \leq \varepsilon \leq \varepsilon_1$  and some  $\varepsilon_1$  sufficiently small.

Indeed the condition  $\Psi[e^{-\rho_j t} F_j(t)] = 0$ ,  $j = 1, \dots, n$ , is necessarily satisfied since  $|\rho_j(\varepsilon) - im\omega| > \delta > 0$ , for all  $j$ 's and  $m$ 's as before. Hence, system (3.21) can be reduced to another analogous to (3.20). A little reflection shows that the method of successive approximations can actually be defined in terms of system (3.21) by taking for  $y^{(0)}(t) = z(t)$  the unique periodic solution of the linear system  $y' = A(\varepsilon)y + F(t)$ , and for  $y^{(m)}(t)$  the unique periodic solution of the

linear system  $y' = A(\varepsilon)y + F(t) + \varepsilon q[y^{(m-1)}](t), t, \varepsilon]$ ,  $m = 1, 2, \dots$ , in other words

$$y^{(0)}(t) = z(t) = e^{iAt} \int e^{-iAu} F(u) du,$$

$$y^{(m)}(t) = z(t) + \varepsilon e^{iAt} \int e^{-iAu} q[y^{(m-1)}](u), u, \varepsilon] du.$$

Extension of (3.1v) in the lines of the remarks  $d_1, d_2$  are the following ones.

(3.v). Consider the system

$$(3.22) \quad dy/dt = A(\varepsilon)y + q(y, t, \varepsilon) + g(y, t, \varepsilon),$$

where  $y, q, g$  are  $n$ -vectors, where  $A, q, g$  satisfy conditions  $(\alpha)$  and  $(K)$  with  $v = 0$ , and, in addition,  $q$  satisfies condition (3.16) and  $g$  condition (3.18). Then there is a periodic solution  $y(t)$  of (3.22) for all  $0 \leq \varepsilon \leq \varepsilon_1$  and some  $\varepsilon_1 > 0$  sufficiently small.

Finally, we may replace  $A(\varepsilon)$  in (3.22) by a  $n \times n$  matrix  $A(t, \varepsilon)$  whose elements are continuous functions of  $t$  and  $\varepsilon$ ,  $-\infty < t < +\infty$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , periodic in  $t$  of period  $2\pi/\omega$  (the same  $\omega$  as for  $q$  and  $g$ ), and whose characteristic exponents, say  $\lambda_j(\varepsilon)$ ,  $j = 1, \dots, n$ , (all defined mod  $\omega 1$ ) verify the relations  $\lambda_j(0) \neq 0 \pmod{\omega 1}$ ,  $j = 1, \dots, n$ .

The last extension contains, as particular cases, results of Antosiewicz and Diliberto, since the former assumed  $R(\lambda_j) < 0$ , and the latter  $R(\lambda_j) \neq 0$ ,  $j = 1, \dots, n$ . Furthermore we assume that the functions  $q$  and  $g$  are periodic and  $L$ -integrable in the period (in the sense of condition  $K$ ) instead of periodic and continuous (S. P. Diliberto and M. D. Marcus, On systems of ordinary differentiable equations. These Contributions, Vol. 3, 1956, 237-241).

An extension in a different direction can be obtained as follows. Suppose that system (3.21) satisfies conditions  $(\alpha)$  and  $(L)$  with  $v = 0$  and  $|R(\rho_j)| \geq \delta > 0$ ,  $j = 1, \dots, n$ , suppose that  $F(t) \in QP$ , where  $QP$  is the class of all vector functions whose components  $F_j(t)$  are  $L^2$ -quasi periodic, i.e.,

$$F_j(t) \sim \sum c_{jm} e^{i\lambda_m t}, \quad \sum |c_{jm}|^2 < +\infty, \quad j = 1, \dots, n.$$

We shall suppose also that  $q_j(y, t, \varepsilon) \in QP$  and  $q_j[y(t), t, \varepsilon] \in QP$ . This condition replaces the periodicity condition (p) of (K) as mentioned in §2.

Note that we have now  $z(t) \equiv 0$  and

$$y_j^0(t) \sim \sum (\rho_j + i\lambda_m)^{-1} c_{jm} e^{i\lambda_m t}, \quad \sum |(\rho_m + i\lambda_m)^{-1} c_{jm}|^2 \leq \delta^{-2} \sum |c_{jm}|^2 < +\infty,$$

i.e.,  $y^0(t) \in QP$ . The same argument proves that system (3.21) has a solution  $y(t)$  of class  $QP$ . [See J. J. Stoker, Nonlinear vibrations, App. 2., Interscience, 1950].

(f). The analytic case. Though not needed in the sequel the following remark may be of some interest. The method of successive approximations defined by relations (2.18) is the one initially proposed by the author [4b] in 1940 for linear systems, and successively used by J. K. Hale, R. A. Gambill [9b, 7abc] for linear systems, and by W. R. Fuller [10] for Lipschitzian nonlinear systems. The corresponding series

$$(3.23) \quad y_j(t) = y_j^{(0)}(t) + \sum_{m=0}^{\infty} [y_j^{(m+1)}(t) - y_j^{(m)}(t)], \quad j = 1, \dots, n,$$

are not power series in  $\varepsilon$  even in the case where the functions  $q_j(y, t, \varepsilon)$  are analytic in  $y_1, \dots, y_n, \varepsilon$ . For this case a modification of the process (2.18) has been used in [9c, 8, 6b]. In harmony with these papers, suppose

$$q_j(y, t, \varepsilon) = \sum_{k=0}^{\infty} \varepsilon^k q_j^{(k)}(y, t), \quad q_j^{(k)}(y, t) \in C_{\omega}, \quad 0 \leq \varepsilon \leq \varepsilon_0,$$

$q_j^{(k)}(y, t)$  analytic in  $y_1, \dots, y_n$ , and let  $s_j^{(r,k)}(t)$  denote the coefficient of  $\varepsilon^{r-1}$ ,  $r = 1, 2, \dots$ , when  $y$  in  $q_j^{(k)}(y, t)$  is replaced by

$$(3.24) \quad y = z^{(0)}(t) + \varepsilon z^{(1)}(t) + \varepsilon^2 z^{(2)}(t) + \dots,$$

where each  $z^{(k)}(t)$  is independent of  $\varepsilon$ . Moreover, let the corresponding coefficient of  $\varepsilon^{r-1}$  in  $q_j(y, t, \varepsilon)$  be denoted by  $s_j^{(r)}(t)$ . Let

$$\begin{aligned}
 c_j s_j^{(r,k)} &= \mathbb{M} \left\{ e^{-i\tau_j t} s_j^{(r,k)}(t) \right\}, & j = 1, \dots, v, \\
 (3.25) \quad c_j s_j^{(r)} &= \mathbb{M} \left\{ e^{-i\tau_j t} s_j^{(r)}(t) \right\}, & j = 1, \dots, v, \\
 S^{(r)} &= \text{diag} (s_1^{(r)}, \dots, s_v^{(r)}, 0, \dots, 0),
 \end{aligned}$$

where

$$\begin{aligned}
 s_j^{(r)} &= s_j^{(r,0)} + s_j^{(r-1,1)} + \dots + s_j^{(1,r-1)}, \\
 S_j^{(r)} &= S_j^{(r,0)} + S_j^{(r-1,1)} + \dots + S_j^{(1,r-1)}, \quad j = 1, \dots, v.
 \end{aligned}$$

The modified method of successive approximations is then defined by

$$y^{(0)}(t) = z(t) = z^{(0)}(t) = (c_1 e^{i\tau_1 t}, \dots, c_v e^{i\tau_v t}, 0, \dots, 0),$$

$$y^{(r)}(t) = z^{(0)}(t) + e^{Bt} \int e^{-Bu} \left[ \sum_{k=1}^r \varepsilon^k s^{(k)}(u) - \right.$$

$$(3.26) \quad \left. - \left( \sum_{k=1}^r \varepsilon^k s^{(k)}(u) \right) z^{(k-1)}(u) \right] du,$$

$$\text{mod } \varepsilon^{r+1}, \quad r = 1, 2, \dots,$$

with  $s^{(k)} = (s_1^{(k)}, \dots, s_v^{(k)}, 0, \dots, 0)$ ,  $y^{(r)} = z^{(0)} + \varepsilon z^{(1)} + \dots + \varepsilon^r z^{(r)}$ , and exactly the same conventions above. In [8, 9c] it was proved directly that this method converges and that  $y(t) = \lim_{r \rightarrow \infty} y^{(r)}(t)$  as  $r \rightarrow \infty$ , satisfies the differential system  $dy/dt = (B - \varepsilon h)y + \varepsilon q(y, t, \varepsilon)$ , where  $h = h(a, b, c, \omega, \varepsilon) = S^{(1)} + \varepsilon S^{(2)} + \varepsilon^2 S^{(3)} + \dots$ . Thus  $h$  may replace  $d$  in the analytic case.

All the proofs in the present paper are given in terms of the transformation  $\mathfrak{z}$  and a fixed point theorem. In the Lipschitzian case, some could be given in terms of the process of successive approximations (2.18) and, in the analytic case, even in terms of the process (3.24, 25, 26). The differences are mostly notational.

(g). A remark. In this section we have supposed all  $c_j \neq 0$  and between positive bounds,  $0 < r_1 \leq |c_j| \leq r_2 < R$ ,  $j = 1, \dots, v$ . There are cases where this requirement can be weakened or omitted.

A first situation where we may consider some  $c_j$  arbitrarily small, say  $0 \leq |c_j| \leq r_2 < R$ , is when, in a convenient class  $n$ , the corresponding mean value (2.11) is identically zero. Then we may assume  $d_j = 0$ ,  $c_j$  arbitrary. This situation will occur in §6.

A second analogous situation will occur when we can prove that the corresponding mean value (2.11) is  $O(c_j)$  as  $|c_j| \rightarrow 0$ . Then  $d_j = O(1)$ ,  $|d_j| \leq M$ , and the previous evaluations are simplified. This situation will occur in §4 with applications in §6.

Another analogous situation would occur if for  $c_j = O(\varepsilon^k)$  as  $\varepsilon \rightarrow 0$  for some  $j$  and  $k > 0$ , we could prove that also the corresponding mean values (2.11) are  $O(\varepsilon^k)$ . We do not discuss this case in the present paper.

#### §4. PROPERTIES OF THE FIXED ELEMENT

(a). The general case. Given system (2.1) under hypotheses ( $\alpha$ ) and (K), and the two sets of integers  $a = (a_1, \dots, a_v)$ ,  $b = (b_1, \dots, b_v)$ ,  $a_j \geq 0$ ,  $b_j > 0$ ,  $j = 1, \dots, v$ , we have determined the  $n$ -vector  $y(t) = y(t, a, b, c, \omega, \varepsilon) = (y_1, \dots, y_n)$  and the relative  $v$ -vector of numbers  $d_j$ ,  $D = D[y] = D(a, b, c, \omega, \varepsilon) = (d_1, \dots, d_v, 0, \dots, 0)$  with  $-\infty < t < +\infty$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ ,  $c = (c_1, \dots, c_v)$ ,  $0 < r_1 \leq |c_j| \leq r_2 < R$ ,  $j = 1, \dots, v$ . As mentioned in §2, in the periodic case  $\omega$  is fixed, since the relations

$$\rho_j(0) = 1a_j\omega/b_j, \quad j = 1, \dots, v,$$

are satisfied, and the functions  $q_j$  are periodic functions of  $t$  of period  $T = 2\pi/\omega$ . In the autonomous case we shall denote by  $\omega = \omega_0$  the value of  $\omega$  for which the  $v$  relations above are verified and then  $\omega$  is thought of as a parameter varying in an interval  $U = [\omega_{01}, \omega_{02}]$ ,  $0 < \omega_{01} < \omega_0 < \omega_{02} < +\infty$ . In this section we shall discuss the properties of  $y$  as a function of  $t$ , as well as the properties of  $y$  and  $D$  as functions of  $c$ ,  $\varepsilon$  and (in the autonomous case) of  $\omega$ . For the sake of brevity we must treat the two cases together: system (2.1) periodic, and system (2.1) autonomous. To do this we shall suppose, in all that follows, that in the periodic case the interval  $U$  is reduced to the single point  $\omega_0$ , thus  $\omega_{01} = \omega_0 = \omega_{02} = \omega$ . We shall take for the  $v$ -vectors  $c$  the norm  $\|c\| = \max |c_j|$ ,  $j = 1, \dots, v$ . The vector function  $y(t)$  is periodic in  $t$  of period  $2\pi/\omega$  and verifies the integral equation (2.15) and the differential equation (2.16), thus

$$y'(t) = (B - \varepsilon D)y(t) + \varepsilon q(y, t, \varepsilon),$$

where  $A = \text{diag} [\rho_j(\varepsilon), j = 1, \dots, n]$ , and  $B = \text{diag} [\rho_j(0), j = 1, \dots, v; \rho_j(\varepsilon), j = v + 1, \dots, n]$ . Let  $M$  be large enough in order that  $|\rho_j(\varepsilon)| \leq M, j = 1, \dots, n, 0 \leq \varepsilon \leq \varepsilon_1$ . Since  $|y_j| \leq R, |q_j| \leq \psi(t)$ , and  $|d_j| \leq r_1^{-1}K$  (§3) we have, as in §3,

$$(4.1) \quad |\dot{y}_j'(t)| \leq MR + \varepsilon_1 r_1^{-1}KR + \varepsilon_1 \psi(t).$$

Thus, we have, for all  $t^1, t^2$ ,

$$|y_j(t^2) - y_j(t^1)| \leq (M + \varepsilon_1 r_1^{-1}K)R |t^2 - t^1| + \varepsilon_1 \int_{t^1}^{t^2} \psi(t) dt,$$

i.e., the functions  $y_j(t), j = 1, \dots, n$ , are equiabsolutely continuous in  $t$  (with respect to  $c$  and  $\varepsilon$ ). In the autonomous case, we have  $\psi(t) = K$ , and

$$(4.2) \quad |\dot{y}_j'(t)| \leq MR + \varepsilon_1 r_1^{-1}KR + \varepsilon_1 K,$$

i.e., the functions  $y_j(t), j = 1, \dots, n$ , are uniformly Lipschitzian in  $t$  (with respect to  $c, \varepsilon$ , and  $\omega$ ).

We could now prove a number of theorems stating that, under condition (L),  $y(t, a, b, c, \omega, \varepsilon), D(a, b, c, \omega, \varepsilon)$  are continuous functions of  $c, \varepsilon$ , and (in the autonomous case) of  $\omega$ , and that further smoothness conditions on the  $q_j$  imply analogous properties for  $y$  and  $D$ . We will prove only the following theorem of a mixed type (which cannot be deduced from general theorems. Cfr. J. Schauder and J. Leray, loc. cit. in §5).

(4.1). Suppose hypotheses  $(\alpha)$  and (L) are satisfied, that  $a = (a_1, \dots, a_v), b = (b_1, \dots, b_v), 0 < r_1 < r_2 < R, 0 \leq \varepsilon_1 \leq \varepsilon_0$ , and  $n_0$  are given as in (3.1), and let  $V \geq 2\pi b_0/\omega_0, b_0 = b_1 \dots b_v$  be any number. Then, there are numbers  $\varepsilon_2, M_2, 0 < \varepsilon_2 \leq \varepsilon_1, M_2 > 0$ , and, given  $\xi > 0$  arbitrary, there is another number  $\xi > 0$  such that, for all  $c^s = (c_{js}, j = 1, \dots, v), r_1 \leq |c_{js}| \leq r_2, j = 1, \dots, v, s = 1, 2$ , for all  $\omega^1, \omega^2, \omega_0^1 \leq \omega^1, \omega^2 \leq \omega_0^2$ , for all  $\varepsilon^1, \varepsilon^2, 0 \leq \varepsilon^1, \varepsilon^2 \leq \varepsilon_2, |\varepsilon^2 - \varepsilon^1| \leq \xi$ , we have

$$\left. \begin{aligned} & |y_j(t, a, b, c^1, \omega^1, \varepsilon^1) - y_j(t, a, b, c^2, \omega^2, \varepsilon^2)| \\ & |d_h(a, b, c^1, \omega^1, \varepsilon^1) - d_h(a, b, c^2, \omega^2, \varepsilon^2)| \end{aligned} \right\} \leq$$

$$\leq M_2 \|c^1 - c^2\| + |\omega^1 - \omega^2| + \xi,$$

$j = 1, \dots, n, h = 1, \dots, v, 0 \leq t \leq V$ . Note that for  $\omega^1 = \omega^2$ , the

functions  $y_j$  above are all periodic of the same period  $2\pi b_0/\omega^1 \leq V$ , and hence, the relations above hold for all  $t$ ,  $-\infty < t < +\infty$ . This occurs, in particular, in the periodic case, since then  $\omega^1 = \omega^2 = \omega_0$ .

PROOF. Take  $\zeta^1 = \zeta/N$ , where  $N$  is a constant which will be determined later, and let  $\xi > 0$  be the number defined in (K) in relation to  $\zeta^1$ . We may assume  $\xi \leq \zeta^1$ . Also, we may assume  $\xi$  so small that  $|\rho_j(\varepsilon^1) - \rho_j(\varepsilon^2)| \leq \xi^1$ ,  $j = 1, \dots, n$ , for all  $0 \leq \varepsilon^1, \varepsilon^2 \leq \varepsilon_1$ ,  $|\varepsilon^1 - \varepsilon^2| \leq \xi$ .

We may suppose  $\omega_{01} \leq \omega^1 \leq \omega^2 \leq \omega_{02}$ . Take  $V_1 = 2\pi b_0/\omega_{02}$ ,  $V_2 = 2\pi b_0/\omega_{01}$ ,  $V_1 \leq V_2 \leq V$ . Since  $\omega_{01} \leq \omega \leq \omega_{02}$  and  $T = 2\pi b_0/\omega$ , we have  $V_1 \leq T \leq V_2 \leq V$ . Note that  $|q_j| \leq \psi(t)$ ,  $j = 1, \dots, n$ , and that we took  $K$  in such a way that

$$\int_0^T \psi(t) dt \leq KT.$$

In the autonomous case  $\psi(t) \equiv K$  and this requirement is certainly satisfied. We shall denote by  $y^s, D^s, y_{js}, d_{js}, \tau_{js}, \rho_{js}, T_s$ , etc.,  $s = 1, 2$ , the values of  $y, D, y_j, d_j, \tau_j, \rho_j, T$  for  $c = c^1, \omega = \omega^1, \varepsilon = \varepsilon^1$ , and  $c = c^2, \omega = \omega^2, \varepsilon = \varepsilon^2$ . We shall omit to indicate all the parameters which are not essential. Note that  $V_1 \leq T_2 \leq T_1 \leq V_2 \leq V$ .

For  $j = 1, \dots, v$ , we have by (2.11)

$$d_{j2} - d_{j1} = c_{j2}^{-1} T_2^{-1} \int_0^{T_2} e^{-i\tau_{j2}t} q_{j2}[y^2(t)] dt - c_{j1}^{-1} T_1^{-1} \int_0^{T_1} e^{-i\tau_{j1}t} q_{j1}[y^1(t)] dt,$$

where  $q_{js}[y^s(t)]$  denotes  $q_j[y(t, c^s, \omega^s, \varepsilon^s), t, \varepsilon^s]$ ,  $s = 1, 2$ . By manipulation, by introducing the new variable of integration  $u = \omega_1 t / \omega_2$  in the second integral, and changing  $u$  into  $t$  we have

$$d_{j2} - d_{j1} = (c_{j2}^{-1} - c_{j1}^{-1}) T_2^{-1} \int_0^{T_2} e^{i\tau_{j2}t} q_{j2}[y^2(t)] dt +$$

$$(4.3) \quad + c_{j1}^{-1} T_2^{-1} \int_0^{T_2} e^{-i\tau_{j2}t} \{q_{j2}[y^2(t)] - q_{j1}[y^1(\omega^2 t / \omega^1)]\} dt.$$

Note that in the periodic case  $T_1 = T_2$ ,  $\tau_{j1} = \tau_{j2}$ ,  $\omega_1 = \omega_2$ , and thus the expression in ( ) actually is



$$(4.4) \quad q_j[y(t, c^2, \omega, \varepsilon^2), t, \varepsilon^2] - q_j[y(t, c^1, \omega, \varepsilon^1), t, \varepsilon^1] ,$$

while in the autonomous case it is

$$(4.5) \quad q_j[y(t, c^2, \omega^2, \varepsilon^2), \varepsilon^2] - q_j[y(\omega^2 t / \omega^1, c^1, \omega^1, \varepsilon^1), \varepsilon^1] .$$

We shall denote by  $\Delta$  the number  $\Delta = \max |y_{j2}(t) - y_{j1}(t)|$  for all  $0 \leq t \leq V$ ,  $j = 1, \dots, n$ . For every  $0 \leq t \leq V$  we have  $0 \leq \omega^2 t / \omega^1 \leq \omega^2 V / \omega^1$  and hence  $|t - \omega^2 t / \omega^1| \leq \omega_{01}^{-1} V |\omega^2 - \omega^1|$ . Because of (4.2) [only in the autonomous case we may take  $\omega^1 \neq \omega^2$ ], we have

$$\begin{aligned} & |y_j(t, c^2, \omega^2, \varepsilon^2) - y_j(\omega^2 t / \omega^1, c^1, \omega^1, \varepsilon^1)| \leq \\ & \leq |y_j(t, c^2, \omega^2, \varepsilon^2) - y_j(t, c^1, \omega^1, \varepsilon^1)| + \\ & + |y_j(t, c^1, \omega^1, \varepsilon^1) - y_j(\omega^2 t / \omega^1, c^1, \omega^1, \varepsilon^1)| \leq \\ & \leq \Delta + \omega_{01}^{-1} V |\omega^2 - \omega^1| [MR + \varepsilon_1 K(1 + r_1^{-1} R)] = \\ (4.6) \quad & = \Delta + P |\omega^2 - \omega^1|, \quad 0 \leq t \leq V, \end{aligned}$$

where  $P$  is a constant.

By (L) we can now evaluate the expression in ( ) of (4.3), that is, either (4.4) or (4.5):

$$\begin{aligned} |q_{j2}[y^2(t)] - q_{j1}[y^1(t)]| & \leq \left[ \sum_{\ell=1}^n |y_{\ell 2}(t) - y_{\ell 1}(t)| + \xi^1 \right] \psi(t) \\ (4.7) \quad & \leq [n\Delta + nP |\omega^2 - \omega^1| + \xi^1] \psi(t) \end{aligned}$$

By (4.3) we have then

$$(4.8) \quad |d_{j2} - d_{j1}| \leq r_1^{-1} K(r_1^{-1} \|c^2 - c^1\| + n\Delta + nP |\omega^2 - \omega^1| + \xi^1) .$$

For  $j = 1, \dots, v$ , by (2.9) we have

$$\begin{aligned} y_{j2}(t) - y_{j1}(t) & = c_{j2} e^{1\tau j_2 t} - c_{j1} e^{1\tau j_1 t} + \\ & + \varepsilon^2 e^{1\tau j_2 t} \int e^{-1\tau j_2 u} \{q_{j2}[y^2(u)] - d_{j2} y_{j2}(u)\} du - \\ & - \varepsilon^1 e^{1\tau j_1 t} \int e^{-1\tau j_1 u} \{q_{j1}[y^1(u)] - d_{j1} y_{j1}(u)\} du . \end{aligned}$$

By manipulation, by introducing the variable  $v = \omega^1 u / \omega^2$  in the second integral and replacing  $v$  by  $u$ , we have

$$\begin{aligned}
 y_{j_2}(t) - y_{j_1}(t) &= (c_{j_2} - c_{j_1}) e^{i\tau_{j_1}t} + (1 - e^{i(\tau_{j_1} - \tau_{j_2})t}) y_{j_2}(t) + \\
 &+ (\varepsilon^2 - \varepsilon^1) e^{i\tau_{j_1}t} \int e^{-i\tau_{j_2}u} \{q_{j_2}[y^2(u)] - d_{j_2}y_{j_2}(u)\} du + \\
 &+ \varepsilon^1 e^{i\tau_{j_1}t} \int e^{-i\tau_{j_2}u} \left\{ q_{j_2}[y^2(u)] - \frac{\omega^2}{\omega^1} q_{j_1}\left[y^1\left(\frac{\omega^2}{\omega^1}u\right)\right] - d_{j_2}y_{j_2}(u) + \right. \\
 &+ \left. \frac{\omega^2}{\omega^1} d_{j_1}y_{j_1}\left(\frac{\omega^2}{\omega^1}u\right) \right\} du + \varepsilon^1 \frac{\omega^2}{\omega^1} e^{i\tau_{j_1}t} \int_{u=\omega^1 t/\omega^2}^{u=t} e^{-i\tau_{j_2}u} \left\{ q_{j_1}\left[y^1\left(\frac{\omega^2}{\omega^1}u\right)\right] - \right. \\
 (4.9) \quad &\left. - d_{j_1}y_{j_1}\left(\frac{\omega^2}{\omega^1}u\right) \right\} du = s_1 + \dots + s_5.
 \end{aligned}$$

For  $0 \leq t \leq V$  we have  $|s_1| \leq \|c^2 - c^1\|$ , and

$$|s_2| = \left| y_{j_2}(t) \int_{s=\tau_{j_1}-\tau_{j_2}}^{s=0} (it) e^{ist} ds \right| \leq RV |\tau_{j_1} - \tau_{j_2}| \leq RVQ |\omega^2 - \omega^1|,$$

since  $\tau_{j_s} = i a_j \omega^s / b_j$ ,  $s = 1, 2$ , and we take  $Q = \max a_j / b_j$ ,  $j = 1, \dots, v$ . By  $|\varepsilon^2 - \varepsilon^1| \leq \xi \leq \xi^1$ , and by (1.11) taking  $N = 2$ , and by (3.2), we have

$$|s_3| \leq \xi \cdot 2 \int_0^{T_2} |q_{j_2}[y^2(u)] - d_{j_2}y_{j_2}(u)| \leq 2KT_2(1 + r_1^{-1}R)\xi^1.$$

In  $s_4$  the integrand is periodic of period  $T_2$ , of class  $C_{\omega^2}$  and mean value zero. By (1.11), taking  $N = 2$ , we have

$$\begin{aligned}
 |s_4| &\leq 2\varepsilon^1 \int_0^{T_2} \left\{ q_{j_2}[y^2(u)] - (\omega^2/\omega^1) q_{j_1}y^1(\omega^2 u/\omega^1) + \right. \\
 &+ \left. |d_{j_2} - (\omega^2/\omega^1)d_{j_1}| |y_{j_2}(u) + (\omega^2/\omega^1)| d_{j_1}| |y_{j_2}(u) - y_{j_1}(\omega^2 u/\omega^1)| \right\} du,
 \end{aligned}$$

and, by (4.6), (4.7), (4.8), also

$$\begin{aligned}
 & |q_{j2}[y^2(u)] - (\omega^2/\omega^1)q_{j1}[y^1(\omega^2 u/\omega^1)]| \\
 & \leq |q_{j2}[y^2(u)] - q_{j1}[y^1(\omega^2 u/\omega^1)]| + |1 - \omega^2/\omega^1| |q_{j1}[y^1(\omega^2 u/\omega^1)]| \\
 & \leq [n\Delta + nP|\omega^2 - \omega^1| + \zeta^1] \psi(t) + \omega_{01}^{-1} |\omega^2 - \omega^1| \psi(t), \quad j = 1, \dots, n; \\
 & |d_{j2} - (\omega^2/\omega^1)d_{j1}| \leq r_1^{-1} K(r_1^{-1} \|c^2 - c^1\| + n\Delta + nP|\omega^2 - \omega^1| + \zeta^1) \\
 & \quad + \omega_{01}^{-1} r_1^{-1} K|\omega^2 - \omega^1|, \\
 & |(\omega^2/\omega^1)d_{j1}| |y_{j2}(u) - y_{j1}(\omega^2 u/\omega^1)| \leq \omega_{01}^{-1} \omega_{02} r_1^{-1} K[\Delta + P|\omega^2 - \omega^1|], \\
 & \quad j = 1, \dots, v.
 \end{aligned}$$

Thus

$$|s_4| \leq \varepsilon^1 P^1 \Delta + P'' |\omega^2 - \omega^1| + P''' \|c^2 - c^1\| + P^{1v} \zeta^1, \quad ,$$

for convenient constants  $P^1, P'', P''', P^{1v}$ . Finally,  $s_5$  may be not zero only in the autonomous case, and

$$|s_5| \leq \varepsilon^1 \omega_{01}^{-1} \omega_{02} |t - \omega^1 t/\omega^2| (K + r_1^{-1} KR) \quad .$$

By (4.9) and the present evaluations, we have

$$\begin{aligned}
 & |y_{j2}(t) - y_{j1}(t)| \leq |s_1| + |s_2| + |s_3| + |s_4| + |s_5| \\
 (4.10) \quad & \leq \varepsilon^1 Q^1 \Delta + Q'' |\omega^2 - \omega^1| + Q''' \|c^2 - c^1\| + Q^{1v} \zeta^1, \\
 & \quad j = 1, \dots, v,
 \end{aligned}$$

for convenient constants  $Q$ .

For  $j = v + 1, \dots, n$ , by (2.10) we have

$$\begin{aligned}
 & y_{j2}(t) - y_{j1}(t) = \varepsilon^2 e^{\rho j_2 t} \int_0^{-\rho j_2 t} q_{j2}[y^2(u)] du - \varepsilon^1 e^{\rho j_1 t} \int_0^{-\rho j_1 t} q_{j1}[y^1(u)] du = \\
 & = (\varepsilon^2 - \varepsilon^1) e^{\rho j_2 t} \int_0^{-\rho j_2 t} q_{j2}[y^2(u)] du + \varepsilon^1 (e^{\rho j_2 t} - e^{\rho j_1 t}) \int_0^{-\rho j_2 t} q_{j2}[y^2(u)] du + \\
 & + \varepsilon^1 e^{\rho j_1 t} \left\{ \int_0^{-\rho j_2 t} q_{j2}[y^2(u)] du - \int_0^{-\rho j_1 t} q_{j2}[y^2(u)] du \right\} + \\
 & + \varepsilon^1 e^{\rho j_1 t} \int_0^{-\rho j_1 t} \{q_{j2}[y^2(u)] - q_{j1}[y^1(u)]\} du = \sigma_1 + \dots + \sigma_4.
 \end{aligned}$$

For  $0 \leq t \leq V$ , by (1.111), (4.7), and for convenient constants  $N$ ,  $H = e^{\gamma V}$ ,  $\gamma = \max_{j=1, \dots, n} |\rho_j(\varepsilon)|$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ ,  $j = 1, \dots, n$ , we have

$$|\sigma_1| \leq \varepsilon^1_{HNKT},$$

$$|\sigma_4| \leq \varepsilon^1_{HNKT} [n\Delta + nP |\omega^2 - \omega^1| + \zeta^1],$$

$$|\sigma_2| \leq \varepsilon^1_{NKT} |e^{\rho_j 2^t} - e^{\rho_j 1^t}| \leq \varepsilon^1_{HNKT V} |\rho_{j2} - \rho_{j1}| \leq \varepsilon^1_{HNKT V \zeta^1}.$$

Finally, by (1.1v), we have

$$|\sigma_3| \leq \varepsilon^1_{HC} |\rho_{2j} - \rho_{1j}| \int_0^T |q_{j2}[y^2(u)]| du \leq \varepsilon^1_{HCKT \zeta^1}.$$

Thus, for  $0 \leq t \leq V$ , we have

$$(4.11) \quad |y_{j2}(t) - y_{j1}(t)| \leq \varepsilon^1_{N' \Delta} + N'' |\omega^2 - \omega^1| + N''' \zeta^1, \quad j = v+1, \dots, n,$$

for convenient constants  $N'$ ,  $N''$ ,  $N'''$ .

By (4.10) and (4.11) we have now

$$|y_{j2}(t) - y_{j1}(t)| \leq \varepsilon^1_{M' \Delta} + M'' |\omega^2 - \omega^1| + M''' \|c^2 - c^1\| + M^{1v}_{\zeta^1},$$

for all  $j = 1, \dots, n$ ,  $0 \leq t \leq V$ , and hence, if we take  $\varepsilon_2 = \min[\varepsilon_1, 1/2M']$ , for  $0 \leq \varepsilon^1, \varepsilon^2 \leq \varepsilon_2$ , we have

$$\Delta \leq (1/2)\Delta + M'' |\omega^2 - \omega^1| + M''' \|c^2 - c^1\| + M^{1v}_{\zeta^1}$$

and finally

$$\Delta \leq 2M'' |\omega^2 - \omega^1| + 2M''' \|c^2 - c^1\| + 2M^{1v}_{\zeta^1}.$$

If we suppose  $N = 2M^{1v}$  and  $M = \max[2M'', 2M''']$ , we have  $2M^{1v}_{\zeta^1} = N\zeta^1 = \zeta$  and

$$(4.12) \quad |y_{j2}(t) - y_{j1}(t)| \leq \Delta \leq M[|\omega^2 - \omega^1| + \|c^2 - c^1\|] + \zeta$$

for all  $0 \leq t \leq V$ ,  $j = 1, \dots, n$ . This proves (4.1).

(c). Further properties of  $y$  in the autonomous case and  
 $v = 2 \leq n$ . The theorems (3.1), (4.1) admit of a stronger form under additional hypotheses. We consider here only the autonomous case with  
 $v = 2$ ,  $-a_2 = a_1 = a = 1$ ,  $b_1 = b_2 = b = 1$ ,  $-c_2 = c_1 = c > 0$ ,  
 $-\tau_2 = \tau_1 = \tau = aw/b$ . We need the following additional hypothesis:

$$(K') \quad q_j(0, 0, x_3, \dots, x_n, \varepsilon) = 0, \quad j = 1, 2.$$

The following statement is stronger than (3.1):

(4.11). In the situation above and under hypotheses  $(\alpha)$ , (L),  $(K')$ , given  $a, b$ , and any  $r_2$ ,  $0 < r_2 < R$ , there is an  $\varepsilon_1$ ,  $0 < \varepsilon_1 \leq \varepsilon_0$ , such that for every  $c$ ,  $0 < c \leq r_2$  and  $0 \leq \varepsilon \leq \varepsilon_1$ ,  $\tau | \Omega_0$  is a contraction and into, where  $\Omega_0$  is a convenient closed subset of  $\Omega$ .

PROOF. The proof is the same as for (3.1) with the following modifications. First, by virtue of (L) and  $(K')$ , we have

$$|q_j(x, \varepsilon) - 0| \leq K(|x_1| + |x_2|), \quad j = 1, 2.$$

For  $r_2/3 \leq c \leq r_2$  Theorem (3.1) holds. Suppose  $0 < c \leq r_2/3$ , and define  $\Omega_0$  as the closed subset of  $\Omega$  with  $|\varphi_j(t) - z_j(t)| \leq 2c$ ,  $j = 1, 2$ ,  $|\varphi_j(t)| \leq r_2$ ,  $j = 3, \dots, n$ , where  $z(t) = (ce^{1-t}, -ce^{-1-t}, 0, \dots, 0)$ . Thus  $|\varphi_j(t)| \leq |z_j(t)| + 2c \leq 3c \leq r_2 < R$ ,  $j = 1, 2$ . Instead of (3.2) we have now

$$\begin{aligned} |d_j| &\leq c^{-1}T^{-1} \int_0^T |q_j[\varphi(u), \varepsilon]| \, du \\ &\leq c^{-1}T^{-1}K \int_0^T [|\varphi_1(u)| + |\varphi_2(u)|] \, du \leq 6K, \quad j = 1, 2. \end{aligned}$$

Relation (3.3) is now replaced by

$$|\psi_j(t) - z_j(t)| \leq 2\varepsilon[6KTc + 6K^2T^2c] = 48\varepsilon K^2Tc, \quad j = 1, 2,$$

while relation (3.4) remains the same

$$|\psi_j(t)| \leq \varepsilon HNK^2T, \quad j = 3, \dots, n.$$

If  $M = \max [24KT, HNK^2T]$ ,  $\varepsilon_1 = \min [\varepsilon_0, 1/M, r_2/M]$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ , we have

$$|\psi_j(t) - z_j(t)| \leq 2\epsilon M c \leq 2c, \quad j = 1, 2,$$

$$|\psi_j(t)| \leq \epsilon M \leq r_2, \quad j = 3, \dots, n,$$

i.e.,  $\tau \cap_0 \subset \cap_0$ . For any two  $\varphi^s \in \cap_0$ ,  $s = 1, 2$ , we have, instead of (3.5L)

$$c(d_{1j} - d_{2j}) = \mathfrak{M} \left[ e^{\frac{1}{\tau} j u} \left\{ q_j[\varphi^1(u), \epsilon] - q_j[\varphi^2(u), \epsilon] \right\} \right],$$

$$c|d_{1j} - d_{2j}| \leq T^{-1} \int_0^T K \sum_{\ell=1}^n |\varphi_\ell^1(u) - \varphi_\ell^2(u)| du \leq nK \cdot \mathfrak{M}[\varphi^1 - \varphi^2],$$

$$|(d_{1j} - d_{2j})\varphi_{2j}(u)| \leq 3c |d_{1j} - d_{2j}| \leq 3nK \cdot \mathfrak{M}[\varphi^1 - \varphi^2], \quad j = 1, 2.$$

Finally (3.6L) is replaced by

$$\begin{aligned} |\psi_{1j}(t) - \psi_{2j}(t)| &\leq 2\epsilon[nKT + 3nKT + 3KnT] \mathfrak{M}[\varphi^1 - \varphi^2] \\ &\leq 14\epsilon nKT \mathfrak{M}[\varphi^1 - \varphi^2], \quad j = 1, 2, \end{aligned}$$

while (3.7L) remains the same. For  $M_2 = \max[14nKT, nHNKT]$ ,  $\epsilon_1^1 = \min[\epsilon_1, 1/2M_2]$ ,  $0 \leq \epsilon \leq \epsilon_1^1$ , we have

$$\mathfrak{M}[\psi^1 - \psi^2] \leq \epsilon M_2 \mathfrak{M}[\varphi^1 - \varphi^2] \leq 2^{-1} \mathfrak{M}[\varphi^1 - \varphi^2],$$

and thus  $\tau \cap_0$  is a contraction. Thereby, (4.11) is proved.

In analogy with (3.12) we have now

$$|\gamma_1(t)|, |\gamma_2(t)| \leq 3c, \quad |\gamma_j(t)| \leq r_2, \quad j = 3, \dots, n,$$

$$|\gamma_1(t) - ce^{1\tau t}|, |\gamma_2(t) + ce^{-1\tau t}| \leq 2c,$$

$$|\gamma_j(t)| \leq \epsilon HNKT, \quad j = 3, \dots, n.$$

If we suppose: (K'') the requirement (K') holds for all  $q_j$ ,  $j = 1, \dots, n$ , then we have also

$$|\gamma_j(t)| \leq 3HNKnc, \quad j = 3, \dots, n.$$

The statements of this section become, under hypotheses  $(\alpha)$ ,  $(L)$ ,  $(K')$ :

(4.111). Under hypotheses  $(\alpha)$ ,  $(L)$  and  $(K')$ , given  $a$ ,  $b$ ,  $r_2$ , and  $\alpha_0$  as in (4.11), there is an  $\varepsilon_2$ ,  $0 < \varepsilon_2 < \varepsilon_1$ , and  $M > 0$  such that, for all  $c^s$ ,  $0 < c^s \leq r_2$ ,  $s = 1, 2$ ,  $0 \leq \varepsilon \leq \varepsilon_2$ , the conclusions of (4.1) hold.

## §5. FIRST EXISTENCE THEOREMS FOR REAL SYSTEMS

Consider the real system, of order  $N = \mu + n$ ,

$$\begin{aligned} x_j'' + 2\alpha_j x_j' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', t, \varepsilon) & j = 1, \dots, \mu, \\ x_j' + \beta_j x_j &= \varepsilon f_j(x, x', t, \varepsilon), & j = \mu + 1, \dots, n, \end{aligned} \quad (5.1)$$

where  $(') = d/dt$ ,  $-\infty < t < +\infty$ ,  $0 \leq \mu \leq n$ ,  $x = (x_1, \dots, x_n)$ ,  $x' = (x_1', \dots, x_n')$ ,  $\sigma_j > 0$ ,  $\alpha_j$ ,  $\beta_j$  real constants, or continuous functions of  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ , and where  $f_j(u, v, t, \varepsilon)$ ,  $j = 1, \dots, n$ , are real functions of the  $n$ -vector  $u$ , of the  $\mu$ -vector  $v$ , of  $-\infty < t < +\infty$ , and of  $\varepsilon \geq 0$ .

For the sake of simplicity we shall assume  $\alpha_j^2 < \sigma_j^2$ ,  $j = 1, \dots, \mu$ , and put  $\gamma_j = (\sigma_j^2 - \alpha_j^2)^{1/2} > 0$ ,  $\rho_{j1}(\varepsilon) = \rho_{j2}(\varepsilon) = -\alpha_j \pm i\gamma_j$ ,  $j = 1, \dots, \mu$ .

We shall order the first  $\mu$  equations (5.1) in such a way that for some fixed  $\omega > 0$  we have  $\alpha_j(0) = 0$ ,  $\sigma_j(0) = a_j\omega/b_j$ ,  $a_j, b_j > 0$  integers,  $j = 1, \dots, \nu$ , and  $\rho_{j1}(0), \rho_{j2}(0) \neq i\omega/b_0$ ,  $j = \nu + 1, \dots, \mu$ ,  $m = 0, \pm 1, \pm 2, \dots$ . The latter can be expressed by saying that either  $\alpha_j(0) \neq 0$ , or  $\alpha_j(0) = 0$  and  $\sigma_j \neq m\omega/b_0$ ,  $b_0 = b_1 \dots b_\nu$ ,  $m = 0, 1, 2, \dots$ , ( $j = \nu + 1, \dots, \mu$ ). We shall order the last  $n - \mu$  equations (5.1) in such a way that  $\beta_j(0) \neq 0$  for  $j = \mu + 1, \dots, r$ , and  $\beta_j(0) = 0$  for  $j = r + 1, \dots, n$ , with  $\mu \leq r \leq n$ .

Thus we have  $0 \leq \nu \leq \mu \leq r \leq n$ .

We shall finally assume that the functions  $f_j$ ,  $j = 1, \dots, n$ , satisfy condition  $(L)$  (with respect to the  $n + \mu$  vector  $(x_1, \dots, x_n, x_1', \dots, x_\mu')$ ).

Let us introduce in (5.1) the new variables  $y_1, \dots, y_N$ ,  $N = n + \mu$ , by the usual relations

$$\begin{aligned}
 (5.2) \quad y_{2j-1} &= -\rho_{j2}x_j + x_j^1, \quad y_{2j} = \rho_{j1}x_j - x_j^1, & j &= 1, \dots, \mu, \\
 y_{\mu+j} &= x_j, & j &= \mu+1, \dots, n,
 \end{aligned}$$

and hence

$$(5.3) \quad x_j = (2i\gamma_j)^{-1}(y_{2j-1} + y_{2j}), \quad x_j^1 = (2i\gamma_j)^{-1}(\rho_{j1}y_{2j-1} + \rho_{j2}y_{2j}), \\
 j = 1, \dots, \mu.$$

Then system (5.1) is transformed into the system of  $N = \mu + n$  first order differential equations

$$(5.4) \quad y' = Ay + \varepsilon q(y, t, \varepsilon),$$

where  $y = (y_1, \dots, y_N)$ ,  $q = (q_1, \dots, q_N)$ ,  $A$  is the diagonal matrix of the  $N$  elements

$$A = \text{diag}(\rho_{j1}, \rho_{j2}, j = 1, \dots, \mu, -\beta_j, j = \mu+1, \dots, n),$$

and

$$q_{2j-1} = f_j, \quad q_{2j} = -f_j, \quad j = 1, \dots, \mu, \quad q_{\mu+j} = f_j, \quad j = \mu+1, \dots, n,$$

where the  $N+2$  arguments of the  $q$ 's are, in order,  $y_1, \dots, y_N; t, \varepsilon$ , and the  $N+2$  arguments of the  $f$ 's are, in order

$$\begin{aligned}
 (2i\gamma_j)^{-1}(y_{2j-1} + y_{2j}), \quad j &= 1, \dots, \mu; \quad y_{\mu+j}, \quad j = \mu+1, \dots, n; \\
 (2i\gamma_j)^{-1}(\rho_{j1}y_{2j-1} + \rho_{j2}y_{2j}), \quad j &= 1, \dots, \mu; \quad t, \varepsilon.
 \end{aligned}$$

System (5.4) is of the type considered in §2, where the first  $2v$  equations and the last  $n-r$  equations replace the first  $v$  equations of §2; hence  $2v + (n-r)$  replaces  $v$ ,  $N = n + \mu$  replaces  $n$ , and  $n + \mu - [2v + (n-r)] = \mu + r - 2v$  replaces  $n - v$ . The numbers  $\rho_j(0) = i\tau_j$ ,  $j = 1, \dots, v$ , of §2 are now replaced by the following  $2v + (n-r)$  numbers

$$i\tau_1, -i\tau_1, \dots, i\tau_v, -i\tau_v, 0, \dots, 0,$$

(the zero repeated  $n-r$  times), where  $\tau_j = a_j\omega/b_j$ ,  $j = 1, \dots, v$ . In other words, for the numbers  $a_j, b_j$  of §2, say  $a_j^1, b_j^1$ , we have now the



integers  $a'_{2j-1} = a_j$ ,  $a'_{2j} = -a_j$ ,  $b'_{2j-1} = b'_{2j} = b_j$ ,  $j = 1, \dots, \nu$ ,  $a'_j = 0$ ,  $b'_j = 1$ ,  $j = \mu + 1, \dots, n$ . Let us take for the  $\nu$  numbers  $c$  of §2 the following  $2\nu + (n - \mu)$  numbers

$$c'_{2j-1} = c_j, c'_{2j} = -\bar{c}_j, c_j \neq 0, \text{ complex}, j = 1, \dots, \nu,$$

$$c_j \neq 0, c_j \text{ real}, j = \mu + 1, \dots, \mu + n.$$

Then the vector  $z(t)$  of §2 is now the  $N$ -vector

$$z(t) = (c_j e^{i\tau_j t}, -\bar{c}_j e^{-i\tau_j t},$$

$$j = 1, \dots, \nu; 0, \dots, 0; 0, \dots, 0; c_j, j = \mu + 1, \dots, \mu + n),$$

where the zero is repeated  $(2\mu - 2\nu) + (r - \mu)$  times. In the present situation we may consider the space  $\Omega$ , analogous to the one of §2, of continuous periodic  $N$ -vector functions  $\varphi(t) = (\varphi_1, \dots, \varphi_N)$  of period  $T = 2\pi b_0/\omega$ ,  $b_0 = b_1 \dots b_\nu$ , with  $\varphi_{2j-1} = -\bar{\varphi}_{2j}$ ,  $j = 1, \dots, \mu$ ,  $\varphi_j$  real,  $j = j = 2\mu + 1, \dots, \mu + n$ , and

$$\forall \{e^{-i\tau_j t} \varphi_{2j-1}(t)\} = c_j, \forall \{e^{i\tau_j t} \varphi_{2j}(t)\} = -\bar{c}_j, \quad j = 1, \dots, \nu,$$

$$m(\varphi_j(t)) = c_j, \quad j = \mu + 1, \dots, \mu + n.$$

Then  $z(t) \in \Omega$ , and we may note that, for every  $\varphi \in \Omega$ , the  $N$  expressions  $(2i\tau_j)^{-1}(\varphi_{2j-1} + \varphi_{2j})$ ,  $(2i\tau_j)^{-1}(\rho_{j1}\varphi_{2j-1} + \rho_{j2}\varphi_{2j})$ ,  $j = 1, \dots, \mu$ ;  $\varphi_j$ ,  $j = 2\mu + 1, \dots, \mu + n$ , are all real, and thus the functions  $q_j(\varphi(u), u, \varepsilon)$ ,  $j = 1, \dots, N$ , are defined, provided the absolute values of the same expressions are all below some prescribed constant  $R' > 0$ . This certainly occurs if  $|\varphi_j| \leq R$ ,  $j = 1, \dots, N$ , for some  $R > 0$ . The transformation  $\mathfrak{z}$  analogous to the one of §2 is then defined for all  $\varphi \in \Omega_R$ ,  $\Omega_R = \{\varphi \in \Omega, |\varphi_j| \leq R, j = 1, \dots, n + \mu\}$  and  $\mathfrak{z}\Omega_R \subset \Omega$  as in §2. As in §2, there is a sphere  $\Omega_0$  about  $z(t)$  in  $\Omega$  with  $\mathfrak{z}(\Omega_0) \subset \Omega_0$ . The transformation  $\psi = \mathfrak{z}\varphi$  is defined by the following relations:

$$\psi_{2j-1} = c_j e^{i\tau_j t} + \varepsilon e^{i\tau_j t} \int e^{-i\tau_j u} \{q_{2j-1}(\varphi(u), u, \varepsilon) - d_{2j-1}\varphi_{2j-1}(u)\} du,$$

$$\psi_{2j} = -\bar{c}_j e^{-i\tau_j t} + \varepsilon e^{-i\tau_j t} \int e^{i\tau_j u} \{q_{2j}(\varphi(u), u, \varepsilon) - d_{2j}\varphi_{2j}(u)\} du,$$

$$j = 1, \dots, \nu,$$

$$\begin{aligned}
 (5.6) \quad \psi_{2j-1} &= \varepsilon e^{\rho_{j1}t} \int e^{-\rho_{j1}u} q_{2j-1}[\varphi(u), u, \varepsilon] du, \\
 \psi_{2j} &= \varepsilon e^{\rho_{j2}t} \int e^{-\rho_{j2}u} q_{2j}[\varphi(u), u, \varepsilon] du, \quad j = v+1, \dots, \mu,
 \end{aligned}$$

$$(5.7) \quad \psi_j = e^{-\beta_j t} \int e^{\beta_j u} q_j[\varphi(u), u, \varepsilon] du, \quad j = 2\mu+1, \dots, \mu+r,$$

$$(5.8) \quad \psi_j = c_j + \varepsilon \int \{q_j[\varphi(u), u, \varepsilon] - d_j \psi_j(u)\} du, \quad j = \mu+r+1, \dots, \mu+n,$$

where  $D = D[\varphi] = (d_1, \dots, d_{2v}, 0, \dots, 0, d_{\mu+r+1}, \dots, d_{\mu+n})$ , and

$$\begin{aligned}
 c_j d_{2j-1} &= \eta_j \left\{ e^{-1\tau_j u} q_{2j-1}[\varphi(u), u, \varepsilon] \right\}, \\
 (5.9) \quad -\bar{c}_j d_{2j} &= \eta_j \left\{ e^{1\tau_j u} q_{2j}[\varphi(u), u, \varepsilon] \right\}, \quad j = 1, \dots, v, \\
 c_j d_j &= \eta_j q_j[\varphi(u), u, \varepsilon], \quad j = \mu+r+1, \dots, \mu+n.
 \end{aligned}$$

(5.1). LEMMA. For every  $\varphi \in \Omega_R$  we have  $\psi \in \Omega$ ,

and  $d_{2j-1} = \bar{d}_{2j}$ ,  $j = 1, \dots, v$ ,  $d_j$  real,

$j = \mu+r+1, \dots, \mu+n$ .

PROOF. We have  $q_{2j-1} = f_j$ ,  $q_{2j} = -f_j$  real,  $j = 1, \dots, v$ , and hence  $q_{2j-1} = -q_{2j}$ , while  $e^{1\tau_j u}$ ,  $e^{-1\tau_j u}$  are complex conjugates. By (5.9) we conclude that  $d_{2j-1} = \bar{d}_{2j}$ , and then the integrands in (5.5) are complex conjugate and of opposite signs. Thus, also their unique primitives of mean value zero have the same property, and (5.5) implies  $\psi_{2j-1} = -\bar{\psi}_{2j}$ ,  $j = 1, \dots, v$ .

Since  $\rho_{j2} = \bar{\rho}_{j1}$ ,  $j = v+1, \dots, \mu$ , we can repeat on relations (5.6) the same reasoning above and thus  $\psi_{2j-1} = -\bar{\psi}_{2j}$ ,  $j = v+1, \dots, \mu$ . Since  $\beta_j$  is real,  $q_j$  is real,  $j = 2\mu+1, \dots, \mu+r$ , by (5.7) we conclude that  $\psi_j$  is real,  $j = 2\mu+1, \dots, \mu+r$ . Finally, for  $j = \mu+r+1, \dots, \mu+n$ , relations (5.9) and (5.8) assure that  $d_j$  and  $\psi_j$  are real,  $j = \mu+r+1, \dots, \mu+n$ . Thereby, (5.1) is proved.

By (5.1) we conclude that  $\exists \Omega_R \subset \Omega$  and, as in §§2, 3, there is a sphere  $\Omega_0$  about  $z(t)$  in  $\Omega_R$  with  $\Omega_0 \subset \Omega_0$ , for  $\varepsilon > 0$  sufficiently small. A fixed element  $y(t) = (y_1, \dots, y_N)$  exists in  $\Omega_0$ ,  $\exists y = y$ , and

$D[y]$  verifies the same relations of (5.1).

We shall consider now the determining equations of §§1, 2, 3, which are in number of  $2v + (n - r)$ , namely

$$\begin{aligned} \rho_{j1} &= i\tau_j - \varepsilon d_{2j-1}, \quad \rho_{j2}(\varepsilon) = -i\tau_j - \varepsilon d_{2j}, & j &= 1, \dots, v, \\ \beta_j &= \varepsilon d_{\mu+j}, & j &= r+1, \dots, n. \end{aligned}$$

By the remark above we conclude that for every  $j = 1, \dots, v$ , the two equations above are equivalent, and thus we have actually only  $v + (n - r)$  determining equations

$$\begin{aligned} \rho_{j1} &= i\tau_j - \varepsilon d_{2j-1}, & j &= 1, \dots, v, \\ \beta_j &= \varepsilon d_{\mu+j}, & j &= r+1, \dots, n. \end{aligned} \quad (5.10)$$

We shall write  $c_j = \lambda_j e^{i\theta_j}$ ,  $j = 1, \dots, v$ ,  $\eta_j = c_{\mu+r+j}$ ,  $j = 1, \dots, n - r$ ,  $\lambda_j \neq 0$ ,  $\eta_j \neq 0$ ,  $\theta_j$  all real,  $\theta_j \pmod{2\pi}$ ,  $\lambda = (\lambda_1, \dots, \lambda_v)$ ,  $\theta = (\theta_1, \dots, \theta_v)$ ,  $\eta = (\eta_1, \dots, \eta_{n-r})$ ,

$$\begin{aligned} d_{2j-1} &= P_j + iQ_j, \quad P_j, Q_j \text{ real}, & j &= 1, \dots, v, \\ d_{\mu+r+j} &= R_j, \quad R_j \text{ real}, & j &= 1, \dots, n - r, \end{aligned}$$

where  $P_j = P_j(a, b, \lambda, \theta, \eta, \omega, \varepsilon)$ ,  $Q_j = Q_j(\dots)$ ,  $R_j = R_j(\dots)$ . Then equations (5.10) reduce to the following  $2v + (n - r)$  real equations

$$\begin{aligned} \varepsilon P_j &= \alpha_j(\varepsilon), & j &= 1, \dots, v, \\ \varepsilon Q_j &= a_j b_j^{-1} \omega - \gamma_j(\varepsilon), & j &= 1, \dots, v, \\ \varepsilon R_j &= \beta_{r+j}(\varepsilon), & j &= 1, \dots, n - r, \end{aligned} \quad (5.11)$$

where  $\alpha_j(0) = 0$ ,  $a_j b_j^{-1} \omega - \gamma_j(0) = 0$ ,  $j = 1, \dots, v$ ,  $\beta_{r+j}(0) = 0$ ,  $j = 1, \dots, n - r$ . Equations (5.11) are the  $2v + (n - r)$  determining equations in real form. By (5.9) we have

$$\begin{aligned} P_j + iQ_j &= \lambda_j^{-1} e^{-i\theta_j} \left\{ e^{-i\tau_j u} d_{2j-1} \right\}, & j &= 1, \dots, v, \\ R_j &= \eta_j^{-1} d_{\mu+r+j}, & j &= 1, \dots, n - r. \end{aligned}$$

and, by obvious manipulations, also

$$\begin{aligned} P_j &= \lambda_j^{-1} [\cos \theta_j \mathfrak{M} \{f_j \cos a_j b_j^{-1} \omega\} - \sin \theta_j \mathfrak{M} \{f_j \sin a_j b_j^{-1} \omega\}] , \\ Q_j &= \lambda_j^{-1} [-\sin \theta_j \mathfrak{M} \{f_j \cos a_j b_j^{-1} \omega\} - \cos \theta_j \mathfrak{M} \{f_j \sin a_j b_j^{-1} \omega\}] , \\ & j = 1, \dots, \nu , \\ R_j &= \eta_j^{-1} \mathfrak{M} \{f_{r+j}\} , & j = 1, \dots, n - r , \end{aligned}$$

where the  $n + \mu + 2$  arguments of  $f_j$  are

$$\begin{aligned} x_j(u) &= (21\gamma_j)^{-1} (\gamma_{2j-1} + \gamma_{2j}), \quad j = 1, \dots, \mu; \quad x_j(u) = \gamma_{\mu+j} , \\ & j = \mu + 1, \dots, n ; \\ (5.12) \quad x_j^1(u) &= (21\gamma_j)^{-1} (\rho_{j1}\gamma_{2j-1} + \rho_{j2}\gamma_{2j}), \quad j = 1, \dots, \mu; \quad u, \varepsilon . \end{aligned}$$

By the continuity, with respect to  $\varepsilon$  proved in (4.11), we can take the limit of both conditions (5.11) and solution (5.12) as  $\varepsilon \longrightarrow 0$ . By using the notation  $x_j(t, \varepsilon)$ ,  $P_j(\varepsilon)$ , etc., we have

$$\begin{aligned} x_j(t, 0) &= \lambda_j \tau_j^{-1} \sin(\tau_j t + \theta_j), & j = 1, \dots, \nu , \\ (5.13) \quad x_j(t, 0) &= 0, \quad j = \nu + 1, \dots, \mu, \quad \text{and} \quad j = \mu + 1, \dots, r , \\ x_j(t, 0) &= \eta_{j-r} , & j = r + 1, \dots, n , \end{aligned}$$

and

$$\begin{aligned} x_j^1(t, 0) &= \lambda_j \cos(\tau_j t + \theta_j), & j = 1, \dots, \nu , \\ (5.14) \quad x_j^1(t, 0) &= 0, & j = \nu + 1, \dots, \mu . \end{aligned}$$

Also,

$$\begin{aligned} P_{j0} &= P_j(0) = (T\lambda_j)^{-1} \left[ \cos \theta_j \int_0^T f_j(\dots, u, 0) \cos a_j b_j^{-1} \omega u \, du - \sin \theta_j \int_0^T f_j \right. \\ & \quad \left. (\dots, u, 0) \sin a_j b_j^{-1} \omega u \, du \right] , \\ (5.15) \quad Q_{j0} &= Q_j(0) = (T\lambda_j)^{-1} \left[ -\sin \theta_j \int_0^T f_j(\dots, u, 0) \cos a_j b_j^{-1} \omega u \, du - \cos \theta_j \int_0^T f_j \right. \\ & \quad \left. (\dots, u, 0) \sin a_j b_j^{-1} \omega u \, du \right] , \quad j = 1, \dots, \nu, \end{aligned}$$

$$R_{j0} = R_j(0) = (T\eta_j)^{-1} \int_0^T f_{r+j}(\dots, u, 0) du, \quad j = 1, \dots, n-r,$$

where the missing  $n + \mu$  arguments in the  $f$ 's are the ones listed in (5.13) and (5.14). We shall use the notation  $\alpha = (\alpha_1, \dots, \alpha_\nu)$ ,  $\gamma = (\gamma_1, \dots, \gamma_\nu)$ ,  $\beta = (\beta_{r+1}, \dots, \beta_n)$ . The previous considerations prove the following

(5.11). THEOREM. If for some  $a, b, \lambda(\varepsilon), \theta(\varepsilon), \eta(\varepsilon), \alpha(\varepsilon), \gamma(\varepsilon), \beta(\varepsilon)$ , all real and for  $\varepsilon > 0$  sufficiently small, equations (5.11) hold, then system (5.1) has a real periodic solution of the form

$$\begin{aligned} x_j(t, \varepsilon) &= \lambda_j \tau_j^{-1} \sin(\tau_j t + \theta_j) + o(\varepsilon), & j &= 1, \dots, \nu, \\ (5.16) \quad x_j(t, \varepsilon) &= o(\varepsilon), & j &= \nu + 1, \dots, r, \\ x_j(t, \varepsilon) &= \eta_{j-r} + o(\varepsilon), & j &= r + 1, \dots, n. \end{aligned}$$

In particular, for  $\varepsilon \geq 0$  sufficiently small, given  $a, b, \lambda(\varepsilon), \theta(\varepsilon), \eta(\varepsilon)$ , there are always  $\alpha(\varepsilon), \gamma(\varepsilon), \beta(\varepsilon)$  such that equations (5.11) are satisfied and system (5.1) has a real periodic solution of the form (5.16).

We shall now consider the more usual question as to whether given  $\alpha, \gamma, \beta$  it is possible to define  $\lambda = (\lambda_1, \dots, \lambda_\nu)$ ,  $\theta = (\theta_1, \dots, \theta_\nu)$ ,  $\eta = (\eta_1, \dots, \eta_{n-r})$  in such a way that equations (5.11) are satisfied and system (5.1) has a periodic solution of the form (5.16). Let us suppose that the derivatives  $\alpha_j'(0), \gamma_j'(0), \beta_{r+j}'(0)$  exist and are finite. By dividing relations (5.11) by  $\varepsilon$  and taking the limit as  $\varepsilon \rightarrow 0$  we have

$$\begin{aligned} P_{j0}(a, b, \lambda, \theta, \eta, \omega, 0) &= \alpha_j'(0), & j &= 1, \dots, \nu, \\ (5.17) \quad Q_{j0}(a, b, \lambda, \theta, \eta, \omega, 0) &= -\gamma_j'(0), & j &= 1, \dots, \nu, \\ R_{j0}(a, b, \lambda, \theta, \eta, \omega, 0) &= \beta_{r+j}'(0), & j &= 1, \dots, n-r. \end{aligned}$$

We shall suppose that these relations are satisfied by some  $\lambda_0 = (\lambda_{j0}, j = 1, \dots, \nu)$ ,  $\theta_0 = (\theta_{j0}, j = 1, \dots, \nu)$ ,  $\eta_0 = (\eta_{j0}, j = 1, \dots, n-r)$ .

If we suppose that the functions  $f_j$  have continuous first partial derivatives with respect to the variables  $x_1, \dots, x_n, x_1', \dots, x_\mu'$ . Then by (§4, Note 1) the functions  $y_j$  have continuous partial derivatives with respect to all  $\lambda_j, \theta_j, \eta_j$ , and the same holds for the  $P_j, Q_j, R_j$ . If equations (5.17) hold for  $\lambda_0, \theta_0, \eta_0$ , and

$$(5.18) \quad \partial(P, Q, R)/\partial(\lambda, \theta, \eta) \neq 0$$

for the same  $\lambda_0, \theta_0, \eta_0$ , then equations (5.11) have a solution  $\lambda(\varepsilon)$ ,  $\theta(\varepsilon)$ ,  $\eta(\varepsilon)$  in a neighborhood of  $\lambda_0, \theta_0, \eta_0$  for all  $\varepsilon \geq 0$  sufficiently small. This proves the following theorem which is essentially of the type usually proved by Poincaré periodicity conditions.

(5.111). THEOREM. If real  $\lambda_0, \theta_0, \eta_0$  exist such that equations (5.17) and (5.18) hold, then for  $\varepsilon \geq 0$  sufficiently small system (5.1) has a real periodic solution of period  $T = 2\pi b_0/\omega$  of the type

$$(5.19) \quad \begin{aligned} x_j(t, \varepsilon) &= \lambda_j \tau_j^{-1} \sin(\tau_j t + \theta_j) + o(\varepsilon), & j &= 1, \dots, v, \\ x_j(t, \varepsilon) &= o(\varepsilon), & j &= v+1, \dots, r, \\ x_j(t, \varepsilon) &= \eta_{j-r} + o(\varepsilon), & j &= r+1, \dots, n. \end{aligned}$$

REMARK. The differentiability condition concerning the functions  $f_j$  used above to prove (5.111) requires more than needed as we shall see below (cf. (5.iv), (5.vi), etc.). The present first remark may be of interest. For instance, let us require that only  $f(x, t, 0)$  has continuous first partial derivatives with respect to  $x$  (and  $f(x, t, \varepsilon)$  is continuous in  $x$  and  $\varepsilon$ ). Then the functions  $P_{j0}, Q_{j0}, R_{j0}$  have continuous first partial derivatives with respect to  $\lambda, \theta, \eta$ . If the Jacobian of the  $P_{j0}, Q_{j0}, R_{j0}$  with respect to  $\lambda, \theta, \eta$  is  $\neq 0$  at  $(\lambda_0, \theta_0, \eta_0)$ , then, for a conveniently small solid sphere  $\sigma$  of center  $(\lambda_0, \theta_0, \eta_0)$ , the mapping  $u_j = P_{j0} - \alpha_j^1(0)$ ,  $v_j = Q_{j0} + \gamma_j^1(0)$ ,  $w_j = R_{j0} - \beta_{r+j}^1(0)$ ,  $(\lambda, \theta, \eta) \in \sigma$ , has topological index one with respect to the origin. By force of continuity only, this holds true also for the (perturbed) continuous mapping  $U_j = P_j(\lambda, \theta, \eta, \varepsilon) - \varepsilon^{-1} \alpha_j^1(\varepsilon)$ ,  $V_j = Q_j + \varepsilon^{-1}(\gamma_j(\varepsilon) - \alpha_j b_j^{-1} \omega)$ ,  $W_j = R_j - \varepsilon^{-1} \beta_{r+j}^1(\varepsilon)$ ,  $(\lambda, \theta, \eta) \in \sigma$ , provided  $\varepsilon \geq 0$  is sufficiently small. We shall discuss this point in more detail in the papers mentioned in the Introduction. For the present argument, see, e.g., J. Leray and J. Schauder, Topologie et équations fonctionnelles, Ann. Ec. Norm. (3) 51, 45-78, 1934.

Let us consider now real systems (5.1) where the functions  $f_j$  do not depend on  $t$ , i.e., autonomous systems,

$$\begin{aligned} x_j'' + 2\alpha_j x_j' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', \varepsilon), & j &= 1, \dots, \mu, \\ x_j' + \beta_j x_j &= \varepsilon f_j(x, x', \varepsilon), & j &= \mu+1, \dots, n, \end{aligned}$$

satisfying the hypotheses listed at the beginning of §5, where now  $\omega$  in condition (K) is actually arbitrary. As mentioned in §2 and §4 we shall suppose  $\omega \in U$ , where  $U$  is a sufficiently small neighborhood of any number  $\omega_0$  for which  $\sigma_j(0) = a_j \omega_0 / b_j$ ,  $j = 1, \dots, \nu$ ,  $\rho_{j1}(0)$ ,  $\rho_{j2}(0) \neq \text{im} \omega_0 / b_0$ ,  $j = \nu + 1, \dots, n$ ,  $m = 0, \pm 1, \dots$ , and we may expect to satisfy (5.11) by a convenient function  $\omega = \omega(\varepsilon)$  with  $\omega(0) = \omega_0$ . Thus

(5.1v). The same as (5.1i) for system (5.20) with  $\omega(\varepsilon)$  added to the list of parameters.

For every solution  $x(t)$  of (5.20) also  $x(t + \theta)$  is a solution for every arbitrary constant phase  $\theta$ . Thus we may expect that (at least) one of the phases  $\theta_j$ ,  $j = 1, \dots, \nu$ , considered above, say  $\theta_1$ , remains arbitrary and we put  $\theta_1 = 0$ . Note that now the  $2\nu + (n - r)$  expressions  $P, Q, R$  are still functions of  $2\nu + (n - r)$  arbitrary parameters, namely  $\lambda_j$ ,  $j = 1, \dots, \nu$ ,  $\theta_j$ ,  $j = 2, \dots, \nu$ ,  $\eta_j$ ,  $j = 1, \dots, n - r$ , and  $\omega \in U$ . We may now denote by  $\theta$  the  $(\nu - 1)$ -vector  $(\theta_2, \dots, \theta_\nu)$ . Again, as above, let us suppose that the functions  $f_j(x, x', \varepsilon)$  have continuous first partial derivatives with respect to the variables  $x_1, \dots, x_n, x'_1, \dots, x'_\mu$ . If equations (5.17) hold for some  $\lambda_0, \theta_0, \eta_0$  and  $\omega = \omega_0$ , and the Jacobian

$$(5.21) \quad \partial(P, Q, R) / \partial(\lambda, \theta, \mu, \omega) \neq 0$$

for the same  $\lambda_0, \theta_0, \eta_0, \omega_0$ , and  $\varepsilon = 0$ , then equations (5.11) have a solution  $\lambda(\varepsilon), \theta(\varepsilon), \eta(\varepsilon), \omega(\varepsilon)$ , in some neighborhood of  $\lambda_0, \theta_0, \eta_0$  and  $\omega \in U$  for all  $\varepsilon \geq 0$  sufficiently small. In analogy with (5.11i) we have now the Poincaré type theorem for autonomous systems:

(5.v). THEOREM. If real  $\lambda_0, \theta_0, \mu_0, \omega_0$  exist such that equations (5.17), (5.21) hold, then for  $\varepsilon \geq 0$  sufficiently small system (5.20) has a real periodic solution of period  $T = 2\pi b_0 / \omega$  of the type (5.19) with  $\theta_1 = 0$  (and then also all other solutions with  $t$  replaced by  $t + \theta$ ,  $\theta$  arbitrary).

We proceed to prove a theorem of a more general type (others will be given in §6). For the sake of simplicity we shall consider a system (5.20) with  $\nu = 1, r = n, \alpha_1 = 0$ , i.e., an autonomous system of the form

$$(5.22) \quad \begin{aligned} x_1'' + \sigma_1^2 x_1 &= \varepsilon f_1(x, x', \varepsilon), \\ x_j'' + 2\alpha_j x_j + \sigma_j^2 x_j^2 &= \varepsilon f_j(x, x', \varepsilon), & j = 2, \dots, \mu, \\ x_j' + \beta_j x_j &= \varepsilon f_j(x, x', \varepsilon), & j = \mu + 1, \dots, n, \end{aligned}$$

where  $1 = \nu \leq \mu \leq n$ ,  $x = (x_1, \dots, x_n)$ ,  $x' = (x'_1, \dots, x'_\mu)$ ,  $\sigma_j(\varepsilon) > 0$ ,  $\alpha_j(\varepsilon)$ ,  $\beta_j(\varepsilon)$  real functions of  $\varepsilon$ ,  $\beta_j(\varepsilon) \neq 0$ , (or constants),  $0 \leq \varepsilon \leq \varepsilon_0$ . Also, we assume that  $a = b = 1$ , and hence  $\sigma_1(0) = \omega_0$ ,  $\rho_{j1}(0)$ ,  $\rho_{j2}(0) \neq 0$ ,  $j = 2, \dots, \mu$ ,  $m = 0, \pm 1, \pm 2, \dots$ . Thus  $\tau_1 = \omega_0 = \sigma_1(0)$ , and there will be only one  $c$ . The functions  $f_j$  are assumed to satisfy a condition (L). There is only one equation (5.10), namely,  $1\omega - \varepsilon d_1(c, \omega, \varepsilon) = 1\sigma_1(\varepsilon)$ , or, in real form, two equations (5.11),  $P_1 = 0$ ,  $\omega - \varepsilon Q = \sigma_1(\varepsilon)$ . Since there is only one phase  $\theta_1$  it is not restrictive to take it equal to zero, i.e.,  $c = \lambda$ ,  $0 < r_1 \leq \lambda \leq r_2 < R$ ,  $\lambda$  real and positive. Thus the two determining equations are actually

$$(5.23) \quad P_1(\lambda, \omega, \varepsilon) = 0, \quad \omega - \varepsilon Q_1(\lambda, \omega, \varepsilon) = \sigma_1(\varepsilon).$$

Note that

$$P_1(\lambda, \omega, 0) = (T\lambda)^{-1} \int_0^T f_1[\lambda\omega^{-1} \sin \omega t, 0, \dots, 0, \lambda \cos \omega t, 0, \dots, 0; 0] \cos \omega t \, dt,$$

(5.24)

$$Q_1(\lambda, \omega, 0) = - (T\lambda)^{-1} \int_0^T f_1[\lambda\omega^{-1} \sin \omega t, 0, \dots, 0, \lambda \cos \omega t, 0, \dots, 0; 0] \sin \omega t \, dt,$$

where  $T = 2\pi/\omega$ . We shall suppose that for some  $\lambda_0$  we have  $P_1(\lambda, \omega_0, 0) = 0$ , and that for two  $\lambda'$ ,  $\lambda''$ ,  $r_1 \leq \lambda' < \lambda_0 < \lambda'' \leq r_2$ , the two numbers  $P_1(\lambda', \omega_0, 0)$ ,  $P_1(\lambda'', \omega_0, 0)$  have opposite signs.

(5.v1). Consider system (5.22) where the functions  $f_j$  are independent of  $t$ , Lipschitzian in  $x$ ,  $x'$  and continuous in  $\varepsilon$ , for  $|x_j| \leq R$ ,  $j = 1, \dots, n$ ,  $|x'_j| \leq R$ ,  $j = 1, \dots, \mu$ ,  $0 \leq \varepsilon \leq \varepsilon_0$ . Suppose that  $\sigma_j$ ,  $\alpha_j$ ,  $\beta_j$  are continuous functions of  $\varepsilon$  [or constants] and that  $\sigma_j(0) > 0$ ,  $j = 1, \dots, \mu$ ,  $\beta_j(0) \neq 0$ ,  $j = \mu + 1, \dots, n$ , and either  $\alpha_j(0) \neq 0$ , or  $\alpha_j(0) = 0$ ,  $\sigma_j(0) \neq 0 \pmod{\sigma_1(0)}$ ,  $j = 2, \dots, \mu$ . Let  $\omega_0 = \sigma_1(0)$ , or  $\omega \in [\omega_{01}, \omega_{02}]$ ,  $\lambda \in [r_1, r_2]$ , with  $0 < \omega_{01} < \omega_0 < \omega_{02}$ ,  $0 < r_1 < r_2 < R$ . If there exists  $\lambda'$ ,  $\lambda_0$ ,  $\lambda''$ ,  $r_1 \leq \lambda' < \lambda_0 < \lambda'' \leq r_2$  such that  $P_1(\lambda_0, \omega_0, 0) = 0$ , and  $P_1(\lambda', \omega_0, 0)$ ,  $P_1(\lambda'', \omega_0, 0)$  have opposite signs, then there is an  $\varepsilon_1 > 0$  such that, for every  $\varepsilon$ ,  $0 \leq \varepsilon \leq \varepsilon_1$ , system (5.22) has at least one periodic solution of the form

$$(5.25) \quad \begin{aligned} x_1(t, \varepsilon) &= \lambda(\varepsilon)\omega^{-1}(\varepsilon)\sin \omega(\varepsilon)t + o(\varepsilon), \\ x_j(t, \varepsilon) &= o(\varepsilon), \quad j = 2, \dots, n, \end{aligned}$$

for conveniently chosen  $\omega(\varepsilon) \in [\omega_{01}, \omega_{02}]$ ,  $\lambda(\varepsilon) \in [\lambda', \lambda'']$ . System (5.22) has also any other solution we obtain by replacing  $t$  by  $t + \theta$ ,  $\theta$  an



arbitrary constant.

PROOF. Let us use statement (1.vi). Take  $F_1(\lambda, \omega, \varepsilon) = P_1(\lambda, \omega, \varepsilon)$ , where  $\omega \in U$ , say  $U = [\omega', \omega'']$ ,  $\omega' < \omega_0 < \omega''$ ,  $\lambda \in V = [\lambda', \lambda'']$ ,  $\varepsilon \in [0, \varepsilon_0]$ . Thus  $F_1(\lambda', \omega_0, 0)$ ,  $F_1(\lambda'', \omega_0, 0)$  have opposite signs, and, if we suppose  $U = [\omega', \omega'']$  sufficiently small, also  $F_1(\lambda', \omega, 0)$ ,  $F_1(\lambda'', \omega, 0)$ ,  $\omega \in [\omega', \omega'']$ , have opposite constant signs. Take  $F_2(\lambda, \omega, \varepsilon) = \omega - \sigma_1(\varepsilon) - \varepsilon Q_1(\lambda, \omega, \varepsilon)$ , and note that  $F_2(\omega_0, \lambda, 0) = \omega_0 - \sigma_1(0) = 0$ ,  $F_2(\omega', \lambda, 0) < 0$ ,  $F_2(\omega'', \lambda, 0) > 0$  for all  $\lambda \in [\lambda', \lambda'']$ . By (1.vi), with  $M = [0, \varepsilon_2]$ ,  $K = [\lambda', \lambda''] \times [\omega', \omega'']$ , we conclude that there are a neighborhood  $[0, \varepsilon_2]$  of  $\varepsilon = 0$ , and  $\lambda(\varepsilon) \in [\lambda', \lambda'']$ ,  $\omega(\varepsilon) \in [\omega', \omega'']$  for every  $0 \leq \varepsilon \leq \varepsilon_2$ , such that  $F_1 = F_2 = 0$ . Then (5.vi) is a consequence of (5.iv).

REMARK 1. Theorem (5.vi) is more general than the usual theorems which are proved by Poincaré periodicity condition, since no differentiability property is involved (other theorems will be given in §6).

Note that the condition of (5.vi) is certainly satisfied if  $P_1(\lambda, \omega_0, 0)$  happens to have derivative with respect to  $\lambda$  at  $\lambda = \lambda_0$  and  $\partial P_1 / \partial \lambda \neq 0$  at that point.

Finally, under usual conditions where the functions  $f_j$  have continuous first partial derivatives with respect to  $x_1, \dots, x_n, x_1', \dots, x_\mu'$ , then  $F_1, F_2$  have also continuous first partial derivatives with respect to  $\lambda$  and  $\omega$ , and  $\partial(F_1, F_2) / \partial(\lambda, \omega) = F_{1\lambda} F_{2\omega} - F_{1\omega} F_{2\lambda}$ , where  $F_{1\lambda} = P_{1\lambda}$ ,  $F_{2\omega} = 1 + o(\varepsilon)$ ,  $F_{1\omega} = o(1)$ ,  $F_{2\lambda} = o(\varepsilon)$ . Thus we have

$$\partial(F_1, F_2) / \partial(\lambda, \omega) = \partial P_1 / \partial \lambda + o(\varepsilon).$$

The condition just mentioned assures that this Jacobian is  $\neq 0$  for  $\lambda = \lambda_0$ ,  $\omega = \omega_0$ ,  $\varepsilon = 0$ , and finally assures the uniqueness and continuity of  $\omega(\varepsilon)$ ,  $\lambda(\varepsilon)$  for  $0 \leq \varepsilon \leq \varepsilon_1$ , and some  $\varepsilon_1 > 0$  sufficiently small. Note that, under the same conditions, the inequality

$$\int_0^T f_{1x_1'} [\lambda \omega^{-1} \sin \omega t, 0, \dots, 0, \lambda \cos \omega t, 0, \dots, 0; 0] dt < 0,$$

or the equivalent Krylov-Bogolyubov condition  $\partial P_1 / \partial \lambda < 0$  [in both  $\lambda = \lambda_0$ ,  $\omega = \omega_0$ ,  $\varepsilon = 0$ ], together with  $\alpha_j < 0$ ,  $j = 2, \dots, \mu$ ,  $\beta_j < 0$ ,  $j = \mu + 1, \dots, n$ , assure that the same periodic solution is asymptotically orbitally stable. [For these and more general stability conditions see J. K. Hale, 91, and E. W. Thompson, 11].

REMARK 2. The following considerations may help to understand the

generality of (5.vi). The real numbers (or functions of  $\varepsilon$ )  $\beta_j(\varepsilon)$ ,  $j = \mu + 1, \dots, n$ ,  $\alpha_j(\varepsilon)$ ,  $\sigma_j(\varepsilon)$ ,  $j = 2, \dots, \mu$ , have no bearing on  $P_1(\lambda, \omega, 0)$ . We require for them only  $\beta_j \neq 0$ ,  $j = \mu + 1, \dots, n$ ,  $\sigma_j(0) > 0$ ,  $j = 1, \dots, n$ , and either  $\alpha_j(0) \neq 0$ , or  $\alpha_j(0) = 0$ ,  $\sigma_j(0) \neq 0 \pmod{\sigma_1(0)}$ ,  $j = 2, \dots, \mu$ ,  $\sigma_1(0) = \omega_0$ . The functions  $f_j(x, x', \varepsilon)$ ,  $j = 1, \dots, n$ , are supposed to be continuous in  $x, x', \varepsilon$ , and  $f_2, \dots, f_n$  have no bearing on  $P_1(\lambda, \omega, 0)$ . If we put

$$Z_1(x_1, x_1') = f_1(x_1, 0, \dots, 0, x_1', 0, \dots, 0; 0),$$

$$g_1(x, x', \varepsilon) = f_1(x, x', \varepsilon) - Z_1(x_1, x_1'),$$

then  $f_1 = Z_1 + g_1$ ,  $g_1(x_1, 0, \dots, 0, x_1', 0, \dots, 0; 0) = 0$  and  $g_1$  has no bearing on  $P_1(\lambda, \omega, 0)$ . Finally, if we put

$$Z_{11} = 4^{-1} [Z_1(x_1, x_1') + Z_1(-x_1, x_1') - Z_1(x_1, -x_1') - Z_1(-x_1, x_1')],$$

and we define  $Z_{12}, Z_{13}, Z_{14}$  analogously by changing the signs  $(+, -, -)$  into  $(+, +, +)$ ,  $(-, -, +)$ ,  $(-, +, -)$ , then  $Z_{11}[Z_{12}]$  is even in  $x_1$  and odd [even] in  $x_1$ ,  $Z_{13}[Z_{14}]$  is odd in  $x_1$  and odd [even] in  $x_1$ . We have  $f_1 = Z_1 + g_1 = Z_{11} + Z_{12} + Z_{13} + Z_{14} + g_1$ , and  $Z_{12}, Z_{13}, Z_{14}, g_1$  have no bearing on  $P_1(\lambda, \omega, 0)$ . We have

$$P_1(\lambda, \omega, 0) = (T\lambda)^{-1} \int_0^T Z_{11}(\lambda\omega^{-1} \sin \omega t, \cos \omega t) dt,$$

where  $T = 2\pi/\omega$ , and  $\omega_0 = \sigma_1(0)$ . In the particular case where  $Z_{11}$  is a polynomial in  $x_1, x_1'$ , hence necessarily of the form

$$Z_{11} = \sum a_{hk} x_1^{2h} x_1'^{2k-1}, \quad h \geq 0, k \geq 1,$$

then

$$P_1(\lambda, \omega_0, 0) = 2^{-2} \sum a_{hk} \frac{(2h)!(2k)!}{h!k!(h+k)!} \left( \frac{\lambda}{2\sigma_1} \right)^{2h+2k-2}$$

where  $\sigma_1 = \sigma_1(0)$ . If, for instance,  $a_{01}$  and the coefficient of the maximal power of  $\lambda$  are ( $\neq 0$  and) of opposite signs, then certainly (5.22) has a periodic solution [cf. J. K. Hale, 9c, for the analytic case].

EXAMPLE 1. The real system (5.22), without restricting its generality, can be written as follows:

$$\begin{aligned}
 x_1'' + \sigma_1^2 x_1 &= \varepsilon [Z_{11}(x_1, x_1') + Z_{12}(x_1, x_1') + Z_{13}(x_1, x_1') \\
 &\quad + Z_{14}(x_1, x_1') + g_1(x, x', \varepsilon)], \\
 (5.24) \quad x_j'' + 2\alpha_j x_j' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', \varepsilon), \quad j = 2, \dots, \mu, \\
 x_j' + \beta_j x_j &= \varepsilon f_j(x, x', \varepsilon), \quad j = \mu + 1, \dots, n,
 \end{aligned}$$

where  $Z_{11}, \dots, Z_{14}, g_1, f_2, \dots, f_n$  are Lipschitzian in  $x, x', \varepsilon$  where  $Z_{11}[Z_{12}]$  is even in  $x_1$  and odd [even] in  $x_1'$ ,  $Z_{13}[Z_{14}]$  is odd in  $x_1$  and odd [even] in  $x_1'$ , where  $g_1(x_1, 0, \dots, 0, x_1', 0, \dots, 0; 0) = 0$ , where  $\alpha_j(\varepsilon), \sigma_j(\varepsilon), \beta_j(\varepsilon)$  are continuous functions of  $\varepsilon$  [or constants] with  $\sigma_j(0) > 0, j = 1, \dots, \mu, \beta_j(0) \neq 0, j = \mu + 1, \dots, n$ , and either  $\alpha_j(0) \neq 0$ , or  $\alpha_j(0) = 0, \sigma_j(0) \neq 0 \pmod{\sigma_1(0)}, \sigma_1(0) = \omega_0, j = 2, \dots, n$ .

If we take  $Z_{11} = (1 - x_1^2)x_1'$ , then  $P_1 = (1/2)(1 - \lambda^2/\sigma_1^2)$ . Thus for every  $\varepsilon, 0 \leq \varepsilon \leq \varepsilon_1$ , and some  $\varepsilon_1 > 0$ , system (5.24) has a real periodic solution of the form

$$(5.25) \quad x_1 = \lambda \omega^{-1} \sin(\omega t + \theta) + o(\varepsilon), \quad x_j = o(\varepsilon), \quad j = 2, \dots, n,$$

with  $\lambda = 2\sigma_1(0) + o(\varepsilon)$ . For  $Z_{12} = Z_{13} = Z_{14} = 0, g_1 = 0, \mu = n = 1$ , we have, as a particular case, the well known Van der Pol equation  $x_1'' + \sigma_1^2 x_1 = \varepsilon(1 - x_1^2)x_1'$ . By (9.1) we can see that the solution (5.25) is asymptotically orbitally stable.

If we take  $Z_{11} = (1 - x_1^2 - x_1'^2)x_1'$ , then  $P_1 = (1/2)(1 - \lambda^2/\sigma_1^2)$  and (5.24) has a real periodic solution of the form above with  $\lambda = \sigma_1(0) + o(\varepsilon), \omega = \sigma_1(0) + o(\varepsilon)$ .

EXAMPLE 2. As in Example 1 with  $Z_{11} = (1 - |x_1|)x_1'$ . Then

$$P_1 = (T\lambda)^{-1} \int_0^T (1 - \lambda \omega^{-1} |\sin \omega t|) \cos^2 \omega t \, dt,$$

and  $P_1(\lambda, \omega_0, 0) = (1/2)(1 - \lambda/\pi\sigma_1)$ . Hence, system (5.24) has a periodic solution of the form (5.25) with  $\lambda = \pi\sigma_1 + o(\varepsilon), \omega = \sigma_1 + o(\varepsilon)$ .

EXAMPLE 3. The real system

$$x'' + x - \varepsilon(1 - x^2 - y^2)x' = \varepsilon f_1(x, y, y') + \varepsilon g_1(x, x', y, y')y,$$

$$y'' + 2y - \varepsilon(1 - x^2 - y^2)y' = \varepsilon f_2(x, x', y) + \varepsilon g_2(x, x', y, y')x,$$

where  $f_1(-x, y, y') = -f_1(x, y, y'), f_2(x, x', -y) = -f_2(x, x', y)$ .

There are two periodic solutions of this system given by

$$x = \lambda \sin (\omega t + \theta) + o(\varepsilon), \quad y = o(\varepsilon), \quad \omega = 1 + o(\varepsilon), \quad \lambda = 2 + o(\varepsilon) ;$$

$$x = o(\varepsilon), \quad y = \lambda \sin (\omega t + \theta), \quad \omega = 2^{1/2} + o(\varepsilon), \quad \lambda = 2 + o(\varepsilon) .$$

[See J. K. Hale, 9c, for the analytic case. In 9i it is proved that both solutions are asymptotically orbitally stable.]

The following further examples have been studied among others, in the lines above, by R. A. Gambill and J. K. Hale [8] with emphasis on the existence of harmonic and subharmonic solutions. In [1], [3], and [9i] the stability of some of these solutions has been discussed. In all examples  $\varepsilon$  is a small parameter.

$$4. \quad x'' + \sigma^2 x = B \cos 2\omega t + \varepsilon \alpha \cos 2\omega t \cdot x + \varepsilon b x^3 ;$$

$$5. \quad x'' + \sigma^2 y = B \cos \omega t + \varepsilon x^3 ;$$

$$6. \quad x'' + x = \varepsilon(1 - x^{2n})x' + \varepsilon p \omega \cos (\omega t + \alpha) ;$$

$$7. \quad \begin{cases} x'' + \sigma_1^2 x = \varepsilon \alpha x + \varepsilon A \cos t \cdot x + \varepsilon \beta x^3 + \varepsilon \gamma x y^2 \\ y'' + \sigma_2^2 y = \varepsilon \delta y + B \cos \omega t \cdot y + \varepsilon \mu y^3 + \varepsilon \nu x^2 y ; \end{cases}$$

$$8. \quad \begin{cases} x'' + 4x = \cos t + \varepsilon[\cos t \cdot x + k y^3 + b y] , \\ y'' + y = \varepsilon[\cos t \cdot y + c x^3] ; \end{cases}$$

$$9. \quad \begin{cases} x'' + \sigma_1^2 x_1 = \varepsilon(\alpha - \gamma y^2)x' + \beta \cos t , \\ y'' + \sigma_2^2 y = \varepsilon(\delta - \rho x^2)y' + r \cos 2t ; \end{cases}$$

Example 4 is a nonlinear Mathieu equation with a large forcing term, 5 is a Duffing equation, 6 is a generalization of the Van der Pol equation, 7 is a system of nonlinear Mathieu equations, 8 is a system of nonlinear Mathieu equations of which one has a large forcing term, 9 is a system of Van der Pol equations.

#### §6. REAL SYSTEMS PRESENTING SYMMETRIES

We shall consider now real systems (5.1) presenting symmetries as defined below. In addition to the hypotheses listed at the beginning of §5, we shall suppose

$$\begin{aligned}\alpha_j &\equiv 0, & j &= 1, \dots, \mu, \\ 0 \leq \nu \leq \mu = r \leq n, & \beta_j \equiv 0, & j &= \mu + 1, \dots, n.\end{aligned}$$

Then we have  $\gamma_j = \sigma_j$ ,  $\rho_{j1} = 1\sigma_j$ ,  $\rho_{j2} = -1\sigma_j$ ,  $j = 1, \dots, \mu$ , and equations (5.2), (5.3) become

$$(6.2) \quad \begin{aligned}y_{2j-1} &= 1\sigma_j x_j + x_j^1, & y_{2j} &= 1\sigma_j x_j - x_j^1, & j &= 1, \dots, \mu, & y_{\mu+j} &= x_j \\ j &= \mu + 1, \dots, n,\end{aligned}$$

$$(6.3) \quad \begin{aligned}x_j &= (21\sigma_j)^{-1} (y_{2j-1} + y_{2j}), & x_j^1 &= 2^{-1} (y_{2j-1} - y_{2j}), \\ j &= 1, \dots, \mu.\end{aligned}$$

Also, if  $u = (x_1, \dots, x_m)$ ,  $v = (x_{m+1}, \dots, x_\mu)$ ,  $w = (x_{\mu+1}, \dots, x_n)$ , hence  $x = (u, v, w)$ ,  $x' = (u', v')$ , for some  $0 \leq m \leq \nu$ , we have

$$(6.4) \quad \begin{aligned}f_j(u, -v, w, -u', v', -t, \epsilon) &= f_j(u, v, w, u', v', t, \epsilon), & j &= 1, \dots, m, \\ f_j(u, -v, w, -u', v', -t, \epsilon) &= -f_j(u, v, w, u', v', t, \epsilon), & j &= m+1, \dots, \mu, \\ f_j(u, -v, w, -u', v', -t, \epsilon) &= -f_j(u, v, w, u', v', t, \epsilon), & j &= \mu+1, \dots, n.\end{aligned}$$

For  $m = 0$  all  $f_j$  are odd with respect to  $(v, u', t)$ . For  $m > 0$  the first  $m$  of the  $f_j$  are even and the remaining ones are odd with respect to  $(v, u', t)$ .

We shall take  $\theta_j = \pi/2$ ,  $j = 1, \dots, m$ ,  $\theta_j = 0$ ,  $j = m+1, \dots, \nu$ , and hence  $c_j = 1\lambda_j$ ,  $j = 1, \dots, m$ ,  $c_j = \lambda_j$ ,  $j = m+1, \dots, \nu$ ,  $\lambda_j$  real; we shall take  $c_j = \eta_{j-\mu-r}$ ,  $j = \mu+r+1, \dots, \mu+n$ . Then we have

$$(6.5) \quad \begin{aligned}P_j &= - (T\lambda_j)^{-1} \int_0^T f_j \sin \tau_j t \, dt, & Q_j &= - (T\lambda_j)^{-1} \int_0^T f_j \cos \tau_j t \, dt, \\ j &= 1, \dots, m, \\ P_j &= (T\lambda_j)^{-1} \int_0^T f_j \cos \tau_j t \, dt, & Q_j &= - (T\lambda_j)^{-1} \int_0^T f_j \sin \tau_j t \, dt, \\ j &= m+1, \dots, \nu, \\ R_j &= (T\eta_j)^{-1} \int_0^T f_{r+j} dt, & j &= 1, \dots, n-r,\end{aligned}$$

where the arguments of the  $f$ 's, besides  $t$  and  $\epsilon$ , are given by the (5.12) of §5, thus, in the present case, by relations analogous to the (6.3).

We shall restrict the space  $\Omega$  to all  $\varphi = (\varphi_1, \dots, \varphi_{\mu+n})$  as in §5 which besides verifying  $\varphi_{2j-1}(t) = -\bar{\varphi}_{2j}(t)$ ,  $j = 1, \dots, \mu$ ,  $\varphi_j$  real,  $j = 2\mu + 1, \dots, \mu + n$ , have the following properties:

$$\varphi_{2j-1}(-t) = -\bar{\varphi}_{2j-1}(t), \quad \varphi_{2j}(-t) = -\bar{\varphi}_{2j}(t), \quad j = 1, \dots, m,$$

$$\varphi_{2j-1}(-t) = \bar{\varphi}_{2j-1}(t), \quad \varphi_{2j}(-t) = \bar{\varphi}_{2j}(t), \quad j = m + 1, \dots, \mu, \quad (6.6)$$

$$\varphi_j(-t) = \varphi_j(t), \quad j = 2\mu + 1, \dots, \mu + n.$$

This is the same as requesting that for  $j = 1, \dots, m$ , all  $\varphi_{2j-1}$ ,  $\varphi_{2j}$  have odd real parts (r.p.) and even imaginary parts (i.p.), that for  $j = m + 1, \dots, \mu$ , all  $\varphi_{2j-1}$ ,  $\varphi_{2j}$  have even r.p. and odd i.p., and that for  $j = 2\mu + 1, \dots, \mu + n$ , all  $\varphi_j$  are real and even. Also the first two lines of (6.6) are equivalent to requesting that for  $j = 1, \dots, m$ , the expressions  $(2i\sigma_j)^{-1}(\varphi_{2j-1} + \varphi_{2j})$  are real and even, and the expression  $2^{-1}(\varphi_{2j-1} - \varphi_{2j})$  are real and odd, while for  $j = m + 1, \dots, \mu$ , the former are real and odd and the latter are real and even.

Because of (6.4) the integrands for the  $R_j$  and  $P_j$  are all real and odd, hence  $P_j \equiv 0$ ,  $j = 1, \dots, v$ ,  $R_j \equiv 0$ ,  $j = 1, \dots, n - \mu$ . As a consequence, for  $j = 2\mu + 1, \dots, \mu + n$ , we have  $d_j = 0$  and, by (5.8), since  $q_j$  is real and odd, we conclude that  $\psi_j$  is real and even.

Relations (5.7) are not present here since  $r = \mu$ .

For  $j = v + 1, \dots, \mu$ , by (5.6), and since  $q_{2j-1}$  is real and odd, and  $\rho_{j1}$  is purely imaginary, we see that the integrand has odd r.p. and even i.p., hence  $\psi_{2j-1}$  has even r.p. and odd i.p. By §5 the same holds for  $\psi_{2j}$ .

For  $j = 1, \dots, m$ , we have  $c_j = i\lambda_j$ ,  $c_j$  is imaginary,  $q_{2j-1}$  is real and even, and by (6.6), the expression under  $\{ \}$  in (5.9) has even r.p. and odd i.p., hence  $c_j d_{2j-1}$  is real and  $d_{2j-1}$  is imaginary. Then the expression under  $\{ \}$  in (5.5) has even r.p. and odd i.p. and the same holds for the integrand. Hence, the integral has odd r.p. and even i.p. and the same holds for  $\psi_{2j-1}$ . By §5 the same is true for  $\psi_{2j}$ .

For  $j = m + 1, \dots, v$ , we have  $c_j = \lambda_j$ ,  $c_j$  is real,  $q_{2j-1}$  is real and odd, and by (6.6) the expression under  $\{ \}$  in (5.9) has odd r.p. and even i.p., hence  $c_j d_{2j-1}$  is imaginary and  $d_{2j-1}$  is imaginary. Then the expression under  $\{ \}$  in (5.5) has odd r.p. and even i.p. and the same holds for the integrand. Hence, the integral has even r.p. and

odd i.p. and the same holds for  $\psi_{2j-1}$ , and finally for  $\psi_{2j}$ .

In connection with §5 we conclude that  $\psi = \tau \varphi \in \Omega$  for every  $\varphi \in \Omega$ , i.e.,  $\tau \Omega_R \subset \Omega$ .

Also, the first  $\nu$  and the last  $n - r$  equations (5.11) are identically satisfied and the second  $\nu$  equations become

$$(6.7) \quad a_j b_j^{-1} \omega - \varepsilon Q_j(a, b, \lambda, \eta, \omega, \varepsilon) = \sigma_j(\varepsilon), \quad j = 1, \dots, \nu,$$

where, for  $\varepsilon = 0$ , we have

$$(6.8) \quad Q_{j0} = -(\tau \lambda_j)^{-1} \int_0^T f_j(\dots, t, 0) \cos a_j b_j^{-1} \omega t \, dt, \quad j = 1, \dots, m,$$

$$Q_{j0} = -(\tau \lambda_j)^{-1} \int_0^T f_j(\dots, t, 0) \sin a_j b_j^{-1} \omega t \, dt, \quad j = m+1, \dots, \nu,$$

and the missing arguments are

$$(6.9) \quad \begin{aligned} x_j(t, 0) &= \lambda_j \tau_j^{-1} \cos \tau_j t, & x_j'(t, 0) &= -\lambda_j \sin \tau_j t, & j &= 1, \dots, m, \\ x_j(t, 0) &= \lambda_j \tau_j^{-1} \sin \tau_j t, & x_j'(t, 0) &= \lambda_j \cos \tau_j t, & j &= m+1, \dots, \nu, \\ x_j(t, 0) &= 0, & x_j'(t, 0) &= 0, & j &= \nu+1, \dots, \mu, \mu = r, \\ x_j(t, 0) &= \eta_{j-r}, & & & j &= r+1, \dots, n. \end{aligned}$$

In connection with §5 we may summarize the preceding results as follows.

Consider the real system

$$(6.10) \quad \begin{aligned} x_j'' + \sigma_j^2(\varepsilon) x_j &= \varepsilon f_j(x, x', t, \varepsilon), & j &= 1, \dots, \mu, \\ x_j' &= \varepsilon f_j(x, x', t, \varepsilon), & j &= \mu+1, \dots, n, \end{aligned}$$

where  $x = (x_1, \dots, x_n)$ ,  $x' = (x_1', \dots, x_n')$ ,  $\sigma_j(\varepsilon) > 0$  are continuous functions of  $\varepsilon$  for  $0 \leq \varepsilon \leq \varepsilon_0$ , and for some integers  $0 \leq m \leq \nu \leq \mu \leq n$  the following occurs:

$$\sigma_j(0) = a_j \omega / b_j, \quad j = 1, \dots, \nu, \quad a_j, b_j > 0 \text{ integers};$$

$$\sigma_j(0) \neq \text{int} \omega / b_0, \quad j = \nu+1, \dots, \mu, \quad b_0 = b_1 \dots b_\nu, \quad m = 0, 1, 2, \dots$$

The functions  $f_j$ ,  $j = 1, \dots, n$ , satisfy condition (L) and the conditions of symmetry (6.4). As usual we shall put  $a = (a_1, \dots, a_v)$ ,  $b = (b_1, \dots, b_v)$ ,  $\sigma = (\sigma_1, \dots, \sigma_v)$ ,  $\lambda = (\lambda_1, \dots, \lambda_v)$ ,  $\eta = (\eta_1, \dots, \eta_{n-\mu})$ .

(6.1). If for some  $a, b, \lambda, \eta$ , all real, and  $\varepsilon$  sufficiently small the determining equations (6.7) are satisfied, then (6.10) has a real periodic solution of the form

$$(6.11) \quad \begin{aligned} x_j(t, \varepsilon) &= \lambda_j \tau_j^{-1} \cos \tau_j t + o(\varepsilon), & j &= 1, \dots, m, \\ x_j(t, \varepsilon) &= \lambda_j \tau_j^{-1} \sin \tau_j t + o(\varepsilon), & j &= m+1, \dots, v, \\ x_j(t, \varepsilon) &= o(\varepsilon), & j &= v+1, \dots, m, \\ x_j(t, \varepsilon) &= \eta_{j-\mu} + o(\varepsilon), & j &= \mu+1, \dots, n, \end{aligned}$$

and  $x_j(-t, \varepsilon) = x_j(t, \varepsilon)$ ,  $j = 1, \dots, m$ ,  $j = \mu+1, \dots, n$ , and  $x_j(-t, \varepsilon) = -x_j(t, \varepsilon)$ ,  $j = m+1, \dots, \mu$ . In particular, given  $a, b, \mu, \lambda$  there is always a  $\sigma(\varepsilon)$  such that equations (6.7) are satisfied and system (6.10) has a real periodic solution of the form (6.11). If system (6.10) is autonomous, the parameter  $\omega$  has to be added.

(6.11). If the functions  $f_j$  have continuous first order partial derivatives and, given  $a, b, \lambda, \eta$ , there are  $v$  numbers  $\lambda_0 = (\lambda_{10}, \dots, \lambda_{v0})$  such that  $Q_j(a, b, \lambda_0, \mu, \omega, 0) = 0$ ,  $j = 1, \dots, v$ , and the Jacobian  $\partial Q / \partial \lambda \neq 0$  at  $\lambda = \lambda_0$ ,  $\varepsilon = 0$ , then for all  $\varepsilon$  sufficiently small system (6.10) has a periodic solution of the form (6.11) with convenient  $\lambda(\varepsilon)$ ,  $\lambda(0) = \lambda_0$ .

Obviously here the Jacobian of §5 of order  $2v + (n - r)$  may be identically zero while the Jacobian above  $\partial Q / \partial \lambda$  of order  $v$  may well be  $\neq 0$ . In the case (6.10) is autonomous, also the parameter  $\omega$  can be taken into consideration as in §5.

The following result is also of interest since it is an explicit existence theorem of a  $(n - \mu + 2)$ -parameter family of periodic solutions. It concerns autonomous systems (6.10) with  $v = 1$ ,  $m = 0$ , or  $m = 1$ ,  $a_1 = b_1 = 1$ .

(6.111). Consider an autonomous system

$$(6.12) \quad \begin{aligned} x_j'' + \sigma_j^2 x_j &= \varepsilon f_j(x, x', \varepsilon), & j &= 1, \dots, \mu, \\ x_j' &= \varepsilon f_j(x, x', \varepsilon), & j &= \mu+1, \dots, n, \end{aligned}$$

where the functions  $f_j$ ,  $j = 1, \dots, n$ , are Lipschitzian in the  $n + \mu$  variables  $x_1, \dots, x_n, x_1', \dots, x_\mu'$ , and continuous in  $\varepsilon$ , for  $|x_s| \leq R$ ,  $s = 1, \dots, n$ ,  $|x_s'| \leq R$ ,  $s = 1, \dots, \mu$ , and  $0 \leq \varepsilon \leq \varepsilon_0$ . In



addition suppose, for  $f_1$  only, that  $f_1(0, x_2, \dots, x_n, 0, x_2', \dots, x_\mu', \varepsilon) = 0$ . Suppose that either all  $f_j, j = 1, \dots, n$ , are odd in  $(x_1, \dots, x_\mu)$ , or  $f_1$  is even and  $f_2, \dots, f_n$  are odd in  $(x_2, \dots, x_\mu, x_1')$ . Suppose  $\sigma_j(\varepsilon) > 0, j = 1, \dots, \mu$ , are continuous functions of  $\varepsilon$  [or constants] and  $\sigma_j(0) \neq 0, \text{ mod } \sigma_1(0), j = 2, \dots, \mu$ . Put  $\omega_0 = \sigma_1(0)$ , and let  $r_2$  be any number,  $0 < r_2 < R$ . Then there exists an  $\varepsilon_1, 0 < \varepsilon_1 \leq \varepsilon_0$ , such that, for all real  $\varepsilon, \lambda_1, \eta_1, \dots, \eta_{n-\mu}, 0 \leq \lambda_1, \eta_1, \dots, \eta_{n-\mu} \leq r_2, 0 \leq \varepsilon \leq \varepsilon_1$ , system (6.12) has a real periodic solution of the form

$$\begin{aligned} x_1(t, \varepsilon) &= \lambda_1 \omega^{-1} \sin \omega t + o(\varepsilon), \quad \text{or } x_1(t, \varepsilon) = \lambda_1 \omega^{-1} \cos \omega t + o(\varepsilon), \\ (6.13) \quad x_j(t, \varepsilon) &= o(\varepsilon), \quad j = 2, \dots, \mu, \\ x_j(t, \varepsilon) &= \eta_{j-\mu} + o(\varepsilon), \quad j = \mu + 1, \dots, n, \end{aligned}$$

where  $x_1$  is odd or even in  $t, x_2, \dots, x_\mu$  are odd,  $x_{\mu+1}, \dots, x_n$  are even, where  $\omega = \omega(\varepsilon, \lambda_1, \eta_1, \dots, \eta_{n-\mu})$  is a continuous function of the same parameters, and  $\omega = \omega_0$  for  $\varepsilon = 0$ . Also,  $t$  can be replaced by  $t + \theta$  in (6.13),  $\theta$  an arbitrary constant. [See J. K. Hale, 9cf, for the analytic case, and W. R. Fuller, 10, for the present situation with  $\mu = n$ .]

PROOF. By (4.111) we know that the unique  $Q_1 = Q_1(\lambda_1, \mu, \omega, \varepsilon)$  where  $\eta = (\eta_1, \dots, \eta_{n-r})$ , is a uniformly continuous and Lipschitzian function of  $\lambda_1, \eta, \omega$ , and uniformly continuous functions of  $\varepsilon$ . The only determining equation (6.7) is

$$\omega - \varepsilon Q_1(\lambda_1, \eta, \omega, \varepsilon) = \sigma_1(\varepsilon).$$

If  $\omega \in U = [\omega_{01}, \omega_{02}]$  with  $\omega_{01} < \omega_0 < \omega_{02}$ , and  $\omega_{02} - \omega_{01}$  sufficiently small, if  $0 \leq \varepsilon \leq \varepsilon_1$  and  $0 < \varepsilon_1 \leq \varepsilon_0$  sufficiently small, we put

$$F_1(\omega, \varepsilon) = \omega - \varepsilon Q_1(\omega, \varepsilon) - \sigma_1(\varepsilon).$$

For all  $\omega_{01} \leq \omega^1 \leq \omega^2 \leq \omega_{02}, 0 \leq \varepsilon \leq \varepsilon_1$ , we have

$$|Q_1(\omega^2, \varepsilon) - Q_1(\omega^1, \varepsilon)| \leq M(\omega^2 - \omega^1),$$

$$F_1(\omega^2, \varepsilon) - F_1(\omega^1, \varepsilon) \geq (1 - \varepsilon M)(\omega^2 - \omega^1),$$

where  $M$  is a constant, independent of  $\lambda_1, \eta, \varepsilon, \omega$ . For  $0 \leq \varepsilon \leq \varepsilon'$ ,  $\varepsilon' = \min\{\varepsilon_1, 1/2M\}$ , the last expression above is positive. Thus  $F_1$  is an increasing continuous function of  $\omega$  with  $F_1(\omega_0, 0) = 0$ . Hence,

$F_1(\omega_{01}, 0) < 0$ ,  $F(\omega_{02}, 0) > 0$ , and finally  $F_1(\omega_{01}, \varepsilon) < 0$ ,  $F(\omega_{02}, \varepsilon) > 0$  for all  $\lambda_1, \eta$  as above, and  $0 \leq \varepsilon \leq \varepsilon''$ ,  $\varepsilon''$  sufficiently small. Thus, for the same  $\lambda_1, \eta$ ,  $0 \leq \varepsilon \leq \varepsilon''$ , there is a unique  $\omega$  with  $F_1(\omega, \varepsilon) = 0$ . By the usual argument of implicit function theorem we know that  $\omega$  is a continuous function of the same parameters. Now (6.111) follows from (6.1) and the previous considerations.

NOTE. If system (6.12) verifies the conditions of (6.111) when any one of the variables  $x_j$  takes the place of  $x_1$  and correspondingly the  $j^{\text{th}}$  equation replaces the first one,  $j = 1, \dots, \mu$ , then (6.111) assures the existence of  $\mu$  families of periodic solutions, each a  $(n-\mu+2)$ -parameter family.

EXAMPLE 1. The equation  $x'' + x = \varepsilon f(x, x')$  with  $f(0, 0) = 0$ , and either  $f(x, -x') = f(x, x')$ , or  $f(-x, x') = -f(x, x')$ , has a family of periodic solutions of the form  $x = \lambda\omega^{-1}\cos(\omega t + \theta) + O(\varepsilon)$ , or  $x = \lambda\omega^{-1}\sin(\omega t + \theta) + O(\varepsilon)$ , with  $\omega = \omega(\lambda, \varepsilon) = 1 + O(\varepsilon)$ ,  $\lambda, \theta$  arbitrary,  $\varepsilon > 0$  sufficiently small. We may take, for instance,  $f = x + x^2 + x'^2$ , or  $f = |x| + |x'|$ , or  $f = x|x'|$ . Also we may take  $f = (2 + x)(|x'| + x)$ .

EXAMPLE 2. The system  $x'' + x = \varepsilon(1 - |y|)|x'|x$ ,  $y'' + 2y = \varepsilon(1 - |x|)|y'|y$ , has two families of periodic solutions of the forms

$$x = \lambda\omega^{-1}\cos(\omega t + \theta) + O(\varepsilon), y = O(\varepsilon), \omega = \omega(\lambda, \varepsilon) = 1 + O(\varepsilon),$$

$$x = O(\varepsilon), y = \lambda\omega^{-1}\cos(\omega t + \theta) + O(\varepsilon), \omega = \omega(\lambda, \varepsilon) = 2^{1/2} + O(\varepsilon).$$

EXAMPLE 3. The system  $x'' + x = \varepsilon f_1(x, y, z, x')$ ,  $y' = \varepsilon f_2(x, y, z, x')$ ,  $z' = \varepsilon f_3(x, y, z, x')$ , with  $f_1(0, y, z, 0) = 0$  and either

$$f_j(-x, y, z, x') = -f_j(x, y, z, x'), j = 1, 2, 3,$$

or

$$\begin{aligned} f_1(x, y, z, -x') &= f_1(x, y, z, x'), f_j(x, y, z, -x') = \\ &= -f_j(x, y, z, x'), j = 2, 3, \end{aligned}$$

has a four parameter family of periodic solutions, for every  $\varepsilon$  sufficiently small, of the form  $x = \lambda\omega^{-1}\sin(\omega t + \theta) + O(\varepsilon)$ , or

$$x = \lambda\omega^{-1}\cos(\omega t + \theta) + O(\varepsilon), y = \eta_1 + O(\varepsilon), z = \eta_2 + O(\varepsilon),$$

with  $\omega = \omega(\lambda, \eta_1, \eta_2, \varepsilon)$ ,  $\theta, \lambda, \eta_1, \eta_2$  arbitrary.

EXAMPLE 4. A third order equation  $x''' + a_1 x'' + a_2 x' + a_3 x = \varepsilon f(x, x', x'', t, \varepsilon)$ , with  $f$  periodic could be studied in the lines of §§5, 6. For instance, if  $f(x, -x', x'', -t, \varepsilon) = -f(x, x', x'', t, \varepsilon)$ , then equation  $x''' + \sigma^2 x' = \varepsilon f$ ,  $\sigma > 0$ , has periodic solutions of the form  $x(-t, \varepsilon) = x(t, \varepsilon)$ ,  $x(t, 0) = -c_1 a \omega \sigma^{-2} b^{-1} \cos ab^{-1} \omega t + c_2$ ,  $a, b$  integers,  $c_1, c_2 \neq 0$  real, provided  $ab^{-1} \omega - \varepsilon H = \sigma$ , where  $H = H(a, b, c_1, c_2, \omega, \sigma, \varepsilon)$  and

$$H = (c_1 T)^{-1} \int_0^T f[x(t, 0), x'(t, 0), x''(t, 0), t, 0] \sin ab^{-1} \omega t \, dt + o(\varepsilon).$$

If  $\sigma = a\omega/b$ , and there is a  $c_{10} = c_{10}(c_2)$  such that  $H = 0$  and  $\partial H / \partial c_1 \neq 0$  for  $c_1 = c_{10}$ , then the equation has a non-zero solution for every  $c_2$ ; that is, a one-parameter family of periodic solutions [J. K. Hale, 9f].

## BIBLIOGRAPHY

- [1] Bailey, H. R., Harmonics and subharmonics for weakly nonlinear Mathieu type differential equations. To appear.
- [2] Bailey, H. R., and Cesari, L., Boundedness of solutions of linear differential systems with periodic coefficients. *Archive Rat. Mech. Anal.* 1, 1958, 246-271.
- [3] Bailey, H. R., and Gambill, R. A., On stability of periodic solutions of weakly nonlinear differential systems. *J. Math. Mech.* 6, 1957, 655-668.
- [4] Cesari, L., (a) Un nuovo criterio di stabilità per le soluzioni delle equazioni differenziali lineari. *Annali scuola Norm. Sup. Pisa* (2) 9, 1940, 163-186. (b) Sulla stabilità delle soluzioni dei sistemi di equazioni differenziali lineari a coefficienti periodici. *Mem. Accad. Italia* (6) 11, 1941, 633-695.
- [5] Cesari, L., Asymptotic Behavior and Stability Problems in Ordinary Differential Equations. *Ergbn. d. Mathematik und ih. Grenzgebiete*, Heft, 16, 1959, vii, 271.
- [6] Cesari, L., and Hale, J. K., (a) Second order linear differential systems with periodic L-integrable coefficients. *Riv. Mat. Univ. Parma*, 5, 1954, 55-61; 6, 1955, p. 159. (b) A new sufficient condition for periodic solutions of weakly nonlinear differential systems. *Proc. Amer. Math. Soc.* 8, 1957, 757-764.
- [7] Gambill, R. A., (a) Stability criteria for linear differential systems with periodic coefficients. *Riv. Mat. Univ. Parma*, 5, 1954, 169-181. (b) Criteria for parametric instability for linear differential systems with periodic coefficients. *Ibid.* 6, 1955, 37-43. (c) A fundamental system of real solutions for linear differential systems with periodic coefficients. *Ibid.* 7, 1956, 311-319.

- [8] Gambill, R. A., and Hale, J. K., Subharmonic and ultraharmonic solutions for weakly nonlinear systems. *J. Rat. Mech. Anal.* 2, 1956, 353-394.
- [9] Hale, J. K., (a) Evaluations concerning products of exponential and periodic functions. *Riv. Mat. Univ. Parma*, 2, 1954, 63-81. (b) On boundedness of the solutions of linear differential systems with periodic coefficients. *Ibid.* 5, 1954, 137-167. (c) Periodic solutions of nonlinear systems of differential equations. *Ibid.* 2, 1954, 281-311. (d) On a class of linear differential equations with periodic coefficients. *Illinois J. Math.* 1, 1957, 98-104. (e) Linear systems of first and second order differential equations with periodic coefficients. *Ibid.* 2, 1958, 586-592. (f) Sufficient conditions for the existence of periodic solutions of systems of weakly nonlinear first and second order differential equations. *Journ. Math. Mech.* 7, 1958, 163-172. (g) A short proof of a boundedness theorem for linear differential systems with periodic coefficients. *Archive Rat. Mech. Anal.* 2, 1959, 429-434. (h) On the behavior of the solutions of linear periodic differential systems near resonance points. *These Contributions*, 5, 1959, (i) On the stability of periodic solutions of weakly nonlinear periodic and autonomous differential systems. *These contributions*, 5, 1959.
- [10] Fuller, W. R., Existence theorems for periodic solutions of systems of differential and differential-difference equations. Ph.D. Thesis, Purdue University, Lafayette, Indiana, 1957.
- [11] Thompson, E. W., On stability of periodic solutions of autonomous differential systems. (To appear).

VII. THE APPLICATIONS OF A FIXED POINT THEOREM TO A  
VARIETY OF NON-LINEAR STABILITY PROBLEMS

Arnold Stokes

§1. INTRODUCTION

We use a fixed point theorem for locally convex linear spaces, due to Tychonoff<sup>1</sup> to reduce the study of the boundedness and stability of certain  $n$ -dimensional vector differential equations to the study of the corresponding properties of related first-order equations. In this manner, we very simply obtain a variety of known results, and in certain cases, we are able to clarify some of these results.

Fixed point theorems have been used to study the stability of systems of differential equations before, notably by Hukuwara<sup>2</sup>, and Bellman<sup>3</sup>, but the present approach, by using Tychonoff's theorem rather than Schauder's result<sup>4</sup> gives many more results in a greatly simplified fashion.

§2. DEFINITIONS

We will let  $E$  be the real vector space of all continuous functions from the non-negative reals into  $R^n$ , the  $n$ -dimensional vector space

---

\* The results presented here are based on material contained in the author's thesis, written as partial fulfillment of the requirements for the Ph.D. degree at the University of Notre Dame. The author wishes to thank Dr. J. P. LaSalle for his valuable suggestions and constant encouragement during the development of this study. The author also is grateful to the Office of Naval Research for their assistance while at the University of Notre Dame.

<sup>1</sup> A. Tychonoff, Ein Fixpunktsatz, Math. Ann. 111, 767-776, 1935.

<sup>2</sup> M. Hukuwara, Sur les Points Singuliers des Equations Differentielles Lineaires, J. Faculty of Sci., Hokkaido Imp. Univ. Ser. 1, Math. 2, 13-88, 1934-36.

<sup>3</sup> R. Bellman, On the Boundedness of Solutions of Non-Linear Difference and Differential Equations, Trans. Amer. Math. Soc., 62, 357-386, 1947.

<sup>4</sup> J. Schauder, Der Fixpunktsatz in Funktionalraum, Studia Math. 2, 171-181, 1930.

over the real field. The topology on  $E$  shall be that induced by the family of pseudo-norms  $(p_n)_{n=1}^{\infty}$ , where for  $x \in E$ ,

$$p_n(x) = \sup_{0 \leq t \leq n} \|x(t)\|$$

where  $\|x(t)\|$  is any vector norm. A fundamental system of neighborhoods is then given by  $(V_n)_{n=1}^{\infty}$ , where  $V_n = \{x \in E \mid p_n(x) \leq 1\}$ . Under this topology,  $E$  becomes a complete, locally convex linear space. For such a space, we have a theorem of Tychonoff's:

THEOREM 1. Let  $E$  be a complete, locally convex linear space. Let  $T : E \longrightarrow E$  be continuous, and let  $A$  be a closed convex subset of  $E$ . If  $T(A) \subset A$  and  $\overline{T(A)}$  is compact, then there exists a fixed point of  $T$  in  $A$ .<sup>5</sup>

If  $T$  is a compact (completely continuous) operator and if  $A$  is bounded, then  $\overline{T(A)}$  is always compact. We then obtain:

COROLLARY 1. Let  $E$  be as in Theorem 1. Let  $T : E \longrightarrow E$  be continuous and compact, and let  $A$  be a closed, convex, bounded subset of  $E$ . If  $T(A) \subset A$ , then there exists a fixed point of  $T$  in  $A$ .

### §3. REDUCTION TO A FIRST-ORDER EQUATION

We shall consider these types of  $n$ -dimensional systems here:

- (1)  $\dot{x} = f(t, x)$
- (2)  $\dot{x} = A(t)x + f(t, x)$ , where  $\|X(t)\| \leq K$ ,  
 $\|X(t)X^{-1}(s)\| \leq K, K > 0$
- (3)  $\dot{x} = A(t)x + f(t, x)$ , where  $\|X(t)\| \leq Ke^{-\sigma t}$   
 $\|X(t)X^{-1}(s)\| \leq Ke^{-\sigma(t-s)}, K, \sigma > 0$ ,

<sup>5</sup> This theorem, while not explicitly stated by Tychonoff, follows from a theorem given in No. 1 in the same manner that the second fixed point theorem of Schauder's in No. 4 follows from his first theorem.

where by  $X(t)$  is meant the principal matrix solution of the system  $\dot{x} = A(t)x$ , where  $A(t)$  is a continuous function of  $t$  for  $t \geq 0$ . If  $A(t)$  is a constant matrix, or periodic in  $t$ , examples of systems satisfying the inequalities in (2) or (3) are well-known.

In addition we shall assume that  $f$  is continuous on  $R^{n+1}$  into  $R^n$ , and that

$$(4) \quad \|f(t, x)\| \leq G(t, \|x\|), \quad \text{for } t \geq 0, x \in D \subset R^n$$

where  $G(t, r)$  is piecewise continuous on  $R^2$ , positive for  $t, r \geq 0$ , and nondecreasing in  $r$  for fixed  $t$ ; and  $D$  is some subset of  $R^n$ .

With (1) above, we shall associate the integral operator

$$(5) \quad T_b(x)(t) = b + \int_0^t f(s, x(s))ds, \quad$$

and with (2) and (3) above we shall associate the operator

$$(6) \quad T_b(x)(t) = X(t)b + \int_0^t X(t)X^{-1}(s)f(s, x(s))ds, \quad$$

where  $b$  is a vector in  $R^n$ .

Clearly fixed points of these operators correspond to solutions of the associated equations, the solution passing through  $b$  at  $t = 0$ . Also it is evident that both operators are compact in the topology of our function space  $E$ .

To apply Corollary 1, we shall proceed as follows: Let  $B$  be a bounded subset of  $R^n$ , and let  $A$  be a subset of  $E$  defined by a positive real-valued function  $g(t)$ , continuous for  $t \geq 0$ , that is,

$$(7) \quad A = \{x \in E \mid \|x(t)\| \leq g(t)\}.$$

For such a  $g$ ,  $A$  is closed, convex, and bounded in the topology given above on  $E$ . We will further assume that for  $x \in A$ ,  $x(t) \in D$  for all  $t \geq 0$ , so that (4) may be used.

Now take  $b \in B$ ,  $x \in A$ . To apply Corollary 1 to either of the operators in (5) or (6), we must show  $T_b(A) \subset A$ , or, by (7),  $\|T_b(x)(t)\| \leq g(t)$ . Using the operator in (5), we obtain

$$\begin{aligned}
\|T_b(x)(t)\| &\leq \|b\| + \int_0^t \|f(s, x(s))\| ds \\
&\leq \|b\| + \int_0^t G(s, \|x(s)\|) ds, \quad \text{by (4)} , \\
&\leq \|b\| + \int_0^t G(s, g(s)) ds, \quad \text{by the definition of } A, \\
&\hspace{15em} \text{as } G \text{ is non-decreasing.}
\end{aligned}$$

Thus, for (5), we have  $T_b(A) \subset A$  if  $g$  satisfies

$$(8) \quad \|b\| + \int_0^t G(s, g(s)) ds \leq g(t) \quad \text{for } b \in B, t \geq 0 .$$

Now consider (6), and let  $A(t)$  be such that  $X(t)$  satisfies the inequalities in (2). In precisely the same fashion as before, we have  $T_b(A) \subset A$  if  $g$  satisfies

$$(9) \quad K \|b\| + \int_0^t KG(s, g(s)) ds \leq g(t), \quad \text{for } b \in B, t \geq 0 .$$

If  $X(t)$  satisfies the inequalities in (3), we have  $T_b(A) \subset A$  if  $g$  satisfies

$$(10) \quad K \|b\| e^{-\sigma t} + \int_0^t K e^{-\sigma(t-s)} G(s, g(s)) ds \leq g(t), \quad \text{for } b \in B, t \geq 0 .$$

Equations (8) and (9) may be combined into

$$(11) \quad K_1 \|b\| + \int_0^t K_1 G(s, g(s)) ds \leq g(t), \quad \text{for } b \in B, t \geq 0 .$$

where  $K_1 = 1$  if equation (1) is being considered, or  $K$ , if (2) is being discussed.

Now  $g$  will clearly satisfy (11) if it satisfies the differential inequality:



$$(12) \quad \dot{g}(t) \geq K_1 G(t, g(t)), \quad g(0) \geq K_1 \|b\|, \quad \text{for } b \in B, t \geq 0,$$

and  $g$  will satisfy (10) if it satisfies the differential inequality:

$$(13) \quad \dot{g}(t) \geq -\sigma g(t) + K G(t, g(t)), \quad g(0) \geq K \|b\|, \quad \text{for } b \in B, t \geq 0,$$

where by a solution is meant a differentiable function  $g$  satisfying (12) or (13) wherever  $G$  is continuous.

Thus, to demonstrate the existence of a solution to any of the equations (1), (2), or (3) which satisfies  $\|x(t)\| \leq g(t)$ , we must choose  $B \subset R^n$  and a  $g$  satisfying (12) or (13), (and, of course,  $g$  must possess the property that  $\|x(t)\| \leq g(t)$  implies  $x(t) \in D$ ).

#### §4. STATEMENT OF THE THEOREMS

(1) On existence in the large. We will consider equation (1) and assume that  $f$  is dominated by  $G$  everywhere, that is,  $D = R^n$ .

**THEOREM 2.** If the equation  $\dot{r} = G(t, r)$  possesses the property that for any  $r_0 > 0$  there exists a solution defined on  $[0, \infty)$ , passing through  $r_0$  at  $t = 0$ , then for an arbitrary vector  $b \in R^n$ , there exists a solution of (1) defined on  $[0, \infty)$  which passes through  $b$  at  $t = 0$ .

**PROOF.** Take  $b \in R^n$ , let  $B = \{b\}$ . Choose  $g$  to be a solution of  $\dot{r} = G(t, r)$  with  $g(0) \geq \|b\|$  which is defined on  $[0, \infty)$ . Clearly such a  $g$  satisfies (12), so the operator in (5) maps  $A$  into  $A$ , where  $A$  is defined as in (7). So by Corollary 1,  $T_b$  has a fixed point in  $A$ ; that is, there exists a solution of (1) defined on  $[0, \infty)$  passing through  $b$  at  $t = 0$ .

**COROLLARY 2.** Assume  $\|f(t, x)\| \leq M(t)L(\|x\|)$ , for  $t \geq 0$ ,  $x \in R^n$ , where  $M, L$  are piecewise continuous, positive, and  $L$  is non-decreasing. If

$$\int_{r_0}^{\infty} \frac{1}{L(s)} ds = \infty,$$

for positive  $r_0$ , then for any  $b \in R^n$ , there exists a solution of (1) passing through  $b$  at  $t = 0$ .

PROOF. Observe that the equation  $\dot{r} = M(t)L(r)$ ,  $r(0) = r_0$  may be solved for any  $r_0 > 0$ , for if we write

$$\int_{r_0}^r \frac{1}{L(s)} ds = \int_0^t M(s) ds, \quad$$

note that the function of  $r$  on the left is strictly increasing, so the inverse function exists, and by our assumption concerning  $L$ , the domain of the inverse functions is  $[0, \infty)$ , so the solution through  $r_0$  at  $t = 0$  is defined on  $[0, \infty)$  for all  $r_0 > 0$ . The result follows from Theorem 2.

Corollary 2 is a theorem due to Wintner<sup>6</sup>, which was later improved by him by removing the requirement that  $L$  be non-decreasing<sup>7</sup>. Later Conti<sup>8</sup> generalized this result still further, by replacing the norm function by a more general function. The above proof is considerably simpler than Wintner's original proof, however.

For an example of a function dominated as in Corollary 2, consider any function  $f(t, x)$  such that  $\|f(t, x)\| = o(\|x\|)$  as  $\|x\| \rightarrow \infty$  for fixed  $t$ .

(ii) On Boundedness. We will consider Equations (1) or (2), and again assume that  $f$  is dominated by  $G$  everywhere, or  $D = \mathbb{R}^n$ .

THEOREM 3. If the equation  $\dot{r} = G(t, r)$  possesses the property that for any  $r_0 > 0$  there exists a solution defined and bounded on  $[0, \infty)$ , passing through  $r_0$  at  $t = 0$ , then for an arbitrary vector  $b \in \mathbb{R}^n$  there exists a bounded solution of (1) or (2) defined on  $[0, \infty)$  which passes through  $b$  at  $t = 0$ .

PROOF. The proof is precisely the same as for Theorem 1, where  $B = \{b\}$ , for  $b \in \mathbb{R}^n$ , and here  $g$  is taken as any bounded solution to  $\dot{r} = G(t, r)$ , with  $g(0) \geq K_1 \|b\|$ .

<sup>6</sup> A. Wintner, The Non-Local Existence Problem of Ordinary Differential Equations, Amer. J. of Math., 67, 277-284, 1945.

<sup>7</sup> A. Wintner, The Infinities in the Non-Local Existence Problem of Ordinary Differential Equations, Amer. J. of Math., 68, 173-178, 1946.

<sup>8</sup> R. Conti, Limitazioni 'in Ampiezza' della Soluzioni di un Sistema di Equazioni Differenziali e Applicazioni, Bollettino della Unione Mat. Ital., Ser. 3, 11, 344-350, 1956.

COROLLARY 3. Assume  $\|f(t, x)\| < M(t)L(\|x\|)$  for  $t \geq 0$ ,  $x \in R^n$ , where  $M, L$  are piecewise continuous, positive, and  $L$  is non-decreasing. If

$$\int_{r_0}^{\infty} \frac{1}{L(s)} ds = \infty,$$

for positive  $r_0$ , and

$$\int_0^{\infty} M(s)ds < \infty,$$

then for any  $b \in R^n$ , there exists a bounded solution of (1) or (2) defined on  $[0, \infty)$  passing through  $b$  at  $t = 0$ .

PROOF. The proof follows from Theorem 3, if we observe that, as in Corollary 2, our assumption on  $L$  implies that for any  $r_0 > 0$ , there exists a solution to  $\dot{r} = M(t)L(r)$  defined on  $[0, \infty)$ , passing through  $r_0$  at  $t = 0$ , and further that

$$\int_0^{\infty} M(s)ds < \infty$$

implies that this solution is bounded.

With reference to the following theorem, observe that the above bounds in general depend upon the initial conditions.

For an example of a function dominated as in Corollary 3, let  $f(t, x) = B(t)x$ , where  $B(t)$  is an  $n \times n$  matrix, continuous in  $t$ , such that

$$\int_0^{\infty} \|B(t)\|dt < \infty.$$

Such a theorem appears in Bellman<sup>9</sup>, where the proof uses Gronwall's inequality. The above corollary shows that the linearity of  $L$  is not essential, but rather that it suffices that

$$\int_{r_0}^r \frac{1}{L(s)} ds$$

<sup>9</sup> R. Bellman, Stability Theory of Differential Equations, (McGraw-Hill Inc., New York, 1953).

diverges as  $r \rightarrow \infty$ .

(iii) On Ultimate Boundedness. Here we use ultimate boundedness in the sense of Yoshizawa<sup>10</sup>; that is, through every point there passes a solution which is not only bounded but as  $t \rightarrow \infty$ , enters a bounded region, which is independent of the initial conditions. We will now consider Equation (3), and again we assume  $f$  is dominated by  $G$  everywhere.

THEOREM 4. If  $G$  is such that for every positive initial condition  $r_0$ , there exists a solution to the inequality  $\dot{r} \geq -\sigma r + KG(t, r)$  passing through  $r_0$  at  $t = 0$  which is ultimately bounded, then for any  $b \in R^n$ , there exists a solution of (3) passing through  $b$  at  $t = 0$  which is ultimately bounded.

PROOF. The proof is the same as for the above two theorems, where  $B = \{b\}$ , for  $b \in R^n$ , and here  $g$  is taken as any ultimately bounded solution to  $\dot{r} \geq -\sigma r + KG(t, r)$ , with  $g(0) \geq K \|b\|$ .

COROLLARY 4. For any  $\epsilon > 0$ , assume there exists  $R(\epsilon) > 0$  such that  $\|f(t, x)\| \leq N$ ,  $\|x\| \leq R$ ,  $t \geq 0$ , and  $\|f(t, x)\| \leq \epsilon \|x\|$ ,  $\|x\| \geq R$ ,  $t \geq 0$ . Then all solutions to (3) are ultimately bounded.

PROOF. Here let

$$G(t, r) = \begin{cases} N, & r \leq R \\ \epsilon r, & r \geq R \end{cases}.$$

By Theorem 4, all we must show is that for  $\epsilon$  sufficiently small, there exist ultimately bounded solutions to  $\dot{r} \geq -\sigma r + KG(t, r)$  passing through  $r_0$  at  $t = 0$ , where  $r_0 \geq K \|b\|$ , for  $b$  an arbitrary vector in  $R^n$ .

Now choose  $\epsilon > 0$  such that  $\epsilon < \sigma/K$ , and choose  $\beta$  such that  $0 < \beta < 1$ . Let  $g(t) = K \|b\| e^{\beta(\epsilon K - \sigma)t} + NK/\sigma$ . By our choice of  $\epsilon$ ,  $g \rightarrow NK/\sigma$  as  $t \rightarrow \infty$ , and so  $g$  is ultimately bounded. And  $g(0) \geq K \|b\|$ , so all that remains is to show that  $g$  satisfies the differential inequality:

$$\dot{g}(t) \geq -\sigma g(t) + K \begin{cases} N, & g(t) \leq R \\ \epsilon g(t), & g(t) \geq R \end{cases}.$$

<sup>10</sup> T. Yoshizawa, Note on the Boundedness and the Ultimate Boundedness of Solutions of  $\dot{x} = F(t, x)$ , Mem. of the Coll. of Sci., Univ. of Kyoto, Ser. A, 29, Math. No. 3, 275-291, 1955.

Now  $\dot{g}(t) = \beta(\epsilon K - \sigma)K \|b\| e^{\beta(\epsilon K - \sigma)t}$ , and

$$(14) \quad \dot{g}(t) + \sigma g(t) = K \|b\| e^{\beta(\epsilon K - \sigma)t} [\beta \epsilon K + \sigma(1 - \beta)] + NK.$$

By our choice of  $\beta$ , this is clearly  $\geq NK$  for all  $t \geq 0$ , and so half the inequality is satisfied. Now  $Kg(t) = K^2 \|b\| e^{\beta(\epsilon K - \sigma)t} + NK^2/\sigma$ , and upon comparison with (14), we see that the inequality

$$\dot{g}(t) \geq -\sigma g(t) + \epsilon K g(t)$$

reduces to the two inequalities

$$\beta \in K + \sigma(1 - \beta) \geq \epsilon K$$

and

$$NK \geq \epsilon K NK/\sigma.$$

But the first of these is equivalent to  $\sigma(1 - \beta) \geq \epsilon K(1 - \beta)$ , or  $\epsilon \leq \sigma/K$ , and the second reduces to  $\epsilon K/\sigma \leq 1$ , or  $\epsilon \leq \sigma/K$ . So by choosing  $\epsilon < \sigma/K$ , we see that the conditions of Theorem 4 are satisfied, and so all solutions to (3) are ultimately bounded.

For an example of a function  $f(t, x)$  dominated as in Corollary 4, it suffices to choose  $f$  so that  $\|f(t, x)\| = o(\|x\|)$  as  $\|x\| \rightarrow \infty$ , uniformly in  $t$ .

(iv) On Stability of a Critical Point. Here we will suppose (1) or (2) has a critical point at the origin, that is,  $f(t, 0) = 0$  for all  $t \geq 0$ , and we wish to examine the stability of this point. We will say the origin is stable if given any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that for any  $b \in R^n$  such that  $\|b\| \leq \delta$ , there exists a solution through  $b$  at  $t = 0$  which is bounded in norm by  $\epsilon$  for  $t \geq 0$ . As this is a local problem, we will assume  $f$  to be dominated by  $G$  in some neighborhood of the origin, that is  $D = \{x \in R^n \mid \|x\| \leq \eta\}$ , where  $\eta > 0$  is given.

**THEOREM 5.** If the equation  $\dot{r} = K_1 G(t, r)$  possesses a stable critical point at the origin, then (1) or (2) also possesses a stable critical point at the origin.

**PROOF.** Take  $\epsilon > 0$ , and let  $\delta > 0$  be such that if  $0 < r_0 \leq \delta$ , there exists a solution to  $\dot{r} = K_1 G(t, r)$  which passes through  $r_0$  at  $t = 0$  and is bounded by  $\min(\epsilon, \eta)$  for  $t \geq 0$ . To apply Corollary 1, let  $B = \{b \in R^n \mid \|b\| \leq \delta\}$ , and let  $g$  be a solution to  $\dot{r} = K_1 G(t, r)$

which is bounded by  $\min(\epsilon, \eta)$  for  $t \geq 0$ , and  $g(0) = \delta$ . Thus for  $b \in B$ ,  $T_b(A) \subset A$ , or there exists a solution to (1) or (2) through  $b$  at  $t = 0$  which lies in  $A$ . The theorem follows.

COROLLARY 5. Let  $\|f(t, x)\| \leq M(t)L(\|x\|)$ , for  $t \geq 0$ ,  $\|x\| \leq \eta$ , where  $M$  and  $L$  are continuous, and for  $r > 0$ ,  $L$  is positive and non-decreasing. If  $L(0) = 0$ , and

$$\lim_{r_0 \rightarrow 0^+} \int_{r_0}^r \frac{1}{L(s)} ds = \infty$$

for  $r$  small and positive, and

$$\int_0^\infty M(s) ds < \infty,$$

then (1) or (2) possesses a stable critical point at the origin.

PROOF. By Theorem 5, if the equation  $\dot{r} = K_1 M(t)L(r)$  possesses a stable critical point at the origin, the result follows. But by considering the equation

$$\int_{r_0}^r \frac{1}{L(s)} ds = \int_0^t M(s) ds,$$

and by the above assumptions on  $L$  and  $M$ , clearly given any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that

$$\int_\delta^\epsilon \frac{1}{L(s)} ds > \int_0^\infty M(s) ds,$$

or, there exists a solution through  $r_0$  at  $t = 0$  which, for  $r_0 \leq \delta$ , is bounded by  $\epsilon$  for  $t \geq 0$ . The corollary follows.

A result of this type, where  $L(r) = r$ , is given by Bellman<sup>9</sup>, where again Gronwall's inequality is applied. As before, Corollary 5 demonstrates that the linearity of  $L$  is not important.

For an example of a function of this type, consider  $f(t, x)$

<sup>9</sup> R. Bellman, Stability Theory of Differential Equations, (McGraw-Hill Inc., New York, 1953).

where the  $i$ -th component is the product of a polynomial in the components of  $x$ , without a constant term, with a function  $\varphi_i(t)$ , where

$$\int_0^{\infty} |\varphi_i(s)| ds < \infty.$$

(v). On Asymptotic Stability of a Critical Point. Here we shall consider (3), and assume that (3) has a critical point at the origin, that is,  $f(t, 0) = 0$ . By asymptotic stability we mean, of course, that the origin is stable, and in addition, there exists a  $\delta > 0$  such that through any  $b$ ,  $\|b\| \leq \delta$ , there passes a solution which approaches 0 as  $t \rightarrow \infty$ . As this is again a local problem, we will assume  $f$  to be dominated in  $G$  in some neighborhood of the origin, that is,  $D = \{x \in \mathbb{R}^n \mid \|x\| \leq \eta\}$ . For our purposes, here we wish to take  $\eta$  as a function of  $\epsilon$ , where  $\epsilon$  is a small parameter, and we will also assume that  $G(t, r) = G(t, r, \epsilon)$  is also a function of this parameter.

THEOREM 6. If the inequality  $\dot{r} \geq -\sigma r + Kg(t, r, \epsilon)$  possesses an asymptotically stable point at the origin for  $\epsilon$  sufficiently small, then (3) possesses an asymptotically stable critical point at the origin.

PROOF. Take  $\epsilon > 0$  so small that the differential inequality above is asymptotically stable. Then there exists a  $\delta > 0$  such that for  $0 < r_0 \leq \delta$ , there exists a solution to the inequality which passes through  $r_0$  at  $t = 0$ , tends to 0 as  $t \rightarrow \infty$ , and for all  $t \geq 0$ , is bounded by  $\eta(\epsilon)$ . Then, to apply Corollary 1, let  $B = \{b \in \mathbb{R}^n \mid \|b\| \leq \delta\}$ , and choose  $g$  to be a solution to the inequality having the above properties, with  $g(0) = \delta$ . The theorem follows.

COROLLARY 6. Assume that for any  $\epsilon > 0$ , we have  $\|f(t, x)\| \leq \epsilon \|x\|$ , for  $t \geq 0$ ,  $\|x\| \leq \eta(\epsilon)$ . Then in (3), the origin is asymptotically stable.

PROOF. By Theorem 6, we must show only that for sufficiently small  $\epsilon$  a solution  $g(t)$  can be found to the differential inequality  $\dot{g}(t) \geq -\sigma g(t) + \epsilon Kg(t)$  which is bounded by  $\eta(\epsilon)$  and tends to 0 as  $t \rightarrow \infty$ . But for  $\epsilon < \sigma/K$ , we may choose  $g(t) = \eta(\epsilon)e^{(\epsilon K - \sigma)t}$ , and the corollary is proved.

As an example, we may take any  $f(t, x)$  for which  $\|f(t, x)\| = o(\|x\|)$  as  $\|x\| \rightarrow 0$  uniformly in  $t$ . This result, by now classical, is originally due to Perron.

COROLLARY 7. Assume that for any  $\epsilon > 0$ , we have  $T(\epsilon) > 0$  such that

$$\|f(t, x)\| \leq \begin{cases} k \|x\| + t^b \|x\|^{1+a}, & 0 \leq t \leq T(\epsilon), \\ \epsilon \|x\| + t^b \|x\|^{1+a}, & t \geq T(\epsilon), \end{cases}$$

for  $\|x\| \leq \eta(\epsilon)$ , where  $k, b, \geq 0, a > 0$ . Then in (3), the origin is asymptotically stable.

PROOF. Again by Theorem 6, we must show only that for sufficiently small  $\epsilon$ , a solution  $g(t)$  can be found to the inequality

$$\dot{g}(t) \geq -\sigma g(t) + K \begin{cases} kg(t) + t^b g(t)^{1+a}, & 0 \leq t \leq T_1, \\ \epsilon g(t) + t^b g(t)^{1+a}, & t \geq T_1, \end{cases}$$

where  $T_1 \geq T(\epsilon)$ , which is bounded by  $\eta(\epsilon)$  and tends to 0 as  $t \rightarrow \infty$ .

Now, for  $0 \leq r \leq 1, 0 \leq t \leq T_1, -\sigma r + K(kr + t^b r^{1+a}) \leq Cr$ , for some constant  $C$ , so clearly we can find a solution to the inequality over  $[0, T_1]$ , for any  $T_1$ , with an arbitrarily small bound on  $[0, T_1]$ , simply by choosing  $g(0)$  sufficiently small. So we may assume  $g(t) \leq \eta(\epsilon)$  on  $[0, T_1]$ . Let  $g_0 = g(T_1)$ . We now wish to define  $g$  for  $t \geq T_1$ .

Take  $\epsilon > 0$  so that  $\epsilon < \sigma/K$ , and choose  $\beta$  so that  $0 < \beta < 1$ . Take  $T_1 \geq T(\epsilon)$ , and let  $g(t) = g_0 e^{\beta(\epsilon K - \sigma)(t - T_1)}$ . Clearly, by our choice of  $\epsilon, g \rightarrow 0$  as  $t \rightarrow \infty$ , and as  $g_0 \leq \eta(\epsilon), g(t) \leq \eta(\epsilon)$  for all  $t \geq 0$ . We now wish to show that for  $T_1$  sufficiently large, and  $g_0$  sufficiently small,  $g$  satisfies the differential inequality  $\dot{g}(t) \geq -\sigma g(t) + \epsilon K g(t) + K t^b g(t)^{1+a}$  for  $t \geq T_1$ . Now

$$\dot{g}(t) - (\epsilon K - \sigma)g(t) = (\beta - 1)(\epsilon K - \sigma)g_0 e^{\beta(\epsilon K - \sigma)(t - T_1)} = \alpha g(t),$$

where  $\alpha = (\beta - 1)(\epsilon K - \sigma) > 0$  by our choice of  $\beta$  and  $\epsilon$ . So we must show that  $\alpha g(t) \geq K t^b g(t)^{1+a}$ , or, as  $g$  is positive,  $\alpha \geq K t^b g(t)^a$ .

But

$$K t^b g(t)^a = g_0^a K e^{-a\beta(\epsilon K - \sigma)T_1} \cdot t^b e^{a\beta(\epsilon K - \sigma)t}.$$

Now choose  $T_1$  so large that  $t^b e^{a\beta(\epsilon K - \sigma)t} < \sqrt{\alpha}$  for  $t \geq T_1$ , and choose  $g_0$  so small that  $g_0^a K e^{-a\beta(\epsilon K - \sigma)T_1} \leq \sqrt{\alpha}$ . Clearly both these choices can be made, and the corollary is proved.

Corollary 7 appears in Coddington and Levinson<sup>11</sup>. There we also have an example of a system to which Corollary 7 applies.



## VIII. QUADRATIC DIFFERENTIAL EQUATIONS AND NON-ASSOCIATIVE ALGEBRAS

Lawrence Markus

### §1. STATEMENT OF THE PROBLEM AND THE RESULTS

In order to obtain a thorough knowledge of the qualitative behavior of the solutions for a class of non-linear differential equations, we classify and analyse differential systems which have quadratic polynomials as coefficients. We find all such differential systems in the plane, and an interesting collection of such systems in higher dimensional spaces.

The method of classification is based on an algebraic technique and thus differs from the geometric methods which are customarily used in the qualitative theory of differential equations. While the algebraic method is somewhat more intricate than the geometric analysis in dimension two, it is available in higher dimensions where the geometry is more difficult to manage.

To each quadratic differential system we attach a certain non-associative, but commutative, real linear algebra. The problem of affine equivalence of differential systems is shown to be the same as the isomorphism problem for the corresponding algebras. This is analogous to the classification of linear differential systems by canonical forms for the coefficient matrix, that is, by means of certain linear endomorphisms of a vector space.

Thus we first find all real, commutative, two-dimensional, linear algebras, see Theorems 6, 7, and 8. This yields all quadratic differential systems, up to affine equivalence, in the affine plane. It is easy to show, Theorem 9, that there are just six geometrical types of quadratic differential systems with isolated critical points in the plane. Representation for these six geometric types are pictured in Figures I through VI. Finally there is a brief discussion of quadratic differential systems in the real projective plane and in higher dimensional spaces. The standard summation notation is used throughout.

## §2. THE ALGEBRA CORRESPONDING TO A QUADRATIC DIFFERENTIAL SYSTEM

DEFINITION. A quadratic differential system is

$$\rho') \quad \frac{dx^1}{dt} = a_{jk}^1 x^j x^k, \quad i = 1, 2, \dots, n,$$

where the  $n^3$  real constants  $a_{jk}^1$  are normalized so that  $a_{jk}^1 = a_{kj}^1$ .

DEFINITION. The related real linear algebra  $\mathfrak{A}$  of the quadratic differential system  $\rho'$  is defined by the multiplication table for a basis  $u_1, u_2, \dots, u_n$  as  $u_j \cdot u_k = a_{jk}^1 u_1$ . Clearly the  $n$ -dimensional algebra  $\mathfrak{A}$  is commutative but it may not be associative.

THEOREM 1. Two quadratic differential systems

$$\rho') \quad \dot{x}^1 = a_{jk}^1 x^j x^k$$

and

$$\hat{\rho}') \quad \dot{y}^1 = \hat{a}_{jk}^1 y^j y^k \quad i, j, k = 1, 2, \dots, n$$

are equivalent under a non-singular linear transformation

$$x^1 = b_\ell^1 y^\ell$$

if and only if their related algebras  $\mathfrak{A}$  and  $\hat{\mathfrak{A}}$ , respectively, are isomorphic.

PROOF. Suppose  $x^1 = b_\ell^1 y^\ell$  carries the system  $\rho'$  into  $\hat{\rho}'$ . Write  $B_1^\ell b_k^1 = b_1^\ell B_k^1 = \delta_k^\ell$ , the Kronecker symbol, so  $y^\ell = B_1^\ell x^1$ . Then

$$\dot{y}^\ell = B_1^\ell \dot{x}^1 = B_1^\ell a_{rs}^1 b_j^r b_k^s y^j y^k = \hat{a}_{jk}^\ell y^j y^k.$$

Thus

$$\hat{a}_{jk}^\ell = B_1^\ell b_j^r b_k^s a_{rs}^1 \quad \text{or} \quad \hat{a}_{jk}^\ell b_\ell^1 = b_j^r b_k^s a_{rs}^1.$$

Now consider the related algebras with multiplication tables

$$\mathfrak{A}) \quad u_j \cdot u_k = a_{jk}^1 u_1$$

and

$$\hat{\mathfrak{A}}) \quad \hat{u}_j \cdot \hat{u}_k = \hat{a}_{jk}^1 \hat{u}_1.$$

The linear transformation of  $\hat{\mathfrak{N}}$  onto  $\mathfrak{N}$  defined by  $\hat{u}_\ell \longrightarrow b_{\ell 1}^1 u_1$  is now shown to be an isomorphism. For

$$\hat{u}_j \cdot \hat{u}_k = \hat{a}_{jk}^1 \hat{u}_1 \longrightarrow \hat{a}_{jk}^1 b_{1\ell}^1 u_\ell$$

and

$$b_{j1}^r u_r \cdot b_{k1}^s u_s = b_{jk}^r b_{rs}^s u_\ell \quad .$$

But

$$\hat{a}_{jk}^1 b_{1\ell}^1 = b_{jk}^r b_{rs}^s a_{\ell}^1 \quad .$$

Conversely, if there is an isomorphism between  $\mathfrak{N}$  and  $\hat{\mathfrak{N}}$ , which is expressed in terms of the given bases by  $\hat{u}_\ell \longrightarrow b_{\ell 1}^1 u_1$ , then the linear transformation  $x^1 = b_{\ell j}^1 x^j$  carries  $\phi'$  into  $\hat{\phi}'$ , as required.

Here we say that  $\phi'$  and  $\hat{\phi}'$  are affinely (or linearly) equivalent.

It is of interest to relate the algebraic properties of  $\mathfrak{N}$  to the behavior of the solution curves of  $\phi'$ . The next three theorems give certain general results in this direction.

THEOREM 2. The quadratic differential system

$$\phi') \quad \dot{x}^1 = a_{jk}^1 x^j x^k \quad 1, j, k = 1, 2, \dots, n$$

has an isolated critical point at the origin if and only if the related algebra  $\mathfrak{N}$  has no nilpotent element  $e \neq 0$  for which  $e \cdot e = 0$ .

PROOF. If  $\phi'$  has another critical point  $P$  than the origin  $O$ ; then the line  $OP$  consists of critical points since  $\phi'$  is homogeneous. Take  $OP$  to be the  $x^1$ -axis and then the coefficients of  $\phi'$  (in the new coordinates) satisfy  $a_{11}^1 = 0$  for  $1 = 1, 2, \dots, n$ . But then the first basis vector  $u_1$ , for the related algebra  $\mathfrak{N}$ , satisfies  $u_1 \cdot u_1 = 0$ .

Conversely if  $e \cdot e = 0$ , take a linear transformation of the  $x$ -space so that  $e = u_1$ , the first basis vector of  $\mathfrak{N}$ . Then  $a_{11}^1 = 0$  and the  $x^1$ -axis consists of critical points of  $\phi'$ . Q.E.D.

THEOREM 3. The quadratic differential system

$$\phi') \quad \dot{x}^1 = a_{jk}^1 x^j x^k \quad 1, j, k = 1, 2, \dots, n$$

has a solution which is a ray to or from the origin

if and only if the related algebra  $\mathfrak{N}$  has a non-zero idempotent (i.e., an element  $e \neq 0$  for which  $e \cdot e = e$ ).

PROOF. If  $\phi'$  has a ray solution, say the  $x^1$ -axis, then  $a_{11}^1 = 0$  for  $i = 2, 3, \dots, n$ . Thus the basis vector  $u_1$  in  $\mathfrak{N}$  satisfies  $u_1 \cdot u_1 = \lambda u_1$  for some  $\lambda \neq 0$ . Take  $e = 1/\lambda u_1$ . Conversely if  $\mathfrak{N}$  has an idempotent, say  $u_1 \cdot u_1 = u_1$ , then  $a_{11}^1 = 0$  for  $i = 2, 3, \dots, n$  and  $a_{11}^1 = 1$  so the  $x^1$ -axis is a ray solution of  $\phi'$ . Q.E.D.

COROLLARY. If  $n = 1, 3, 5, 7, \dots$  is odd, and if  $\phi'$  has an isolated critical point at the origin ( $\mathfrak{N}$  has no non-zero element  $e \cdot e = 0$ ), then  $\phi'$  has a solution ray ( $\mathfrak{N}$  has an idempotent element).

PROOF. This is an immediate consequence of the topological result that the sphere  $S^{n-1}$  cannot support a continuous, non-vanishing, tangent vector field. Q.E.D.

THEOREM 4. The quadratic differential system

$$\phi') \quad \dot{x}^i = a_{jk}^i x^j x^k \quad i, j, k = 1, 2, \dots, n$$

has an invariant  $r$ -plane,  $1 \leq r < n$ , through the origin if and only if the related algebra  $\mathfrak{N}$  has an  $r$ -dimensional subalgebra.

PROOF. Say the  $r$ -plane spanned by the coordinate axes of  $x^1, x^2, \dots, x^r$  is invariant under the flow of  $\phi'$ . Then  $a_{jk}^i = 0$  for  $r+1 \leq i \leq n$  and  $j, k \leq r$ . Thus the basis vectors  $u_1, u_2, \dots, u_r$  of  $\mathfrak{N}$  span an  $r$ -dimensional algebra. The converse also follows from the same observation. Q.E.D.

If  $\mathfrak{N}$  has an  $r$ -dimensional ideal then, in a certain sense,  $\phi'$  has a projection on an  $(n-r)$ -dimensional quadratic differential system. However if  $\mathfrak{N}$  splits as a direct product, then so does  $\phi'$ , as is indicated in the next theorem.

THEOREM 5. Let  $\phi'$  be a quadratic differential system with related algebra  $\mathfrak{N}$ . If there exist ideals  $\mathfrak{N}_1$  and  $\mathfrak{N}_2$  intersecting only in zero and spanning  $\mathfrak{N}$ , then  $\phi'$  is affinely equivalent to a product differential system of the form

$$\dot{x}^1 = a_{jk}^1 x^j x^k \quad 1, j, k = 1, 2, \dots, r$$

and

$$\dot{x}^1 = a_{jk}^1 x^j x^k \quad 1, j, k = r+1, \dots, n.$$

PROOF. We require that  $a_{jk}^1 = 0$  for  $1 \leq r$  with  $j > r$  or  $k > r$ ; and for  $1 > r$  with  $j < r$  or  $k < r$ . Choose a basis  $u_1, u_2, \dots, u_r, u_{r+1}, \dots, u_n$  of  $\mathfrak{u}$  such that  $u_1, \dots, u_r$  is a basis for the ideal  $\mathfrak{u}$  and  $u_{r+1}, \dots, u_n$  is a basis for the ideal  $\mathfrak{c}$ . Then the multiplication tensor  $a_{jk}^1$  has the desired properties. Q.E.D.

### §3. EXAMPLES AND APPLICATION OF QUADRATIC DIFFERENTIAL SYSTEMS

In the study of the potential flow from a doublet we are lead to the differential equation

$$\frac{dy}{dx} = \frac{y^2 - x^2}{2xy}$$

whose solution curves describe the streamlines. The equi-potential lines are the orthogonal trajectories which thereby satisfy

$$\frac{dy}{dx} = \frac{-2xy}{y^2 - x^2}.$$

The quadratic differential system

$$\dot{x} = 2xy, \quad \dot{y} = y^2 - x^2$$

corresponds to the algebra 4) of Theorem 8, which is the complex numbers.

Next consider the quadratic differential system

$$\begin{aligned} \dot{x} &= yz - x^2 \\ \dot{y} &= xz - y^2 \\ \dot{z} &= xy - z^2 \end{aligned}$$

which occurs in a solution of the Einstein gravitational field equations, cf. [9]. The corresponding algebra has a basis of idempotents  $e_1, e_2, e_3$  with

$$\begin{aligned} e_1 \cdot e_1 &= e_1, & e_2 \cdot e_2 &= e_2, & e_3 \cdot e_3 &= e_3 \\ e_1 \cdot e_2 &= -\frac{1}{2} e_3, & e_1 \cdot e_3 &= -\frac{1}{2} e_2, & e_2 \cdot e_3 &= -\frac{1}{2} e_1. \end{aligned}$$

There is just one (independent) nilpotent of index two, namely,  
 $e_1 + e_2 + e_3$ .

Next consider the Jacobi elliptic functions<sup>1</sup>  $u = \operatorname{sn} t$ ,  $v = \operatorname{cn} t$ ,  
 $w = \operatorname{dn} t$  for a fixed parameter value of  $k^2$  on  $0 < k < 1$ . Then

$$\frac{du}{dt} = vw, \quad \frac{dv}{dt} = -uw, \quad \frac{dw}{dt} = -k^2 uv.$$

This corresponds to an algebra with basis  $e_1, e_2, e_3$  of nilpotents such that

$$\begin{aligned} e_1 \cdot e_1 &= 0, & e_2 \cdot e_2 &= 0, & e_3 \cdot e_3 &= 0 \\ e_1 \cdot e_2 &= -\frac{1}{2} k^2 e_3, & e_1 \cdot e_3 &= -\frac{1}{2} e_2, & e_2 \cdot e_3 &= \frac{1}{2} e_1. \end{aligned}$$

Let

$$E_1 = \frac{1}{k} e_1, \quad E_2 = \frac{1}{k} e_2, \quad E_3 = e_3$$

and then

$$E_1 \cdot E_1 = 0, \quad E_2 \cdot E_2 = 0, \quad E_3 \cdot E_3 = 0$$

and

$$E_1 \cdot E_2 = -\frac{1}{2} E_3, \quad E_1 \cdot E_3 = -\frac{1}{2} E_2, \quad E_2 \cdot E_3 = \frac{1}{2} E_1.$$

Thus for all  $k$ , the corresponding quadratic differential systems for the elliptic functions are linearly equivalent.

One of the major applications of quadratic differential systems occurs in interaction processes which depend on collisions of entities. For example, second order chemical processes, or biological interactions are often of this nature.

Following Volterra [18], we consider  $n \geq 2$  species  $\sigma_1, \sigma_2, \dots, \sigma_n$  of creatures with populations  $N_1, N_2, \dots, N_n$  respectively. Say that the members of  $\sigma_1$  each have mass  $\beta_1$  and also assume that each species

<sup>1</sup> The author wishes to thank Professor G. Birkhoff for calling attention to this example and for discussing with him the topic of quadratic differential equations.

has a natural net birth-death rate  $\epsilon_1$ . Assume that when individuals of species  $\sigma_1$  and  $\sigma_j$  meet, there is a certain probability that the member of  $\sigma_1$  will destroy and eat the member of  $\sigma_j$ . Then the fluctuations of the populations with time  $t$  can be studied by the differential system

$$\frac{dN_1}{dt} = \left( \epsilon_1 + \sum_{j=1}^n \frac{\beta_j}{\beta_1} a_{1j} N_j \right) N_1$$

for  $i = 1, 2, \dots, n$ . Here the  $a_{ij}$  are real biological constants and the conservation of mass requires that

$$\frac{a_{ij}}{\beta_i} = - \frac{a_{ji}}{\beta_j}.$$

One computes easily  $\beta_1 \dot{N}_1 + \beta_2 \dot{N}_2 + \dots + \beta_n \dot{N}_n = \epsilon_1 \beta_1 N_1 + \dots + \epsilon_n \beta_n N_n$ , which leads to a first integral in certain important cases. For example, if  $\epsilon_1 = \epsilon_2 = \dots = \epsilon_n = 0$  we have a quadratic differential system with an integral  $\beta_1 N_1 + \dots + \beta_n N_n = \text{const}$ , which is a hyperplane.

To be more definite consider the case of two interacting species,

$$\begin{aligned} \dot{N}_1 &= \left( \epsilon_1 + \frac{\beta_2 a_{12}}{\beta_1} N_2 \right) N_1 \\ \dot{N}_2 &= \left( \epsilon_2 + \frac{\beta_1 a_{21}}{\beta_2} N_1 \right) N_2. \end{aligned}$$

Then an integral is

$$N_1^{\epsilon_2} \exp \left\{ \frac{\beta_1 a_{21} N_1}{\beta_2} \right\} N_2^{-\epsilon_1} \exp \left\{ \frac{-\beta_2 a_{12} N_2}{\beta_1} \right\} = \text{const}.$$

In case  $\epsilon_1 = \epsilon_2 = 0$  we have the quadratic differential system

$$\dot{N}_1 = \beta_2 \gamma N_1 N_2, \quad \dot{N}_2 = -\beta_1 \gamma N_1 N_2$$

for

$$\beta_1 > 0, \beta_2 > 0, \gamma = \frac{a_{12}}{\beta_1} > 0.$$

This corresponds to the algebra 3) of Theorem 6.

As a second example of Volterra's theory consider the problem of the three fish. There are three species  $\sigma_1, \sigma_2, \sigma_3$  with populations  $N_1, N_2, N_3$ , respectively. We assume that the net birth-death rates  $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = 0$ , that is, each species is in equilibrium with the environment whenever the other two species are removed. Assume that the first type of fish eats the second type, which eats the third, which eats the first. Then  $a_{12} > 0, a_{23} > 0, a_{31} > 0$ . The resulting quadratic differential system corresponds to an algebra with a basis of nilpotents  $e_1, e_2, e_3$  such that

$$\begin{aligned} e_1 \cdot e_1 &= 0, & e_2 \cdot e_2 &= 0, & e_3 \cdot e_3 &= 0 \\ e_1 \cdot e_2 &= Ae_1 - Ce_2, & e_1 \cdot e_3 &= -Be_1 + Ee_3, & e_2 \cdot e_3 &= De_2 - Fe_3. \end{aligned}$$

Here the positive constants  $A, B, C, D, E, F$  are given by

$$\begin{aligned} 2A &= \frac{\beta_2 a_{12}}{\beta_1}, & -2B &= \frac{\beta_3 a_{13}}{\beta_1}, & -2C &= \frac{\beta_1 a_{21}}{\beta_2} \\ 2D &= \frac{\beta_3 a_{21}}{\beta_2}, & 2E &= \frac{\beta_1 a_{31}}{\beta_3}, & -2F &= \frac{\beta_2 a_{32}}{\beta_3}. \end{aligned}$$

Volterra's theory can be modified to apply to military situations, where it is termed Lanchester's law [5], but here the conservation of mass is no longer required since civilized warriors do not (as yet) eat one another.

Finally we point out that by increasing the dimension of the space of dependent variables we can make a differential system with non-homogeneous quadratic polynomial coefficients into a quadratic differential system. For example, consider the damped harmonic oscillator with the equations of motion  $\ddot{x} + h\dot{x} + \omega^2 x = 0$ , for positive constants  $h$  and  $\omega$ . In the affine phase space this is written

$$\dot{x} = y, \quad \dot{y} = -\omega^2 x - hy.$$

Consider the differential system in the real projective plane, with homogeneous coordinates  $(x, y, z)$  and write the differential equations in an affine plane "at infinity" where  $x \neq 0$ . In this affine plane use the coordinates

$$v = \frac{y}{x}, \quad z = \frac{1}{x}$$



and then

$$\dot{z} = -zv, \quad \dot{v} = -\omega^2 - 2hv - v^2.$$

Now introduce a new coordinate  $w$  and consider the quadratic differential system in  $R^3$ ,

$$\dot{z} = -zv, \quad \dot{v} = -\omega^2 w^2 - 2hvw - v^2, \quad \dot{w} = 0.$$

The solutions of this enlarged differential system, in the plane  $w = 1$ , are solutions for the harmonic oscillator.

#### §4. REAL COMMUTATIVE ALGEBRAS OF DIMENSION TWO

LEMMA. Let  $\mathfrak{A}$  be a real 2-dimensional commutative linear algebra. If  $\mathfrak{A}$  does not contain a (non-zero) nilpotent element with square zero, then  $\mathfrak{A}$  contains an idempotent  $e$  with  $e \cdot e = e \neq 0$ .

PROOF. For some basis  $u_1, u_2$  of  $\mathfrak{A}$  we have

$$u_1 \cdot u_1 = a_{11}^1 u_1 + a_{11}^2 u_2, \quad u_1 \cdot u_2 = a_{12}^1 u_1 + a_{12}^2 u_2, \quad u_2 \cdot u_2 = a_{22}^1 u_1 + a_{22}^2 u_2.$$

If  $a_{11}^2 = 0$ , then  $a_{11}^1 \neq 0$  and take

$$e = \frac{1}{a_{11}^1} u_1$$

which is idempotent. If  $a_{11}^2 = 0$ , then  $a_{22}^2 \neq 0$  and one could take

$$e = \frac{1}{a_{22}^2} u_2$$

which is idempotent.

Thus suppose  $a_{11}^2 \neq 0$  and  $a_{22}^1 \neq 0$ . Consider the real cubic polynomial  $a_{22}^1 x^3 + (2a_{12}^1 - a_{22}^2)x^2 + (a_{11}^1 - 2a_{12}^2)x - a_{11}^2$ . This has a real non-zero root  $x_0$ . Thus

$$a_{22}^2 x_0^2 + 2a_{12}^2 x_0 + a_{11}^1 = x_0 [a_{22}^1 x_0^2 + 2a_{12}^1 x_0 + a_{11}^1].$$

If  $a_{22}^2 x_0^2 + 2a_{12}^2 x_0 + a_{11}^1 = a_{22}^1 x_0^2 + 2a_{12}^1 x_0 + a_{11}^1 = 0$ , then  $E = u_1 + x_0 u_2$  is nilpotent which is impossible. Then  $E \cdot E = \lambda E$  where  $\lambda = a_{11}^1 + 2a_{12}^1 x_0 + a_{22}^1 x_0^2 \neq 0$ . Define the idempotent  $e = 1/\lambda E$ . Q.E.D.

We first classify the quadratic differential systems in the plane where the origin is not an isolated critical point. Here the related algebra contains a non-zero nilpotent element with zero square, that is, a nilpotent of index two.

**THEOREM 6.** There are ten (types of) real commutative 2-dimensional algebras which contain a nilpotent of index two.

- 1)  $e_1 \cdot e_1 = 0, e_2 \cdot e_2 = 0, e_1 \cdot e_2 = 0$
- 2)  $e_1 \cdot e_1 = 0, e_2 \cdot e_2 = 0, e_1 \cdot e_2 = e_2$
- 3)  $e_1 \cdot e_1 = 0, e_2 \cdot e_2 = 0, e_1 \cdot e_2 = e_1 + e_2$
- 4)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = 0, e_2 \cdot e_2 = e_2$
- 5)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = 0, e_2 \cdot e_2 = e_1$
- 6)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_1, e_2 \cdot e_2 = ke_2, \text{ real } k \neq 0$
- 7)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_1, e_2 \cdot e_2 = e_1 + 2e_2$
- 8)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_2, e_2 \cdot e_2 = e_1$
- 9)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_2, e_2 \cdot e_2 = -e_1$
- 10)  $e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_2, e_2 \cdot e_2 = ke_1 + e_2, \text{ real } k \neq 0.$

The first three algebras each have a basis of nilpotent elements of index two whereas the last seven algebras each have just one independent nilpotent or index two. In Cases 6) and 10) the algebras are distinct for distinct values of the parameter  $k$ .

**PROOF.** First assume that the algebra has a basis of nilpotents of index two, say  $e_1$  and  $e_2$ . Then we need only determine the real constants  $c, d$  in

$$e_1 \cdot e_2 = ce_1 + de_2.$$

If  $c = 0, d = 0$  we have the first Case 1). If  $c = 0, d \neq 0$ , then a scalar change in  $e_1$  yields the algebra 2). If  $c \neq 0, d = 0$  we again obtain an algebra isomorphic with 2). If  $c \neq 0, d \neq 0$  let

$$E_1 = \frac{1}{d} e_1, \quad E_2 = \frac{1}{c} e_2$$

and compute  $E_1 \cdot E_2 = E_1 + E_2$ , which yields the Case 3).

The Case 1) has a zero multiplication and hence is different from 2) or 3). In the algebras 2) and 3) the only nilpotents of index 2

are scalar multiples of  $e_1$  and  $e_2$ . The algebra 3) is distinguished from 3) by the property that the product of two independent nilpotents is a nilpotent of index 2.

From now on assume that the algebra has just one independent nilpotent  $e_1$  which we take as the first vector of a basis,  $e_1$  and  $e_2$ . Write  $e_1 \cdot e_2 = ce_1 + de_2$ . If  $c = d = 0$ , then  $e_1 \cdot e_2 = 0$ . If  $c = 0$ ,  $d \neq 0$ , replace  $e_1$  by  $1/d e_1$  and then  $e_1 \cdot e_2 = e_2$ . If  $c \neq 0$ ,  $d = 0$  replace  $e_2$  by  $1/c e_2$  to obtain  $e_1 \cdot e_2 = e_1$ . If  $c \neq 0$  and  $d \neq 0$  let  $E_1 = 1/d e_1$ ,  $E_2 = c/d e_1 + e_2$  to obtain  $E_1 \cdot E_1 = 0$ ,  $E_1 \cdot E_2 = E_2$ . Thus one can always select a basis  $e_1, e_2$  for which  $e_1 \cdot e_1 = 0$  and  $e_1 \cdot e_2 = 0$ , or  $e_1 \cdot e_2 = e_1$ , or  $e_1 \cdot e_2 = e_2$ . Exactly one of these alternatives must hold. For only in the first case does  $e_1$  (which is the unique nilpotent up to a scalar multiple) annihilate the algebra upon left multiplication; in the second case  $e_1$  generates a 1-dimensional ideal and this is not so in the third alternative.

Now take a basis in the algebra  $\mathfrak{N}$  with  $e_1 \cdot e_1 = 0$ ,  $e_1 \cdot e_2 = 0$ . Write  $e_2 \cdot e_2 = fe_1 + ge_2$  for real constants  $f, g$ . By a change to a new basis, still preserving the properties  $e_1 \cdot e_1 = 0$ ,  $e_1 \cdot e_2 = 0$  we can always obtain  $e_2 \cdot e_2 = e_2$  or  $e_2 \cdot e_2 = e_1$ . These yield the algebras 4) and 5) above. They differ in that 4) contains an idempotent whereas 5) does not.

Now take a basis in the algebra  $\mathfrak{N}$  with  $e_1 \cdot e_1 = 0$ ,  $e_1 \cdot e_2 = e_1$ . Take a new basis  $E_1 = \alpha e_1$ ,  $\alpha \neq 0$ , and  $E_2 = \lambda e_1 + e_2$  so that  $E_1 \cdot E_1 = 0$  and  $E_1 \cdot E_2 = E_1$ . Write  $e_2 \cdot e_2 = fe_1 + ge_2$  and then

$$E_2 \cdot E_2 = \frac{(2-g)\lambda + f}{\alpha} E_1 + gE_2.$$

If  $g = 0$ , choose  $\alpha = 1$ ,  $\lambda = 1/2 f$  which makes  $E_2$  nilpotent and this is not allowed here. Thus  $g \neq 0$ . If  $g \neq 2$ , we can obtain  $E_2 \cdot E_2 = gE_2$ . If  $g = 2$ , one obtains either  $E_2 \cdot E_2 = E_1 + 2E_2$  or  $E_2 \cdot E_2 = 2E_2$ . These yield the algebras 6) and 7) above. The algebra 7) has no 1-dimensional subalgebra, other than that generated by a nilpotent element, and so 7) is different from every algebra in 6). If  $k = 2$  in 6) then every element generates a 1-dimensional algebra whereas if  $k \neq 2$  there is only one (up to scalar multiples) non-nilpotent element which generates a 1-dimensional subalgebra. Now consider the Case 6) with  $k \neq 2$ . Here the 1-dimensional space generated by  $e_2$  is distinguished. But then  $e_1 \cdot e_2 = e_1$  normalizes  $e_2$  so that  $k$  is an invariant of the algebra and distinct values of  $k$  determine non-isomorphic algebras.

Finally take a basis in  $\mathfrak{M}$  corresponding to the third alternative:

$$e_1 \cdot e_1 = 0, \quad e_1 \cdot e_2 = e_2.$$

Write  $e_2 \cdot e_2 = fe_1 + ge_2$ . Replace  $e_2$  by  $E_2 = \mu e_2$ ,  $\mu \neq 0$ . Then  $E_2 \cdot E_2 = \mu^2 fe_1 + \mu g E_2$ . If  $g = 0$ , we can obtain algebras 8) or 9). If  $g \neq 0$ , we obtain algebra 10). In this last case we exclude  $k = 0$  since this yields an algebra with a basis of nilpotents.

In the algebras 8), 9) and 10) the idempotents  $\alpha e_1$  each define a linear transformation upon multiplication of  $\mathfrak{M}$ . Moreover only  $e_1$  produces an eigenvalue of  $+1$  and that for the eigenvectors  $be_2$ ,  $b \neq 0$ . Thus if there is an isomorphism of algebra 8) onto 9) or 10) it must carry  $e_1$  to  $e_1$  and  $e_2$  to  $be_2$ . This distinguishes between algebras 8), 9), and 10). Also for distinct values of  $k$  in the algebras 10) we obtain non-isomorphic algebras. Q.E.D.

The only algebras occurring in this Theorem 6 which are associative are 1), 4), 5), and 6), for  $k = 1$ . The algebra 6) for  $k = 2$  is power associative and all the remaining algebras of Theorem 6 are not power associative. The necessary and sufficient condition for the power associativity of real commutative algebras is the identity  $x^2 \cdot x^2 = (x^2 \cdot x) \cdot x$ , cf. [1], and a direct computation yields the desired result.

**THEOREM 7.** Every real commutative 2-dimensional algebra  $\mathfrak{M}$ , containing no nilpotents of index two, and with a basis of idempotents, is isomorphic with exactly one of the following:

- 1)  $e_1 \cdot e_1 = e_1, e_2 \cdot e_2 = e_2, e_1 \cdot e_2 = 0$
- 2a)  $e_1 \cdot e_1 = e_1, e_2 \cdot e_2 = e_2, e_1 \cdot e_2 = e_2$
- b)  $e_1 \cdot e_1 = 2e_1, e_2 \cdot e_2 = e_2, e_1 \cdot e_2 = e_2$
- c)  $e_1 \cdot e_1 = \alpha e_1, e_2 \cdot e_2 = e_2, e_1 \cdot e_2 = e_2$ ,  
for  $0 < \alpha < 1$  and  $\alpha > 2$
- 3a)  $e_1 \cdot e_1 = 2e_1, e_2 \cdot e_2 = \beta e_2, e_1 \cdot e_2 = e_1 + e_2$ ,  
with  $\beta \neq 0, \beta \neq 2$
- b)  $e_1 \cdot e_1 = \alpha e_1, e_2 \cdot e_2 = \beta e_2, e_1 \cdot e_2 = e_1 + e_2$ ,  
with  $\alpha \neq 0, \alpha \neq 2, \beta \neq 0, \beta \neq 2, \alpha \beta \neq 4, \alpha + \beta \neq \alpha\beta$ ,  
and  $\frac{2}{\alpha - 1} \leq \beta \leq \frac{2 + \alpha}{\alpha}$  on  $\alpha < -1$  and on  $\alpha > 2$ .

Distinct values of the parameters yield non-isomorphic algebras.

PROOF. Take a basis  $E_1$  and  $E_2$  of idempotents in  $\mathcal{M}$ .

Then  $E_1 \cdot E_2 = cE_1 + dE_2$ .

If  $c = d = 0$ , we have Case 1) of the theorem.

If  $c = 0$  and  $d \neq 0$  write

$$e_1 = \frac{1}{d} E_1, \quad e_2 = E_2$$

to obtain an algebra of the form, Case 2,  $e_1 \cdot e_1 = \alpha e_1$ ,  $e_2 \cdot e_2 = e_2$ ,  $e_1 \cdot e_2 = e_2$ . The parameter restrictions on  $\alpha$  will be explained later.

If  $c \neq 0$  and  $d = 0$ , the algebras are isomorphic to those of Case 2.

If  $c \neq 0$  and  $d \neq 0$ , let

$$e_1 = \frac{1}{d} E_1, \quad e_2 = \frac{1}{c} E_2$$

to obtain an algebra of the form, Case 3,

$$e_1 \cdot e_1 = \alpha e_1, \quad e_2 \cdot e_2 = \beta e_2, \quad e_1 \cdot e_2 = e_1 + e_2.$$

The condition that there are no nilpotents imposes no restriction on Case 1, but requires  $\alpha \neq 0$  in Case 2, and  $\alpha \neq 0$ ,  $\beta \neq 0$ ,  $\alpha\beta \neq 4$  in Case 3.

Next we show that no algebra of one of the above three types 1), 2), or 3) is isomorphic with an algebra of a different type.

In Case 1) the only idempotents are  $e_1$  and  $e_2$  and their product is zero. This characterizes Case 1). In Case 2) the only idempotents are

$$\frac{1}{\alpha} e_1, \quad e_2,$$

and

$$\frac{1}{\alpha} e_1 + (1 - \frac{2}{\alpha})e_2.$$

Here the idempotent  $e_2$  generates a 1-dimensional ideal, yet it does not annihilate the algebra upon multiplication. Thuse Case 2) differs from Case 1). In Case 2) we distinguish the subcases 2a) where  $\alpha = 1$ , for only in this case does the corresponding idempotent  $e_1 - e_2$  also generate an ideal. Also the subcase 2b), where  $\alpha = 2$ , is the only algebra of 2) which contains just two idempotents.

In Case 3), the idempotents are

$$\frac{1}{\alpha} e_1, \quad \frac{1}{\beta} e_2,$$

and

$$\frac{2-\beta}{4-\alpha\beta} e_1 + \frac{2-\alpha}{4-\alpha\beta} e_2.$$

If  $\alpha + \beta \neq \alpha\beta$ , it is easy to see that no idempotent generates a 1-dimensional ideal. Thus Case 3) differs from 1) and 2). Finally we show that  $e_1 \cdot e_1 = \alpha e_1$ ,  $e_2 \cdot e_2 = \beta e_2$ ,  $e_1 \cdot e_2 = e_1 + e_2$  with  $\alpha \neq 0$ ,  $\beta \neq 0$ ,  $\alpha\beta \neq 4$  is an algebra of Case 2) in case  $\alpha + \beta = \alpha\beta$ . Here there are idempotents

$$\bar{E}_1 = \frac{1}{\alpha} e_1, \quad \bar{E}_2 = \frac{2-\beta}{4-\alpha\beta} e_1 + \frac{2-\alpha}{4-\alpha\beta} e_2.$$

If  $\alpha = 2$  and  $\alpha + \beta = \alpha\beta$ , then  $\beta = 2$  so  $\alpha\beta = 4$  which is impossible. Thus  $\alpha \neq 2$  and also  $\beta \neq 2$ . Compute  $\bar{E}_1 \cdot \bar{E}_2 \neq 0$ . But  $\bar{E}_2$  generates a 1-dimensional ideal and thus we have an algebra of Case 2). The subcase 3a),  $\alpha = 2$ , is distinguished in that there are just two idempotents. If  $\beta = 2$  (so  $\alpha \neq 2$ ) we again have Case 3a).

We next study the restrictions on the parameter  $\alpha$  in Case 2) and  $(\alpha, \beta)$  in Case 3), so as to have a unique form for each possible algebra.

Thus consider  $e_1 \cdot e_1 = \alpha e_1$ ,  $e_2 \cdot e_2 = e_2$ ,  $e_1 \cdot e_2 = e_2$  for  $\alpha \neq 0$ ,  $\alpha \neq 1$ ,  $\alpha \neq 2$ . Here  $e_2$  is the unique idempotent which generates an ideal. Using

$$\frac{1}{\alpha} e_1 + \left(1 - \frac{2}{\alpha}\right) e_2$$

and  $e_2$  as a base for the algebra we find that the algebras with values  $\alpha$  and

$$\frac{\alpha}{\alpha-1}$$

are isomorphic and that no other value of the parameter yields an isomorphic algebra. Thus we pick out a single representative for each algebra of Case 2c) by demanding

$$\alpha > \frac{\alpha}{\alpha-1}.$$

Thus  $0 < \alpha < 1$  and  $\alpha > 2$  enforces a one-to-one correspondence between parameter values  $\alpha$  and isomorphism classes of algebras in 2c).

Consider Case 3a) with  $\alpha = 2$  (so  $\beta \neq 2$ ). There are just two idempotents

$$\frac{1}{2} e_1 \quad \text{and} \quad \frac{1}{\beta} e_2.$$

The product of these idempotents is

$$\frac{1}{\beta} \left( \frac{1}{2} e_1 \right) + \frac{1}{2} \left( \frac{1}{\beta} e_2 \right) .$$

Thus isomorphic algebras in 3a) must yield the same value for

$$\frac{1}{\beta} + \frac{1}{2} .$$

Thus  $\beta$  is different for different algebras. The case  $\beta = 2$  (so  $\alpha \neq 2$ ) is isomorphic to an algebra with  $\alpha = 2$  and  $\beta \neq 2$ .

Now turn to Case 3b):

$$e_1 \cdot e_1 = \alpha e_1, \quad e_2 \cdot e_2 = \beta e_2, \quad e_1 \cdot e_2 = e_1 + e_2$$

with

$$\alpha \neq 0, \alpha \neq 2, \beta \neq 0, \beta \neq 2, \alpha\beta \neq 4, \alpha + \beta \neq \alpha\beta .$$

There are three idempotents

$$E_1 = \frac{1}{\alpha} e_1, \quad E_2 = \frac{1}{\beta} e_2 ,$$

and

$$E_3 = \frac{2 - \beta}{4 - \alpha\beta} e_1 + \frac{2 - \alpha}{4 - \alpha\beta} e_2 .$$

The table

$$E_1 \cdot E_2 = \frac{1}{\beta} E_1 + \frac{1}{\alpha} E_2 ,$$

$$E_1 \cdot E_3 = \frac{\alpha + \beta - \alpha\beta}{4 - \alpha\beta} E_1 + \frac{1}{\alpha} E_3 ,$$

$$E_2 \cdot E_3 = \frac{1}{\beta} E_3 + \frac{\alpha + \beta - \alpha\beta}{4 - \alpha\beta} E_2 ,$$

shows that the algebras of 3b) with parameter values  $(\bar{\alpha}, \bar{\beta}) = (\alpha, \beta)$ , or  $(\beta, \alpha)$ , or

$$\left( \alpha, \frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta} \right), \quad \text{or} \quad \left( \frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta}, \alpha \right)$$

or

$$\left( \frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta}, \beta \right), \quad \text{or} \quad \left( \beta, \frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta} \right)$$

are isomorphic. Also these are the only isomorphic parameter sets.

To obtain a unique parametrization of the algebras in 3b), we must find a fundamental domain, in the  $(\alpha, \beta)$ -plane with the deletions  $\alpha \neq 0$ ,  $\alpha \neq 2$ ,  $\beta \neq 0$ ,  $\beta \neq 2$ ,  $\alpha\beta \neq 4$ ,  $\alpha + \beta \neq \alpha\beta$  under the group generated by the transformations

$$T : (\alpha, \beta) \longrightarrow \left( \alpha, \frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta} \right)$$

and

$$R : (\alpha, \beta) \longrightarrow (\beta, \alpha) .$$

We compute  $T^2 = R^2 = I$ , the identity. Also  $RTR = TRT$ . Call  $Z = RT$  (first  $T$  then  $R$ ) and  $Z^3 = I$ ,  $RZ = Z^2R$ . Thus the group consists of six elements  $I, Z, Z^2, R, ZR, \text{ and } Z^2R$ .

Use the fact that

$$\frac{4 - \alpha\beta}{\alpha + \beta - \alpha\beta}$$

is a monotonic decreasing function of  $\beta$ , for each fixed  $\alpha$ . A careful study of the geometry of the six transformations of the group  $I, Z, Z^2, R, ZR, Z^2R$  shows that every allowable point of the  $(\alpha, \beta)$ -plane is equivalent to one and only one point of the fundamental domain

$$\frac{2}{\alpha - 1} \leq \beta \leq \frac{2 + \alpha}{\alpha}$$

with  $\alpha > 2$ , and

$$\frac{2}{\alpha - 1} \leq \beta \leq \frac{2 + \alpha}{\alpha}$$

for  $\alpha < -1$ . Q.E.D.

A long computation based on the identity  $x^2 \cdot x^2 = (x^2 \cdot x) \cdot x$  shows that algebras 1) and 2a) of Theorem 7 are associative but that none of the other algebras of Theorem 7 are power associative.

**THEOREM 8.** A real commutative two-dimensional algebra  $\mathfrak{A}$ , with just one idempotent and no nilpotent elements of index two, is isomorphic with exactly one of the following:

- 1)  $e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = 0, e_2 \cdot e_2 = e_1 + g e_2$ ,  
with  $0 \leq g < 2$



- 2)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = e_1 + e_2$ ,  $e_2 \cdot e_2 = fe_1$ ,  
with  $f < -1$
- 3a)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = \frac{1}{2} e_2$ ,  $e_2 \cdot e_2 = e_1$
- b)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = de_2$ ,  $e_2 \cdot e_2 = e_1 + ge_2$ ,  
with  $d \neq 0$ ,  $g \geq 0$ ,  $g^2 < 4(1 - 2d)$
- c)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = \frac{1}{2} e_2$ ,  $e_2 \cdot e_2 = -e_1$
- d)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = de_2$ ,  $e_2 \cdot e_2 = -e_1 + ge_2$ ,  
with  $d \neq 0$ ,  $d \neq 1$ ,  $g^2 \neq 4d^2$ ,  $g^2 < 4(2d - 1)$ ,  $g \geq 0$ .
- 4)  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = e_2$ ,  $e_2 \cdot e_2 = -e_1$ .

Distinct values of the parameters yield non-isomorphic algebras.

PROOF. Let  $e_1$  be the idempotent element of  $\mathfrak{N}$ . Then there is a basis  $e_1, e_2$  in  $\mathfrak{N}$  such that exactly one of the following three cases holds:

CASE 1:  $e_1 \cdot e_2 = 0$

CASE 2:  $e_1 \cdot e_2 = e_1 + e_2$

CASE 3:  $e_1 \cdot e_2 = de_2$ , for  $d \neq 0$ .

To see this write  $e_1 \cdot e_2 = ce_1 + de_2$ . If  $c = d = 0$  we have Case 1. If  $c \neq 0$ ,  $d = 0$  replace  $e_2$  by  $E_2 = -ce_1 + e_2$  to obtain Case 1. If  $c = 0$ ,  $d \neq 0$  we have Case 3. Finally suppose  $c \neq 0$ , and  $d \neq 0$ . Here replace  $e_2$  by

$$E_2 = \frac{c}{1-d} e_1 - e_2$$

(if  $d \neq 1$ ) to obtain Case 3 again. If  $d = 1$ ,  $c \neq 0$  use

$$E_2 = \frac{1}{c} e_2$$

to obtain Case 2. Thus every  $\mathfrak{N}$  lies in Case 1, 2, or 3.

Also an algebra  $\mathfrak{N}$  can fall in just one of these three cases. For in  $\mathfrak{N}$  the element  $e_1$  is distinguished as the unique idempotent. The linear transformation  $T_1: \mathfrak{N} \rightarrow \mathfrak{N}$  of multiplication by  $e_1$  has a null space in just Case 1. In Case 2  $T_1$  has just one independent eigenvector,  $e_1$ . In Case 3  $T_1$  has two independent eigenvectors.

Now consider an algebra  $\mathfrak{N}$  of Case 1. We consider three subcases designated by:

- 1A)  $e_2 \cdot e_2 = ge_2$  for  $g \neq 0$

$$1B) \quad e_2 \cdot e_2 = fe_1 \quad \text{for } f > 0$$

$$1C) \quad e_2 \cdot e_2 = \pm e_1 + ge_2 \quad \text{for } g \neq 0.$$

Suppose  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = 0$ ,  $e_2 \cdot e_2 = fe_1 + ge_2$ . Sub-case 1A) is inadmissible since it allows a basis of idempotents in  $\mathfrak{M}$ . If  $g = 0$ ,  $f \neq 0$  then there are nilpotents unless  $f > 0$ . If  $g \neq 0$ ,  $f \neq 0$  we have Case 1C), upon replacing  $e_2$  by  $E_2 = |f|^{-1/2}e_2$ .

In Case 1B) replace  $e_2$  by  $E_2 = f^{-1/2}e_2$  to obtain the result  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot E_2 = 0$ ,  $E_2 \cdot E_2 = e_1$  which is listed in the theorem. In Case 1C) the possibility  $e_2 \cdot e_2 = -e_1 + ge_2$  is not allowed since it admits a basis of idempotents. Also in case  $e_2 \cdot e_2 = e_1 + ge_2$  we must have  $g^2 < 4$  to prevent a basis of idempotents. But the values  $+g$  and  $-g$  yield isomorphic algebras under the automorphism  $e_1 \longrightarrow e_1$ ,  $e_2 \longrightarrow -e_2$ . Therefore every algebra  $\mathfrak{M}$  of Case 1 is listed under 1) in the theorem.

It is easy to check that each algebra of 1) has exactly one idempotent and no nilpotent of index two. Also in an algebra  $\mathfrak{M}$  of 1) the element  $e_1$  is distinguished by  $e_1 \cdot e_1 = e_1$ . Also the subspace  $\lambda e_2$  is the null space of  $T_1$ , multiplication by  $e_1$ . The condition  $e_2 \cdot e_2 = e_1 + ge_2$  fixes the pair  $(e_2, -e_2)$ . But then the restriction  $g \geq 0$  shows that no algebra in 1) corresponds to two distinct admissible values of  $g$ .

Now consider Case 2),  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = e_1 + e_2$ ,  $e_2 \cdot e_2 = fe_1 + ge_2$ . If  $g \neq 0$  we replace  $e_2$  by

$$E_2 = -\frac{g}{2}e_1 + e_2$$

to obtain  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot E_2 = e_1 + E_2$ ,  $E_2 \cdot E_2 = he_1$ . Therefore we can always take  $g = 0$  in Case 2. If  $f \geq -1$  there are two independent idempotents in  $\mathfrak{M}$  and so we are lead to the Case 2) listed in the theorem.

Each algebra  $\mathfrak{M}$  of Case 2) has no nilpotent of index two and only one idempotent. In an algebra  $\mathfrak{M}$  of 2) the element  $e_1$  is distinguished by  $e_1 \cdot e_1 = e_1$ . The affine space  $\{e_2 + \alpha e_1\}$  is distinguished by the relation  $e_1 \cdot e_2 = e_1 + e_2$ . But the only member of  $\{e_2 + \alpha e_1\}$  whose square is a scalar multiple of  $e_1$  is  $e_2$ , which is thereby distinguished in  $\mathfrak{M}$ . Thus distinct values of  $f$  yield non-isomorphic algebras in Case 2).

Now consider Case 3,  $e_1 \cdot e_1 = e_1$ ,  $e_1 \cdot e_2 = de_2$ ,  $e_2 \cdot e_2 = fe_1 + ge_2$ , with  $d \neq 0$ . If  $f = 0$  there is either a nilpotent of index two or two idempotents in  $\mathfrak{M}$  so we must have  $f \neq 0$ . Replace  $e_2$  by  $E_2 = |f|^{-1/2}e_2$  and thereby we can assume that  $f = +1$  or

$f = -1$ . Further we separate the case where the operator  $T_1$ , of multiplication by  $e_1$ , is the identity, that is  $d = 1$ . We are then lead to three subcases of 3:

- I)  $e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = e_2, e_2 \cdot e_2 = \pm e_1 + ge_2$   
 II)  $e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = de_2, e_2 \cdot e_2 = e_1 + ge_2,$   
 $d \neq 0, d \neq 1$   
 III)  $e_1 \cdot e_1 = e_1, e_1 \cdot e_2 = de_2, e_2 \cdot e_2 = -e_1 + ge_2,$   
 $d \neq 0, d \neq 1$ .

An algebra  $\mathbb{M}$  of Case 3 has a distinguished element  $e_1$  with  $e_1 \cdot e_1 = e_1$ . Thus the operator  $T_1$  is an algebraic invariant and in 3I) it is the identity. Case 3 II) is distinguished from 3 III) in that the eigenvector  $e_2$  has a square with  $e_2 \cdot e_2 = e_1 + ge_2$ , instead of  $e_2 \cdot e_2 = -e_1 + ge_2$ .

In 3 I) the possibility  $e_2 \cdot e_2 = e_1 + ge_2$  is eliminated since such an algebra always has a basis of idempotents. Also the possibility  $e_2 \cdot e_2 = -e_1 + ge_2$  is restricted since if  $g^2 = 4$  there are nilpotents whereas if  $g^2 > 4$  there are two idempotents.

Thus take  $g^2 < 4$  and define new basis vectors  $E_1 = e_1$ ,  $E_2 = \lambda e_1 + \mu e_2$  where  $\mu > 0$  and  $\lambda$  are defined by

$$\lambda = -\frac{\mu g}{2}, \quad \mu^2 = \frac{4}{4 - g^2}.$$

Then one computes directly  $E_1 \cdot E_1 = E_1, E_1 \cdot E_2 = E_2, E_2 \cdot E_2 = -E_1$ , which is algebra 4) in the theorem. This algebra is the complex numbers.

We next show that in 3 II) the admissible values of the parameters are  $g^2 < 4(1 - 2d)$  and  $g = 0, d = \frac{1}{2}$ . Use  $(\alpha e_1 + \beta e_2)^2 = (\alpha^2 + \beta^2)e_1 + (2\alpha\beta d + \beta^2 g)e_2$  to show that there are neither nilpotents nor idempotents, with  $\beta \neq 0$ , for  $g^2 < 4(1 - 2d)$ . But if  $g^2 > 4(1 - 2d)$  then there are two idempotents and so this is not admissible. If  $g^2 = 4(1 - 2d)$  there are two idempotents in  $\mathbb{M}$  unless  $g = 0, d = \frac{1}{2}$ . Now the interchange of  $e_2$  and  $-e_2$  replaces  $g$  by  $-g$  and so we can take  $g \geq 0$ . Thus algebras of 3 II) are those of Cases 3a) and 3b) in the theorem.

In Case 3b) the arithmetical condition  $g^2 < 4(1 - 2d)$  shows that  $d \neq 1, d \neq \frac{1}{2}$ . Thus 3a) is different from any algebra in 3b). In an algebra  $\mathbb{M}$  of 3b) we can distinguish the idempotent  $e_1$ , then the pair  $\{e_2, -e_2\}$  by  $e_2 \cdot e_2 = e_1 + ge_2$ . Since  $g \geq 0$  there is just one algebra for each admissible parameter value.

Next consider the cases 3 III). We shall show that the admissible

values are exactly those where  $d \neq 0$ ,  $d \neq 1$ ,  $g \neq \pm 2d$ ,  $g^2 < 4(2d - 1)$  and also  $g = 0$ ,  $d = \frac{1}{2}$ . Use  $(\alpha e_1 + \beta e_2)^2 = (\alpha^2 - \beta^2)e_1 + (2\alpha\beta d + \beta^2 g)e_2$ . Since  $g \neq \pm 2d$ , there are no nilpotents of index two. Since  $g^2 < 4(2d - 1)$ , or  $g = 0$ ,  $d = \frac{1}{2}$ , there are no idempotents other than  $e_1$ . Now suppose  $d \neq 0$ ,  $d \neq 1$  but  $g^2 = 4d^2$  or  $g^2 \geq 4(2d - 1)$  (excepting  $g = 0$ ,  $d = \frac{1}{2}$ ). If  $g^2 = 4d^2$  there are nilpotents of index two. If  $g^2 \geq 4(2d - 1)$  then there are idempotents other than  $e_2$  (provided we exempt  $g = 0$ ,  $d = \frac{1}{2}$ ). Therefore we have the algebras of Cases 3c) and 3d) in the theorem.

Finally we must note that two parameter values in 3d) define non-isomorphic algebras. For such an algebra  $\mathfrak{N}$  first distinguish the idempotent  $e_1$ , then the eigenspace of  $e_2$ , then the pair  $(e_2, -e_2)$ . Interchanging  $e_2$  and  $-e_2$  replaces  $g$  by  $-g$  and since  $g \geq 0$  there is a unique algebra for each admissible parameter value. Q. E. D.

Only the algebra 4) in Theorem 8 is associative. Moreover none of the other of these algebras in Theorem 8 is even power associative.

In Theorems 6, 7, and 8 we have found all real commutative algebras of dimension two. An interesting by-product of our classification is the following result.

COROLLARY. The only real commutative two-dimensional algebra which is power associative, yet not associative is:

$$e_1 \cdot e_1 = 0, e_1 \cdot e_2 = e_1, e_2 \cdot e_2 = 2e_2.$$

This is a Jordan algebra, that is, it satisfies the identity  $(uv)u^2 = u(vu^2)$ .

#### §5. QUALITATIVE BEHAVIOR OF QUADRATIC DIFFERENTIAL EQUATIONS IN THE AFFINE PLANE

We shall study the topological behavior of the solution curve families for quadratic differential systems in the plane, with an isolated critical point at the origin. Thus we shall utilize the algebras listed in Theorems 7 and 8.

Consider the algebra 1) of Theorem 7:

$$e_1 \cdot e_1 = e_1, e_2 \cdot e_2 = e_2, e_1 \cdot e_2 = 0.$$

This corresponds to the quadratic differential system

$$\dot{x} = x^2, \quad \dot{y} = y^2.$$

Corresponding to the three idempotents  $e_1$ ,  $e_1 + e_2$ , and  $e_2$  there are three lines through the origin which define solution curves. The computation

$$\dot{\theta} = \frac{x\dot{y} - y\dot{x}}{x^2 + y^2} = \frac{xy}{r^2} (y - x)$$

where  $\theta = \arctan y/x$ , displays these ray solutions along the lines  $x = 0$ ,  $y = 0$ , and  $y - x = 0$ . We designate a ray solution by  $+$  if the radial velocity is positive and  $-$  if the radial velocity is negative. For each sector between the ray solutions we write an arrow to the right  $\longrightarrow$  if the angular velocity is positive and an arrow to the left  $\longleftarrow$  if the angular velocity is negative. Thus the differential system

$$\dot{x} = x^2, \quad \dot{y} = y^2$$

is represented by the combinatorial scheme

$$\longrightarrow + \longleftarrow + \longrightarrow + \longleftarrow - \longrightarrow - \longleftarrow - \longrightarrow,$$

where the first  $+$  (in the order of reading left to right) represents the positive half-axis solution.

The combinatorial scheme of each quadratic differential system, with an isolated critical point and a finite number of solution rays in the plane, is defined similarly. The combinatorial scheme is only specified up to a cyclic ordering; thus

$$\longleftarrow + \longrightarrow + \longleftarrow - \longrightarrow - \longleftarrow - \longrightarrow + \longleftarrow$$

also represents

$$\dot{x} = x^2, \quad \dot{y} = y^2.$$

**DEFINITION.** Two quadratic differential systems,  $q_1$  and  $q_2$ , with isolated critical points in the plane, are called geometrically equivalent in case  $q_1$  has the same cyclic combinatorial scheme as  $q_2$ , or as  $q_2$  transformed by an orientation reversing reflection.

**DEFINITION.** Two autonomous differential systems  $s_1$  and  $s_2$  on a differentiable manifold  $M^n$  are called topologically equivalent in case there exists a homeomorphism of the  $M^n$  onto itself carrying each critical point of  $s_1$  onto one of  $s_2$ , and each directed (but non-

parametrized) solution curve of  $\delta_1$  onto one of  $\delta_2$ , and conversely carrying the solution curves of  $\delta_2$  onto those of  $\delta_1$ . If  $M^n$  is orientable and the homeomorphism is orientation preserving, then  $\delta_1$  and  $\delta_2$  are o-topologically equivalent.

LEMMA. Let  $\phi_1$  and  $\phi_2$  be quadratic differential systems, with isolated critical points and finitely many ray solutions in the affine plane. Let  $\sigma_1$  and  $\sigma_2$  be their corresponding cyclic combinatorial schemes, as above. Then  $\phi_1$  and  $\phi_2$  are o-topologically equivalent if  $\sigma_1$  and  $\sigma_2$  are the same.

PROOF. This is an immediate consequence of the general theory of separatrix configurations developed in Theorem 7.1 of [14]. Q.E.D.

It should be noted that topologically equivalent differential systems can fail to be equivalent under a differentiable homeomorphism.

THEOREM 9. Each quadratic differential system, with an isolated critical point in the affine plane, is geometrically equivalent to exactly one of the following six differential systems:

- I.  $\dot{x} = x^2$ ,  $\dot{y} = y^2$  which represent Cases 1), 2a), 2c), and 3b) for  $\alpha > 2$ ,  $0 < \beta < 2$ ,  $4 - \alpha\beta < 0$ , and 3b) for  $\alpha < -1$ ,  $-1 < \beta < 0$  in Theorem 7.
- II.  $\dot{x} = 3x^2 + 2xy$ ,  $\dot{y} = y^2 + 2xy$ , which represents Cases 3b) for  $\alpha > 2$ ,  $0 < \beta < 2$ ,  $4 - \alpha\beta > 0$ , and 3b) for  $\alpha < -1$ ,  $0 < \beta < 1$ , in Theorem 7.
- III.  $\dot{x} = 2x^2 + 2xy$ ,  $\dot{y} = -y^2 + 2xy$ , which represents Cases 2b), 3a) for  $\beta < 0$ , and 3a) for  $\beta > 2$ , in Theorem 7.
- IV.  $\dot{x} = 2x^2 + 2xy$ ,  $\dot{y} = y^2 + 2xy$ , which represents Cases 3a) for  $0 < \beta < 2$  in Theorem 7.

All of the above have four or six directions of approach to the origin, and next there are systems with just two directions of approach.

- V.  $\dot{x} = x^2 + y^2$ ,  $\dot{y} = y^2$  which represents Cases 1), 3a), and 3b) of Theorem 8.
- VI.  $\dot{x} = x^2 - 2y^2 + 2xy$ ,  $\dot{y} = 2xy$  which represents Cases 2), 3c), 3d) and 4) of Theorem 8.

PROOF. By examination of

$$\dot{\theta} = \frac{x\dot{y} - y\dot{x}}{x^2 + y^2}$$

we sketch the solution curves and find the schemes for the quadratic differential systems as follows.

In Theorem 7, we find

- 1)  $\longrightarrow + \langle \longrightarrow + \longrightarrow \rangle + \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle$
- 2a)  $\langle \longrightarrow + \longrightarrow \rangle + \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle + \langle \longrightarrow$
- 2b)  $\longrightarrow + \longrightarrow + \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle$
- 2c)  $\longrightarrow + \langle \longrightarrow + \longrightarrow \rangle + \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle$  for all  $\alpha$
- 3a)  $\longrightarrow + \langle \longrightarrow + \longrightarrow \rangle - \longrightarrow$  for  $\beta < 0$   
 $\langle \longrightarrow + \longrightarrow \rangle - \longrightarrow - \langle \longrightarrow$  for  $0 < \beta < 2$   
 $\longrightarrow + \longrightarrow + \langle \longrightarrow - \longrightarrow \rangle$  for  $\beta > 2$
- 3b)  $\langle \longrightarrow + \longrightarrow \rangle + \langle \longrightarrow + \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow$   
for  $\alpha > 2$ ,  $0 < \beta < 2$ ,  $4 - \alpha\beta > 0$ , and for  
 $\alpha < -1$ ,  $0 < \beta < 1$   
 $\longrightarrow + \langle \longrightarrow + \longrightarrow \rangle + \langle \longrightarrow - \longrightarrow \rangle - \langle \longrightarrow - \longrightarrow \rangle$   
for  $\alpha > 2$ ,  $0 < \beta < 2$ ,  $4 - \alpha\beta < 0$  and for  
 $\alpha < -1$ ,  $-1 < \beta < 0$ .

In Theorem 8, we find

- 1)  $\longrightarrow + \langle \longrightarrow - \longrightarrow \rangle$  for all  $g$
- 2)  $\langle \longrightarrow + \longrightarrow \rangle - \langle \longrightarrow$  for all  $f$
- 3a)  $\longrightarrow + \langle \longrightarrow - \longrightarrow \rangle$
- 3b)  $\longrightarrow + \langle \longrightarrow - \longrightarrow \rangle$  for all  $d$  and  $g$
- 3c)  $\langle \longrightarrow + \longrightarrow \rangle - \langle \longrightarrow$
- 3d)  $\langle \longrightarrow + \longrightarrow \rangle - \langle \longrightarrow$  for all  $d$  and  $g$
- 4)  $\langle \longrightarrow + \longrightarrow \rangle - \langle \longrightarrow$

An examination of these combinatorial schemes, and those corresponding to the above differential systems after an orientation reversing map  $x \longrightarrow -x$ ,  $y \longrightarrow y$ , yields the geometrical classification stated in the theorem. Q.E.D.

A necessary and sufficient condition that quadratic differential equations  $\mathcal{Q}_1$  and  $\mathcal{Q}_2$ , with isolated critical points in the plane, be topologically equivalent is that they have the same separatrix configurations, cf. [14]. An examination of Figures I through VI shows that I and




Fig. I.




Fig. II.




Fig. III.




Fig. IV.



Fig. V.




Fig. VI.



III are topologically equivalent and also II and IV are topologically equivalent. Thus there are only four topologically different planar quadratic differential systems having isolated critical points.

# §6. QUADRATIC DIFFERENTIAL EQUATIONS AND LINE ELEMENT FIELDS IN REAL PROJECTIVE SPACE

Consider a quadratic differential system in the affine  $n$ -space  $R^n$

$$\varphi) \quad \frac{dx^i}{dt} = \sum_{j,k=1}^n a_{jk}^i x^j x^k \quad i = 1, 2, \dots, n.$$

Now  $\varphi$  defines a non-directed line-element field  $\mathcal{L}$  in  $R^n$ , with the critical points of  $\varphi$  deleted. We shall only consider  $\varphi$  with isolated critical points in  $R^n$  in this section.

Now  $\mathcal{L}$  can be extended to a unique differentiable line element field on the real projective space  $P^n$ , with a minimal set of critical points occurring on the  $P^{n-1}$  at infinity. We define linear equivalence of two line-element fields  $\mathcal{L}_1$  and  $\mathcal{L}_2$  on  $R^n$ , and topological equivalence of  $\mathcal{L}_1$  and  $\mathcal{L}_2$  on either  $R^n$  or  $P^n$  just as was done for differential systems (except that here the solution curves have no direction or sense).

THEOREM 10. Let  $\varphi_1$  and  $\varphi_2$  be quadratic differential systems, with isolated critical points in  $R^n$ . Let the corresponding line-element fields be  $\mathcal{L}_1$  and  $\mathcal{L}_2$  in  $R^n$ . If  $\varphi_1$  and  $\varphi_2$  are linearly equivalent in  $R^n$ , then so are  $\mathcal{L}_1$  and  $\mathcal{L}_2$ . Conversely if  $\mathcal{L}_1$  and  $\mathcal{L}_2$  are linearly equivalent in  $R^n$ , then  $\varphi_1$  is linearly equivalent to  $\varphi_2$ , or to the quadratic differential system  $c\varphi_2$  obtained by multiplying each coefficient of  $\varphi_2$  by the fixed positive number  $c$ .

PROOF. If  $\varphi_1$  and  $\varphi_2$  are linearly equivalent in  $R^n$ , then certainly  $\mathcal{L}_1$  and  $\mathcal{L}_2$  are linearly equivalent in  $R^n$ .

Next suppose that after a linear transformation of  $R^n$ ,  $\mathcal{L}_1$  coincides with  $\mathcal{L}_2$ . Then there are two quadratic differential systems

$$\varphi_2) \quad \dot{x}^i = p^i(x), \quad i = 1, 2, \dots, n,$$

and the transform of  $\varphi_1$ , say

$$\hat{\varphi}_1) \quad \hat{x}^1 = \hat{P}^1(x) \quad , \quad 1 = 1, 2, \dots, n \quad ,$$

which yield the same line-element field. Therefore  $P^1(x) = \lambda(x)\hat{P}^1(x)$  where  $\lambda(x) \neq 0$  in  $R^n - \{0\}$ , the space  $R^n$  with the origin deleted. Since

$$\lambda(x)^2 = \sum_{i=1}^n [P^i(x)]^2 \bigg/ \sum_{i=1}^n [\hat{P}^i(x)]^2 \quad ,$$

we see that  $\lambda(x)$  is a real analytic function on  $R^n - \{0\}$ . By the identity theorem for power series,  $\lambda(x)$  is a constant in a neighborhood of each point and is therefore everywhere constant in  $R^n - \{0\}$ . Thus  $\lambda(x) = \lambda_0$ .

Since  $\varphi_1$  and  $-\varphi_1$  are linearly equivalent, we can take  $\lambda_0 = c > 0$ . Q.E.D.

**THEOREM 11.** Let  $\varphi_1$  and  $\varphi_2$  be quadratic differential systems, with isolated critical points in  $R^n$ . Let  $\mathcal{L}_1$  and  $\mathcal{L}_2$  be the corresponding line-element fields in the real projective space  $P^n$ . If  $\varphi_1$  and  $\varphi_2$  are linearly equivalent in  $R^n$ , then  $\mathcal{L}_1$  and  $\mathcal{L}_2$  are topologically equivalent in  $P^n$ .

**PROOF.** A linear transformation of  $R^n$  onto itself can be extended to a projective transformation of  $P^n$  onto itself. This projective transformation is a homeomorphism of  $P^n$  onto itself which carries the solution curves of  $\mathcal{L}_1$  onto those of  $\mathcal{L}_2$ , since  $\mathcal{L}$  is a unique extension of  $\varphi$ . Q.E.D.

It may happen that a topological equivalence of  $\mathcal{L}_1$  and  $\mathcal{L}_2$  in  $R^n$  can not be extended to a topological equivalent in  $P^n$ . We shall not classify line-element fields in  $P^n$ ; not even in the projective plane  $P^2$ .

## §7. QUADRATIC DIFFERENTIAL EQUATIONS IN 3-SPACE $R^3$

We shall only investigate real commutative 3-dimensional algebras which are the direct product of a 2-dimensional algebra and the real numbers  $R$ .

**REMARK.** Let  $\mathfrak{A}$  be a real commutative 2-dimensional algebra

THE LINE-ELEMENT FIELD FOR  $\dot{x} = x^2, \dot{y} = y^2$  IN  $P^2$




FIGURE VII

and  $R$  the real numbers. Then  $\mathfrak{N} \times R$  contains a nilpotent of index two if and only if  $\mathfrak{N}$  does. This is easily seen from the multiplication  $(e_1, \lambda_1) \cdot (e_2, \lambda_2) = (e_1 \cdot e_2, \lambda_1 \lambda_2)$  in  $\mathfrak{N} \times R$ .

**THEOREM 12.** Let  $\mathfrak{N}_1$  and  $\mathfrak{N}_2$  be real commutative 2-dimensional algebras containing no nilpotents of index two. Then  $\mathfrak{N}_1 \times R$  is isomorphic with  $\mathfrak{N}_2 \times R$  if and only if  $\mathfrak{N}_1$  is isomorphic with  $\mathfrak{N}_2$ .

**PROOF.** Assume  $\mathfrak{N}_1 \times R$  is isomorphic with  $\mathfrak{N}_2 \times R$ . Now  $\mathfrak{N}_1$  (or  $\mathfrak{N}_1 \times 1$ ) is a 2-dimensional ideal in  $\mathfrak{N}_1 \times R$ . We shall consider the image  $\mathfrak{N}_1'$  of  $\mathfrak{N}_1$  in  $\mathfrak{N}_2 \times R$ .

Let  $e_1, e_2, e_3$  be a basis for  $\mathfrak{N}_2 \times R$  such that  $e_1, e_2$  are a basis for  $\mathfrak{N}_2$ . Take  $e_1, e_2$  so that  $\mathfrak{N}_2$  has one of the canonical forms for commutative 2-dimensional algebras found in Theorems 7 and 8,

and also demand  $e_1 \cdot e_3 = e_2 \cdot e_3 = 0$ ,  $e_3 \cdot e_3 = e_3$ .

If  $\mathbb{M}_1$  and  $\mathbb{M}_2$  are each isomorphic with the algebra  $\mathcal{U}$ , having a basis of orthogonal idempotents (algebra 1) in Theorem 7), then the theorem holds.

Consider the 2-dimensional subalgebra  $\mathbb{M}'_1$  in  $\mathbb{M}_2 \times \mathbb{R}$ . If  $\mathbb{M}'_1 = \mathbb{M}_2 \times e_3$  the theorem holds. Otherwise choose a basis  $u = \alpha e_1 + \beta e_2 + \gamma e_3$  and  $v = \xi e_1 + \eta e_2 + \zeta e_3$  of  $\mathbb{M}'_1$ . Since  $\mathbb{M}'_1$  is an ideal  $u \cdot e_3 = \gamma e_3$  must lie in  $\mathbb{M}'_1$ . Then

1)  $\gamma = \xi = 0$  in which case  $\mathbb{M}'_1 = \mathbb{M}_2$  and the theorem holds, or

2)  $\gamma \neq 0$  (or  $\xi \neq 0$ ) and  $e_3$  is in  $\mathbb{M}'_1$ .

In the second case choose a basis  $w = \lambda e_1 + \mu e_2$  and  $e_3$  for  $\mathbb{M}'_1$ . In fact  $w \cdot w = vw$  for some real  $v \neq 0$ , so  $\mathbb{M}'_1 \cong \mathcal{U}$ . In this case  $\mathbb{M}_1 \cong \mathcal{U}$ .

Now repeat the argument for the image  $\mathbb{M}'_2$  of  $\mathbb{M}_2$  in  $\mathbb{M}_1 \times \mathbb{R}$  to show that either  $\mathbb{M}'_2 = \mathbb{M}_1 \times 1$  or else  $\mathbb{M}'_2 \cong \mathcal{U}$ . In any case  $\mathbb{M}_1 \cong \mathbb{M}_2$ . Q.E.D.

Thus we have constructed an infinite class of quadratic differential systems in  $\mathbb{R}^3$ , with isolated critical points, such that no two of them are linearly equivalent.

#### BIBLIOGRAPHY

- [1] ALBERT, A. A., "Power associative rings," Trans. Am. Math. Soc. 64 (1948), pp. 552-593.
- [2] BROUWER, L. E. J., "Over de loodrechte trajectorien der baankrommen eener vlakke eenladige projectieve groep," Nieuw. Archief. (2) 11, (1915), pp. 265-290.
- [3] COLEMAN, C., Homogeneous differential equations, Princeton University thesis, 1955.
- [4] DULAC, H., Points singuliers des équations différentielles, Mém. Sci. Math. Fasc. 61, Paris (1934).
- [5] ENGEL, J. H., "A verification of Lanchester's Law," J. of Op. Research Soc. of Am. (1954), pp. 163-171.
- [6] FORSTER, H., "Über das Verhalten der Integralkurven einer gewöhnlichen Differentialgleichung erster Ordnung in der Umgebung eines singulären Punktes," Math. Zeitsch. 43, (1938), pp. 271-320.
- [7] FROMMER, M., "Die Integralkurven einer gewöhnlichen Differenzialgleichung in der Umgebung rationaler Unbestimmtheitsstellen," Math. Ann. 99, (1928), pp. 222-272.
- [8] GAVURIN, M., "On systems of differential equations of the form  $y' = Ay^2 - 2By + C$ ," Doklady Akad. Nauk, SSSR 84, (1952), p. 205.

- [9] KASNER, E., "Solutions of the Einstein equations involving functions of only one variable," Trans. Am. Math. Soc. 27, (1925), pp. 155-162.
- [10] KESTIN, J. and ZAREMBA, A., "Geometrical methods in the analysis of ordinary differential equations," Appl. Sci. Research Bull. 3, (1953), pp. 149-189.
- [11] LEFSCHETZ, S., "Notes on Differential Equations," Contributions II, Princeton Annals Studies 29, (1952), pp. 61-73.
- [12] ———, "The ambiguous case in planar differential systems," Bol. de la Soc. Mat. Mexicana (1957), pp. 63-74.
- [13] LIAGHINA, L., "The integral curves of the equations  $y' = \frac{ax^2+bxy+cy^2}{dx^2+exy+fy^2}$ ," Uspeki Matem. Nauk, VI, 2(42), (1951), pp. 171-183.
- [14] MARKUS, L., "Global structure of ordinary differential equations in the plane," Trans. Am. Math. Soc. 76, (1954), pp. 127-148.
- [15] NEMICKII, V. and STEPANOV, V., Qualitative Theory of Differential Equations, Princeton, 1959, esp. Ch. II.
- [16] SANSONE, G. and CONTI, R., Equazioni Differenziali Non Lineari, Rome, 1956, esp. Chapter II.
- [17] STEBAKOV, S., "Qualitative investigation of the system  $\dot{x} = P(x, y)$ ,  $\dot{y} = Q(x, y)$  by means of isoclines," Doklady Akad. Nauk. SSSR, 82, (1952), pp. 677-680.
- [18] VOLTERRA, V., Leçons sur la théorie mathématique de la lutte pour la vie, Paris, 1931.

University of Minnesota  
Minneapolis, Minnesota



# IX. SUR UNE PROPRIÉTÉ DE L'ENSEMBLE DES TRAJECTOIRES BORNÉES DE CERTAINS SYSTÈMES DYNAMIQUES

Georges Reeb

Nous exposerons la remarque (cf. théorème 1) qui fait l'objet de la présente note sur un cas particulier, les généralisations dont cette remarque est susceptible étant immédiates.

Considérons la variété à bord  $S_n \times I$  (où  $I$  est l'intervalle  $[-1, 1]$  et où  $S_n$  est la sphère à  $n$  dimensions munie de la structure d'espace de Riemann usuelle) et désignons par  $E$  un champ de vecteurs sur  $S_n \times I$  (de classe  $C_1$ ) vérifiant les propriétés suivantes:

- (i)  $E$  ne s'annule en aucun point de  $S_n \times I$ .
- (ii) Sur  $S_n \times \{-1\}$  le champ  $E$  est dirigé vers l'intérieur de  $S_n \times I$ , tandis que sur  $S_n \times \{1\}$  il est dirigé vers l'extérieur de  $S_n \times I$ .

Il est clair que de tels champs  $E$  existent. Parmi les trajectoires de  $E$  nous pouvons distinguer les trajectoires entrantes (c. à d. celles qui rencontrent  $S_n \times \{-1\}$ ) et les trajectoires sortantes (celles qui rencontrent  $S_n \times \{1\}$ ). Désignons la réunion des trajectoires entrantes par  $R$  et la réunion des trajectoires sortantes par  $S$ . Les ensembles  $S$  et  $R$  sont ouverts et homéomorphes à  $S_n \times [0, 1]$  si  $S \cap R = \emptyset$ .

Nous désignerons par  $H$  l'hypothèse suivante:

$$(H) \quad S \cap R = \emptyset.$$

Le complémentaire compact de  $S \cap R$  est noté  $A$ . L'ensemble  $A$  mérite d'être appelé l'ensemble des trajectoires bornées de  $E$ .

Si  $H$  est vérifiée, l'ensemble  $A$  jouit d'un certain nombre de propriétés dont voici quelques unes des plus importantes:

$$(1) \quad H(A) \cong H(S_n) \quad (\text{en homologie de Čech})$$

(La propriété (1) est évidente si on remarque que  $S$  et  $R$  sont homéomorphes à  $S_n \times [0, 1]$  et que l'homologie de  $S_n/A$  vérifie l'excision complète).

(2) l'ensemble  $A$  admet des transformations, homotopes à l'identité, sans points fixes.

(La propriété (2) est évidente puisque  $A$  est un ensemble de trajectoires bornées).

La remarque que nous avons en vue résulte de (1) et (2):

THÉORÈME 1. Si  $n$  est pair et si  $H$  est vérifiée, alors  $A$  n'est pas homéomorphe à un polyèdre, ni localement connexe.

En effet si  $A$  était localement connexe le théorème de points fixes lui serait applicable donc  $\chi(A) = 0$  et il y aurait contradiction avec  $\chi(A) = \chi(S_n) = 2$ .

Le théorème met en évidence une propriété topologique de  $A$ , assez intéressante sous réserve d'établir les deux points suivants:

- (i)  $H$  est vérifiée par certains champs  $E$  même si  $n$  est pair.
- (ii) si  $n$  est impair, l'ensemble  $A$  peut-être un polyèdre.

Nous allons maintenant construire deux exemples qui apporteront la preuve de ces deux points.

1. Exemple montrant que si  $n$  est impair il se peut que  $A$  soit un polyèdre

Un tel exemple est très facile à fabriquer et correspond à un phénomène bien familier. Soit  $u$  un champ de vecteurs de classe  $C_1$  sur  $S_n$ , ne s'annulant en aucun point de  $S_n$ . On définira  $E(x, t)$  par ses deux composantes dans  $S_n \times I$ , à savoir:

$$E(x, t) = [u(x), |t|] \quad .$$

Ici l'ensemble  $A$  se réduit à  $S_n \times \{0\}$ .

2. Exemple montrant que  $H$  peut-être vérifiée même si  $n$  est pair

Nous allons décrire cet exemple dans le cas particulier où  $n = 2$ , le cas général pouvant être calqué assez simplement sur le cas où



$n = 2$ . A cet effet on considérera  $S_2 \times I$  comme l'espace obtenu en recollant convenablement (le bord  $S_n \times \{1\}$  étant recollé sur  $S_n \times \{-1\}$ ) quatre exemplaires de  $S_2 \times I$ ; nous définirons  $E$  dans chacun de ces exemplaires, d'où le champ  $E$  total.

Mais au préalable introduisons dans  $S_2$  un axe polaire et un système de coordonnées sphériques ( $\theta$  latitude,  $\varphi$  longitude). Les calottes définies par

$$1 \leq \theta \leq \frac{\pi}{2} \text{ et } -\frac{\pi}{2} \leq \theta \leq -1$$

seront désignées respectivement par  $\Gamma_0$  et  $\Gamma_1$ ; la zone  $-1 \leq \theta \leq 1$  est notée  $B$ . On désignera par  $e$  le champ de vecteurs de composantes  $(\sigma, \cos \theta)$ ; ce champ est analytique et admet les parallèles de  $S_2$  comme lignes intégrales. Une rotation de  $\frac{\pi}{2}$  autour d'un diamètre équatorial de  $S_2$  transformera les coordonnées  $(\theta, \varphi)$  et les éléments  $\Gamma_0, \Gamma_1, B, e$  en nouveau système de coordonnées et de nouveaux éléments qui seront notés  $(\bar{\theta}, \bar{\varphi}), \bar{\Gamma}_0, \bar{\Gamma}_1, \bar{B}, \bar{e}$ . Nous définissons maintenant  $E$  dans les quatres exemplaires de  $S_2 \times I$ :

(i) Premier exemplaire:

Les composantes de  $E$  sont:

$$\begin{aligned} (e_0, 1 - t) & \quad \text{s1 } (\theta, \varphi) \in B \\ (e_0, (1 - t) + (\theta - 1)) & \quad \text{s1 } (\theta, \varphi) \in \Gamma_0 \\ (e_0, (1 - t) - (\theta + 1)) & \quad \text{s1 } (\theta, \varphi) \in \Gamma_1 \end{aligned}$$

On remarque que sur  $S_2 \times \{-1\}$  le champ  $E$  est dirigé vers l'intérieur de  $S_2 \times I$ . Les trajectoires de  $E$  ne traversent pas  $B \times \{1\}$  mais elles traversent  $\Gamma_0 \times \{1\}$  et  $\Gamma_1 \times \{1\}$ .

(ii) Deuxième exemplaire:

Les composantes de  $E$  sont:

$$\begin{aligned} (e_0, \frac{(1-t)}{2}, \frac{t+1}{2}) & \quad \text{s1 } (\theta, \varphi) \in B \\ (e_0, \frac{(1-t)}{2}), ((1+t) + (\theta - 1)) (\frac{(1-t)}{2} + \frac{1+t}{2}) & \quad \text{s1 } (\theta, \varphi) \in \Gamma_0 \\ (e_0, \frac{(1-t)}{2}), ((1+t) - (\theta + 1)) \frac{1-t}{2} + \frac{1+t}{2} & \quad \text{s1 } (\theta, \varphi) \in \Gamma_1 \end{aligned}$$

On remarquera que  $t = 1$  implique  $E(x, 1) = (\sigma, 1)$ .

(iii) Troisième exemplaire:

$$((t+1)e_1, \frac{1-t}{2}) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \bar{B}$$

$$((t+1)e_1, \frac{1-t}{2} + (\bar{\theta}-1)(1+t)) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \Gamma_0$$

$$((t+1)e_1, \frac{1-t}{2} - (\bar{\theta}+1)(1+t)) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \Gamma_1$$

(iv) Quatrième exemplaire:

$$(2e_1, \frac{t+1}{2}) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \bar{B}$$

$$(2e_1, 2(\bar{\theta}-1) + (t+1)) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \bar{\Gamma}_0$$

$$(2e_1, -2(\bar{\theta}+1) + (t+1)) \text{ si } (\bar{\theta}, \bar{\varphi}) \in \bar{\Gamma}_1 .$$

Le champ  $E$  que nous venons de définir n'est pas exactement de classe  $C_1$  (en effet sur les zones de recollement et sur  $B \cap (\Gamma_0 \cup \Gamma_1) \times I$ ) le champ  $E$  est seulement de classe  $C_0$ ). Cependant notre exemple conserve sa valeur démonstrative; il serait d'ailleurs facile, en modifiant légèrement les définitions analytiques, de donner effectivement un exemple de classe  $C_\infty$  (Voir également la dernière phase de cette note).

Pour voir que notre exemple établit effectivement la propriété voulue il suffit de remarquer que les trajectoires de  $E$  qui entrent par  $B \times \{-1\}$  sont arrêtées dès le bord du premier exemplaire de  $S_2 \times I$ , tandis que les trajectoires qui entrent par  $(\Gamma_0 \cup \Gamma_1) \times \{-1\}$  traversent les deux premiers exemplaires de  $S_2 \times I$  mais sont arrêtées au bord du troisième (parce que les calottes  $\Gamma_0$  et  $\Gamma_1$  ont été choisies suffisamment petites).

Il reste une dernière remarque: les exemples ci-dessus sont peut-être critiquables parce qu'une légère modification de  $E$  peut entraîner la non-validité de  $H$ ; on peut facilement parer à ce dernier inconvénient comme le suggère dans le cas où  $n$  est impair l'exemple du champ:

$$(u(x)t, (t + \frac{1}{2})(t - \frac{1}{2})) .$$

Si  $n$  est pair on peut également faire une construction analogue. Cette dernière remarque, combinée avec le théorème d'approximation de Weierstrass établit le fait suivant: le théorème 1 reste valable même si le champ  $E$  est supposé analytique.

# X. ON LAGRANGE STABLE MOTIONS IN THE NEIGHBORHOOD OF CRITICAL POINTS

Pinchas Mendelson

1. We consider an autonomous system of ordinary differential equations

$$(1.1) \quad \frac{dx_i}{dt} = f_i(x_1, \dots, x_n), \quad (i = 1, 2, \dots, n),$$

defined in some region  $\Omega$  in the  $n$ -dimensional Euclidean space  $E^n$  and having a unique solution there.

We shall employ the standard vector notation: the  $n$ -tuple  $(x_1, \dots, x_n)$  will be denoted by  $x$ .

Let  $P$  be a point of  $\Omega$ . The (unique) trajectory or orbit, of system (1.1), which passes through  $P$ , is denoted by  $\Gamma_P$ ; its parametrization by (the time)  $t$  is assumed to be chosen in such a way that  $\Gamma_P(0) = P$ . The positive semi-orbit  $\{\Gamma_P(t) \mid t \geq 0\}$  and the negative semi-orbit  $\{\Gamma_P(t) \mid t \leq 0\}$  are designated by  $\Gamma_P^+$  and  $\Gamma_P^-$ , respectively.

Let  $\omega$  be an open subset of  $\Omega$  and let  $H(\omega)$  be the boundary of  $\omega$  in  $\Omega$ . A point  $p \in H(\omega)$  is said to be a point of egress from the set  $\omega$  (with respect to the system (1.1) and the set  $\Omega$ ) if there exists an  $\eta_1 > 0$  such that  $\Gamma_P(t) \in \omega$  for all  $-\eta_1 < t < 0$ . If, moreover, there exists an  $\eta_2 > 0$  such that  $\Gamma_P(t) \in \Omega - \bar{\omega}$  for all  $0 < t < \eta_2$ , then  $P$  is said to be a point of strict egress from  $\omega$ .

Consider  $\omega$  fixed. The set of all points of egress from  $\omega$  will be called  $E$ . The set of all points of strict egress from  $\omega$  will be called  $E^*$ . Clearly  $E^* \subset E$ .

If  $P \in \omega$  and  $\Gamma_P^+ \not\subset \omega$ , then  $\Gamma_P^+$  must intersect  $E \subset H(\omega)$ ; we denote the first such point of intersection by  $\tilde{P}$  (note that the open arc of the semi-orbit  $\Gamma_P^+$  between  $P$  and  $\tilde{P}$  is contained in  $\omega$ ).

2. The following theorem, due to T. Ważewski ([3], [4]) will be used in the sequel:

THEOREM (T. Ważewski). Let every point of egress from  $\omega$  (with respect to the system (1.1) and the set  $\Omega$ ) be a point of strict egress, and let a certain set  $Z$  satisfy the following conditions:

- (2.1)            (i)  $Z \subset \omega \cup E$   
                   (ii)  $Z \cap E$  is a deformation retract of  $E$   
                   (iii)  $Z \cap E$  is not a deformation retract of  $Z$ ,

then there exists a point  $P_0 \in \omega \cap Z$  such that  $\Gamma_{P_0}^+ \subset \omega$ .

The above theorem, sometimes referred to as Ważewski's principle, was formulated by A. Pliś [2] in a somewhat more general form. His statement of the principle is as follows:

THEOREM (A. Pliś). Let  $E = E^*$  and let certain sets  $Z, E_1$  satisfy the following conditions:

- (2.2)            (i)  $E_1 \subset E$   
                   (ii)  $Z \subset \omega \cup E_1, Z \neq \emptyset$   
                   (iii)  $E_1$  is not a quasi-isotopic deformation retract<sup>1</sup> of  $Z \cup E_1$  in  $\omega \cup E_1$ ,

then there exists a point  $P_0 \in \omega \cap Z$  such that either  $\Gamma_{P_0}^+ \subset \omega$  or  $\tilde{P}_0 \in E - E_1$ .

The results contained in this note will be obtained by a suitable application of Ważewski's principle in the neighborhood of an isolated critical point of system (1.1).

3. Let  $\Phi$  be a system such as (1.1) and let  $O \in \Omega$  be a (not necessarily isolated) critical point of  $\Phi$ . Let  $\omega$  be an open topological  $n$ -sphere containing  $O$  in its interior; we assume  $\omega$  small enough so

<sup>1</sup> Let  $I$  denote the closed unit interval  $[0, 1]$ . A set  $A$  is a quasi-isotopic deformation retract of a set  $B$  in the topological space  $E$  if there exists a continuous mapping  $\varphi: B \times I \rightarrow E$  having the following properties:

- (i)  $\varphi(P, 0) = P$  for all  $P \in B$ .  
 (ii)  $\varphi(P, s) = P$  for all  $P \in A$  and any  $0 \leq s \leq 1$ .  
 (iii)  $\varphi(P, 1) \in A$  for all  $P \in B$ .  
 (iv)  $\varphi_s: B \rightarrow E$  defined by  $\varphi_s(P) = \varphi(P, s), P \in B, s$  fixed, is a homeomorphism for all  $0 \leq s < 1$ .

that  $\omega \subset \Omega$ . The sets  $E \subset H(\omega)$  and  $E^* \subset H(\omega)$  are comprised, as above, of the points of egress and the points of strict egress from  $\omega$ , respectively.

**THEOREM.** The equality  $E = E^*$  is a sufficient condition for the existence of a semi-orbit (other than 0) wholly contained in  $\omega$ .

**PROOF.** We always have  $E^* \subset E$ . If  $E$  is empty the equality  $E^* = E$  holds trivially. In this case it follows from the definition of  $E$  that no motion which starts in  $\omega$  can intersect  $H(\omega)$  when continued in the positive direction. In other words, every orbit which passes through a point  $Q \in \omega$  satisfies  $r_Q^+ \subset \omega$ .

Similarly, if  $H(\omega) - E$  is empty and  $E = E^*$ , every point of  $H(\omega)$  is a point of strict egress from  $\omega$ ; therefore no motion which starts in  $\omega$  can intersect  $H(\omega)$  when continued in the negative direction. Every orbit which passes through  $Q \in \omega$  satisfies  $r_Q^- \subset \omega$ .

We may therefore assume that  $E = E^*$  is a non-empty proper subset of  $H(\omega)$ .

The set  $E$  may be connected, or it may be disconnected; in the latter case we denote its components by  $E_\alpha$  where  $\alpha$  ranges over some index set  $\mathfrak{M}$ . We shall treat these cases separately.

Suppose first that  $E$  is disconnected. Then  $E = \bigcup_{\alpha \in \mathfrak{M}} E_\alpha$  and the cardinality of  $\mathfrak{M}$  is at least 2.

Let  $P$  be any point of  $\omega$  other than 0. Let  $P_\alpha$  be an arbitrary point of  $E_\alpha$  and let  $\pi_\alpha$  be a polygonal path with the following properties:

- (i)  $\pi_\alpha$  has  $P$  for its initial point and  $P_\alpha$  for its terminal point.
- (ii) The set  $\pi_\alpha$ , except for its terminal point  $P_\alpha$ , is contained in  $\omega - 0$ .

Let  $Z = \bigcup_{\alpha \in \mathfrak{M}} (\pi_\alpha \cup E_\alpha)$ . Clearly the set  $Z$  is connected. Furthermore, it follows from the above construction that  $Z \cap E = \bigcup_{\alpha \in \mathfrak{M}} E_\alpha$  and that  $0 \notin Z$ .

Thus  $Z \cap E = E$  is (trivially) a deformation retract of  $E$ . On the other hand,  $Z \cap E$  — being disconnected — is clearly not a deformation retract of  $Z$ , which is connected. Ważewski's theorem may therefore be applied to the set  $Z$  and we conclude that there exists a point  $Q \in Z$  such that the semi-orbit  $r_Q^+ \subset \omega$ . This completes the proof in the case when  $E$  is disconnected.

Assume next that  $E$  has only one component. Let  $P$  be an arbitrary point of  $E$ . Let  $H_0(0)$  be the surface of a sphere of radius  $\delta$  and center  $O$ ;  $\delta$  is assumed small enough so that  $H_0(0) \subset \omega$ . Let  $\pi$  be a polygonal path with the following properties:

- (i) The initial point of  $\pi$  is on  $H_0(0)$ ; its terminal point is  $P$ ,
- (ii) The set  $\pi$ , except for its terminal point  $P$  is contained in  $\omega - O$ .

Let  $Z = H_0(0) \cup \pi \cup E$ . Clearly  $Z \cap E = E$  and  $Z \neq \emptyset$ . Let  $\omega_1 = \omega - O$ . Then  $H(\omega_1) = H(\omega) \cup O$ . Since  $O$  is a critical point,  $O$  is not a point of egress from  $\omega_1$  (with respect to the system  $\phi$  and the set  $\Omega$ ). Hence the set  $E'$  of points of egress from  $\omega_1$  is identical with the set  $E$ . Clearly  $E$ , as a consequence of the Jordan-Brouwer Separation Theorem, is not a quasi-isotopic deformation retract of  $Z \cup E$  in  $\omega_1 \cup E$ . Conditions (2.2) are satisfied, with  $E_1 = E$ ; there exists a point  $Q \in \omega_1 \cap Z$  such that  $\Gamma_Q^+ \subset \omega_1 \subset \omega$ . This completes the proof of the theorem.

We recall that a trajectory  $\Gamma_P$  is said to be positively stable in the sense of Lagrange, or simply  $L^+$ -stable if  $\overline{\Gamma_P^+}$  is compact; it is  $L^-$ -stable if  $\overline{\Gamma_P^-}$  is compact. Thus the theorem proved above furnishes a sufficient condition for the existence of an  $L^+$  or  $L^-$ -stable motion in the neighborhood of the critical point  $O$ . It is of interest to contrast this local condition with the global one derived by the author [1], namely that the system  $\phi$  have no improper saddle points<sup>2</sup>.

4. In the event that system (1.1) is of class  $C^2$  we can employ the theorem proved above to derive sufficient conditions, expressed analytically, for the existence of Lagrange stable motions in the neighborhood of the critical point  $O$ . The particular analytical expressions obtained will depend on the choice of the set  $\omega$ . Two examples will be worked out by way of illustration.

(4.1) Let  $\omega$  be the open sphere of radius  $\lambda$  and center  $O$ .

<sup>2</sup> The system  $\phi$  is said to have an improper saddle point if there exists a sequence of points  $\{P_n\} \subset \Omega$  and two sequences  $\{\tau_n\}, \{t_n\}$  of real numbers satisfying the following conditions:

- (i)  $0 < \tau_n < t_n$ , ( $n = 1, 2, \dots$ ).
- (ii)  $\tau_n \longrightarrow +\infty$  as  $n \longrightarrow +\infty$ .
- (iii)  $P_n \longrightarrow P \in \Omega$  as  $n \longrightarrow +\infty$ .
- (iv)  $\Gamma_{P_n}(t_n) \longrightarrow Q \in \Omega$  as  $n \longrightarrow +\infty$ .
- (v)  $\{\Gamma_{P_n}(\tau_n)\}$  has no limit point in  $\Omega$ .

We assume, without loss of generality, that  $0$  is at the origin. Let  $\rho = (x_1^2 + \dots + x_n^2)^{1/2}$ . The condition  $E = E^*$  means that every point of egress from  $\omega$  is a point of strict egress; in other words,  $\rho = \rho(t)$  along any orbit which intersects  $\omega$  cannot have a relative maximum on the surface  $\rho = \lambda$ . This, in turn, is certainly the case if for every

$P \in H(\omega)$  such that:

(a)  $P$  is not a critical point; and

(b)  $\left. \frac{d\rho}{dt} \right|_P = 0$ ,

we would have  $\left. \frac{d^2\rho}{dt^2} \right|_P > 0$ . In fact, this last statement is actually stronger than the statement  $E = E^*$  and could be weakened somewhat if necessary.

In terms of the notation employed for system (1.1) we have:

$$(4.11) \quad \rho^2 = \sum_{i=1}^n x_i^2$$

$$(4.12) \quad \frac{d\rho}{dt} = \dot{\rho} = \frac{1}{\rho} \sum_{i=1}^n x_i f_i$$

$$(4.13) \quad \frac{d^2\rho}{dt^2} = \ddot{\rho} = \frac{1}{\rho} \left( \sum_{i=1}^n f_i^2 + \sum_{i,j=1}^n \frac{\partial f_i}{\partial x_j} f_j x_i - \dot{\rho}^2 \right) .$$

Thus the sufficient condition under consideration obtains the following form:

COROLLARY 1. Let  $\lambda > 0$  be arbitrary. If for every regular point  $P(\bar{x}_1, \dots, \bar{x}_n) = P(\bar{x})$  such that

$$\rho_P^2 = \sum_{i=1}^n \bar{x}_i^2 = \lambda^2$$

and at which

$$\rho \dot{\rho} = \sum_{i=1}^n \bar{x}_i f_i(\bar{x}) = 0 ,$$

there holds the inequality

$$(4.14) \quad \sum_{i=1}^n f_1^2(\bar{x}) + \sum_{i,j=1}^n \bar{x}_i f_j(\bar{x}) \frac{\partial f_1}{\partial x_j} \bigg|_{\bar{x}} > 0,$$

then there exists a semi-orbit wholly contained in the open sphere of radius  $\lambda$  and center  $O$ .

(4.2) Next let  $\omega$  be an open box with sides parallel to the coordinate axes; i.e., the open set defined by the inequalities  $-\lambda_1 < x_1 < \mu_1$ ,  $\lambda_1 > 0$ ,  $\mu_1 > 0$ , ( $i = 1, \dots, n$ ). Following the same line of reasoning as in (4.1) we obtain

COROLLARY 2. Let  $\lambda_1 > 0$ ,  $\mu_1 > 0$ , ( $i = 1, \dots, n$ ), be arbitrary. If for every regular point  $P(\bar{x}_1, \dots, \bar{x}_n) = P(\bar{x})$  lying on the  $i^{\text{th}}$  face of the boundary of  $\omega$  (i.e., such that  $\bar{x}_1 = -\lambda_1$  or  $\bar{x}_1 = \mu_1$ ) at which  $f_1(\bar{x}_1, \dots, \bar{x}_n) = 0$ , there holds the inequality

$$(4.21) \quad \sum_{j=1}^n \frac{\partial f_1}{\partial x_j} f_j > 0,$$

then there exists a semi-orbit wholly contained within the box  $-\lambda_1 < x_1 < \mu_1$ , ( $i = 1, \dots, n$ ).

Other topologically equivalent choices of  $\omega$  will result in still further analytical expressions for the above sufficient condition.

Columbia University

#### BIBLIOGRAPHY

- [1] MENDELSON, P., "On dynamical systems without improper saddle points," to appear.
- [2] FLIŚ, A., "On a topological method of the study of the behavior of the integrals of ordinary differential equations," Bull. Acad. Polon. des Sciences Cl. III, Vol. II, No. 9 (1954), pp. 415-418.
- [3] WĄŻEWSKI, T., "Sur un principe topologique de l'allure asymptotique des intégrales des équations différentielles," Ann. Soc. Polon. de Math., Vol. XX (1947), pp. 279-313.
- [4] WĄŻEWSKI, T., "Sur une méthode topologique de l'examen de l'allure asymptotique des intégrales des équations différentielles," Proceedings of the International Congress of Mathematicians (1954), Vol. 3, pp. 132-139.



## XI. THE LOCAL THEORY OF PIECEWISE CONTINUOUS DIFFERENTIAL EQUATIONS

### I. IDEAL SYSTEMS \*

J. André and P. Seibert

#### INTRODUCTION

The theory of automatic control, which is an important field of modern engineering, gave rise to the investigation of a class of differential equations which, until very recently, have not been studied by mathematicians; these equations involve piecewise continuous functions. Among the previous contributions to the subject we mention a paper of Solncev [10] which gives a stability theory for two-dimensional piecewise continuous systems, the monograph of Flügge-Lotz [5] who investigates various piecewise linear systems and stresses mainly the engineering point of view, two notes by the authors [1a], [1b] concerned with n-dimensional piecewise linear systems, a number of recent papers on the problem of optimal control (e.g., those of Bellman, Glicksberg, Gross [2], Boltyanskii, Gamkrelidze, Pontryagin [3], [6], [9], Krasovskii [8], Bushaw [4]), and a paper of Krasovskii [7] on stability in the large of piecewise continuous systems.

The present paper gives a generalization of the two above-mentioned notes by the authors. The systems under consideration are piecewise of type  $C^2$  and the set of discontinuity is assumed to consist of certain hypersurfaces of class  $C^3$ . The latter are called the "switching spaces" of the system, their points "switching points".<sup>†</sup> The precise hypotheses and the definition of a solution will be given in §1. In §2 we study the

---

<sup>†</sup> It will be noted that in most of the papers on optimal control ([2], [3], [6], [8], [9]) the right hand sides of the differential equations considered depend discontinuously on  $t$  and not, as in the other papers (including the present one), on  $x$ . However, since the problem treated in these papers is to find optimal switching times for a given initial point, the solution of this problem for all initial points is actually equivalent to that of finding certain switching spaces in the sense considered here (provided, the uncontrolled system is autonomous). The systems studied by Krasovskii in [7] (which are generalizations of those occurring in the control problems) have right hand sides which also depend discontinuously on  $t$ .

behavior of the solutions near the switching spaces and give a classification of the switching points. The latter consist of three principal types: transition points (i.e., points at which a solution traverses the switching space), end-points (at which two solutions "end", (i.e., are not continuable beyond the point), and starting points (at which two solutions "start", i.e., are not continuable into the past). At switching points of some of the other classes (which are less frequent) solutions may fuse or fiburcate. In §5 we describe qualitatively the entire sets of solutions passing through given closed subsets of a switching space. For this purpose we introduce "local flows" (in §4) which are defined on compact subsets of the phase space and can be considered as "local dynamical systems" in analogy to the concept of local groups. The system of solutions around the switching spaces can then be conceived as a complex of local flows, connected with each other by certain neighborhood relations.

The second part of the paper will be devoted to systems with switching delay. These formally belong to a class of difference-differential equations.<sup>1</sup> They are distinguished from the systems considered in the present part of the paper (also called "ideal systems") by the property that the discontinuous function changes sign shortly after the trajectory has reached the switching space, rather than at the exact moment of transition. They usually represent a better approximation to physical reality than the ideal systems and, in contrast with the latter, their solutions are continuable indefinitely into the future, i.e., they have no end-points.

#### §1. THE CONCEPT OF PIECEWISE CONTINUOUS DIFFERENTIAL EQUATIONS AND THEIR SOLUTIONS

1. The system (S). Consider  $m$  real single-valued functions  $s_\mu(x)$ , defined in  $E^n$  ( $n$ -dimensional euclidean space) and satisfying the following conditions.

(a) The functions  $s_\mu(x)$  are of class  $C^3$  (i.e., they possess continuous partial derivatives of first, second and third order).

(b) At no point of  $E^n$  does any function  $s_\mu(x)$  vanish simultaneously with its gradient

$$\left( \frac{\partial s_\mu}{\partial x_1}, \dots, \frac{\partial s_\mu}{\partial x_n} \right).$$

<sup>1</sup> Strictly speaking, this is true only for systems with "constant time lag", not for those with "threshold".

Under these conditions the sets

$$(1.1) \quad S_\mu : s_\mu(x) = 0$$

( $\mu = 1, \dots, m$ ) are smooth hypersurfaces of  $E^n$ . We introduce the following notations:

$$\underline{S} = \bigcup_{\mu=1}^m S_\mu, \quad$$

$$s(x) = (s_1(x), \dots, s_m(x)) .$$

We now consider the domains into which the space is decomposed by the hypersurfaces  $S_\mu$ . Denoting by  $\text{sgn}(\alpha_1, \dots, \alpha_m)$  the vector  $(\text{sgn } \alpha_1, \dots, \text{sgn } \alpha_m) = (\alpha_1/|\alpha_1|, \dots, \alpha_m/|\alpha_m|)$  ( $\alpha_\mu$  real,  $\neq 0$ ), we can associate to every point  $x \in E^n - \underline{S}$  a vector  $e(x)$  by setting

$$(1.2) \quad e(x) = \text{sgn } s(x) .$$

Furthermore, we associate to every vector  $e$  with coordinates  $\pm 1$  an (open) domain<sup>1</sup>

$$(1.3) \quad D_e = \{x \in E^n - \underline{S} : e(x) = e\} .$$

Apparently, every such domain is bounded by hypersurfaces  $S_\mu$ .

Finally, we associate to every vector  $e$  (for which  $D_e$  is  $\neq \emptyset$ ) an  $n$ -dimensional vector function  $f(x, e)$  which is defined and of class  $C^2$  throughout  $E^n$ <sup>2</sup> and which does not vanish on the boundary of  $D_e$ .

After these preparations we consider the differential equation

$$(1.4) \quad \dot{x} = f(x) \quad (x = (x_1, \dots, x_n), \quad \dot{\phantom{x}} = d/dt)$$

where the vector function  $f(x)$  is given by

$$(1.5) \quad f(x) = f(x, e) \quad \text{for } x \in D_e$$

<sup>1</sup> which may be empty or disconnected.

<sup>2</sup> As long as we restrict ourselves to ideal systems, it is sufficient that  $f(x, e)$  is defined in an open set containing the closure  $\bar{D}_e$  of  $D_e$ .

and undefined on  $\underline{S}$ . Using (1.2) and (1.5), we can write (1.4) in the form

$$(S) \quad \dot{x} = f(x, \operatorname{sgn} s(x)) \quad .$$

2. The Concept of Solution. Given a point  $p$  of a domain  $D_e$ , there exists exactly one solution of the differential equation

$$(S, e) \quad \dot{x} = f(x, e)$$

which passes through  $p$  at the time  $t = 0$ . We denote it by  $x(t, p)$ . Since the system (S) coincides with  $(S, e)$  for all  $x \in D_e$ , the function  $x(t, p)$  is also a solution of (S) if it is restricted to a  $t$ -interval  $I$  for which<sup>1</sup>

$$x(I, p) \subset D_e$$

holds. We now extend the concept of a solution of (S) by the following definition:

DEFINITION 1.1. We call a (single valued) vector function  $x(t)$ , defined in an interval  $I$ , a solution of the system (S) if it satisfies the following conditions:

- 1° It is continuous throughout  $I$ .
- 2° For every  $t \in I$ , such that  $x(t) \in D_e$  [which is equivalent to  $e(x(t)) = \operatorname{sgn} s(x(t)) = e$ ], the function  $x(t)$  is differentiable and satisfies equation  $(S, e)$ .
- 3° The set of values  $t \in I$  for which  $x(t) \in \underline{S}$  holds, has no cluster point in  $I$ .

A solution according to this definition satisfies equation (S) in the ordinary sense for almost all values  $t \in I$ . In the second part of the paper we will extend the notion of solution in order to include also certain paths contained entirely in  $\underline{S}$  which are related to a phenomenon in the theory of discontinuous control (after end-point motions).

## §2. SWITCHING POINTS AND THEIR CLASSIFICATION

3. Topological classification of switching points. The hypersurfaces  $S_\mu$  are called the switching spaces of the system (S). Points in a switching space, or switching points, will usually be denoted by  $u$ .

<sup>1</sup>  $x(I, p)$  denotes, as usual, the set  $\{x(t, p)\}_{t \in I}$ .

Unless the contrary is stated, it will always be assumed that  $u$  belongs to exactly one switching space  $S_\mu$ , and we will therefore drop the index  $\mu$ . Under this assumption  $u$  belongs to the boundaries of exactly two domains  $D_e$  which we denote by  $D_e^+$  and  $D_e^-$ , respectively. The corresponding vectors  $e^+$  and  $e^-$  differ only in their  $\mu$ -th coordinate which we assume to be  $+1$  in  $e^+$  and  $-1$  in  $e^-$ . The solutions of the equations

$$\dot{x} = f(x, e^\pm)$$

with the initial value  $u$  will be denoted by  $x^\pm(t, u)$ .

We first classify the switching points  $u$  with respect to each of the cruves  $x^+(t, u)$  and  $x^-(t, u)$  by introducing the following definitions: If the limits

$$(3.1) \quad \gamma_1^\pm = \lim_{t \rightarrow -0} e_\mu(x^\pm(t, u)), \quad \gamma_2^\pm = \lim_{t \rightarrow +0} e_\mu(x^\pm(t, u))$$

both exist, the point  $u$  will be called

an A-point (with respect to  $x^\pm(t, u)$ ) if  $\gamma_1^\pm = \mp 1$ ,  $\gamma_2^\pm = \pm 1$ ,

an E-point if  $\gamma_1^\pm = \pm 1$ ,  $\gamma_2^\pm = \mp 1$ ,

an L-point if  $\gamma_1^\pm = \gamma_2^\pm = \mp 1$ ,

an R-point if  $\gamma_1^\pm = \gamma_2^\pm = \pm 1$ .

In the case of an A-point there exists a  $t$ -interval  $B = (0, \beta)$  for which  $e_\mu(x^\pm(t, u)) = +1$ , and consequently  $x^\pm(B, u) \subset D_{e^+}$ . Since  $x^\pm(t, u)$  satisfies  $(S, e^\pm)$  and this equation coincides with  $(S)$  for  $x \in D_{e^+}$ , the function  $x^\pm(t, u)$ , restricted to the interval  $B$ , is a solution of  $(S)$ . Analogously we find that for some interval  $(-\alpha, 0)$  the same function does not satisfy  $(S)$ .

In the case of an E-point the situation is vice versa, i.e.,  $x^\pm(t, u)$  satisfies  $(S)$  for small negative values of  $t$ , but not for small positive values.

If  $u$  is an R-point,  $x^\pm(t, u)$  is a solution of  $(S)$  in a certain open neighborhood of  $t = 0$ , while in the case of an L-point there exists an interval around 0 in which  $x^\pm(t, u)$  does not satisfy  $(S)$ .

Consider now the case where some of the limits (3.1) fail to exist. This situation occurs whenever the intersections of one or both of the curves  $x^\pm(t, u)$  with  $S$  cluster at  $u$ . We call  $u$

an  $A^*$ -point if  $\gamma_2^+$  exists and is equal to  $\pm 1$ ,

an  $E^*$ -point if  $\gamma_1^+$  exists and is equal to  $\pm 1$ ,

an  $L^*$ -point in all other cases.

It is easy to see that  $A^*$ -,  $E^*$ -,  $L^*$ -points have the above stated properties of  $A$ -,  $E$ -,  $L$ -points, respectively, except that  $x^\pm(t, u)$  usually satisfies equation (S) in certain intervals clustering at  $t = 0$  and contained (in the case of  $A^*$  -  $[E^*]$ -points) in the semi-neighborhoods which are void of solutions in the case of  $A$ -  $[E]$ -points.

We now extend the classes  $A, E, L$  by the sets  $A^*, E^*, L^*$ , respectively. The set of all  $X$ -points  $u$  ( $X = A, E, L, R$ ) with respect to  $x^\pm(t, u)$  will be denoted by  $X^+$ . Then the complete topological classification of the switching points is obtained by considering the intersections of the sets  $X^+$  and  $Y^-(X, Y = A, E, L, R)$ . Since the points of the sets  $X^+ \cap Y^-$  and  $X^- \cap Y^+$  are of the same topological character with respect to the solutions passing through them, the sets of topologically equivalent switching points are given by the expressions

$$(3.2) \quad XY = (X^+ \cap Y^-) \cup (X^- \cap Y^+)$$

( $X, Y = A, E, L, R$ ). The symmetry relation  $XY = YX$  reduces the number of these classes to 10.

4. Behavior of solutions at switching points. If  $u$  is of type  $AE$ , there exists a unique solution with  $u$  as its initial value which is defined in an open interval around  $t = 0$ . This solution, which we denote by  $x(t, u)$ , is given by

$$x(t, u) = \begin{cases} x^+(t, u) & \text{for } t \geq 0 \\ x^-(t, u) & \text{for } t \leq 0 \end{cases}$$

in the case  $a \in A^+ \cap E^-$ , and vice versa if  $u \in A^- \cap E^+$ . At the point  $u$  the solution crosses the switching space;  $u$  is therefore called a transition point. In the topological behavior of the solution  $x(t, u)$  there is no difference between a transition point and a point at which  $f(x)$  is continuous. (Vid. Figure 1, also for the following cases.)<sup>1</sup>

If  $u$  is an  $AA$  -  $[EE]$  point, the positive [negative] half

<sup>1</sup> Dotted lines represent half trajectories which are not solution of (S).

trajectories<sup>1</sup> of  $x^{\pm}(t, u)$  are both solutions of (S), so that the uniqueness theorem fails to hold in this case. Apparently, these two half trajectories cannot be continued beyond  $u$  into the past [future]. An  $AL - [EL-]$  point is the initial point of exactly one solution of (S) which is defined only for  $t \geq 0$  [ $t \leq 0$ ]. We call the  $AA-$  and  $AL-$  points starting points, the  $EE-$  and  $EL-$  points end-points.

If  $u$  is an  $AR - [ER-]$  point, both  $x^+(t, u)$  and  $x^-(t, u)$  are solutions of (S), one being defined only for  $t \leq 0$  [ $t \geq 0$ ], the other for positive and negative  $t$ -values. This implies the existence of two solutions actually passing through  $u$ . Assume, e.g.,  $u \in A^+ \cap R^-$  [ $u \in E^+ \cap R^-$ ]. (In the case  $u \in A^- \cap R^+$  [ $u \in E^- \cap R^+$ ] the situation is analogous.) Then

$$x_1(t, u) = x^-(t, u)$$

and

$$x_2(t, u) = \begin{cases} x^-(t, u) & \text{for } t \leq 0 \text{ } [t \geq 0] , \\ x^+(t, u) & \text{for } t \geq 0 \text{ } [t \leq 0] , \end{cases}$$

are both solutions of (S) passing through  $u$ . Since for  $t \leq 0$  [ $t \geq 0$ ] these two solutions are identical, the situation can be described as a bifurcation [fusion] of trajectories.<sup>2</sup>

Through an  $IL-$  point there apparently exists no solution of (S). - If  $u$  is an  $IR-$  point, there is exactly one solution [either  $x^+(t, u)$  or  $x^-(t, u)$ ] with  $u$  as its initial value. It is defined for positive and negative values of  $t$  and does not cross the switching space. Apart from the transition ( $AE-$ ) points, the  $IR-$  points are the only switching points at which the existence and uniqueness theorem holds in the strict sense. - If, finally,  $u$  belongs to the set  $RR$ , both  $x^+(t, u)$  and  $x^-(t, u)$  are solutions of (S) around  $t = 0$ . This implies the existence of altogether four solutions passing through  $u$ , the latter therefore being a point of fusion and of bifurcation simultaneously.

5. Normal and exceptional switching points. Beside the topological qualities of a switching point which we analysed in the preceding setion, it is of significance whether the curves  $x^{\pm}(t, u)$  associated to the point  $u$  are tangent to  $S$  or not. We introduce the following notions:

<sup>1</sup> i.e., the restrictions of  $x^{\pm}(t, u)$  to  $t \geq 0$  [ $t \leq 0$ ].

<sup>2</sup> The concept of solutions introduced in the paper André-Seibert [1a] was slightly less general than the one we use here. According to the former, there exist fusions of trajectories, but no bifurcations.




Fig. 1.



A point  $u \in S$  will be called a normal switching point if neither of the curves  $x^\pm(t, u)$  are tangent to the switching space  $S$  at  $u$ , otherwise an exceptional point. In particular, all points of the types  $A^*$ ,  $E^*$ ,  $L^*$  are apparently exceptional. The same is true for all points of the types  $LX$ ,  $RX$  ( $X = A, E, L, R$ ), since for all of these at least one of the curves  $x^\pm(t, u)$  fails to traverse  $S$  at the point  $u$ . Denoting the set of all normal points by  $S^0$ , we therefore have the inclusion

$$(5.1) \quad S^0 \subset AA \cup AE \cup EE.$$

The sets of normal starting, end-, and transition points will be denoted by  $A$ ,  $E$ ,  $T$ , respectively:

$$(5.2) \quad A = AA \cap S^0, \quad E = EE \cap S^0, \quad T = AE \cap S^0.$$

We call an exceptional point  $u$  of first order, if only one of the curves  $x^\pm(t, u)$  is tangent to  $S$  at  $u$  and the contact is of 1<sup>st</sup> order; in all other cases we say  $u$  is of higher order. By  $S'$  we denote the set of all exceptional points, by  $S''$  that of all exceptional points of higher order, and by  $S^1 = S' - S''$  the set of exceptional points of first order. The intersections of any subset  $Q$  of  $S$  with  $S^0$ ,  $S'$ ,  $S''$ ,  $S^1$  will be denoted by  $Q^0$ ,  $Q'$ ,  $Q''$ ,  $Q^1$ .<sup>1</sup> Since at points of types  $LL$ ,  $LR$ ,  $RR$  both curves  $x^\pm(t, u)$  are tangent to  $S$ , we have the relation

$$(5.3) \quad LL \cup LR \cup RR \subset S''.$$

6. Determination of the type of a switching point. In order to state the criteria for the type of a given switching point  $u \in S = S_\mu$  it is sufficient to assume  $s_\mu \in C^2$ ,  $f \in C^1$  instead of the stronger conditions formulated in §1.1. Then the functions  $x^\pm(t, u)$  which satisfy the equations  $\dot{x} = f(x, e^\pm)$  are of type  $C^2$  and we can apply the following Taylor expansion:

$$(6.1) \quad \begin{aligned} s_\mu(x^\pm(t, u)) &= \frac{d}{d\tau} s_\mu(x^\pm(\tau, u)) \Big|_{\tau=0} t + 1/2 \frac{d^2}{d\tau^2} s_\mu(x^\pm(\tau, u)) \Big|_{\tau=0} t^2 = \\ &= c_1^\pm(u)t + 1/2 c_2^\pm(st, u)t^2 \quad (0 < \theta < 1). \end{aligned}$$

<sup>1</sup> In the case of the sets  $A$ ,  $E$ ,  $T$  [vid. (5.2)] we omit the superscript 0. -- It should be noted that the sets  $S^0$ ,  $S'$ ,  $S''$ ,  $S^1$  comprise only that part of  $S$  which is not contained in any other switching space.

Here

$$c_1^+(t, u) = \frac{d}{dt} s_\mu(x^+(t, u))$$

and

$$c_1^+(u) = c_1^+(0, u) \quad (i = 1, 2) \quad .$$

For the present purpose it is sufficient to examine the signs of the coefficients  $c_1^+(u)$  and  $c_2^+(u)$ . An obvious calculation yields the following formulas:<sup>1</sup>

$$(6.2) \quad c_1^+(u) = f(u, e^+) \frac{d}{dx} s_\mu(u) \quad ,$$

$$(6.3) \quad c_2^+(u) = f(u, e^+) \left( \frac{d^2}{dx^2} s_\mu(u) \right) f(u, e^+)' \\ + f(u, e^+) \left( \frac{d}{dx} f(u, e^+) \right)' \frac{d}{dx} s_\mu(u) \quad ;$$

Here  $\frac{d}{dx}$  denotes the gradient when applied to a scalar and the Jacobian matrix when applied to a vector:

$$\frac{ds_\mu}{dx} = \begin{pmatrix} \frac{\partial s_\mu}{\partial x_1} \\ \vdots \\ \frac{\partial s_\mu}{\partial x_n} \end{pmatrix}, \quad \frac{df}{dx} = \frac{d}{dx} (f_1, \dots, f_n) = \begin{pmatrix} \frac{\partial f_1}{\partial x_1}, \dots, \frac{\partial f_n}{\partial x_1} \\ \dots \\ \frac{\partial f_1}{\partial x_n}, \dots, \frac{\partial f_n}{\partial x_n} \end{pmatrix}$$

The operator  $\frac{d^2}{dx^2}$ , applied to a scalar, denotes the Jacobian matrix of the gradient. From the formula (6.1) we immediately obtain the following theorem:

**THEOREM 6.1.** If  $u$  is a point of the switching space  $S$  (and of no other switching space), the following implications hold:<sup>2</sup>

<sup>1</sup> By ' we denote the operation of transposition.

<sup>2</sup> The upper [lower] signs +, - in the superscripts corresponds to the upper [lower] signs <, >.

$$c_1^+(u) > 0 \quad \text{implies} \quad u \in A^+,$$

$$c_1^-(u) < 0 \quad \text{implies} \quad u \in E^+$$

$$c_1^+(u) = 0 \quad \text{and} \quad c_2^+(u) > 0 \quad \text{together imply} \quad u \in R^+,$$

$$c_1^-(u) = 0 \quad \text{and} \quad c_2^-(u) < 0 \quad \text{together imply} \quad u \in L^+.$$

The coefficients  $c_1^+(u)$ ,  $c_2^+(u)$  are given by the formulas (6.2) and (6.3).

In particular,  $u$  is

a normal transition point if  $c_1^+(u)$  and  $c_1^-(u)$  are either both positive or both negative,

a normal starting point if  $c_1^+(u) > 0$  and  $c_1^-(u) < 0$ ,

a normal endpoint if  $c_1^+(u) < 0$  and  $c_1^-(u) > 0$ .

Unconsidered in this theorem remain only those exceptional points of higher order which involve contacts of higher than first order [i.e., at least 3-point contact].

The three last implications require only the hypotheses  $f \in C^0$ ,  $s \in C^1$ . If these are satisfied, the functions  $c_1^\pm$  are continuous and we obtain the following corollary:<sup>1</sup>

COROLLARY 6.1. The sets  $A$ ,  $E$ ,  $T$  of normal switching points are open relative to  $S$ .

Since every point of  $S^1$  involves a contact of first order, we conclude from the third and fourth implication in Theorem 6.1 and from the continuity of  $c_2^\pm(u)$  (assuming again  $f \in C^1$ ,  $s \in C^2$ ):

COROLLARY 6.2.<sup>2</sup> The set  $S^1$  of exceptional points of first order consists of the sets  $(AL)^1$ ,  $(EL)^1$ ,  $(AR)^1$ ,  $(ER)^1$ , all of which are open relative to  $S^1$ .

<sup>1</sup> The boundaries of the sets  $A$ ,  $E$ ,  $T$  may contain parts of intersections of  $S$  with other switching spaces. The latter, however, being closed sets, do not affect the validity of Corollary 6.1.

<sup>2</sup> All sets with superscript 1 consist of exceptional points of first order.

### '§3. TOPOLOGICAL PROPERTIES OF THE CLASSES OF SWITCHING POINTS

7. The exceptional classes. Let  $u$  be an exceptional point of  $S$  ( $= S_\mu$ ) [vid. §2.5]. Then the conditions

$$(7.1) \quad \begin{cases} s_\mu(u) = 0 \\ c_1^\pm(u) = f(u, e^\pm) \frac{d}{dx} s_\mu(u) = 0 \end{cases}$$

hold, the second either for  $e^+$  or  $e^-$  [vid. (6.1), (6.2)]. The functions  $s_\mu(u)$  and  $c_1^\pm(u)$  are of class  $C^1$ . Therefore, if the Jacobian matrix

$$J_1^\pm(u) = \frac{d}{dx} (s_\mu(u), c_1^\pm(u))$$

$$\left[ = \frac{d}{dx} (s_\mu(x), c_1^\pm(x)) \right]_{x=u}$$

of the system (7.1) is of rank 2, it follows from the implicit function theorem that the set of all exceptional points in a certain neighborhood of  $u$  constitute an  $(n-2)$ -dimensional differentiable submanifold of  $S$  [or, if  $c_1^\pm(u)$  both vanish, the union of two such manifolds]. We introduce the following terminology:

If the matrices  $J_1^\pm(u)$  are of rank 2 at all points of  $S$  [or of a part  $Q$  of  $S$ ] satisfying the corresponding conditions (7.1), we say, the system (S) satisfies condition A on the switching space  $S$  [or on the subset  $Q$ ].

We can then state the following theorem:

**THEOREM 7.1.** If condition A is satisfied on a subset  $Q^1$  of the switching space  $S$ , the set  $Q^1$  of exceptional points in  $Q$  consists of  $(n-2)$ -dimensional differentiable submanifolds of  $S$  if it is not empty. Every point  $u \in Q^1$  possesses a neighborhood  $N$  such that  $Q^1 \cap N$  is contained in the union of at most two of the connected manifolds constituting  $Q^1$  (namely, one (connected) component of each of

---

<sup>1</sup> It is assumed that  $Q$  contains no points of switching spaces other than  $S$ .

the sets  $\{u \in Q' : c_1^+(u) = 0\}$  and  $\{u \in Q' : c_1^-(u) = 0\}$ .  
 If, in particular,  $U$  is an exceptional point of  
 first order (vid. §2.5), the set  $Q' \cap N$  is contained  
 in a single (connected) manifold.

We now establish a similar theorem for the set  $S''$  of exceptional points of higher order [vid. §2.5]. Every point  $u \in S''$  satisfies one of the two following sets of equations:

$$(7.2) \quad s_\mu(u) = c_1^+(u) = c_1^-(u) = 0, \quad .$$

$$(7.3) \quad s_\mu(u) = c_1^+(u) = c_2^+(u) = 0$$

(the latter either for  $c_1^+$  or  $c_1^-$ ). Denote by  $J_2(u)$ ,  $J_3^+(u)$  the Jacobian matrices of these systems:

$$J_2(u) = \frac{d}{dx} (s_\mu(u), c_1^+(u), c_1^-(u)) ,$$

$$J_3^+(u) = \frac{d}{dx} (s_\mu(u), c_1^+(u), c_2^+(u)) .$$

We say, condition B is satisfied on  $S$  [or  $Q \subset S$ ], if the matrices  $J_2(u)$ ,  $J_3^+(u)$  are of rank 3 at every point of  $S$  [or  $Q$ ] satisfying the corresponding set of conditions (7.2), (7.3).

Then, in analogy to Theorem 7.1, we obtain

**THEOREM 7.2.** If condition B is satisfied on a subset<sup>1</sup>  $Q$  of the switching space  $S$ , the set  $Q''$  of exceptional points of higher order in  $Q$  consists of  $(n-3)$ -dimensional differentiable submanifolds of  $Q'$  if it is not empty. Every point  $u \in Q''$  possesses a neighborhood  $N$  such that  $Q'' \cap N$  is contained in the union of at most three of the connected manifolds constituting  $Q''$ .

**REMARK 7.1.** By comparing the number of (scalar) equations entering into the conditions A and B with the number of independent variables, it is easy to see that, in general, both conditions hold in the entire switching space.

<sup>1</sup> Vid. Footnote on p. 236

REMARK 7.2. Since  $S''$  is of lower dimension than  $S'$  (provided that condition B is satisfied on  $S'$  and the latter is not empty),  $S' [= S' - S'']$  is apparently dense in  $S'$ :

$$(7.4) \quad \overline{S'} = S'.$$

8. The classes of normal switching points. For the following considerations we assume conditions A and B to be satisfied throughout the switching space  $S$ . Then the following lemma holds:

LEMMA 8.1. If  $c_1^+(u) = 0$  and  $c_2^+(u) \neq 0$  hold, the function  $c_1^+$  assumes positive and negative values in every neighborhood of  $u$ .

PROOF. Under the assumptions  $c_1^+(u) = 0$ ,  $c_2^+(u) > 0$  the curve  $x^+(t, u)$  is contained in the set  $\bar{D}_{0+}$  for  $|t| < \delta$  and  $\delta$  sufficiently small [vid. (6.1)]. Now assume that  $c_1^+$  is non-negative in a neighborhood  $N$  of  $u$  relative to  $S$ :

$$(8.1) \quad c_1^+(v) \geq 0 \quad \text{for } v \in N,$$

and take  $N$  so small that

$$(8.2) \quad c_2^+(v) > 0 \quad \text{for } v \in N.$$

Then, due to the continuous dependence of  $x^+$  on the initial values, there exists a neighborhood  $N_1 \subset N$  of  $u$  such that, if  $I$  denotes the interval  $(0, \delta)$  and  $\delta$  is chosen small enough, the set  $x^+(I, N_1)$  contains no point of  $S$  outside  $N$ . Then (8.1) and (8.2) imply

$$x^+(I, N_1) \subset \bar{D}_{0+}.$$

Since  $x^+(I, N_1)$  is a homeomorphic image of the product set  $N_1 \times I$  and  $N_1$  is an open set on the hypersurface  $S$ , the set  $x^+(I, N_1)$  apparently contains a semi-neighborhood  $\mathcal{N} \cap D_{0+}$  of  $u$  ( $\mathcal{N}$ : neighborhood of  $u$ ).

On the other hand, it follows from the hypotheses (and (6.1)) that  $x^+(t, u) \subset D_{0+}$  tends to  $u$  for  $t \rightarrow -0$ , which leads to a contradiction.

If  $c_1^+ < 0$  is assumed in  $N$  instead of  $c_1^+ > 0$ , the preceding argument needs only to be modified by considering the interval  $(-\delta, 0)$  instead of  $(0, \delta)$ . - For the coefficients  $c_1^-, c_2^-$  the proof is exactly the same.

We now consider the boundaries relative to  $S$  of the sets  $A$ ,  $E$ ,  $T$  and denote them by  $\partial A$ ,  $\partial E$ ,  $\partial T$ .

From Theorem 7.1 and Lemma 8.1 we easily obtain the following pair of inclusions:

$$(8.3) \quad (AL)^1 \cup (AR)^1 \subseteq \partial A \cap \partial T,$$

$$(8.4) \quad (EL)^1 \cup (ER)^1 \subseteq \partial E \cap \partial T.$$

Consider, e.g., the case  $u \in (A^+ \cap L^-)^1$ . [The other cases are treated in obvious analogy.] Here  $c_1^+ > 0$ ,  $c_1^- = 0$ ,  $c_2^- > 0$  [Theorem 6.1]. According to the lemma there exist points  $v, w \in S$  arbitrarily near  $u$  with  $c_1^+(v) > 0$ ,  $c_1^-(v) > 0$ ;  $c_1^+(w) > 0$ ,  $c_1^-(w) < 0$ . This implies  $v \in T$ ,  $w \in A$  [Theorem 6.1].

After these preparations we can prove the following theorem:

**THEOREM 8.1.** If the conditions  $A$  and  $B$  are satisfied on the switching space  $S$ , one of the following statements is true, provided that  $S$  intersects no other switching space:

- (a)  $S$  consists entirely of normal switching points of one type, i.e.,  $S = A$ ,  $E$ , or  $T$ .
- (b) The sets  $T$  and  $A \cup E$  are both non-empty and their boundaries  $\partial T$  and  $\partial(A \cup E)$  coincide. Thus the closure  $\bar{T}$  of  $T$  separates  $A$  from  $E$  and the components of  $A$  and  $E$  from each other and, conversely, the components of  $T$  are separated from each other by  $\overline{A \cup E}$ .

If  $S$  intersects other switching spaces  $S_\lambda$ , the preceding statements hold for every component of  $S - \bigcup_\lambda (S_\lambda \cap S)$ .

**PROOF.** We first show that the set of exceptional points  $S^1$  is empty if all normal points are of the same type. (The converse is obvious because  $\partial A$ ,  $\partial E$ ,  $\partial T \subseteq S^1$ .) Corollary 6.2 yields

$$(8.5) \quad S^1 \subseteq (AL)^1 \cup (AR)^1 \cup (EL)^1 \cup (ER)^1.$$

From (8.3), (8.4) it follows that all sets on the right-hand side of (8.5) are empty, hence  $S^1 = \emptyset$ . Finally, (7.4) implies  $S^1 = \emptyset$ , so that we have

the case (a).

Suppose now that  $S'$  and therefore also  $S^1$  are non-empty. Then it follows immediately from (8.3), (8.4), and (8.5) that both  $T$  and  $A \cup E$  are non-empty. The same relations imply

$$(8.6) \quad (\partial A)^1 \subset \partial T, (\partial E)^1 \subset \partial T, (\partial T)^1 \subset \partial(A \cup E).$$

Observing that  $\partial A$ ,  $\partial E$ ,  $\partial T$  are closed subsets of  $S$  and using (7.4) and (8.6), we obtain

$$\partial A = \partial A \cap S' \subset \partial A \cap \overline{S^1} = \overline{\partial A \cap S^1} = \overline{(\partial A)^1} \subset \overline{\partial T} = \partial T$$

and the analogous relations

$$\partial E \subset \partial T \quad \text{and} \quad \partial T \subset \partial(A \cup E) .$$

Hence,

$$(8.7) \quad \partial(A \cup E) = \partial T ,$$

which yields the separation property of case (b).

The extension to the case where  $S$  intersects other switching spaces is immediate.

#### §4. LOCAL FLOWS

9. Definitions and elementary properties. As we have seen in §2.4, the qualitative behavior of the solution of a system (S) differs in many respects from that of a dynamical system in the usual sense, defined, e.g., by the solutions of differential equations satisfying a Lipschitz condition. In order to study families of solutions of (S) qualitatively, we introduce the notion of "local flows".

DEFINITION 9.1. Let  $Y$  be a compact connected topological space. Denote by  $t_0$ ,  $t_1$  two continuous maps of  $Y$  into the negative and positive halves of the real line (both including 0) respectively, and by  $I(y)$  the interval  $[t_0(y), t_1(y)]$ . Then a continuous mapping  $\varphi$  of the set  $\Omega = \{(y, t) : y \in Y, t \in I(y)\}$  onto  $Y$  defines a local flow  $\mathcal{F} = (Y, I, \varphi)$  if  $\varphi$  has the following properties:

$$(a) \quad y \in Y, t \in I(y) \text{ implies } I(\varphi(t, y)) = I(y) - t = [t_0(y) - t, t_1(y) - t].$$

$$(b) \quad \varphi(0, y) = y \text{ holds for all } y \in Y.$$



- (c) If  $y \in Y$ ,  $t \in I(y)$ ,  $t + t' \in I(y)$ , the relation  $\varphi(t + t', y) = \varphi(t', \varphi(t, y))$  holds.
- (d)  $\varphi(t, y)$  is a topological mapping of the set  $\{y \in Y : t \in I(y)\}$  for every fixed  $t$ .

In the case where  $Y$  is a  $k$ -dimensional manifold with boundary<sup>1</sup> we call  $F$  a local  $k$ -flow.

The sets  $\varphi(I(y), y)$  [ $y$  fixed] are called the paths of the local flow; we denote them by  $\gamma(y)$ . The set  $\bigcup_{p \in P} \gamma(p)$  ( $P \subset Y$ ) will be denoted by  $\gamma(P)$ . It follows from (a) and (c) that

$$q \in \gamma(p) \text{ implies } \gamma(q) = \gamma(p) .$$

To every point  $y \in Y$  we associate two points

$$y_0 = \varphi(t_0(y), y), \quad y_1 = \varphi(t_1(y), y)$$

and define a pair of sets  $Y_0, Y_1$  by

$$Y_0 = \{y_0(y)\}_{y \in Y}, \quad Y_1 = \{y_1(y)\}_{y \in Y} .$$

We call  $Y_0$  and  $Y_1$  the entrance and exit set of  $\mathcal{F}$ . Apparently, the paths of  $\mathcal{F}$  are in one-one correspondence with the points of each of the sets  $Y_0, Y_1$ ;  $y_0(y)$  and  $y_1(y)$  are the end-points of the path through  $y$ .

PROPOSITION 9.1. The entrance and exit set  $Y_0, Y_1$  of a local flow are connected sets.

PROOF. Since the mappings  $t_0, t_1, \varphi$  are continuous, so are  $y_0$  and  $y_1$ . Therefore the sets  $Y_0, Y_1$ , which are images under  $y_0$  and  $y_1$  of the connected set  $Y$ , are connected.

PROPOSITION 9.2. The mapping  $\varphi(t, y)$  is univalent (as a function of  $t$  for every point  $y \in Y_0$ ).<sup>2</sup>

<sup>1</sup> I.e., a Hausdorff space with the property that each of its points possesses a neighborhood which is homeomorphic either to an open  $k$ -sphere or to a semi- $k$ -sphere  $x_1^2 + \dots + x_k^2 < 1, x_1 \geq 0$ .

<sup>2</sup> It should be noted that Definition 9.1 excludes the case where  $I(y)$  is the whole real line. Therefore, in particular, critical points  $[\varphi(t, y) = y]$  are excluded due to condition (a).

PROOF. Suppose that  $\varphi(t^1, y) = \varphi(t^2, y)$ . Then obviously  $\varphi(-t^1, \varphi(t^1, y)) = \varphi(-t^1, \varphi(t^2, y))$  holds. By applying (b) and (c) we obtain

$$\varphi(0, y) = y = \varphi(t^2 - t^1, y) .$$

Finally (a) yields  $I(y) = I(y) - (t^2 - t^1)$ , i.e.,  $t^1 = t^2$ .

PROPOSITION 9.3. Consider a dynamical system<sup>1</sup> [or flow] defined on a finite-dimensional manifold  $X$  by the mapping function  $\Phi$ . Let  $C$  be a compact connected subset of a hypersurface without contact with the paths of the flow.<sup>2</sup> Then there exists a number  $T > 0$  such that  $\Phi$  defines a local flow on the set  $Y = \Phi([0, T], C)$ .

PROOF. Choosing  $T$  so small that  $Y$  is contained in a neighborhood  $H$  of the kind mentioned in footnote 2, it is clear that every point  $y \in Y$  possesses a unique representation  $y = \Phi(t, y_0)$  ( $y_0 \in C$ ,  $t \in [0, T]$ ). Then, if we define  $I(y) = [-t, T - t]$ , the conditions of Definition 9.1 are obviously satisfied.

So far only interior properties of local flows have been considered, i.e., properties which do not depend on an imbedding of  $Y$  into a larger space. Now we are going to study properties which depend substantially on such an imbedding. We therefore assume  $Y$  to be a subspace of a topological space  $X$ . Closure, interior, and boundary of  $Y$  will always be understood relative to  $X$ .

DEFINITION 9.2. A local flow  $\mathcal{F}' = (Y', I', \varphi')$  is called a subflow of the local flow  $\mathcal{F} = (Y, I, \varphi)$  if  $Y' \subseteq Y \subseteq X$ ,  $I'(y) = \{t \in I(y) : \varphi(t, y) \in Y'\}$  for all  $y \in Y'$  and  $\varphi'$  is the restriction of  $\varphi$  to the set  $n' = \{t, y : y \in Y', t \in I'(y)\}$ .

If  $\mathcal{F}'$  is a subflow of  $\mathcal{F}$  we write

$$\mathcal{F}' \subseteq \mathcal{F}.$$

PROPOSITION 9.4. Consider a local  $k$ -flow  $\mathcal{F} = (Y, I, \varphi)$

<sup>1</sup> I.e., a continuous mapping of the product space  $X \times I$  ( $I$ : real line) onto  $X$  which satisfies the conditions (b), (c), (d) of Definition 9.1 in which  $Y$  and  $I(Y)$  are replaced by  $X$  and  $I$ .

<sup>2</sup> In this context we mean by a "hypersurface without contact" a hypersurface  $H$  of  $X$  with a neighborhood  $N$  such that  $H$  does not contain more than one point of any component of the intersection of any path with  $N$ .

and denote by  $\partial Y$  the boundary of  $Y$ . If a path  $\gamma$  of  $\mathcal{F}$  contains a point  $p \in \partial Y$  not lying in  $Y_0$  or  $Y_1$ , the entire path  $\gamma$  is contained in  $\partial Y$ .

PROOF. To every point  $p \in Y$  we associate the set<sup>1</sup>  $Q_p = \varphi(-t_0(p), Y_0)$  [which is a  $(k-1)$ -dimensional manifold with boundary]. Since  $t_0$  and  $t_1$  are continuous, given two numbers  $\alpha \in (t_0(p), 0)$ ,  $\beta \in (0, t_1(p))$ , there exists a neighborhood  $Q_p^I$  of  $p$  relative to  $Q_p$  such that  $\varphi(t, p')$  is defined for all  $t \in I = [\alpha, \beta]$  and all points  $p' \in Q_p^I$ . If  $p$  is an interior point of  $Q_p^I$  (i.e., possesses a neighborhood relative to  $Q_p^I$  which is homeomorphic to a  $(k-1)$ -sphere), it is also an interior point of the set  $\varphi(I, Q_p^I)$  (since the latter is a homeomorphic image of the product set  $I \times Q_p^I$ ) and therefore of  $Y$ . The converse is also true. Indeed, consider a sequence of points  $p^n \in Y$  tending to  $p$ . Then (again using the continuity of  $t_0$ )  $t_0(p^n) - t_0(p) \in I$  for sufficiently large  $n$ , while, due to the continuity of  $\varphi$ , the points  $\varphi(t_0(p^n) - t_0(p), p^n)$  ultimately belong to  $Q_p^I$ . Thus  $p^n \in \varphi(I, Q_p^I)$ . Therefore, the intersections of  $Y$  and  $\varphi(I, Q_p^I)$  with a sufficiently small neighborhood of  $p$  coincide. This implies the equivalence of  $p \in \partial Y$  and  $p \in \partial Q_p$ .

Now consider two points  $p \in \partial Y$  and  $q = \varphi(t_q, p)$  ( $t_q \in I(p)$ ). Then  $p \in \partial Q_p$  and, choosing  $Q_p^I$  as before (with  $t_q \in I$ ),  $\varphi(t_q, \gamma)$  maps  $Q_p^I$  homeomorphically onto  $Q_q^I = \varphi(t_q, Q_p^I)$ , so that  $q \in \partial Q_q^I$ . This implies  $q \in \partial Y$  which proves the proposition.

DEFINITION 9.3. A subflow  $\mathcal{F}' = (Y', I', \varphi')$  of a local  $k$ -flow  $\mathcal{F} = (Y, I, \varphi)$  is called a face of  $\mathcal{F}$  if  $Y'$  is contained in the boundary of  $Y$ . We also say,  $\mathcal{F}'$  is incident with  $\mathcal{F}$  and use the notations<sup>2</sup>

$$\mathcal{F}' \subseteq \partial \mathcal{F}$$

and

$$\mathcal{F}' \subseteq \partial \mathcal{F}_1 \cap \partial \mathcal{F}_2 \text{ if } \mathcal{F}' \subseteq \partial \mathcal{F}_1 \text{ and } \mathcal{F}' \subseteq \partial \mathcal{F}_2.$$

Apparently, a face of a local  $k$ -flow is a local  $(k-1)$ -flow.

PROPOSITION 9.5. Consider a local  $k$ -flow  $\mathcal{F} = (Y, I, \varphi)$

<sup>1</sup> If  $\varphi(-t_0(p), \gamma)$  is defined only on a subset  $Y_0'$  of  $Y_0$ , we mean by  $\varphi(-t_0(p), Y_0)$  the set  $\varphi(-t_0(p), Y_0')$ .

<sup>2</sup> The symbol  $\partial \mathcal{F}$  denotes the collection of all faces of  $\mathcal{F}$ .

and a local  $(k-1)$ -flow  $\mathcal{F}' = (Y', I', \varphi')$  which satisfy the conditions:

- (1)  $I'(y) = \{t \in I(y) : \varphi(t, y) \in Y'\}$   
for all  $y \in Y' \cap Y$ .
- (2) The restrictions of  $\varphi$  and  $\varphi'$  to the set  $\{t, y : y \in Y \cap Y', t \in I'(y)\}$  are identical.
- (3) The entrance [exit] set  $Y'_0[Y'_1]$  of  $\mathcal{F}'$  is contained in the boundary of the entrance [exit] set  $Y_0[Y_1]$  of  $\mathcal{F}$ .

Then

$$\mathcal{F}' \subset \partial \mathcal{F}.$$

PROOF. Assume condition (3) to be satisfied, e.g., for  $Y_0$  and  $Y'_0$ . Then it follows by an argument analogous so that in the proof of Proposition 9.4 that

$$\gamma(Y'_0) = Y' \subseteq \partial Y \subset Y.$$

Conditions (1) and (2), therefore, imply that  $\mathcal{F}'$  is a subflow of  $\mathcal{F}$  and the proposition follows.

DEFINITION 9.4. We call two local  $k$ -flows  $\mathcal{F}_1 = (Y_1, I_1, \varphi_1)$ ,  $\mathcal{F}_2 = (Y_2, I_2, \varphi_2)$  coherent if  $Y_1 \cap Y_2 \neq \emptyset$  and there exists a  $(k-1)$ -flow  $\mathcal{F}_{12} = (Y_1 \cap Y_2, I_{12}, \varphi_{12})$  such that

$$\mathcal{F}_{12} \subseteq \partial \mathcal{F}_1 \cap \partial \mathcal{F}_2$$

and  $I_{12}$  does not reduce to a single point for any  $y$ .

Apparently,  $Y_1 \cap Y_2 = \partial Y_1 \cap \partial Y_2$ ,  $I'_{12}(y) = \{t \in I_1(y) : \varphi_1(t, y) \in Y_1 \cap Y_2\} = \{t \in I_2(y) : \varphi_2(t, y) \in Y_1 \cap Y_2\}$ , and  $\varphi_1 = \varphi_2 = \varphi_{12}$  in the set  $\{t, y : y \in Y_1 \cap Y_2, t \in I'_{12}(y)\}$  [vid. Definitions 9.2 and 9.3].

DEFINITION 9.5. A local flow  $\mathcal{F} = (Y, I, \varphi)$  is called the union of the flows  $\mathcal{F}' = (Y', I', \varphi')$  and  $\mathcal{F}'' = (Y'', I'', \varphi'')$ , if  $\mathcal{F}'$  and  $\mathcal{F}''$  are both subflows of  $\mathcal{F}$  and  $Y$  is the union of  $Y'$  and  $Y''$ . We use the notation

$$\mathcal{F} = \mathcal{F}' \cup \mathcal{F}''.$$

Obviously the union of local flows is associative in the sense

that if one of the sums  $\mathcal{F}_1 \cup (\mathcal{F}_2 \cup \mathcal{F}_3)$ ,  $\mathcal{F}_2 \cup (\mathcal{F}_1 \cup \mathcal{F}_3)$ ,  $\mathcal{F}_3 \cup (\mathcal{F}_1 \cup \mathcal{F}_2)$  exist, they all exist and are equal.

PROPOSITION 9.6. The necessary and sufficient condition in order that two coherent local flows  $\mathcal{F}_1 = (Y_1, I_1, \varphi_1)$  and  $\mathcal{F}_2 = (Y_2, I_2, \varphi_2)$  possess a union is that

$$(9.1) \quad \gamma_1(Y_1 \cap Y_2) = \gamma_2(Y_1 \cap Y_2) = Y_1 \cap Y_2.$$

Here  $\gamma_1, \gamma_2$  denote the paths of  $\mathcal{F}_1, \mathcal{F}_2$ .

PROOF. Apparently, (9.1) implies

$$(9.2) \quad I_1 = I_2 \quad \text{on} \quad Y_1 \cap Y_2.$$

If (9.1) is satisfied, we define  $I$  and  $\varphi$  on  $Y = Y_1 \cap Y_2$  by

$$I(y) = \begin{cases} I_1(y) & \text{for } y \in Y_1, \\ I_2(y) & \text{for } y \in Y_2 \end{cases}$$

$$\varphi(t, y) = \begin{cases} \varphi_1(t, y) & \text{for } y \in Y_1, t \in I_1(y), \\ \varphi_2(t, y) & \text{for } y \in Y_2, t \in I_2(y). \end{cases}$$

[On  $Y_1 \cap Y_2$  the equality  $\varphi_1 = \varphi_2$  holds due to the coherence property (vid. the remark following Definition 9.4).] Then  $(Y, I, \varphi)$  obviously satisfies the conditions (a), (b), (c) of Definition 9.1. Condition (d) follows easily from the following lemma:

Consider two compact sets  $A_1, A_2$  in a topological space  $X$  and two homeomorphic mappings  $\varphi_1 : A_1 \longrightarrow A_1', \varphi_2 : A_2 \longrightarrow A_2' (A_1', A_2' \subseteq X)$ . Assume  $\varphi_1 = \varphi_2$  on  $A_1 \cap A_2$ . Then the mapping

$$\varphi = \begin{cases} \varphi_1 & \text{on } A_1 \\ \varphi_2 & \text{on } A_2 \end{cases}$$

is homeomorphic if and only if

$$(9.3) \quad A_1' \cap A_2' = \varphi(A_1 \cap A_2).$$

Indeed, (9.3) is necessary and sufficient in order that  $\varphi$  is one-one. The continuity of  $\varphi$  is obvious; therefore, since  $A_1$  and  $A_2$

are compact,  $\varphi$  is homeomorphic if (9.3) is satisfied.

For every  $t$  the mappings  $\varphi_j(y) = \varphi_j(t, y)$  ( $j = 1, 2$ ) of the sets  $A_j = \{y \in Y_j : t \in I_j(y)\}$  satisfy the hypotheses of the lemma and condition (9.3) due to (9.1), so that  $\varphi$  satisfies (d).

We now assume the existence of a union  $\mathcal{F} = (Y, I, \varphi) = \mathcal{F}_1 \cup \mathcal{F}_2$ . Consider an interior point  $y$  of  $Y_1$  and suppose that  $I(y) \supset I_1(y)$ . Then, by Definition 9.2, one endpoint  $y_0$  of  $\gamma_1(y)$  would belong to  $Y_1 \cap Y_2 \subset \partial Y_1$ . Consider the path  $\gamma$  through  $y_0$  of the common face of  $\mathcal{F}_1$  and  $\mathcal{F}_2$  [vid. Definition 9.4]. Since  $\gamma$  is contained in  $\partial Y_1 \cap \gamma_1(y)$  and does not reduce to a point, a whole arc of  $\gamma_1(y)$  would lie on  $\partial Y_1$ , in contradiction to Proposition 9.4. Thus  $I(y) = I_1(y)$ .

Now consider a point  $y \in Y_1 \cap Y_2 \subset \partial Y_1$ . Every neighborhood of  $y$  contains an interior point  $y_1$  of  $Y_1$ . Since  $I(y_1) = I_1(y_1)$  and the endpoints of  $I$  and  $I_1$  depend continuously on  $y$ , we conclude that also in this case  $I(y) = I_1(y)$  holds and, in the same manner,  $I(y) = I_2(y)$ , so that (9.2) follows.

Now consider two points  $p \in Y_1 \cap Y_2$ ,  $q = \varphi_1(t, p)$ . Then, since  $\varphi_1$  and  $\varphi_2$  are both restrictions of  $\varphi$ ,  $\varphi_2(t, p) = q$  holds. Thus  $q \in Y_1 \cap Y_2$  and (9.1) follows.

PROPOSITION 9.7. If two local flows  $(Y', I', \varphi')$  and  $(Y'', I'', \varphi'')$  are linked by the relation

$$Y'_1 = Y''_0,$$

they possess a union  $(Y, I, \varphi)$ , where  $Y, I, \varphi$  are defined by

$$Y = Y' \cup Y'' ,$$

$$I(y) = \begin{cases} [t'_0(y), t'_1(y) + t'_1(y'_1(y))] & \text{for } y \in Y' , \\ [t''_0(y) + t'_0(y''_0(y)), t''_1(y)] & \text{for } y \in Y'' , \end{cases}$$

$$\varphi(t, y) = \begin{cases} \varphi'(t, y) & \text{for } y \in Y' , \\ \varphi''(t, y) & \text{for } y \in Y'' . \end{cases}$$

The proof follows easily from the definitions of the sets  $Y_0, Y_1$ .

# §5. THE QUALITATIVE PICTURE OF THE SOLUTIONS NEAR A SWITCHING SPACE

10. Local flows around sets of normal switching points. Throughout this chapter  $U$  will denote a compact connected subset of  $S - S''$  [vid. §2.5], containing no points of switching spaces other than  $S$ .

Consider first the case  $U \subset T$ . Since  $U$  is compact, there exist two numbers  $\alpha (< 0)$ ,  $\beta (> 0)$  such that the functions  $x^\pm(t, u)$  satisfy (S) in the intervals  $I^+ = [0, \beta]$  and  $I^- = [\alpha, 0]$ , respectively, and define local flows  $\varphi^\pm$  on the sets  $x(I^\pm, U)$ . Moreover,  $x(t, u)$  defines a local flow  $\varphi = \varphi^+ \cup \varphi^-$  on  $x(I, U)$  ( $I = I^- \cup I^+$ ) [vid. Proposition 9.7].

In the case  $U \subset A \cup E$  the set  $U$  is the common entrance [exit] set of two local flows defined by solutions of (S) on the sets  $x^+(I^+, U)$  [ $x^-(I^-, U)$ ], respectively, where  $I^+ = [0, \beta]$ ,  $I^- = [\alpha, 0]$ ;  $\alpha, \beta > 0$  and sufficiently small.

The ambiguity occurring in connection with starting and end points can be eliminated by a modification of the phase space  $E^n$ , namely by "cutting" it along the sets  $A, E$  and replacing every point  $u \in A, E$  by a pair of points  $u^\pm$ , situated on either "edge" of the cut. More precisely, this procedure can be described as follows: Denote by  $V$  the set

$$V = A \cup E$$

and consider an open set  $X \subset E^n$  satisfying the condition  $X \cap S = V$ . Then complete the set  $X - V$  (with respect to the euclidean metric), and finally replace  $E^n$  by the space

$$D^* = (E^n - X) \cup \overline{(X - V)},$$

where the closure is to be understood in the sense of the new topology, defined by the Cauchy sequences in  $X - V$ . Obviously, the construction of the space  $D^*$  does not depend on the choice of  $X$ . Denote by  $\varphi$  the continuous mapping of  $D^*$  onto  $E^n$  which leaves every point of  $E^n - V$  fixed. Each point  $v \in V$  possesses two images under  $\varphi = \varphi^{-1}$  which we denote by  $\varphi^\pm(v) = v^\pm$  and which are distinguished from each other by the property

$$v^\pm \in \bar{D}_e^\pm.$$

Apparently, through every point  $p \in D^*$  there exists exactly one solution of (S). This we denote by  $x(t, p)$ .

11. Local flows around the sets  $(AL)^1$  and  $(EL)^1$ . If  $U$  is a subset of  $(AL)^1$ , it is easy to see that it is contained in one of the sets  $A^+ \cap L^-$  and  $A^- \cap L^+$ . Indeed, in the opposite case the connectedness of  $U$  would imply the existence of a point

$$u \in U \cap \overline{A^+ \cap L^-} \cap \overline{A^- \cap L^+},$$

which we could, e.g., assume to belong to  $A^+ \cap L^-$ . This would imply  $c_1^+ > 0$  [vid. Theorem 6.1]. Since, however, every neighborhood of  $u$  would contain points of  $A^- \cap L^+$ , on which  $c_1^+ = 0$ ,  $c_1^+$  could not be continuous. Therefore, without loss of generality, we may assume

$$U \subset A^+ \cap L^-.$$

Relation (8.3) implies  $U \subset \partial A \cap \partial T$ . Moreover,  $U$  is contained in the intersection of the boundaries of two components  $A_1, T_1$  of  $A$  and  $T$ :

$$(11.1) \quad U \subset \partial A_1 \cap \partial T_1.$$

This follows immediately from the connectedness of  $U$  and from Theorem 7.1,<sup>1</sup> according to which every point  $u \in U \subset S^1$  lies on the boundary of exactly two components of  $S^0$ .

For every point  $u \in U$  the relation

$$(11.2) \quad x^-(t, u) \in D_{\theta}^+ \quad (0 < |t| \leq t')$$

holds for sufficiently small  $t'$  [Theorem 6.1]. If  $v \in A_1$ , we have

$$(11.3) \quad x^-(t, v) \in D_{\theta}^- \quad (0 < t < t_1(v))$$

for small  $t_1(v)$ . The continuous dependence of  $x^-$  on the initial value, together with (11.2), implies the existence of a neighborhood  $N$  of  $u$  such that

$$(11.4) \quad x^-(t', N) \subset D_{\theta}^+$$

<sup>1</sup> We again assume condition A to be satisfied.



holds. We now restrict  $v$  to  $A_1 \cap N$ . Because of (11.3) and (11.4), we can choose  $t_1(v)$  in such a way that  $x^-(t_1, v) \in S$  and (11.3) holds for  $v \in N \cap A_1$ . Since  $U$  is compact, there exist a connected compact set  $\mathcal{A}$ , containing  $U$  and contained in  $U \cup A_1$ , and a (continuous) function  $t_1$  defined on  $\mathcal{A}$  such that

$$(11.5) \quad x^-(t, v) \in \begin{cases} A_1 & \text{for } t = 0, \\ D_0 & \text{for } 0 < t < t_1(v), \\ T_1 & \text{for } t = t_1(v) \end{cases},$$

holds for all  $v \in \mathcal{A} - U$ . [It is sufficient that  $\mathcal{A}$  is contained in the union of a finite number of sufficiently small open sets  $N_K$  (satisfying, in particular, (11.4)) which together cover  $U$ .] Moreover, we can assume  $\mathcal{A}$  to be a  $(n-1)$ -dimensional manifold with boundary (because  $A_1$  is an open subset of the hypersurface  $S$ ). Since  $U \subseteq (AL)^1$  and  $\mathcal{A} \subseteq U \cup A_1$ , (11.1) implies

$$U \subseteq \partial T_1 \subseteq \bar{T}_1 \subseteq \overline{S - (A_1 \cup (AL)^1)} \subseteq \overline{S - (A_1 \cup U)} \subseteq \overline{S - \mathcal{A}}.$$

This, together with  $U \subseteq \mathcal{A}$ , yields

$$(11.6) \quad U \subseteq \partial \mathcal{A}.$$

By cutting the space along  $A$ , as described in the preceding section, the set  $\mathcal{A}$  is split into two "sheets"  $\mathcal{A}^\pm = \psi^\pm(\mathcal{A})$ . Since  $\mathcal{A}$  is compact, the mappings  $\psi^\pm : \mathcal{A} \rightarrow \mathcal{A}^\pm$  are homeomorphic. This implies the compactness of  $\mathcal{A}^\pm$  and, together with (11.6), the relation

$$(11.7) \quad U \subseteq \partial \mathcal{A}^+ \cap \partial \mathcal{A}^-.$$

Then it follows immediately that

$$(11.8) \quad \mathcal{A}^+ \cap \mathcal{A}^- = U.$$

After these preparations it is easy to analyse the local flows around  $U$ . According to (11.5), the solutions of (S) define a local  $n$ -flow  $\mathcal{F}^- = (Y^-, I^-, x^-)$  on the set

$$Y^- = \{x(t, v) \mid t \in I^-(v), v \in \mathcal{A}^-\}, \quad I^-(v) = [0, t_1(v)] \text{ for } v \in \mathcal{A}^-.$$

Using the notations of §4 (and indicating by a superscript  $-$  the relatedness to  $\mathcal{F}^-$ ),  $t_1(v) = t_1^-(v)$  holds for  $v \in \mathcal{A}^-$ . Since  $t_1(v) \rightarrow 0$

with  $v \longrightarrow U$ , it follows that

$$(11.9) \quad t_0^-(y) \longrightarrow 0 \text{ with } y \longrightarrow U.$$

Since  $x^+(t, v) \in D_0^+$  holds for all  $v \in U \cup A_1 \cup T_1$  and small  $t$  and  $\mathcal{X} \cup Y_1^-$  is a compact subset of  $U \cup A_1 \cup T_1$ , there exists an interval  $I_0 [0, t^*]$  ( $t^* > 0$ ) such that

$$x^+(I_0, \mathcal{X} \cup Y_1^-) \subset D_0^+.$$

As a consequence of (11.9), the inequality  $t_0^-(v) < t^*$  holds for all  $v \in Y_1^-$  if  $\mathcal{X}$  is sufficiently small. Therefore, the solutions  $x(t, v)$  of (S) define a local flow  $\mathcal{F}^+$  with the entrance set  $\mathcal{X}^+ \cup Y_1^- = Y_0^+$  and the intervals of definition

$$(11.10) \quad I^+(v) = \begin{cases} [0, t^*] & \text{for } y \in \mathcal{X}^+, \\ [0, t^* + t_0^-(v)] & \text{for } v \in Y_1^-. \end{cases}$$

Now we split up the local flow  $\mathcal{F}^+$  into two subflows  $\mathcal{F}^1$  and  $\mathcal{F}^2$ , defined on the sets  $x(I^+, \mathcal{X}^+)$  and  $\bigcup_{v \in Y_1^-} x(I^+(v), v)$ , respectively. Then  $Y_1^- = Y_0^2$  holds and, due to Proposition 9.7,  $\mathcal{F}^- \cup \mathcal{F}^2$  is a local flow which we denote by  $\mathcal{F}' = (x(I_0, \mathcal{X}^-), I', x)$ . Since  $\mathcal{X}$  is an  $(n-1)$ -dimensional manifold with boundary,  $\mathcal{F}^1$  and  $\mathcal{F}'$  are both local  $n$ -flows. Denoting by  $\mathcal{F}^{12}$  the  $(n-1)$ -flow defined on  $x(I_0, U)$ , (11.7) together with Proposition 9.5 implies

$$\mathcal{F}^{12} \subset \partial \mathcal{F}^1 \cap \partial \mathcal{F}'.$$

From this and (11.8) it follows that  $\mathcal{F}^1$  and  $\mathcal{F}'$  are coherent.

Finally, if  $v \in Y_1^-$ , Proposition 9.7 and (11.10) yield

$$I'(v) = [t_0^-(v), t^* + t_0^-(v)]$$

which, together with (11.9), implies

$$I'(u) = [0, t^*] = I_0 \quad (u \in U).$$

Since also  $I^1(u) = I_0$ , condition (9.1) of Proposition 9.6 is satisfied for the flows  $\mathcal{F}^1$  and  $\mathcal{F}'$ , their union thus being a local flow:<sup>1</sup>

<sup>1</sup> Vid. remark following Definition 9.5.

$$\begin{aligned} \mathcal{X} &= \mathcal{X}^1 \cup \mathcal{X}^- = \mathcal{X}^1 \cup (\mathcal{X}^2 \cup \mathcal{X}^-) = (\mathcal{X}^1 \cup \mathcal{X}^2) \cup \mathcal{X}^- \\ &= \mathcal{X}^+ \cup \mathcal{X}^- . \end{aligned}$$

This yields the following theorem:

**THEOREM 11.1.** Consider a connected compact subset  $U$  of  $(AL)^1$  [ $= AL \cap S^1$ ; vid. §2, Sections 4 and 5], on which condition A is satisfied [vid. §3.7]. Assume the phase space to be cut along the set  $A$  of starting points. Then  $U$  is contained in a compact connected set  $Y$  which is an  $n$ -dimensional manifold with boundary and has the property that all solutions of (S) which contain points of  $Y$  form a single local flow  $\mathcal{X}$  on  $Y$ . The entrance set of  $\mathcal{X}$  consists of  $U$  and the two "edges"  $\mathcal{X}^+$  of the cut along  $A \cap Y$ . The paths originating on  $\mathcal{X}^-$  [or on  $\mathcal{X}^+$  in the case  $U \subset A^- \cap L^+$ ] traverse the switching space exactly once. The exit set is contained in  $D_{e^+} \cup D_{e^-}$ . (Vid. Figures 2, 3).

The case  $U \subset (EL)^1$  is exactly analogous and can be reduced to the one considered above by reversing the sign of  $t$ .

**12. Local flows around the sets  $(AR)^1$  and  $(ER)^1$ .** If  $U$  is a subset of  $(AR)^1$  it can be assumed without loss of generality that it is contained in  $A^+ \cap R^-$ . [The proof is analogous to the corresponding one in the preceding section.] Denote again by  $A_1, T_1$  the components of  $A, T$  which are adjacent to  $U$ . Let  $\mathcal{X}$  and  $\mathcal{T}$  be compact connected sets satisfying the conditions

$$U \subset \mathcal{X} \subset U \cup A_1, \quad U \subset \mathcal{T} \subset U \cup T_1 .$$

We can assume  $\mathcal{X}$  and  $\mathcal{T}$  to be  $(n-1)$ -dimensional manifolds with boundary;  $\mathcal{X}$  will again be replaced by two "sheets"  $\mathcal{X}^\pm$ .

Through every point  $u \in U$  there exist exactly two solutions  $x_1(t, u), x_2(t, u)$  of (S). Since  $U$  is compact, the inclusions

$$(12.1) \quad x_1(t, U) \subset D_{e^-} \quad (-t_1 < t < t_2; t \neq 0)$$

and




Fig. 2.




Fig. 3.




Fig. 4.

$$(12.2) \quad x_2(t, U) \subset \begin{cases} D_{\theta}^- & \text{for } -t_1 < t < 0, \\ D_{\theta}^+ & \text{for } 0 < t < t_3 \end{cases}$$

hold for sufficiently small positive numbers  $t_1, t_2, t_3$  [vid. §2.6]. Due to the continuous dependence of  $x^\pm$  on the initial values, it follows from (12.1) and (12.2) that the solutions of (S) define three local flows  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_4$  on the sets  $x(I_1, \mathcal{A}^+)$ ,  $x(I_2, \mathcal{T})$ ,  $x(I_4, \mathcal{A}^-)$ , respectively, where  $I_1 = [0, t_3]$ ,  $I_2 = [-t_1', t_3']$ ,  $I_4 = [0, t_2']$ ,  $0 < t_j' < t_j$  ( $j = 1, 2, 3$ ) and  $\mathcal{A}$  and  $\mathcal{T}$  are sufficiently small. The local  $(n-1)$ -flows defined on the sets  $x_2(I_1, U)$ ,  $x_1([-t_1', 0] \cup U)$ ,  $x_1(I_4, U)$  we denote by  $\mathcal{F}_{12}, \mathcal{F}_{23}, \mathcal{F}_{34}$ . The relations (11.1) and (11.7) obviously hold also in the case under consideration. Since (11.1) implies  $U \subset \partial \mathcal{T}$ , Proposition 9.5 yields the incidences

$$\mathcal{F}_{12} \subset \partial \mathcal{F}_1 \cap \partial \mathcal{F}_2, \quad \mathcal{F}_{23} \subset \partial \mathcal{F}_2, \quad \mathcal{F}_{34} \subset \partial \mathcal{F}_4.$$

The first of these implies the coherence of the local flows  $\mathcal{F}_1$  and  $\mathcal{F}_2$ .<sup>1</sup> Moreover, due to Proposition 9.7, the unions  $\mathcal{F}_{23} \cup \mathcal{F}_{12}$  and  $\mathcal{F}_{23} \cup \mathcal{F}_{34}$  exist.<sup>2</sup>

Finally, a local flow which is incident with the  $(n-1)$ -flows  $\mathcal{F}_{23}$  and  $\mathcal{F}_{34}$  is obtained by considering a hypersurface  $H$  without contact<sup>3</sup> with the solutions of  $(S, e^-)$ , which contains  $U$ ,<sup>4</sup> and a compact connected subset  $C$  of  $H \cap \bar{D}_{\theta}^-$  the boundary of which contains  $U$ . Then, denoting by  $I_3$  the interval  $[-t_1', t_2']$ , every arc  $x^-(I_3, p)$ , where  $p \in C$ , intersects  $C$  only at  $p$ . Consequently, the solutions of  $(S)$  define a local flow  $\mathcal{F}_3$  on the set  $x^-(I_3, C)$ . Since  $U \subset \partial C$ , the incidence relations

$$\mathcal{F}_{23} \subset \partial \mathcal{F}_3, \quad \mathcal{F}_{34} \subset \partial \mathcal{F}_3$$

follow. Therefore,  $\mathcal{F}_2$  and  $\mathcal{F}_3$  as well as  $\mathcal{F}_3$  and  $\mathcal{F}_4$  are coherent.<sup>5</sup> We therefore have the result:

<sup>1</sup> However, the union of  $\mathcal{F}_1$  and  $\mathcal{F}_2$  is not a flow [vid. Proposition 9.6].

<sup>2</sup> This situation can be described as a "bifurcation" of the flow  $\mathcal{F}_{23}$  into  $\mathcal{F}_{12}$  and  $\mathcal{F}_{34}$ .

<sup>3</sup> Vid. p. , footnote 1.

<sup>4</sup> The set  $x^+(I, U)$ , e.g., has this property if  $I$  is a suitable interval around  $t = 0$ .

<sup>5</sup> Again (due to Proposition 9.6) neither of the unions is a local flow.

THEOREM 12.1. If  $U$  is a compact connected subset of  $(AR)^1 [= AR \cap S^1]$ , satisfying condition A, and the space is cut along the set of starting points, splitting it into two "sheets"  $\mathcal{A}^+$ , there exists a compact connected set  $Y$  which is an  $n$ -dimensional manifold with boundary containing  $U$  and has the following property: The solutions of (S) which contain points of  $Y$  constitute a complex of four local flows  $\mathcal{F}_1, \dots, \mathcal{F}_4$ , separated from each other by  $(n-1)$ -flows  $\mathcal{F}_{12}, \mathcal{F}_{23}, \mathcal{F}_{34}$  such that  $\mathcal{F}_{\mu, \mu+1}$  ( $\mu = 1, 2, 3$ ) is incident with  $\mathcal{F}_\mu$  and  $\mathcal{F}_{\mu+1}$  and the two latter are coherent. Upon reaching the set  $U$ , the flow  $\mathcal{F}_{23}$  bifurcates into  $\mathcal{F}_{12}$  and  $\mathcal{F}_{34}$  in the sense that the exit set of  $\mathcal{F}_{23}$  and the entrance sets of  $\mathcal{F}_{12}$  and  $\mathcal{F}_{34}$  are all equal to  $U$ . The local flows  $\mathcal{F}_1$  and  $\mathcal{F}_4$  originate on  $\mathcal{A}^+$  and  $\mathcal{A}^-$ , respectively. The paths of  $\mathcal{F}_2$  (or, if  $U \subset A^- \cap R^+$ , of  $\mathcal{F}_3$ ) and only these traverse the switching space.

Figure 4 gives a symbolic picture of the local flows and their interrelations.

The analogous result for the case  $U \subset (ER)^1$  is again obtained by reversing the orientation of the paths.

#### BIBLIOGRAPHY

- [1a] André, J. and Seibert, P., "Über stückweise lineare Differentialgleichungen, die bei Regelungsproblemen auftreten I, II," Archiv d. Math., Vol. 7 (1956), p. 148-156, 157-164.
- [1b]
- [2] Bellman, R., Glicksberg, I., and Gross, O., "On the "bang-bang" control problem," Quart. Appl. Math., Vol. 14 (1956), p. 11-18.
- [3] Boltyanskii, V. G., Gamkrelidze, R. V., Pontryagin, L. S., "On the theory of optimal processes," (Russian), Dokl. Akad. Nauk SSSR (N.S.) Vol. 110 (1956), p. 7-10.
- [4] Bushaw, D., "Optimal discontinuous forcing terms," these Contributions, Vol. IV, Annals of Math. Studies 41 (1958), p. 29-52.
- [5] Flügge-Lotz, I., Discontinuous automatic control, Princeton, 1953.
- [6] Gamkrelidze, R. V., "Theory of time-optimal processes in linear systems," (Russian), Izvest. Akad. Nauk SSSR, Ser. Mat., Vol. 22 (1958), p. 449-474.

- [7] Krasovskiy, N. N., "On stability in the case of large initial perturbations," (Russian), Prikl. Mat. Meh., Vol. 21 (1957), p. 309-319.
- [8] \_\_\_\_\_, "On the theory of optimal control," (Russian), Avtomat. i Telemekh., Vol. 18 (1957), p. 960-970.
- [9] Pontryagin, L. S., "Optimal control processes," (Russian), Uspehi Mat. Nauk, Vol. 14 (1959), p. 3-20.
- [10] Solncev, Yu. K., "On stability according to Lyapounov of the equilibrium positions of a system of two differential equations in the case of discontinuous right hand sides," (Russian), Moscov. Gos. Univ. Uchenye Zapiski, Vol. 148, Mat. 4 (1951), p. 144-180.





## XII. ASYMPTOTIC STABILITY IN 3-SPACE

Courtney Coleman

1. Consider the system,

$$(1) \quad \frac{dx}{dt} = f^{(m)}(x) + g(x, t) \quad ,$$

where  $f^{(m)}(x)$  and  $g(x, t)$  satisfy the following conditions:

- a) Both are  $n$ -vector functions of the  $n$ -vector  $x$ .  
 $g$  may depend on  $t$ .  $f^{(m)}(0) = g(0, t) = 0$ .
- b) In some neighborhood,  $D$ , of  $x = 0$  and for all  $t$  greater than some  $t_0$ ,  $f^{(m)}(x)$  and  $g(x, t)$  are continuous in  $x$  and  $t$  and satisfy a Lipschitz condition in  $x$ .
- c)  $f^{(m)}(x)$  is a homogeneous function of degree  $m$  —  
 $(m$  an integer  $> 1$ ) in  $D$ ; i.e.,  $f^{(m)}(x) = c^m f(x)$  for any constant  $c > 0$  and for all  $x$  in  $D$ .
- d)  $g(x, t) = o(\|x\|^m)$  uniformly in  $t$  in  $D$ .

Malkin [1] and Massera [2] have shown that if the trivial solution,  $x = 0$ , of the system of first approximation,

$$(2) \quad \frac{dx}{dt} = f^{(m)}(x) \quad ,$$

is asymptotically stable, then so is the trivial solution of (1). The problem, then, is to find conditions ensuring the asymptotic stability of the trivial solution of (2). This will be done for  $n = 3$  (see the theorem below) using some of the results obtained by the author [3]. For  $n = 2$  such conditions can be found implicitly in a number of papers -- e.g., see Forster [4]. The conditions are simply that the origin be an attracting focus if the function  $h(x_1, x_2) = x_2 f_1^{(m)} - x_1 f_2^{(m)}$  has no real zeros, and that

$$\frac{dr}{dt} < 0 \quad (r^2 = x_1^2 + x_2^2)$$

on the zeros of  $h$  if they exist.

It should be noted that for  $m = 1$  and for any  $n$  the theory of characteristic roots holds. Clearly, if  $m$  is even, there can be no asymptotic stability. Hence,  $m$  is assumed odd.

2. Let  $n = 3, m > 1$ . Let  $r = (x_1^2 + x_2^2 + x_3^2)^{\frac{1}{2}}, y = \frac{1}{r} x$ . Then (2) becomes in  $r, y$ ,

$$\begin{aligned} \text{a.} \quad \frac{dr}{dt} &= (y, f^{(m)}(y))r^m \\ \text{b.} \quad \frac{dy}{dt} &= (f^{(m)}(y) - (y, f^{(m)}(y))y)r^{m-1}, \end{aligned} \quad (3)$$

where  $(, )$  indicates scalar product. Let  $r^{m-1}(t)dt = d\tau$ . Then (3) becomes,

$$\begin{aligned} \text{a.} \quad \frac{dr}{d\tau} &= (y, f^{(m)}(y))r \\ \text{b.} \quad \frac{dy}{d\tau} &= f^{(m)}(y) - (y, f^{(m)}(y))y. \end{aligned} \quad (4)$$

If  $y(\tau)$  is a solution of (4b) and  $r(\tau)$  is a solution of (4a) in which  $y$  stands for  $y(\tau)$  (the solution of 4b), then  $(r(\tau), y(\tau))$  is a solution of (4). Similarly for (3a, b). System (4) is more suitable for our purposes since no solution of (4) may have a finite "escape time" while a solution of (3) may. Ignoring the differing parametrizations, the trajectories in 3-space of (3) coincide with those of (4); for the only difference between the two systems is the factor  $r^{m-1}$ , which is positive everywhere except at the origin. This implies in addition that the trajectories of the two systems agree in the sense of parametric description. If all the trajectories of (4) close enough to 0 tend to 0 as limit with increasing  $\tau$  and if none tend to 0 with decreasing  $\tau$ , then the same is true for the trajectories of (3) and hence of (2). Since systems (2), (3), (4) are autonomous, to prove asymptotic stability it is sufficient to show that there is a neighborhood  $D^* \subset D$  such that for any trajectory  $r$  of (4) which cuts  $D^*$   $\Omega(r)$  (the omega limit set of  $r$ ) contains 0 only and  $A(r)$  (the alpha limit set of  $r$ ) has no points in  $D^*$ . Actually for systems (2), (3) and (4)  $D$  and  $D^*$  might as well be all of 3-space.

The following properties of system (4) and its trajectories were observed in [3].

- P I. Each trajectory  $\gamma : y = y(\tau, y_0, \tau_0)$  of (4b) lies on the unit sphere  $S^2$ . The Poincaré-Bendixson theory can then be applied to  $\Omega(\gamma)$  and  $A(\gamma)$ .
- P II. Each trajectory  $\gamma$  of (4b) determines a cone  $S(\gamma)$  generated by rays through 0 and the points of  $\gamma$ .  $S(\gamma)$  is an integral surface of (4).
- P III. On  $S(\gamma)$  each  $\Gamma$  of (4) is given by

$$\left[ K \cdot \exp \int_{\tau_0}^{\tau} (y, f^{(m)}(y)) d\tau \right] \cdot y(\tau) ,$$

where  $K$  is an arbitrary positive constant.

- P IV. If  $\lambda(p)$  is a ray on  $S(\gamma)$  through 0 and  $p$ , then all trajectories of (4) on  $S(\gamma)$  cut  $\lambda(p)$  at the same non-zero angle in the same sense (except in the trivial case that  $\gamma$  is a singular point of (4b)).

The following general properties of a system of autonomous differential equations,

$$(5) \quad \frac{dx}{dt} = h(x), \quad x \text{ an } n\text{-vector},$$

and its trajectories will be needed. All functions are assumed analytic.

- P Va. If  $N(p)$  is a small enough neighborhood of the non-critical point  $p$  in the normal hyperplane to the trajectory of (5) through  $p$ , then the trajectories of (5) through the points of  $N(p)$  cross  $N(p)$  from one side to the other in the same sense.
- P Vb. If  $S$  is a smooth surface in  $x$ -space and  $C$  a simple smooth arc on  $S$  such that all trajectories of (5) cutting  $C$  do so crossing  $C$  from one side of  $S$  to the other and not tangent to  $S$ , then there is a neighborhood  $N(C)$  in  $S$  of  $C$  such that the same is true for all points of  $N(C)$ .

P VI. The  $A$  limit set ( $\Omega$  limit set) of a bounded trajectory of (5) is closed, non-empty and connected. If a point belongs to a limit set of some trajectory, then so does the full trajectory through the point.

In the proof of the theorem that follows frequent use will be made of PI - PVI. Unless required for clarity specific reference will not be made.

THEOREM. Suppose

$$a) \quad \begin{aligned} & (y, f(y)) < 0 \text{ on all zeros of} \\ & f^{(m)}(y) - (y, f^{(m)}(y))y \end{aligned}$$

and

$$b) \quad \int_0^T (y, f^{(m)}(y)) d\tau < 0 \text{ for each periodic}$$

solution  $y(\tau)$  of (4b) ( $T$  is the period of  $y(\tau)$ ).

Then the solution  $x = 0$  of (1) for  $n = 3$  is asymptotically stable.

PROOF. It is sufficient to show that under assumptions a) and b)  $\Omega(\Gamma)$  contains only 0 and  $A(\Gamma)$  has no finite points for any trajectory  $\Gamma$  of (4).

In the following  $\gamma, \gamma^*, \gamma_i, i = 1, \dots, k$ , denote trajectories of (4b) on  $S^2$  and  $P, P_i, i = 1, \dots, k$  critical points of (4b) on  $S^2$ .  $S(\gamma_i), S(P_i), \dots$  denote the integral surfaces determined by  $\gamma_i, P_i, \dots$ .  $\Gamma, \Gamma^*, \Gamma_i, i = 1, \dots, k$  denote trajectories of (4) on the integral surfaces  $S(\gamma), S(\gamma^*), S(P), S(\gamma_i), \dots$  respectively. The Poincaré-Bendixson theory states that a limit set of a trajectory  $\gamma$  of (4b) on  $S^2$  is one of the following:

(L 1a.)  $\gamma$  itself, where  $\gamma$  is a single point;

(L 2b.)  $\gamma$  itself, where  $\gamma$  is a closed curve;

(L 2a.) a critical point  $P$ , not  $\gamma$ ;

(L 2b.) a closed curve  $\gamma^*$ , not  $\gamma$ ;

(L 3) a closed graph  $G : P_1 \cup \gamma_1 \cup \dots \cup P_k \cup \gamma_k \cup P_{k+1}$ ,  
where  $\Omega(\gamma_1) = P_1, A(\gamma_1) = P_{1+1}, P_{k+1} = P_1$ .

The various combinations of these possibilities for  $A(\gamma)$  and  $\Omega(\gamma)$  must now be considered.

- A. (L 1a.) for  $A(\gamma)$  and hence for  $\alpha(\gamma)$ . Hypothesis a) ensures that for the ray trajectory  $\Gamma$  of (4) through  $\gamma$ ,  $\alpha(\Gamma)$  contains 0 alone and  $A(\Gamma)$  is infinite. (See P III).
- B. (L 1b.) for  $A(\gamma)$  and hence for  $\alpha(\gamma)$ . Here hypothesis b) ensures that  $\alpha(\Gamma)$  consists only of 0 and  $A(\Gamma)$  is infinite (See P III and [3]).
- C. (L 2a.) for  $A(\gamma)$  and for  $\alpha(\gamma)$ . Whether or not  $A(\gamma) = \alpha(\gamma)$ ,  $A(\Gamma)$  is infinite and  $\alpha(\Gamma)$  contains just 0. (See P III, Iv and [3]).
- D. (L 2a.) for  $A(\gamma)$  and (L 2b.) for  $\alpha(\gamma)$ . Let  $A(\gamma) = P$ . Then  $\Gamma_P$  is as in A above. By (P VI) for any  $\Gamma$  on  $S(\gamma)$ ,  $A(\Gamma)$  cannot contain any point of  $\Gamma_P$  unless it contains  $\Gamma_P$  itself. This is impossible by (P Va.) which also implies that  $A(\Gamma)$  cannot contain 0 alone. Hence  $A(\Gamma)$  has no finite points. On  $S(\gamma^*)$  B. above obtains. Hence  $\alpha(\Gamma)$  has no finite points, or contains 0 and at least one  $\Gamma^*$  on  $S(\gamma^*)$ , or contains 0 alone. Let  $p$  be any point of  $\gamma^*$ . Let

$$\int_0^T (y^*, f(y^*)) d\tau = -K < 0,$$

where  $\gamma^*$  is given by  $y = y^*(\tau)$ ,  $y^*(\tau + T) = y^*(\tau)$ . Let  $\sigma$  be an arc (bounded on one end by  $p$ ) of the great circle on  $S^2$  through  $p$  normal to  $\gamma^*$ . Let  $\gamma$  be given by  $y = y(\tau)$  and let  $d$  denote Euclidean distance. If  $y(\tau)$  is on  $\sigma$  for some  $\tau$ , denote by  $T_\tau$  the smallest positive number such that  $y(\tau + T_\tau)$  is on  $\sigma$ . If the length of  $\sigma$  is small enough,  $T_\tau$  is defined for all  $\tau$  for which  $y(\tau)$  is on  $\sigma$ ; and if  $y(\tau_0)$  is on  $\sigma$ , then

$$\max_{\tau \geq \tau_0} d(y(\tau), \gamma^*)$$

(for all  $\tau \geq \tau_0$ ) and

$$T_\tau, \tau \geq \tau_0 \quad |T - T_\tau| \quad (y(\tau) \text{ on } \sigma)$$


are so small that

$$\int_{\tau}^{\tau+T_{\tau}} (y, f(y)) d\tau \leq -K/2 < 0 \quad (y(\tau) \text{ on } \sigma) .$$

But this can be done for every point of  $\gamma^*$ . Hence,  $\alpha(r)$  can contain 0 alone (recalling that  $r$  is given by

$$\left[ K \exp \int_{\tau_0}^{\tau} (y, f(y)) d\tau \right] \cdot y(\tau) .$$

- E. (L 2a.) for  $A(\gamma)$  and (L 3.) for  $\alpha(r)$ . As in D, for any  $r$  on  $S(\gamma)$   $A(r)$  contains no finite points. We must show that  $\alpha(r)$  contains 0 alone. In the first place for any  $r_1$  or  $r_p$  on the cone  $S(G)$ ,  $A(r_1)$  and  $A(r_{p_1})$  have no finite points and  $\alpha(r_1)$  and  $\alpha(r_{p_1})$  contain 0 alone. For, each  $S(\gamma_1)$  or  $S(p_1)$  is covered by cases A. or C. above (Figure 1). Hence, as in D, for any  $r$  on  $S(\gamma)$ ,  $\alpha(r)$  has no finite points, contains 0 and at least one  $r_1$  or  $r_{p_1}$  on  $S(G)$ , or else contains 0 alone. The first two alternatives must be excluded.



First a simple closed curve without contact  $C$  is constructed on  $S(G)$  such that one of the components of the complement of  $C$  on  $S(G)$  has 0 as a boundary point and is bounded. Let  $q_0$  be any point on  $S(\gamma_1)$ . Suppose each  $r_1$  on  $S(\gamma_1)$  tends to 0 with increasing  $\tau$  tangent to  $r_{P_1}$  and to infinity parallel to  $r_{P_2}$  with decreasing  $\tau$  (see C. above and [3]). Let

$$y(\tau_0) = \frac{1}{\|q_0\|} \cdot q_0 .$$

FIGURE 1

Considering all rays as being directed toward 0, let  $0 < \theta(q) < \pi$  be the angle defined by the field vector at  $q$  and the directed ray  $\lambda(q)$  for each  $q$  on  $S(\gamma_1)$ .  $\theta(q)$  is a continuous function of  $q$  as  $q$  varies along any continuous curve on  $S(\gamma_1)$ . Define  $\varepsilon(\tau)$  as any positive continuous function of  $\tau$  such that

$$\lim_{\tau \rightarrow \infty} \int_{\tau_0}^{\tau} [(y, f^{(m)}(y)) + \varepsilon(\tau)] d\tau = \ln \frac{\|q_1\|}{\|q_0\|}$$

where  $q_1$  is some point of  $\Gamma_{P_1}$ . There are many ways of defining  $\varepsilon(\tau)$ . For example, since by hypothesis (a)  $(y, f(y))$  is negative and bounded away from 0 for  $\tau$  large enough,  $\varepsilon(\tau)$  could be taken as

$$\frac{1}{(\tau+1-\tau_0)^2} \cdot \ln \frac{\|q_1\|}{\|q_0\|} - (y, f(y))$$

for  $\tau$  large enough.

Define the curve  $C_1^+(q_0, q_1, \tau)$  as

$$q_1 U \left\{ \|q_0\| \exp \int_{\tau_0}^{\tau} [(y, f(y)) + \varepsilon(\tau)] d\tau \right\} \cdot y(\tau)$$

for  $\tau \geq \tau_0$ .  $C_1^+$  is then a smooth arc bounded by  $q_0$  and  $q_1$ .  $\frac{dr}{d\tau}$  at any point  $q$ ,  $q \neq q_0, \neq q_1$ , of  $C_1^+(q_0, q_1, \tau)$  is greater by the positive quantity  $\varepsilon(\tau)$  than  $\frac{dr}{d\tau}$  given by (4a). Hence the angle  $\theta(q)$  defined by the directed tangent to  $C_1^+(q_0, q_1, \tau)$  at  $q$  and the directed ray  $\lambda(q)$  is greater than  $\theta(q)$ . Then  $C_1^+(q_0, q_1, \tau)$  is an arc without contact. In a similar way,  $C_1^-(q_0, q_2, \tau)$  for  $\tau \leq \tau_0$ ,  $q_2$  on  $\Gamma_{P_2}$ , can be defined as a smooth curve without contact bounded by  $q_0$  and  $q_2$ . After small changes in  $C_1^-$  and  $C_1^+$  near  $q_0$ , if necessary, define  $C_1(q_1, q_2)$  as  $C_1^- U C_1^+$ .  $C_1(q_1, q_2)$  is then a smooth curve without contact on  $S(\gamma_1)$  bounded by  $q_1$  on  $\Gamma_{P_1}$  and  $q_2$  on  $\Gamma_{P_2}$ . All trajectories on  $S(\gamma_1)$  cross  $C_1(q_1, q_2)$  into the region on  $S(\gamma_1)$  bounded by 0,  $\Gamma_{P_1}$ ,  $\Gamma_{P_2}$ , and  $C_1(q_1, q_2)$ . Note that  $q_1$  and  $q_2$  are arbitrary points on  $\Gamma_{P_1}$  and  $\Gamma_{P_2}$ .

In like fashion a curve without contact  $C_1$  is constructed on each  $S(\gamma_i)$ . Since the boundary points of each  $C_1$  may be chosen arbitrarily, the  $C_1$ 's may be connected to form a simple closed curve  $C$  on  $S(G)$  with the desired properties. (See Figure 2.)




FIGURE 2

Now, let  $N(C, \sigma)$  be the surface generated by the normals of length  $\sigma$  to  $S(G)$  along  $C$ . The normals are taken in the component of the complement of  $S(G) \cup O$  which contains  $r$ . By (P Vb) and a compactness argument, for  $\sigma$  small enough  $N(C, \sigma)$  is a smooth (except on the boundaries) non-self-intersecting surface without contact bounded by  $C$  and the curve generated by the ends of the normals.

Let  $G'$  be a smooth simple closed curve on  $S^2$  enclosing  $G$  and cut by  $\gamma$ . There is a  $\tau_1$  such that for all  $\tau \geq \tau_1$   $\gamma(\tau)$  lies in the region on  $S^2$  bounded by  $G$  and

by  $G'$ . Let the maximal distance between  $G$  and  $G'$  be so small that the cone (not an integral surface),  $S(G')$  determined by  $G'$  and  $O$  is cut by  $N(C, \sigma)$  in a simple closed curve. For example,  $G'$  might be constructed by using arcs of fixed length (sufficiently small) of great circles of  $S^2$  normal to the  $\gamma_1$  and then joining the subarcs around the  $P_1$  in some smooth fashion.

Let  $R$  be the region bounded by  $S(G)$ ,  $S(G')$ , and  $O$ . Let  $R^*$  be the subregion of  $R$  bounded by  $S(G)$ ,  $S(G')$ ,  $O$  and  $N(C, \sigma)$ . For all

$$\tau \geq \tau_1, \Gamma : \left\{ r(\tau_0) \exp \int_{\tau_0}^{\tau} (y, f(y)) d\tau \right\} \cdot \gamma(\tau)$$

is in  $R$ . Furthermore, if  $\Gamma$  is in  $R^*$  for some  $\tau$ , it is in  $R^*$  for all greater  $\tau$ . For evidently  $\Gamma$  could not leave  $R^*$  via  $S(G')$ ,  $S(G)$  or  $O$ . It could not leave through  $N(C, \sigma)$  because by construction of  $N(C, \sigma)$  all trajectories cutting the interior of  $N(C, \sigma)$  enter  $R^*$ . Hence in this case using (P VI)  $\Omega(\Gamma)$  contains only  $O$ . Suppose it is not known that  $\Gamma$  intersects  $R^*$ . Let  $p$  be a point of  $\Gamma$  corresponding to some  $\tau_2$ ,  $\tau_2 \geq \tau_1$ . Then for some  $k$ ,  $0 < k < 1$ ,  $k \cdot p$  is in  $R^*$ . Then the trajectory  $k \cdot \Gamma$ :

$$\left\{ k \cdot r(\tau_0) \cdot \exp \int_{\tau_0}^{\tau} (y, f(y)) d\tau \right\} \cdot \gamma(\tau)$$



must lie in  $R^*$  for all  $\tau \geq \tau_2$  and  $n(k \cdot r)$  contains 0 alone. By (P III) this implies that  $n(r)$  consists of just 0 (see Figure 3).




FIGURE 3

F-I. The remaining combinations of possibilities for  $A(\gamma)$  and  $n(\gamma)$  are not discussed since in each case the argument is similar to the preceding.

Thus, in every case  $A(\gamma)$  and  $n(\gamma)$  have the desired properties and the theorem is proved.

### 3. EXAMPLE

Consider the system

$$\begin{aligned} \frac{dx_1}{dt} &= -\mu(x_1^3/3 - x_1x_3^2) + ax_1x_3^2 + bx_1x_2^2 + x_2x_3^2 + g_1(x, t) = f_1(x) + g_1(x, t) \\ (5) \quad \frac{dx_2}{dt} &= -x_1x_3^2 + ax_2x_3^2 + bx_2^3 + g_2(x, t) = f_2(x) + g_2(x, t) \\ \frac{dx_3}{dt} &= ax_3^3 + bx_2^2x_3 + g_3(x, t) = f_3(x) + g_3(x, t) \end{aligned}$$

where  $a < 0$ ,  $b < 0$ ,  $\mu > 0$ ,  $x = (x_1, x_2, x_3)$ . Suppose that  $g = (g_1, g_2, g_3)$  satisfies the hypotheses given in Section 1 for  $m = 3$ .

In  $r, y$  coordinates (5) become

$$(a) \quad \frac{dr}{d\tau} = r(y, f(y))$$

(6)

$$(b) \quad \frac{dy}{d\tau} = f(y) - (y, f(y))y, \quad$$

where  $f = (f_1, f_2, f_3)$  and  $r^2 dt = d\tau$ . It is not hard to verify that the only roots of  $f(y) - (y, f(y))y = 0$  are  $(\pm 1, 0, 0)$ ,  $(0, \pm 1, 0)$ ,  $(0, 0, \pm 1)$  and that  $(y, f(y)) < 0$  on each of these roots ( $a$  and  $b$  must be negative). Projecting the hemisphere  $y_3 > 0$  from the origin onto the plane  $y_3 = 1$ , we obtain

$$\frac{dz_1}{d\tau} = z_2 - \mu(z_1^3/3 - z_1)$$

(7)

$$\frac{dz_2}{d\tau} = -z_1, \quad$$

where

$$z_1 = y_1(1 - y_1^2 - y_2^2)^{-1/2}, \quad z_2 = y_2(1 - y_1^2 - y_2^2)^{-1/2}.$$

This is just Van der Pol's equation, and there is a unique periodic solution, for  $\mu$  small enough, which is near the circle:  $z_1^2 + z_2^2 = 4$ . Thus, there is a unique periodic solution to (6b) on the hemisphere  $y_3 > 0$  (and, of course, another on the hemisphere  $y_3 < 0$ ). This solution is near the circle  $y_3 = 5^{-1/2}$  on the two sphere. But then  $(y, f(y))$  is approximately  $-\mu y_1^2(y_1^2/3 - 1/5) + 1/5a + by_2^2$  which is negative for  $|a|$  large enough,  $a < 0$ . Hence,

$$\int_0^T (y, f(y)) d\tau < 0$$

where  $T$  is the period of the periodic solution. Similarly for the periodic solution on the hemisphere  $y_3 < 0$ .

The theorem then applies, and the solution  $x = 0$  of (5) is

asymptotically stable.

4. REMARKS

- A. The two conditions given in the theorem for asymptotic stability are usually easy to apply. The first is always easy, and the second condition can be verified without knowing either a precise expression for a periodic solution of (4b) or its exact period -- as was seen in the above example.
- B. The two conditions are in a certain sense necessary for asymptotic stability. That is, if either of the inequalities is reversed, the critical point of (2) at the origin is not even stable. It is evident that if one of the two inequalities is replaced by an equality, the trivial solution of (1) may or may not be asymptotically stable, depending on the form of  $g(x, t)$ .
- C. If  $m = 1$ , the linear case, the two conditions are equivalent to the condition that the characteristic roots of the matrix of coefficients of the linear terms have negative real parts. The proof of this fact is not difficult, but is a little long because of the necessity of considering the various possible canonical forms for the matrix. The proof will be omitted.
- D. Zubov [5] has given conditions for asymptotic stability for  $m = p/q$  ( $p, q$  integers,  $q$  odd) where  $n$  is arbitrary. His conditions do not seem easy to use inasmuch as one must have a priori bounds on the solutions  $x(t)$  of (1) for which  $\|x(0)\| = 1$ , namely  $\|x(t)\| < At^{-\alpha}$ ,  $A, \alpha$  positive constants.

RIAS, Baltimore

BIBLIOGRAPHY

- [1] Malkin, I. G., "A theorem on stability in the first approximation," (in Russian), Doklady Akademii Nauk SSSR, 76, (1951), pp. 783-784.
- [2] Massera, J. L., "Contributions to stability theory," Annals of Math., 64, (1956), pp. 182-205.
- [3] Coleman, C. S., "A certain class of integral curves in 3-space," Annals of Math., 69, May, 1959.

- [4] Forster, H., "Über das Verhalten der Integralkurven einer Gewöhnlichen Differentialgleichung erster Ordnung in der Umgebung eines singulären Punktes," *Mathematische Zeitschrift*, 43, (1937), pp. 271-320.
- [5] Zubov, V. I., "An investigation of the stability problem for a system of equations with homogeneous first terms," *Doklady Akademii Nauk* (in Russian), 114, (1957), pp. 942-944.

### XIII. EXISTENCE AND UNIQUENESS OF THE PERIODIC SOLUTION OF AN EQUATION FOR AUTONOMOUS OSCILLATIONS

Rui Pacheco de Figueiredo

#### I. PRELIMINARY REMARKS

Consider the differential equation

$$(1) \quad \ddot{x} + R(\dot{x}) + x = 0$$

where  $R(\dot{x})$  is a real-valued function having a piecewise continuous derivative\*\* and from now on the dot on a variable denotes differentiation with respect to  $t$ . Both (1) and its alternate form

$$(2) \quad \ddot{y} + r(y)\dot{y} + y = 0$$

where  $y = \dot{x}$  and  $r(y) = R'(y)$ , have been widely used to represent the behavior of physical systems undergoing self-sustained oscillatory motion.

In this paper we propose a general set of conditions for the existence and uniqueness of the periodic solution of (1).

---

\* The results of this paper are from a part of the author's doctorate thesis at Harvard University and were presented at the annual meeting of the American Mathematical Society in Philadelphia on January 20, 1959. It is a pleasure to thank Professor P. Le Corbeiller of Harvard for his very valuable advice throughout this research, and Dr. F. Huckemann, now at the Mathematical Institute of the University of Giessen, Giessen-Lahn, Germany, for his careful reading of, and very useful remarks on the original manuscript of this thesis. This work was supported in part by a grant from the Junta de Investigações do Ultramar, Portugal.

\*\* For such a function, it can be readily shown (cf. the author's thesis) that the two-dimensional system equivalent to (1):  $\dot{x} = y$ ,  $\dot{y} = -R(y) - x$  possesses locally a unique solution everywhere on the  $(x, y)$  plane. This and the continuity of the right-sides of this system assure (see reference [1]) that the Poincaré-Bendixon theorem is applicable to this system under the above assumption.

Noting that  $r(\dot{x})$  ( $= R'(\dot{x})$ ) represents the damping in the physical system, we may describe our existence conditions simply as follows:

(1) has at least one periodic solution if, for small  $|\dot{x}|$ , the damping in the system is negative and provided that, for large  $|\dot{x}|$ , the damping is positive for positive  $\dot{x}$  and greater than a negative minimum for negative  $\dot{x}$  (or vice-versa). We further show that the periodic solution is unique (except for translation in  $t$ ) if  $R(\dot{x})$  satisfies some additional restrictions.

Among the previous contributions on this subject, notable are those of Liénard [2] and Levinson and Smith\* [3]. In the present conditions, we remove the restriction of these authors that, for large  $\dot{x}$ , the damping in the system be positive for both positive and negative  $\dot{x}$ , and in particular we do not require  $R(\dot{x})$  to be odd.

It is convenient to replace (1) by the equivalent system

$$(3a, b) \quad \dot{x} = y, \quad \dot{y} = -R(y) - x$$

which is a special case of the two-dimensional autonomous system

$$(4a, b) \quad \dot{x} = X(x, y), \quad \dot{y} = Y(x, y)$$

where  $X$  and  $Y$  are real continuous functions defined on the real  $(x, y)$  plane and such that (4) is assured of possessing locally a unique solution everywhere on this plane. Some of the well known definitions and results of the Poincaré-Bendixson theory [1] relative to (4) will be now briefly recalled, the terms defined being assumed to refer to (4) unless otherwise stated.

A point of the  $(x, y)$  plane is said to be critical if both the right sides of (4) vanish there and it is said to be regular otherwise. If  $x(t)$  and  $y(t)$  constitute a solution of (4) then for a given  $t$  they define the  $x$ - and  $y$ -coordinates of a representative point  $P(t)$  on the  $(x, y)$  plane for this solution. The set  $\{x(t), y(t)\}$  for all real  $t$ , that is the curve on the  $(x, y)$  plane which is the locus of all the representative points for this solution, is called an orbit. Uniqueness of the solutions of (4) implies that no two orbits can intersect. A closed orbit (that is an orbit which is represented by a closed curve in the  $(x, y)$  plane) made of regular points only is periodic (i.e., it is such that only a finite change in  $t$  is required for a representative point to describe a complete circuit on it), and vice-versa. In particular a

---

\* Levinson and Smith actually considered the more general equation  $\ddot{y} + r(y, \dot{y})\dot{y} + g(y) = 0$ . Their results may be readily particularized to (2).

periodic orbit is said to be a stable (unstable) limit cycle if it is approached asymptotically by its neighboring orbits as  $t \rightarrow \infty$  ( $-\infty$ ). If these remarks are particularized to the system (3), it is clear that, for the purpose of this paper, it is sufficient to determine the conditions for the existence and uniqueness of a periodic orbit of (3).

In order to establish the existence conditions we shall need the conclusions of the following section and also the following general result [4].

LEMMA 1 (Poincaré-Bendixson). If an orbit of (4) remains for  $t \rightarrow \infty$  in a finite region  $K$  of the  $(x, y)$  plane and if  $K$  is free of critical points of (4), then either this orbit is periodic or it approaches asymptotically a limit cycle as  $t \rightarrow \infty$ .

## 2. AN AUXILIARY AUTONOMOUS SYSTEM OF TWO FOCI OR OF A FOCUS AND A NODE

Consider the piecewise linear autonomous system

$$(5a, b) \quad \dot{x} = y, \quad \dot{y} = -R_1(y) - x$$

where

$$(6a) \quad R_1(y) = \begin{cases} -\rho_1 y & (y \leq 0) \end{cases}$$

$$(6b) \quad \begin{cases} \rho_2 y - \gamma & (y > 0) \end{cases}.$$

Here,  $\rho_1, \rho_2$  and  $\gamma$  are positive constants and  $2 > \rho_1 < \rho_2$ .

The patterns of the orbits of this system for  $\rho_2 < 2$ ,  $\rho_2 = 2$  and  $\rho_2 > 2$  are illustrated respectively in Figures 1(a), (b) and (c), where we make use of the well-known results for the  $(x, y)$  plane solutions of a linear second order differential equation in  $t$  with constant coefficients.

Since  $0 < \rho_1 < 2$ , the origin  $O$  of the  $(x, y)$  plane is an unstable focus for the parts of orbits of (5) lying in the half-plane  $y \leq 0$ . Hence the arc of an orbit of (5) beginning at an arbitrary point  $A$  on the positive  $x$ -axis, when followed in the clockwise (increasing  $t$ ) direction, although getting farther from the origin, must necessarily turn and meet the negative  $x$ -axis at some point  $B$ . Then from the well-known results alluded in the preceding paragraph, the ratio of the distances  $OB$  and  $OA$  is





where

$$(10) \quad \omega_2 = \sqrt{1 - \frac{1}{4} \rho_2^2}$$

From (7) and (9) there is

$$\frac{OC}{OA} = e^{-\frac{\pi}{2} \left( \frac{\rho_2}{\omega_2} - \frac{\rho_1}{\omega_1} \right)} + \gamma \left( e^{-\frac{\pi}{2} \frac{\rho_2}{\omega_2}} + 1 \right) e^{\frac{\pi}{2} \frac{\rho_1}{\omega_1}} \frac{1}{OA}$$

which shows that as OA increases from 0 to  $\infty$ , the ratio OC/OA decreases monotonically from  $\infty$  to the value

$$e^{-\frac{\pi}{2} \left( \frac{\rho_2}{\omega_2} - \frac{\rho_1}{\omega_1} \right)}$$

which is positive and less than unity. Hence there is one and only one position of A, say A', at which A and C coincide, this occurring when the ratio (11) is unity, that is, when

$$(12) \quad OA = OA' = \frac{\gamma \left( e^{-\frac{\pi}{2} \frac{\rho_2}{\omega_2}} + 1 \right)}{e^{-\frac{\pi}{2} \frac{\rho_1}{\omega_1}} - e^{-\frac{\pi}{2} \frac{\rho_2}{\omega_2}}}$$

In virtue of (11), it is clear that the resulting closed orbit A'BA' is a limit cycle of (5) to which all the other orbits of the system tend as  $t \rightarrow \infty$ .

We next assume that  $\rho_2 \geq 2$  (see Figures 1(b) and (c)). Then, whatever the position of the point B on the negative x-axis, the orbit through AB always tends to F when followed clockwise from B, since now F acts as a stable node for the arc BF. Thus as A varies its position on the positive x-axis from 0 to  $\infty$ , there is only one position of A corresponding to a closed orbit of the system, namely when A coincides with F. All other orbits of the system tend to F (and hence approach the closed orbit FBF) as  $t \rightarrow \infty$ .

The above conclusions may be summarized as follows:

LEMMA 2. Under the conditions stated, the system (5) has in the  $(x, y)$  plane one and only one closed

orbit which is approached by all the other orbits of the system as  $t \rightarrow \infty$ .

### 3. ON THE EXISTENCE OF A PERIODIC SOLUTION

We now state the following result for the original equation (1):

**THEOREM 1.** The equation (1) has at least one periodic solution of

$$(a) \quad R(0) = 0$$

$$(b) \quad R'(0) \text{ exists and is negative}$$

and provided there is a  $y_0 > 0$  such that

$$(c) \quad R'(\dot{x}) > 0, \min(\dot{x} \geq y_0)$$

$$(d) \quad 2 > -\min R'(\dot{x}) < R'(-\dot{x}), (\dot{x} \leq -y_0)$$

except for the values of  $\dot{x}$  at which  $R'(\dot{x})$  undergoes simple discontinuities.

**REMARK.** Below it will be necessary to compare the slopes of the elements of orbits, through a point  $P$  in the  $(x, y)$  plane, of two systems  $\Sigma_1$  and  $\Sigma_2$  of the form

$$(13) \quad \Sigma_1 : \dot{x} = y, \quad \dot{y} = -F_1(y) - x$$

$$(14) \quad \Sigma_2 : \dot{x} = y, \quad \dot{y} = -F_2(y) - x$$

where  $F_1$  and  $F_2$  are real, continuous (piecewise differentiable) functions. For this purpose we let  $z$  denote the distance from the origin of the  $(x, y)$  plane to a point on an orbit of  $\Sigma_1$  and  $\Sigma_2$ , i.e.,

$$z = \sqrt{x^2 + y^2},$$

which after differentiation with respect to  $t$  and combination with (13) or (14) gives

$$(15) \quad \dot{z} = -\frac{1}{z} F_1(y)y, \quad i = 1, 2$$

This shows that at  $P$ , if  $y > 0$  and  $F_1 < F_2$ ,  $\dot{z}$  is larger along the element of orbit of  $\Sigma_1$  than along that of  $\Sigma_2$ , in other words, when taken in the direction of increasing  $t$ , the first of these elements intersects the second one outward or away from the origin. The same is

the case if  $y < 0$  and  $F_1 > F_2$ , the opposite being true if  $y > 0$  and  $F_1 > F_2$  or  $y < 0$  and  $F_1 < F_2$ .

PROOF. Consider the two-dimensional system (3) equivalent to (1).

Since  $R(y)$  is continuous on  $-y_0 \leq y \leq y_0$ ,  $|R(y)|$  is bounded by a positive constant  $M$  there.

It will be convenient to introduce the variable

$$(16) \quad u = x + M.$$

The conditions (c) and (d) of the theorem allow us to define the positive constants  $\rho_2$  and  $\rho_1$  satisfying

$$(17) \quad \rho_2 = \min R'(y), \quad (y \geq y_0)$$

$$(18) \quad 2 > \rho_1 < \rho_2$$

$$(19) \quad \rho_1 \geq -\min R'(y), \quad (y \leq -y_0).$$

Finally, let

$$(20) \quad \gamma = \rho_2 y_0 + 2M.$$

$\rho_1$ ,  $\rho_2$  and  $\gamma$  thus defined satisfy the conditions assumed for these constants in (6) and hence by Lemma 2, the system

$$(21) \quad \dot{u} = y, \quad \dot{y} = -R_1(y) - u$$

$$(22a) \quad R_1(y) = \begin{cases} -\rho_1 y, & (y \leq 0) \\ \rho_2 y - \gamma, & (y > 0) \end{cases}$$

$$(22b)$$

has a unique orbit in the  $(u, y)$  plane and therefore in the  $(x, y)$  plane. We label this orbit  $\Gamma_2$ .

Now consider the annular region  $K$  (see Figure 2) in the  $(x, y)$  plane bounded outside by  $\Gamma_2$  and inside by the circle  $\Gamma_1$  defined by

$$(23) \quad x^2 + y^2 = \delta$$

where  $\delta$  is a sufficiently small positive constant such that  $R'(y) < 0$  in the interior of and on  $\Gamma_1$ .




FIGURE 2

Let (21) and (22), after combination with (16) and (20), be rewritten as

$$(24a, b) \quad \dot{x} = y, \quad \dot{y} = -F(y) - x$$

$$(25a) \quad F(y) = \begin{cases} -\rho_1 y + M, & (y \leq 0) \end{cases}$$

$$(25b) \quad F(y) = \begin{cases} \rho_2(y - y_0) - M, & (y > 0) \end{cases}.$$

Then since  $|R(y)| \leq M$  for  $0 \leq |y| \leq y_0$  and according to (17) and (18) we have

$$(26a) \quad \text{For } y < 0: R(y) \leq -\rho_1 y + M = F(y)$$

$$(26b) \quad \text{For } y > 0: R(y) \geq \rho_2(y - y_0) - M = F(y)$$

where the rightmost terms in (26a) and (26b) arise from the definitions (25a) and (25b). It follows, from (26a) and (26b) (when considered together with (3) and (24)) and by virtue of the remark just preceding this proof, that the elements of orbits of (3) on  $\Gamma_2$ , taken in the direction of increasing  $t$ , remain in  $K$ .

Also, according to the conditions (a) and (b) of the theorem,

$R(y) < 0$  for  $y > 0$  and  $R(y) > 0$  for  $y < 0$ . Moreover (23) is a solution of the system  $\dot{x} = y, \dot{y} = -x$ . Hence, by the remark just referred to, the elements of orbits of (3) on  $r_1$ , when taken in the direction of increasing  $t$ , intersect  $r_1$  outward from the origin and hence into  $K$ .

Thus an arc of orbit of (3) starting at a point on  $K$  cannot leave this region for increasing  $t$ . Moreover, the origin of the  $(x, y)$  plane, which is the only critical point of (3), is external to  $K$ . Hence by Lemma 1,  $K$  contains at least one periodic orbit of (3). This concludes the proof.

#### 4. UNIQUENESS OF THE PERIODIC SOLUTION

We now make the following proposition:

**THEOREM 2.** Assume that the conditions of Theorem 1 are satisfied for the equation (1) and let there exist a  $y_1 > 0$  such that\*

- (a)  $R(y_1) = R(0) = 0$
- (b)  $\dot{x} R(\dot{x}) < 0, \quad (0 < |\dot{x}| < y_1)$
- (c)  $R(\dot{x}) > 0, \quad (\dot{x} > y_1)$
- (d)  $R'(\dot{x}) \geq \frac{1}{\dot{x}} R(\dot{x}), \quad (\dot{x} < 0, \dot{x} > y_1)$

except at the values of  $\dot{x}$  where  $R'(\dot{x})$  undergoes simple discontinuities. Then (3) has a periodic solution which is unique except for the translations in  $t$ .

**REMARKS.** As before, instead of (1), the equivalent system (3) will be considered.

Figure 3 illustrates the direction, for increasing  $t$ , of the elements of orbits of (3) under the conditions of this theorem. In this and the following figures,  $-R(y)$  is also shown plotted along the axis of the abscissae. It follows from (3) that, for increasing  $t$ , the elements of orbits of (3) above the  $x$ -axis are directed rightward and those below it leftward; the ones at the left of  $-R(y)$  are directed upward and those at its right, downward. The curve  $-R(y)$  itself is intersected

\* By evident changes in the proof it can be shown that the result of this theorem holds if the conditions (a), (c) and (d) are replaced respectively by (a)  $R(-y_1) = R(0) = 0$ , (c)  $R(\dot{x}) > 0, (\dot{x} < -y_1)$ , (d)  $R'(\dot{x}) \geq \frac{1}{\dot{x}} R(\dot{x}), (\dot{x} < -y_1, \dot{x} > 0)$ .




FIGURE 3




FIGURE 4

by every orbit horizontally and the  $x$ -axis is crossed vertically.

Let  $\Gamma$  in Figure 4 represent an arc of orbit of (3) that describes a complete clockwise turn around the origin, leaving the positive  $x$ -axis at  $A$  and meeting this axis again for the first time at some point  $I$ . Then nowhere on  $\Gamma$  is an element of  $\Gamma$  along a radial direction. In fact, this follows directly from the discussion in the preceding paragraph, for that part of  $\Gamma$  that does not lie in the region between the curve  $-R(y)$  and the  $y$ -axis (indicated by the hatched area in the Figure). To show that the same holds for the arcs  $FG$  and  $BC$  of  $\Gamma$  that lie in this region, consider first  $FG$ . In the  $y$ -interval on which  $FG$  is defined there is a  $N > 0$  such that  $R(y)/y \leq N$ . Since an orbit through  $F$  of the linear system  $\dot{x} = y$ ,  $\dot{y} = -Ny - x$ , when followed clockwise from  $F$ , does not approach a radial direction in this  $y$ -interval, the arc  $FG$  cannot certainly approach a radial direction (since  $R(y) \leq Ny$ ). Similar argument holds for  $BC$ , but in this case one follows  $BC$  counterclockwise from  $C$  to  $B$ .

If  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$  are two points on an orbit of (3), we shall often need to relate their distances  $OP$  and  $OQ$  from the origin of the  $(x, y)$  plane. For this purpose (3) is reduced to a single first-order equation in  $x$  and  $y$  and then integrated to give

$$(27) \quad OQ^2 - OP^2 = (x_Q^2 + y_Q^2) - (x_P^2 + y_P^2) = -2 \int_P^Q R(y) dx$$

where the rightmost term is a line integral computed on the orbit from  $P$  to  $Q$ .

PROOF. Since the conditions of Theorem 1 are satisfied, (3) has at least one periodic orbit. It remains to be proved that this periodic orbit is unique under the present set of restrictions.

Condition (b) of the theorem clearly implies\* that the periodic orbit must be partly above and below the band  $|y| \leq y_1$ .

Consider the arc of an orbit of (3)  $ACDFI$  (see Figure 5) which




FIGURE 5

starts at a point  $A$  on the positive  $x$ -axis and continues clockwise from  $A$  intersecting: the curve  $-R(y)$  for  $y < 0$  at  $C$ , the negative  $x$ -axis at  $D$ , the curve  $-R(y)$  for  $y > 0$  at  $F$ , and again the positive  $x$ -axis at  $I$ . Clearly,  $C$  and  $F$  are respectively the minimum and maximum of  $y$  for this arc.

If  $C$  and  $F$  (the  $y$ -coordinates of which are denoted by  $y_C$

\* This follows from application of (27) to the arcs of orbits of (3) lying in the band  $|y| \leq y_1$ , taking into account the condition (b) of the theorem.

and  $y_F$ ) are inside or at the edges of the band  $|y| \leq y_1$ ,  $A$  is certainly at the left\* of  $I$ . Let  $A_0C_0D_0F_0I_0$  be the arc  $ACDFI$  such that  $\min\{-y_{C_0}, y_{F_0}\} = y_1$ . The proof of the theorem now proceeds via the following lemma.

LEMMA 3. As the position of the point  $A$  moves rightward on the positive  $x$ -axis from  $A_0$ , the ratio of the distances  $OI/OA$  decreases monotonically.

PROOF OF LEMMA 3. Suppose  $A_1$  and  $A_2$  are two points on the positive  $x$ -axis such that  $A_2$  is to the right of  $A_1$  and  $A_1$  is to the right of (or coincides with)  $A_0$ . The corresponding arcs of orbits  $A_1C_1D_1F_1I_1$  and  $A_2C_2D_2F_2I_2$  will be denoted by  $\Gamma_1$  and  $\Gamma_2$  and the quantities associated with these arcs will be indicated respectively by the subscripts 1 and 2. The lemma is proved by showing that

$$(28) \quad \frac{OI_2}{OA_2} \leq \frac{OI_1}{OA_1}.$$

Let the left- and right-side intercepts of  $y = y_1$  with  $\Gamma_1$  be denoted respectively by  $E_1$  and  $H_1$ , and those with  $\Gamma_2$  respectively by  $E_2$  and  $H_2$ . Assume that  $OE_1$  and  $OH_1$  when produced in the directions from  $O$  to  $E_1$  and  $H_1$  intersect  $\Gamma_2$  respectively at  $E_3$  and  $H_3$ .

Now the change in the radial distance of a representative point moving clockwise on  $\Gamma_1$  will be compared to that on  $\Gamma_2$ . For this purpose, each of the parts into which  $\Gamma_1$  is divided by the points  $D_1$ ,  $E_1$  and  $H_1$  will be considered together with the corresponding part on  $\Gamma_2$ .

Consider first the arcs  $A_1C_1D_1$  and  $A_2C_2D_2$ .

Introducing the polar coordinates  $z$  (already defined) and  $\theta$  (which is the angle measured in the clockwise direction from the positive  $x$ -axis), we write

$$(29a) \quad x = z \cos \theta$$

$$(29b) \quad y = -z \sin \theta$$

with which (3) may be expressed as the single first order equation

$$(30) \quad \frac{dz}{d\theta} = \frac{z R \sin \theta}{z + R \cos \theta}$$

---

\* See footnote on preceding page.



where  $R = R(-z \sin \theta)$ .

(30) may be conveniently rewritten in the form

$$(31) \quad \frac{d}{d\theta} (\log z) = \frac{R \sin \theta}{z + R \cos \theta}$$

Consider now two differential elements  $ds_1$  and  $ds_2$  belonging respectively to  $A_1C_1D_1$  and  $A_2C_2D_2$  and lying along the same radial direction, as indicated in Figure 5. For both these elements the terms in  $\theta$  in the right side of (31) are the same.

Partial differentiation of (31) with respect to  $z$  gives

$$(32) \quad \frac{\partial}{\partial z} \frac{d}{d\theta} (\log z) = \frac{\left( \frac{\partial R}{\partial z} z - R \right) \sin \theta}{(z + R \cos \theta)^2}$$

The denominator in the right side of (32) does not vanish at any point on  $A_1C_1D_1$  and  $A_2C_2D_2$ . For if it did, then (30) would imply that the element of  $A_1C_1D_1$  or  $A_2C_2D_2$ , there, is along a radial direction, and, as remarked earlier, this is not possible.

Also, for the elements  $ds_1$  and  $ds_2$

$$(33) \quad \sin \theta > 0$$

and, according to (29b),

$$(34) \quad \frac{\partial R}{\partial z} z - R = R'(y)y - R \leq 0$$

the right-most member of (34) resulting from the condition (d) of the theorem.\*

(33) and (34) together with (32) imply that

$$(35) \quad \frac{\partial}{\partial z} \left[ \frac{d}{d\theta} (\log z) \right] \leq 0$$

and since  $ds_2$  is at a greater radial distance than  $ds_1$ , (35) shows that the right side of (31) is less (or equal) for  $ds_2$  than (and) for  $ds_1$ , or symbolically:

---

\* Note that since  $y < 0$ , multiplication by  $y$  changes the sign of the inequality given in the condition (d) of the theorem.

$$(36) \quad \left\{ \frac{R \sin \theta}{z + R \cos \theta} \right\}_{ds_2} \leq \left\{ \frac{R \sin \theta}{z + R \cos \theta} \right\}_{ds_1}$$

Integrating (31) along  $A_1C_1D_1$  and  $A_2C_2D_2$ , making, in this integration, the differential elements of these arcs along the same radial direction correspond, and using (36), we obtain

$$(37) \quad \int_{OA_2}^{OD_2} d \log z = \int_{A_2C_2D_2}^{\pi} \frac{R \sin \theta}{z + R \cos \theta} d\theta \leq \int_{A_1C_1D_1}^{\pi} \frac{R \sin \theta}{z + R \cos \theta} d\theta = \\ = \int_{OA_1}^{OD_1} d \log z$$

and hence

$$(38) \quad \frac{OD_2}{OA_2} \leq \frac{OD_1}{OA_1}$$

Next we consider the arcs  $D_1E_1$  and  $D_2E_2$  using a modification of Liénard's [2] approach.

Now  $y$  is taken as the independent variable (in the line integration to be performed) and a differential element  $ds_1$  on  $D_1E_1$  is made to correspond to the element  $ds_2$  of  $D_2E_2$  having the same  $y$ -coordinate, as shown in the figure.

Let  $(x', y')$  and  $(x'', y'')$  denote the coordinates of  $ds_1$  and  $ds_2$ . Since  $x'' < x' < 0$  (see Figure 5)),  $y' = y'' = y > 0$ , and according to the condition (b) of the theorem,  $R(y') = R(y'') = R(y) < 0$ , we deduce from (3) (by reducing (3) first to a single order equation in  $x$  and  $y$  and expressing  $dx$  in terms of the other variables):

$$(39) \quad dx' > dx'' \geq 0$$

and

$$(40) \quad R(y) dx' \leq R(y) dx'' \leq 0.$$

Then integrating (40) along  $D_1E_1$  and  $D_2E_2$  in the way indicated in (27), we have:

$$\begin{aligned}
 (41) \quad OE_1^2 - OD_1^2 &= -2 \int_{D_1 E_1} R(y) dx' \geq -2 \int_{D_2 E_2} R(y) dx'' = \\
 &= OE_2^2 - OD_2^2 \geq 0.
 \end{aligned}$$

or simply

$$(42) \quad OE_1^2 - OD_1^2 \geq OE_2^2 - OD_2^2 \geq 0$$

which may be rearranged as

$$(43) \quad OD_1^2 \left( \frac{OE_1^2}{OD_1^2} - 1 \right) \geq OD_2^2 \left( \frac{OE_2^2}{OD_2^2} - 1 \right).$$

Since, according to (42), both sides of (43) are non-negative and because  $OD_2 \geq OD_1$  we derive from (43)

$$(44) \quad \frac{OE_1^2}{OD_1^2} - 1 \geq \frac{OD_2^2}{OD_1^2} \left( \frac{OE_2^2}{OD_2^2} - 1 \right) \geq \frac{OE_2^2}{OD_2^2} - 1$$

and therefore

$$(45) \quad \frac{OE_2}{OD_2} \leq \frac{OE_1}{OD_1}.$$

For the arcs  $E_2 E_3$  and  $H_3 H_2$  of  $r_2$ ,  $R(y) > 0$  (since  $y > y_1$ ). Hence applying (27), separately, to each of these arcs, we get

$$(46) \quad \frac{OE_3}{OE_2} < 1$$

$$(47) \quad \frac{OH_2}{OH_3} < 1.$$

The arcs  $E_1 F_1 H_1$  and  $E_3 F_2 H_3$  are treated in the same way as  $A_1 C_1 D_1$  and  $A_2 C_2 D_2$ . For those arcs (33) is negative and (34) is positive. Hence, the right side of (32) is again negative and therefore, as in (38), there is

$$(48) \quad \frac{OH_3}{OE_3} \leq \frac{OH_1}{OE_1} \quad .$$

Finally, dealing with the arcs  $H_1I_1$  and  $H_2I_2$  in much the same way as with  $D_1E_1$  and  $D_2E_2$ , we deduce

$$(49) \quad \frac{OI_2}{OH_2} \leq \frac{OI_1}{OH_1} \quad .$$

Multiplying together separately the left and right sides of (38), (45), (46), (47), (48) and (49), the result (28) is established.

CONCLUSION OF THE PROOF OF THEOREM 2. It follows from the above lemma that if the points  $A$  and  $I$  coincide, they can do so only once. Since (3) has at least one periodic orbit (since it is assumed to satisfy the conditions of Theorem 1),  $A$  and  $I$  must coincide at least once. In other words, (3) possesses one and only one periodic orbit and this orbit corresponds to the periodic solution of (1). This completes the proof of the theorem.

#### BIBLIOGRAPHY

- [1] Coddington, E. A., and Levinson, N., "Theory of Ordinary Differential Equations," McGraw-Hill, (1955), p. 389.
- [2] Liénard, A., "Étude des Oscillations Entretenues," Rev. Gen. d'Électricité, 23, (1928), pp. 901-912.
- [3] Levinson, N., and Smith, O. K., "A General Equation for Relaxation Oscillations," Duke Math. Jnal., 9, (1941), pp. 382-403.
- [4] Bendixson, I., "Sur les Courbes Définies par des Équations Différentielles," Acta Mathematica, 24, (1901), pp. 1-88.

# APPENDIX

E. Pinney

Errata to "Nonlinear Differential Equations Systems", Contributions to the Theory of Nonlinear Oscillations, vol. III, pp. 31-56. Princeton University Press, 1956.

I. In formulas (2.7), (3.3), (5.1), (8.4), (9.13), (10.1), (10.7), (10.8) replace

$$\frac{\mu_j^1}{D(\mu_j^1)(z_j)} G_1^{(q)}(z_j) \quad \text{by} \quad G_{1j}^{(q)}(z_j) \quad .$$

II. In (2.7) replace the upper limit of the second summation sign by  $\mu_j - 1$ .

III. Replace (2.9) by

$$(2.9) \quad G_{1j}(z) = - (z - z_j)^{\mu_j} D_1(z) / [D(z) \bar{D}(z)]$$

IV. Replace (2.11) and the material above (2.12) by

$$(2.11) \quad y_1(t) = - \frac{1}{2\pi i} \int_C \sum_{\alpha=1}^m \frac{D_{1\alpha}(z)}{D(z)} [e^{zt} v_\alpha + \int_0^t e^{z(t-\tau)} u_\alpha(\tau) d\tau] dz \quad ,$$

where  $C$  is a contour enclosing the zeros of  $D(z)$  but not those of  $\bar{D}(z)$ . Then

$$\begin{aligned} \frac{dy_1}{dt} - \sum_{\beta=1}^m A_{1\beta} y_\beta &= \frac{1}{2\pi i} \int_C \sum_{\alpha=1}^m \sum_{\beta=1}^m \frac{D_{\beta\alpha}(z)}{D(z)} (A_{1\beta} - \delta_{1\beta} z) [e^{zt} v_\alpha \\ &+ \int_0^t e^{z(t-\tau)} u_\alpha(\tau) d\tau] dz - \frac{1}{2\pi i} \sum_{\alpha=1}^m u_\alpha(t) \int_C [D_{1\alpha}(z)/D(z)] dz \quad . \end{aligned}$$

By (2.4) the first term on the right vanishes because its integrand has no poles inside  $C$ . Therefore

V. Replace equation (2.13) by

$$(2.13) \quad S_{1\alpha} = -\frac{1}{2\pi i} \int_C [D_{1\alpha}(z)/D(z)] dz .$$

VI. Replace the material below equation (2.14) and above equation (2.16) by, "To evaluate  $S_{1\alpha}$  note that as  $z \rightarrow \infty$ ,"

VII. Change formula number (2.16) to (2.15).

VIII. Line 11, p. 36 should read, "degree  $m - 1$  or  $\leq m - 2$  according as ... ."

IX. Replace "inserting (2.16) into (2.15)," by "inserting (2.15) into (2.13),"

X. Omit lines 15-18, p. 36.

XI. Replace line 19, p. 36 by "Solving (2.5) for  $D_{1\alpha}(z)$  and inserting in (2.11),"

XII. Replace lines 4 and 5, p. 37 by

$$y_1(t) = -\frac{1}{2\pi i} \int_C \frac{D_1(z)}{D(z)\overline{D}(z)} [e^{zt}B(z) + \int_0^t e^{z(t-\tau)}U(z,\tau)d\tau] dz .$$

The integrand has poles at the characteristic roots only, so by the theory of residues and (2.9),

$$y_1(t) = \sum_{S_j} \frac{1}{(\mu_1 - 1)!} \frac{\partial^{\mu_j - 1}}{\partial z_j^{\mu_j - 1}} \left\{ G_{1j}(z_j) [e^{z_j t} B(z_j) + \int_0^t e^{z_j(t-\tau)} U(z_j, \tau) d\tau] \right\} .$$

















# PRINCETON MATHEMATICAL SERIES

*Edited by Marston Morse and A. W. Tucker*

|                                                                                             |                 |
|---------------------------------------------------------------------------------------------|-----------------|
| 1. The Classical Groups, Their Invariants and Representations<br>By HERMANN WEYL            | 314 pp. \$6.00  |
| 3. An Introduction to Differential Geometry<br>By LUTHER PFAHLER EISENHART                  | 316 pp. \$6.00  |
| 4. Dimension Theory<br>By WITOLD HUREWICZ and HENRY WALLMAN                                 | 174 pp. \$4.50  |
| 5. Analytical Foundations of Celestial Mechanics<br>By AUREL WINTNER                        | 460 pp. \$7.50  |
| 6. The Laplace Transform<br>By DAVID VERNON WIDDER                                          | 416 pp. \$7.50  |
| 7. Integration<br>By EDWARD J. McSHANE                                                      | 400 pp. \$6.00  |
| 8. Theory of Lie Groups: I<br>By CLAUDE CHEVALLEY                                           | 229 pp. \$5.00  |
| 9. Mathematical Methods of Statistics<br>By HARALD CRAMER                                   | 570 pp. \$7.50  |
| 10. Several Complex Variables<br>By S. BOCHNER and W. T. MARTIN                             | 216 pp. \$4.50  |
| 11. Introduction to Topology<br>By S. LEFSCHETZ                                             | 226 pp. \$5.00  |
| 12. Algebraic Geometry and Topology<br>Edited by R. H. FOX, D. C. SPENCER, and A. W. TUCKER | 408 pp. \$7.50  |
| 14. The Topology of Fibre Bundles<br>By NORMAN STEENROD                                     | 232 pp. \$5.00  |
| 15. Foundations of Algebraic Topology<br>By SAMUEL EILENBERG and NORMAN STEENROD            | 342 pp. \$7.50  |
| 16. Functionals of Finite Riemann Surfaces<br>By MENAHEM SCHIFFER and DONALD C. SPENCER     | 464 pp. \$8.00  |
| 17. Introduction to Mathematical Logic, Vol. I<br>By ALONZO CHURCH                          | 384 pp. \$7.50  |
| 19. Homological Algebra<br>By H. CARTAN and S. EILENBERG                                    | 408 pp. \$7.50  |
| 20. The Convolution Transform<br>By I. I. HIRSCHMAN and D. V. WIDDER                        | 280 pp. \$5.50  |
| 21. Geometric Integration Theory<br>By HASSLER WHITNEY                                      | 396 pp. \$8.50  |
| 22. Qualitative Theory of Differential Equations<br>By V. V. NEMICKII and V. V. STEPANOV    | 530 pp. \$12.50 |
| 23. Topological Analysis, REVISED EDITION<br>By GORDON T. WHYBURN                           | 132 pp. \$5.00  |
| 24. Analytic Functions<br>By AHLFORS, BEHNKE and GRAUERT, BERS, <i>et al.</i>               | 204 pp. \$5.00  |
| 25. Continuous Geometry<br>By JOHN VON NEUMANN                                              | 296 pp. \$7.50  |
| 26. Riemann Surfaces<br>By L. AHLFORS and L. SARIO                                          | 393 pp. \$10.00 |

---

PRINCETON UNIVERSITY PRESS  
PRINCETON, NEW JERSEY

---

# ANNALS OF MATHEMATICS STUDIES

*Edited by Robert C. Gunning, John G. Moore, and Marston Morse*

- |                                                                               |         |        |
|-------------------------------------------------------------------------------|---------|--------|
| 1. Algebraic Theory of Numbers                                                |         |        |
| By HERMANN WEYL                                                               | 240 pp. | \$3.00 |
| 3. Consistency of the Continuum Hypothesis                                    |         |        |
| By KURT GÖDEL                                                                 | 75 pp.  | \$3.00 |
| 11. Introduction to Nonlinear Mechanics                                       |         |        |
| By N. KRYLOFF and N. BOCOLIUBOFF                                              | 108 pp. | \$2.95 |
| 16. Transcendental Numbers                                                    |         |        |
| By C. L. SIEGEL                                                               | 110 pp. | \$2.00 |
| 19. Fourier Transforms                                                        |         |        |
| By S. BOCHNER and K. CHANDRASEKHARAN                                          | 223 pp. | \$3.50 |
| 20. Contributions to the Theory of Nonlinear Oscillations, Vol. I             |         |        |
| Edited by S. LEFSCHETZ                                                        | 360 pp. | \$4.50 |
| 21. Functional Operators, Vol. I                                              |         |        |
| By JOHN VON NEUMANN                                                           | 271 pp. | \$5.50 |
| 24. Contributions to the Theory of Games, Vol. I                              |         |        |
| Edited by H. W. KUHN and A. W. TUCKER                                         | 217 pp. | \$3.00 |
| 25. Contributions to Fourier Analysis                                         |         |        |
| Edited by A. ZYGMUND, W. TRANSUE, M. MORSE, A. P. CALDERON,<br>and S. BOCHNER | 192 pp. | \$3.00 |
| 28. Contributions to the Theory of Games, Vol. II                             |         |        |
| Edited by H. W. KUHN and A. W. TUCKER                                         | 404 pp. | \$4.00 |

*Continued on back cover*

# ANNALS OF MATHEMATICS STUDIES

*Continued from inside back cover*

|                                                                                                                        |         |        |
|------------------------------------------------------------------------------------------------------------------------|---------|--------|
| 38. Linear Inequalities and Related Systems<br>Edited by H. W. KUHN and A. W. TUCKER                                   | 352 pp. | \$5.00 |
| 39. Contributions to the Theory of Games, Vol. III<br>Edited by M. DRESHER, A. W. TUCKER and P. WOLFE                  | 448 pp. | \$5.00 |
| 40. Contributions to the Theory of Games, Vol. IV<br>Edited by R. DUNCAN LUCE and A. W. TUCKER                         | 468 pp. | \$6.00 |
| 41. Contributions to the Theory of Nonlinear Oscillations, Vol. IV<br>Edited by S. LEFSCHETZ                           | 224 pp. | \$3.75 |
| 42. Lectures on Fourier Integrals<br>By S. BOCHNER                                                                     | 342 pp. | \$5.00 |
| 43. Ramification Theoretic Methods in Algebraic Geometry<br>By S. ABHYANKAR                                            | 104 pp. | \$2.75 |
| 44. Stationary Processes and Prediction Theory<br>By H. FURSTENBERG                                                    | 300 pp. | \$5.00 |
| 45. Contributions to the Theory of Nonlinear Oscillations, Vol. V<br>Edited by L. CESARI, J. LASALLE, and S. LEFSCHETZ | 296 pp. | \$5.00 |
| 46. Seminar on Transformation Groups<br>By A. BOREL <i>et al.</i>                                                      | 250 pp. | \$4.50 |
| 47. Theory of Formal Systems<br>By R. SMULLYAN                                                                         | 160 pp. | \$3.00 |
| 48. Lectures on Modular Forms<br>By R. C. GUNNING                                                                      | 94 pp.  | \$2.75 |
| 49. Composition Methods in Homotopy Groups of Spheres<br>By H. TODA                                                    | 201 pp. | \$4.50 |
| 50. Cohomology Operations<br>Lectures by N. E. STEENROD, written and revised by D. B. A. EPSTEIN.                      | 137 pp. | \$3.00 |
| 51. Morse Theory<br>By J. W. MILNOR                                                                                    | 160 pp. | \$3.00 |
| 52. Advances in Game Theory<br>Edited by M. DRESHER, L. SHAPLEY, and A. W. TUCKER                                      | 679 pp. | \$8.50 |
| 53. Flows on Homogeneous Spaces<br>By L. AUSLANDER, L. GREEN, F. HAHN, <i>et al.</i>                                   | 120 pp. | \$2.75 |
| 54. Elementary Differential Topology<br>By J. R. MUNKRES                                                               | 120 pp. | \$2.75 |
| 55. Degrees of Unsolvability<br>By G. E. SACKS                                                                         | 176 pp. | \$3.50 |

PRINCETON UNIVERSITY PRESS  
PRINCETON, NEW JERSEY