

UNIVERSAL
LIBRARY

OU_172119

UNIVERSAL
LIBRARY

A MANUAL OF DISEASES OF THE EYE

BY

CHARLES H. MAY, M.D.

CONSULTING OPHTHALMOLOGIST TO THE MT. SINAI HOSPITAL, TO THE
FRENCH HOSPITAL, NEW YORK, AND TO THE
MONMOUTH MEMORIAL HOSPITAL

AND


CLAUD WORTH, F.R.C.S. ENG.

CONSULTING SURGEON TO THE ROYAL LONDON OPHTHALMIC HOSPITAL, MOORFIELDS
CONSULTING SURGEON TO QUEEN MARY'S HOSPITAL FOR THE EAST END

SEVENTH EDITION, EXTENSIVELY REVISED BY

MONTAGUE L. HINE, M.D. LOND., F.R.C.S. ENG.

SURGEON TO ROYAL WESTMINSTER OPHTHALMIC HOSPITAL, OPHTHALMIC SURGEON
TO CHARING CROSS HOSPITAL, CONSULTING OPHTHALMIC SURGEON TO MILLER
GENERAL HOSPITAL FOR SOUTH-EAST LONDON



LONDON

BAILLIÈRE, TINDALL AND COX

7 & 8, HENRIETTA ST., COVENT GARDEN, W.C. 2

1934

First English Edition, April, 1906
Second Edition, May, 1908
Reprinted, January, 1909
Third Edition, October, 1911
Reprinted, April, 1912
Fourth Edition, November, 1914
Reprinted, January, 1917, January, 1918,
February, 1919, March and September,
1920, April, 1922
Fifth Edition, September, 1924
Reprinted, May, 1927
Sixth Edition, February, 1930
Reprinted, September, 1933
Seventh Edition, September, 1934

PRINTED IN GREAT BRITAIN

PREFACE TO THE SEVENTH EDITION

FOR the present edition this manual has been extensively revised and, in parts, rewritten. Much new matter has been incorporated, but the size of the book has only been increased by thirty pages, as it has been found possible to omit some paragraphs and sections included in previous editions.

Two new coloured plates have been added. These illustrate thrombosis of the central retinal vein and a branch, and a tear in a detached retina before and after operation. Many new illustrations, more in keeping with modern practice, have replaced some of those in the last edition. Of these, I am indebted to Mr. C. L. Gimblett for Figs. 21 and 36, to Messrs. Clement Clarke for the illustrations for 'Orthoptic Training,' and to Messrs. T. Hamblin for many others throughout the book.

While numerous changes and additions will be found throughout, the chapters on Diseases of the Lacrymal Apparatus, Glaucoma, Cataract, Squint and Heterophoria, and the section on Detachment of the Retina, have been subject to the widest revision. Contact Glasses and the newer methods of Orthoptic Treatment have been dealt with. The chapter on Colour Vision has been deleted, and a completely rewritten section on the subject added to Chapter XXI. Similarly, with the approval of Mr. S. H. Browning, the chapter on Vaccines in Ophthalmology has been omitted and replaced by a much shorter section added to Chapter XXXI.

More attention has been given to Pathology than formerly.

The co-operation of Dr. C. H. May in the preparation of the present edition is gratefully acknowledged.

While the original objects of this Manual, as set forth in the preface to the first edition, have been kept in mind, it is hoped that the new material which has been added will very definitely increase the usefulness of the book.

MONTAGUE L. HINE.

September, 1934.

PREFACE TO THE FIRST EDITION

IN the following pages the authors have endeavoured to present a concise, practical, and systematic Manual of the Diseases of the Eye, intended for the student and the general practitioner of medicine. An attempt has been made to give the fundamental facts of ophthalmology, and to cover all that is essential in this branch of medicine, always keeping in mind that the book has been written for students and general practitioners. Space, therefore, has been allotted as the necessities of such readers require, estimated by an extended experience in teaching. Thus, rare conditions have merely been mentioned; uncommon affections, of interest chiefly to the specialist, have been dismissed with a few lines; and common diseases, which the general practitioner is most frequently called upon to treat, have been described with comparative fulness.

The illustrations, excepting those showing instruments, are original, and have been inserted wherever it seemed that they would be of value in elucidating the text. Those in colours represent the most common changes in the fundus, a knowledge of which is desirable in the treatment of general diseases, as well as in ophthalmic practice. These plates form a fairly complete ophthalmoscopic atlas.

Since the year 1900 four editions of May's 'Manual of Diseases of the Eye' have been published. It is hoped that the present volume may prove equally popular.

CHARLES H. MAY.
CLAUD WORTH.

March, 1906.

CONTENTS

CHAPTER	PAGE
I. EXAMINATION OF THE EYE - - - -	I
II. SUBJECTIVE OR FUNCTIONAL EXAMINATION OF THE EYE -	II
III. OBJECTIVE EXAMINATION OF THE EYE CONDUCTED IN THE DARK-ROOM - - - - -	23
IV. THE SLIT-LAMP AND CORNEAL MICROSCOPE - -	43
V. AFFECTIONS OF THE EYELIDS - - - -	52
VI. DISEASES OF THE LACRYMAL APPARATUS - - -	77
VII. DISEASES OF THE ORBIT - - - - -	87
VIII. DISEASES OF THE CONJUNCTIVA - - - -	101
IX. DISEASES OF THE CORNEA - - - - -	139
X. DISEASES OF THE SCLERA - - - - -	166
XI. DISEASES OF THE IRIS - - - - -	172
XII. DISEASES OF THE CILIARY BODY - - - -	187
XIII. DISEASES OF THE CHOROID - - - - -	193
XIV. SYMPATHETIC OPHTHALMITIS—PANOPHTHALMITIS -	201
XV. INTRA-OCULAR TUMOURS - - - - -	207
XVI. GLAUCOMA - - - - -	211
XVII. DISEASES OF THE VITREOUS - - - - -	236
XXIII. DISEASES OF THE LENS - - - - -	244
XIX. DISEASES OF THE RETINA - - - - -	272
XX. DISEASES OF THE OPTIC NERVE - - - - -	295
XXI. AMBLYOPIA AND FUNCTIONAL DISEASES OF THE RETINA	306
XXII. GENERAL OPTICAL PRINCIPLES - - - -	321
XXIII. OPTICAL CONSIDERATION OF THE EYE - - -	337
XXIV. ERRORS OF REFRACTION - - - - -	357
XXV. ANOMALIES OF ACCOMMODATION - - - -	388
XXVI. PARALYSES OF EXTERNAL OCULAR MUSCLES -	393
XXVII. COMITANT <u>SQUINT</u> - - - - -	410

CHAPTER	PAGE
XXVIII. HETEROPHORIA - - - - -	424
XXIX. OPERATIONS ON THE EXTERNAL OCULAR MUSCLES -	432
XXX. THE OCULAR MANIFESTATIONS OF GENERAL DISEASES -	441
XXXI. OCULAR THERAPEUTICS: GENERAL RULES FOR OPERATIONS UPON THE EYE - - - - -	455
XXXII. VISUAL REQUIREMENTS FOR BRITISH AND INDIAN PUBLIC SERVICES - - - - -	480
INDEX - - - - -	486

ERRATA

Page 401, sixth line from bottom, *for* abversion
read adversion.

DISEASES OF THE EYE

CHAPTER I

EXAMINATION OF THE EYE

OPHTHALMOLOGY is a branch of general medicine and surgery, from which, in practice, it cannot be entirely separated. For this reason all licensing authorities have wisely decided that he who would practise ophthalmology must first qualify for general practice. The ophthalmologist should keep up his general knowledge and be familiar with recent developments in medicine and surgery, at least in their broad outlines.

By examination of the eye is meant not the investigation of an isolated organ, but the examination of the patient with special reference to that organ and its functions.

Every case should be investigated thoroughly and according to a definite routine, otherwise there would be a danger of the attention being concentrated upon some obvious symptom, while something else, perhaps more important, might be overlooked. A beginner should write a list in their proper order of all the points to be investigated, and should, while examining a patient, consider each point in turn, rejecting those which obviously do not apply to the case in question, and investigating those which may be relevant. With increasing experience it will no longer be necessary for him *consciously* to go through the list. He will have acquired the habit of systematic enquiry.

Before beginning the examination of the eye, it is well to hear what the patient has to say about his symptoms and history. While one is making notes of any parts of his narrative which seem to be of importance, the patient himself should be studied.

The examination of the eye may be divided into Objective and Subjective or Functional.

The objective examination consists in—

Examination of the appendages and the anterior portions of the eyeball by means of inspection and palpation; this part of the examination may be conducted in daylight or artificial light.

Examination of the cornea and of the interior of the eyeball in the dark-room, with artificial light, by means of oblique illumination, transillumination, corneal microscope, ophthalmoscope, etc.

EXTERNAL EXAMINATION OF THE EYE

Inspection.

The patient should be seated facing a window or adequate source of artificial illumination. A general inspection will reveal any obvious symptoms which may be present, such as congestion, discharge, lacrymation, photophobia, undue prominence of the eyes, obvious squint, etc.

Proceeding from the superficial to the deeper parts, we commence with the *lids*, noticing their thickness, colour, and position; the condition of their margins, whether swollen, crusted, or ulcerated; the power of opening and closing; the size of the palpebral aperture, and the position and permeability of the lacrymal puncta. Then passing to the region of the *tear-sac*, we see whether this is swollen, and whether pressure with the tip of the index finger causes escape of secretion. We examine the condition and direction of the *cilia*, and notice whether any are misdirected.

Next we inspect the inner or *conjunctival surfaces of the lids*, observing any change in smoothness, thickness, and secretion of this membrane, and looking for foreign bodies.

• *Exposure of the conjunctiva of the lower lid* is easy: Place the thumb near the margin of the lid, press downward, while the patient looks up (Fig. 2).

Eversion of the upper lid requires a little practice: Grasp the central lashes between the thumb and index finger of the right hand, and draw the lid strongly downward and away from the globe, directing the patient to look down

(Fig. 3); place the left thumb (or a probe held horizontally) at the upper margin of the tarsus, and press downward, at the same time quickly turning the lid. Having turned the lid, it can be kept everted by shifting the left thumb against




FIG. 2.—EVERSION OF THE LOWER LID.




FIG. 3.—FIRST STEP IN EVERSION OF THE UPPER LID.




FIG. 4.—KEEPING THE UPPER LID EVERTED.




FIG. 5.—EXPOSURE OF THE RETROTARSAL FOLD OF THE CONJUNCTIVA OF THE UPPER LID.

the margin, the other fingers of the left hand being applied to the patient's forehead (Fig. 4).

This exposes the tarsal portion of the conjunctiva. If we wish to inspect the *retrotarsal fold* (and this is important

in the examination for trachoma), it is necessary to continue as follows: Press the edge of the everted upper lid firmly against the supra-orbital margin with the thumb of the left hand; then push the lower lid upward over the cornea with the right index finger, at the same time exerting gentle backward pressure upon the eyeball (Fig. 5).

There is a better method of everting the upper lid, which is easily learned. Put the tip of the index finger on the upper lid just above its margin. Put the thumb on the lower lid just below its margin. Tell the patient to look down. Push the upper lid up and back: this tilts its margin away from the globe. With the thumb slide the lower lid under the upper lid. The upper lid is now grasped between the finger and thumb, and is readily everted by a sort of semi-rotary movement of the thumb and finger. The whole act is continuous, and is done quickly, easily, and with a minimum of discomfort to the patient. Only one hand is required. There is no danger of pulling out lashes.

Next we proceed to the *eyeball*, and notice its position in the orbit, whether this is normal or whether the globe is pushed forward (exophthalmos, Fig. 101, p. 89) or sunken (enophthalmos).

The position of the eyeballs in reference to the visual axes should be roughly ascertained; we see whether the two axes meet at the object looked at by directing the patient to gaze at a finger held about a foot in front of the eyes; if they deviate, we investigate if there is loss of movement in any direction (paralysis) or whether there is faulty muscle balance, either latent (heterophoria) or manifest (squint or strabismus), as explained in Chapters XXVI.-XXVIII.

We observe whether there is any œdema of the bulbar conjunctiva (chemosis), or congestion of the interior part of the eyeball. If the latter is present, it should be examined carefully, for the nature of this injection points to the seat of inflammation (pp. 102 and 176).

The *cornea* is next inspected, and may show inflammation, ulceration, vascularization, opacities, or foreign bodies. As an aid we may now use a strong convex lens with which to

concentrate light from a window, but this method (oblique illumination) gives better results in the dark-room with artificial light, and is therefore described in Chapter III.

The *corneal reflex* derived from the window bars gives us information about the curvature and smoothness of this part of the eye. Placido's disc, like a target, 8 or 10 inches in diameter, with alternate black and white circles, a central hole and a handle, may also be used. Distortion of the corneal

reflection of the circles, or of the corneal reflex of the window bars, indicates a change of curvature or roughness.

To bring an abrasion or ulcer of



FIG. 6.—PLACIDO'S DISC.




FIG. 7.—CORNEAL REFLECTION OF PLACIDO'S DISC.

A, Normal; B, distortion caused by foreign body on the cornea.

the cornea more clearly into view, we may instil a drop of a 2 per cent. solution of fluorescein, washing off the excess with water. Wherever the corneal epithelium is absent there will be a green stain. This examination is best made in the dark-room with oblique illumination and corneal magnifier; if available, it is best to use a 'daylight' electric bulb to show up a faint staining.

We often find evidences of previous ulceration of the cornea in the form of opacities. When a corneal opacity is very faint and cloud-like, it is called a nebula; when denser, a macula; and when perfectly opaque and white, a leucoma (Figs. 144, 145, 146, p. 161).

The *sensitiveness* of the cornea may be noted by touching it gently with a thread or piece of soft paper, taking care not to touch the lids or lashes.

When there is much irritation, spasm of the lids (blepharo-

spasm) prevents a proper examination. In such cases, the instillation of a solution of cocaine, 2 per cent. in liquid paraffin, will aid us in exposing the eyeball.

In *infants or young children*, when blepharospasm, swelling, inflammation, or obstinacy prevents us from inspecting the cornea in the usual way, the child is laid upon its back across the nurse's lap, its hands are held, and its head is steadied between the knees of the examiner (Fig. 8). Under such



FIG. 8.—EXAMINATION OF YOUNG CHILD. HEAD BETWEEN SURGEON'S KNEES.

circumstances the lids may usually be everted by pulling upon them at a little distance from the margin. To inspect the eyeball, we part the lids by placing the thumbs or index fingers at the edges, rolling in the latter somewhat, and then separating, keeping close to the surface of the eyeball (Fig. 9). Having exposed the eyeball, we may replace the thumb of the right hand by the index finger of the left, thus leaving the right hand free for other uses. The eye will usually be found turned upward, hence the cornea will be hidden from view; but after a

minute it will appear in the palpebral aperture. Care must be taken not to scrape the cornea and cause an abrasion, nor to exert any pressure upon the eyeball, on account of the danger of perforation in case the cornea has become weakened by ulceration.

It is sometimes necessary to use retractors (Fig. 10) in order to separate the lids under such circumstances, and with these the same caution is required against wounding the cornea or pressing upon the eyeball. If this procedure should prove unsatisfactory, a general anæsthetic must be employed. When forcibly separating the lids

we must remember that pent-up secretions are released suddenly, and may squirt into the eyes of the examiner.

Next we examine the *anterior chamber* and note its depth, whether normal, shallow, or increased, and whether the aqueous humour is clear; if the latter is altered, we observe whether the exudation consists of pus (hypopyon), blood (hyphæma), spongy exudation, or the like. This examination should be repeated with focal illumination in the dark-room.




FIG. 9.—METHOD OF EXPOSING THE EYEBALL.




FIG. 10.—DESMARRE'S LID RETRACTOR.

The *iris* comes next. We note its colour, smoothness, and thickness, whether its markings are clearly defined or blurred (*muddy*), and whether it is steady or tremulous during movements of the eyeball (*iridodonesis*); the latter condition is seen when the iris is not properly supported by the lens—*e.g.*, in absence (*aphakia*), shrinkage, or dislocation of the lens. Adhesions to the cornea (anterior synechiæ) or to the capsule of the lens (posterior synechiæ) are looked for. These may require the instillation of a mydriatic for their detection.

Then we note the characteristics of the *pupil*: size, shape and position; also its reaction to light, and in accommodation and convergence as explained on p. 184. The size varies with exposure to light, with accommodation and convergence, with age, being smaller in advanced years and wider in youth, and with the state of refraction, being smaller in hypermetropes and larger in myopes. Behind the pupil we see the central part of the anterior surface of the *lens*, and observe its transparency or any abnormal condition which may be present,



FIG. 11.—TESTING THE TENSION OF THE EYEBALL.

such as cataract and deposits. To explore the lens fully, dilatation of the pupil and artificial illumination are required.

Palpation.

Palpation gives us information regarding the presence or absence of sensitiveness in the ciliary region, the tension of the eye compared with that of a normal eye, and may possibly reveal the existence of a tumour or swelling in or about the orbit.

Ciliary Tenderness.—By exerting gentle pressure upon the sclera, just behind the cornea (Fig. 11), as described below,

we may discover increased sensitiveness of the ciliary body; this is an important symptom of cyclitis.

Eye-ball Tension.—To ascertain the tension, we direct the patient to look down, and then gently palpate the sclera above the cornea, by means of the two index fingers placed upon the upper lid (Fig. 11), just as in feeling for fluctuation in an abscess, but pressing more downward than backward. We estimate the degree of tension by comparison with the other eye, if normal, or with another healthy eye.

Increase of tension is a prominent symptom of glaucoma; detached retina, perforating wounds, and degenerated conditions of the eyeball cause diminished tension; alternations in tension are sometimes found in cyclitis.

Tension is expressed roughly by the sign T. followed by n. when normal, by + or - when increased or diminished, with numerals indicating the degree of change, as follows:

T.n. = Tension normal.	T. - = Tension diminished.
T. + = Tension increased.	T. - 1 = Appreciable softness.
T. + 1 = Appreciable hardness.	T. - 2 = Decided softness.
T. + 2 = Decided hardness.	T. - 3 = Eyeball very soft.
T. + 3 = Board-like hardness.	

This method serves to estimate and record tension satisfactorily when the surgeon's fingers are educated, but for more accurate measurement and for purposes of record we use a tonometer; that of Schiötz is in most general use; McLean's is sometimes substituted. The tonometer records the resistance offered to definite weights, used to produce an indentation of the cornea, by the movement of a needle on a scale. The eye is anæsthetized with two instillations of 1 per cent. holocaine; cocaine is contra-indicated because it lowers tension of the normal eye slightly and also softens the corneal epithelium, making it more liable to abrasion from contact with the tonometer. The patient should lie on a couch, looking directly up-




FIG. 12.—SCHIÖTZ
TONOMETER, TAND-
BERG PATTERN.

ward; both lids are separated with the fingers of the left hand, without pressing on the eyeball. The tonometer is then rested by its own weight on the upturned cornea, different weights having been placed on the needle, depending on the degree of suspected alteration in tension; the needle of the instrument becomes deflected to a certain number on the scale, which an accompanying chart translates into a definite number of millimetres of mercury. Normal tension varies from 15 to 25 mm. of mercury as registered by the Schiötz tonometer.

For minute inspection of the cornea, anterior chamber, iris and lens, as well as for examination of the vitreous and fundus, we resort to oblique illumination, the ophthalmoscope, transillumination (Chapter III.), and the corneal microscope with slit lamp in the dark-room (Chapter IV.).

CHAPTER II

SUBJECTIVE OR FUNCTIONAL EXAMINATION OF THE EYE

THE *subjective examination*, dependent upon the statements of the patient, comprises the testing of the function (sight) of each eye separately. The functions to be tested are the form sense in direct vision, the extent of the fields of vision, the colour sense, and the light sense.

The *form sense* is the faculty which the eye possesses of perceiving the shape or form of objects in direct vision, and is expressed as acuteness of vision. The *field of vision* of each eye is mapped out by a series of points showing the angular distances from the fixation point at which light can be perceived. The *colour sense* is the power which the eye has of distinguishing light of different wave lengths—*i.e.*, distinguishing colours. The *light sense* is the faculty of perceiving different degrees of intensity of illumination (brightness).

The Acuteness of Vision.

Central or Direct Vision.—When we wish to see an object clearly, we look directly at it so that the image falls upon the macula lutea, the portion of the retina which is adapted for the most acute vision. The acuteness should be tested both for distant and for near vision.

Distant Vision.—In testing for distance a range of 20 feet (6 metres) is selected, since rays of light from a point at this distance are nearly parallel. For this purpose we make use of *Snellen's test types*, which are constructed upon the following principle: Each letter is inscribed within a square (Fig. 13), which subtends a visual angle of 5' at the distance at which

the normal eye should distinguish the letter. The visual angle is included between two lines drawn from the extremities of the object through the nodal point of the eye (Fig. 14).




FIG. 13. — CONSTRUCTION OF SNELLEN'S TEST TYPES. SERIF LETTER.

Each side of the square is subdivided into five equal parts; the smaller squares thus formed subtend a visual angle of $1'$, which is the minimum visual angle for the normal eye—that is, if two black objects on a white ground are separated by a space subtending a smaller angle, they will no longer be seen separate. In order to subtend the same visual angle, the size of the letters must

increase the farther they are removed from the eye (Fig. 14).

Snellen's test types consist of square-shaped serif letters arranged upon a chart, the size of the letters diminishing from above downward. The height of each letter subtends a visual angle of $5'$, the component lines a visual angle of $1'$. The uppermost letter is of such a size that it can be read at 60 metres; then follow rows of letters which should be read at 36, 24, 18, 12, 9, 6, and 5 metres respectively (Fig. 15).

Another type which is becoming popular is Green's (Fig. 16), which is based on a similar principle. Non-serif letters are used,




FIG. 14.—THE ESTIMATION OF THE SIZE OF SNELLEN'S TEST TYPES AT VARIOUS DISTANCES. NON-SERIF LETTERS.

and the total width of each letter subtends an angle of $4'$, the height still subtending $5'$. Letters are cleaner cut, and of type more in everyday use than the serif letters used in Snellen's types.

The *acuteness of vision* is expressed by a fraction, the numerator of which corresponds to the number of metres separating the patient from the chart (preferably 6 metres), and the denominator to the number indicating the distance at which

the smallest letter read by the patient should be read by the normal eye. If the patient's sight is normal, his acuteness of vision will equal $\frac{6}{6}$; this is expressed $V. = \frac{6}{6}$. If he can see only the third line from the top, $V. = \frac{6}{24}$. If he cannot read more than the top letter, $V. = \frac{6}{80}$. If he reads some letters in the $\frac{6}{18}$ line, but not all of this size, $V. = \frac{6}{18} -$ or $\frac{6}{24} +$. Many persons, especially during youth, can read the line which should be read at 5 metres from the chart; the fraction in this case would be $\frac{6}{5}$.




FIG. 15.—SNELLEN'S TEST TYPES.



FIG. 16.—GREEN'S TEST TYPES.




FIG. 17.—TEST TYPES FOR ILLITERATES.

If the patient's vision is less than $\frac{6}{80}$, we reduce the distance from the chart. If he sees the largest letter at 2 metres, $V. = \frac{2}{60}$. If he cannot read the top letter at any distance, we record the distance in feet or inches at which he can correctly count the examiner's fingers held against a dark background; for example, $V. =$ Fingers at 1 foot, or $V. =$ Fingers at 7 inches. If he has less sight than this, we move the hand before the eye, and if he is capable of appreciating such

movements, we say he has 'perception of hand movements.' If his vision is still further reduced, we ascertain whether he has perception of light (P. L.) by alternately shading and exposing the eye. This is done by means of the hand, or by throwing light upon the eye with the ophthalmoscopic mirror in the dark-room, and noting whether he indicates the presence or absence of illumination.

Each eye is tested separately, one eye being covered with a card, or with the opaque disc supported in the trial frame. Daylight may be used for illuminating the chart, but artificial light thrown directly upon the test letters is preferable, being more uniform. The test types are hung opposite a window, at about the level of the patient's eyes, and the patient is placed with his back to the source of illumination. For a child or illiterate patient, we may employ a chart such as Fig. 17; the acuteness of vision is determined by the smallest row in which the patient can correctly tell the number of funnels in each ship.

Near Vision.—When in a state of rest, the eye is adapted for parallel rays coming from a distant object. In order that divergent rays from a near object shall be focussed on the retina, there must be an increase in the refractive power of the eye. This change is known as accommodation; it will be more fully described in Chapter XXIII.

The test types usually employed to determine near vision consist of different sizes of ordinary printer's types; the finest is numbered 1, successive numbers indicating coarser type. They are known as Jaeger's test types (Fig. 18).

The patient should be placed with his back to the light, so that the page is well illuminated, and each eye tested separately. His near vision is expressed by J., followed by the number corresponding to the finest print which he can read; thus, J. 3 means that the patient is able to read the third paragraph.

Types based on the Snellen principle are also used, and are required for the testing of candidates for several public services, such as the Army, Navy, etc. They should be read at 0.5 m., 0.75 m., 1 m., etc.

1.

As she spoke, Moses came slowly on foot, and sweating under the deal box, which he had strapped round his shoulders like a pedlar. "Welcome, welcome, Moses! well, my boy, what have you brought us from the fair?"—"I have brought you myself," cried Moses, with a sly look, and resting

2.

five shillings and twopence is no bad day's work. Come, let us have it then."—"I have brought back no money," cried Moses again. "I have laid it all out in a bargain, and here it is," pulling out a bundle from his breast: "here they are; a

4.

mother," cried the boy, "why won't you listen to reason? I had them a dead bargain, or I should not have brought them. The silver

6.

the rims, for they are not worth sixpence; for I perceive they are only copper varnished over."—"What!"

8.

with copper rims and shagreen cases? A murrain take such trumpery! The block-

10.

the idiot!" returned she, "to bring me such stuff: if I had them I would throw them in

12.

By this time the unfortunate

14.

asked the circumstances

15.

to a tent, under

The Field of Vision.

Peripheral vision is exercised when the image falls upon some part of the retina outside the fovea centralis; such vision is indistinct, but of great importance.

The field of vision represents the limits of peripheral or indirect vision; it is the space within which an object can be seen while the eye remains fixed upon some one point. It usually refers to one eye, the other being covered, and, when not otherwise stated, applies to a white object. The field can be outlined roughly by the hand, more accurately by a piece of chalk upon a blackboard, or a lighted candle, most exactly by means of a perimeter.

The Hand Test.—The patient is turned with his back to the light, and the examiner faces him at a distance of 2 feet. After covering one eye, the patient is directed to fix that eye of the examiner which is opposite. The examiner closes his other eye. The hand with extended fingers is then moved from various parts of the periphery inward, midway between examiner and patient, and the latter indicates when he sees the fingers. In this way the examiner can compare the patient's field with his own; if both be normal, patient and examiner must see the fingers simultaneously. This is a very simple and rapid method, and will reveal any large defect in the field. Instead of the hand, a small white knob upon the end of a rod may be used to measure the field.

The Candle Test.—When the patient is no longer able to see the hand, we make use of a lighted candle, or light reflected from an ophthalmoscopic mirror, in the same manner, in the dark-room.

The Blackboard Test gives us an approximately correct graphic representation. The patient is placed 12 inches in front of a blackboard, upon which we mark a cross to serve as the point of fixation. A piece of chalk is now gradually brought from the periphery towards the centre, and the patient indicates when he sees it in the several directions. These points are marked, and by connecting them an outline of the field is obtained.

The Perimeter (Fig. 19) furnishes the most exact method. It may be very simple and cheap with test object moved by hand, or very complicated, self-recording, and expensive. Accurate results are given by the simpler instruments, but more time is spent in the examination. It consists of a metallic semicircle or quadrant, which can be revolved so as to take the direction of any meridian. This arc is marked




FIG. 19.—LISTER PERIMETER. RADIUS OF ARC 33 CM.

in degrees, '0' corresponding to the middle point and '90' to either extremity. The patient's head is supported upon a chin-rest, one eye covered and the other fixed upon an object placed at the centre of the arc. The test object, a white or coloured disc 5 mm. in diameter, is carried along the inner surface of the arc, and the points where it is first seen in the different meridians are marked upon

diagrams of the normal field; the lines connecting these form the boundary of the field. Larger or smaller test objects are used in special cases; sometimes the sight is not completely lost in an area which will not see a 5 mm. test object, and a field can be obtained with a larger test object. If the larger size is used first a small defect may be missed.

The size and colour of the test object must be noted on the chart; also the character of the illumination, the distance of the eye from fixation point, the date on which the field was taken, and the name of the patient. Many test objects are now graded according to the angle they subtend on the retina at a fixed distance, 1° , 2° , etc. Or the size of the test object and the distance at which it is used may conveniently be expressed as a fraction; thus $\frac{5}{330}$ indicates a 5 mm. test object used at the ordinary perimetric distance of $\frac{1}{3}$ metre or 330 mm.

Extent of the Normal Field of Vision.—For a 5 mm. or 1° white test object this is, toward the temporal side, 90° (or over); in other directions it is less extensive, owing to interference from the nose and brow, and because the percipient layers of the retina do not extend as far forward on the temporal as on the nasal side. On the nasal side the field extends 60° ; above, 60° ; below, 70° (Fig. 20).

The limits vary according to the size and colour of the test object, the intensity of the illumination, the contrast between the test object and its surroundings, the state of adaptation of the eye, and the co-operation of the patient.

Pathological Alterations in the Field of Vision.—These consist of limitation and defects. Limitations may assume the form of contraction evenly in all directions (concentric) (Fig. 240), irregular contraction (Fig. 182), or loss of part of the field, on one side or the other (Fig. 244).

Concentric contraction affects all parts of the periphery alike; when considerable, nothing but central vision may remain (Fig. 240); such contraction with preservation of good central vision is met with especially in retinitis pigmentosa. The contraction may affect only one side of the periphery; in such cases we speak of temporal or nasal contraction, or upper or lower contraction. When one-half of

the field is absent (Fig. 244), it constitutes *hemianopsia* (p. 318). Sector-shaped contractions sometimes exist; the defect then has the shape of a triangle the base of which is peripheral. Certain affections produce characteristic contraction of the visual field; for instance, in atrophy of the optic nerve the contraction is concentric; in glaucoma it is usually greatest on the nasal side.

A *scotoma* is a defect within the visual field. A physiological scotoma is the blind spot, the situation of which is




FIG. 20.—NORMAL FIELDS FOR WHITE AND FOR COLOURS (BLUE, RED, AND GREEN).

about 15° to the temporal side of the point of fixation, corresponding to the entrance of the optic nerve (the small oval ring in Fig. 20). According to their situation, we divide scotomata into central and peripheral. A central scotoma corresponds to the point of fixation (Fig. 239); when marked, it interferes with or abolishes central vision altogether; the scotoma caused by hæmorrhage at the macula furnishes an example. A paracentral scotoma (as in early tobacco amblyopia), is situated to one side of the point of fixation and a *ring or annular scotoma* encircles this point (e.g., retinitis

pigmentosa). Peripheral scotomata cause little disturbance, of sight, and may exist without the patient's knowledge, especially when situated far from the point of fixation; disseminated choroiditis furnishes examples of scotomata of this sort (Fig. 166, p. 194).

Scotomata may be *positive*, when the patient sees them as black spots in his field, or *negative*, when they exist as defects in the visual field, but are not perceived by the patient until the visual field is examined. Positive scotomata are due to changes in the media or in the retina. If the opacities exist in the vitreous, the scotomata are motile; muscæ volitantes represent one variety of defects of this sort. Negative scotomata may be *absolute*, when perception of light is entirely lost over the defective area, or *relative*, when there is only diminished perception of light, or loss of perception of certain colours over this area. Toxic amblyopia gives us an example of a scotoma which is central, relative, and often negative (Fig. 239).

For the detection of central and paracentral defects in the field within a radius of 30° , and for estimating the size of the blind spot, 2 mm. white and coloured test objects are used and the examination conducted with a flat surface instead of a perimeter arc. For this purpose we employ a Bjerrum's screen, or one of its many modifications; sometimes stereoscopic charts (Haitz) are useful for small central defects.

Bjerrum's screen consists of a black curtain, 2 metres square, supported on a framework; the patient is seated at a distance of 2 metres. A smaller screen 1 metre square, with patient seated at 1 metre, may be employed as an alternative. We use a 1 to 3 mm. test object (a white ball on a black rod). With one eye covered, the patient fixes the other on a white spot in the centre of the screen; the test object is moved in from the periphery to the centre along various meridians, and the points at which it becomes invisible and the points at which it is seen again are marked with black-headed pins. The result is then transferred to a chart for record, many devices nowadays making this a simple process. Sinclair's rule, which translates distances from fixation point into the appropriate

degrees, is the best known. The different meridians are also shown by the rule. The greater distance between the eyes and the screen than in examination on a perimeter furnishes a larger projection of the defect and thus makes possible its early detection.

The Colour Sense.

The colour sense as a whole (*i.e.*, the faculty of distinguishing different colours) is investigated by the methods described in Chapter XXI.

Central perception of colour is tested by the methods described on p. 309. The peripheral extent of the colour fields is ascertained by small objects, such as squares of coloured paper or small coloured knobs 5 or 10 mm. in diameter, which are moved from the periphery towards the centre, on the perimeter or in the coarser methods of testing the field.

The field for colours is smaller than that for white, but has the same general shape. It varies also for different colours; that for blue is the largest, next comes red, while green has the smallest field. The limits (given in Fig. 20) correspond to the points at which the colours are recognized, not to those points at which merely the presence of a moving object is perceived. The degree of brightness or colour saturation of the test object markedly affects the size of the field. The examination of the colour fields is of considerable importance, since we frequently find that contraction of the field for colours exists at an earlier period than that for white. It is a more delicate test, and detects diminution of visual power before it has become sufficiently pronounced to affect the field for white. But one must not place too much reliance on the accuracy of such fields, since they are so very dependent on the intelligence of the patient.

The Light Sense.

The light sense is the power of perceiving gradations in the intensity of illumination (brightness). We determine either the lowest limit of illumination with which an object is still

visible (light minimum=L.M.), or the smallest difference in illumination which can be appreciated (light difference=L.D.) In passing from a bright light to darkness or from darkness into a bright light vision is much interfered with, and it takes a certain length of time for vision to become normal; this adjustment is known as adaptation. The light sense is estimated by an apparatus known as a photometer or by a chart upon which grey letters are printed upon a grey background of different density.

Reduction of the light sense is present in diseases of the optic nerve and retina and also in glaucoma. Marked reduction is seen in cases which are accompanied by night blindness—*e.g.*, retinitis pigmentosa. Diminution in the light sense is not always proportionate to reduction in the acuteness of vision.

The examination of the motility of the eye is described in Chapters XXVI., XXVII., and XXVIII.

CHAPTER III

OBJECTIVE EXAMINATION OF THE EYE CONDUCTED IN THE DARK-ROOM

THE examination in the dark-room comprises the following steps:

I. *Oblique illumination*, for the physical examination of the anterior portions of the eyeball.

II. *Examination with the ophthalmoscope at a distance*, for exploring the media of the eyeball, and to determine whether the eyes fix binocularly.

III. *The indirect method of ophthalmoscopy*, for examining the fundus, giving an inverted picture of low magnification.

IV. *The direct method of ophthalmoscopy*, for examining the fundus, giving an erect picture of greater magnification.

V. *Transillumination*.

VI. *Examination with the corneal microscope and the slit-lamp*.

The examining-room should have rather dark walls, and all light should be excluded except that given by the surgeon's lamp. The source of illumination usually preferred is an electric light upon a 'universal bracket,' which permits the light to be placed on either side of the patient, and to be raised and lowered at will. The patient should be seated; the examiner may be either standing or seated.

Oblique Illumination.

Oblique, lateral, or focal illumination furnishes a very valuable means of minutely exploring the cornea, anterior chamber, iris, and lens. By means of a strong convex lens of 2 or 3 inch focus, light is concentrated upon the eye in such a manner that the apex of the cone of light corresponds to the part to be examined (Fig. 21). The source of illumination should be about 18 inches to the side of the patient,

several inches in advance, and on a level with the eye. The lens is grasped by its margin, or handle, between the thumb and index finger, held so that its surfaces are at right angles to the direction from which the light proceeds, and steadied by means of the little finger placed against the side of the patient's




FIG. 21.—OBLIQUE ILLUMINATION AND EXAMINATION WITH LOUPE.

face. After having examined one eye, without removing the supporting finger, we turn the patient's head slightly toward the light and illuminate the opposite eye. The light




FIG. 22.—CORNEAL MAGNIFIER.

may be placed on either side; if on the patient's right, we use the left hand for holding the lens; if on the left, we use the right hand. To avoid being opposite the patient's face during examination with the corneal magnifier, it is often preferable to direct the light obliquely across the nose on to the eye that is being examined; the surgeon then stands to the side of the eye he is examining. After having examined the

cornea, the lens is brought nearer to the eye, so that the apex of the cone of light corresponds to the deeper structures which we wish to explore.

With a second strong convex lens (loupe), magnifying 8 or 10 times, held at its focal distance (1 or 2 inches) in front of the patient's eye, we can magnify the illuminated area, and thus obtain greater detail.

Opacities of the cornea, aqueous, or lens, seen by oblique illumination, appear as grayish or white spots upon the black ground of the pupil (Figs. 28, 30, 32, 34, Plate II.).

The Ophthalmoscope.

The ophthalmoscope was invented by Helmholtz in 1851. Previous to this period we had practically no knowledge of the interior of the eye during life. The importance of this invention lies not only in our ability to explore the interior of the eye, but also in the means of recognizing changes in the fundus which may give valuable information in the diagnosis of disease of the system in general.

There are two types of ophthalmoscope:

1. The ordinary reflecting form in which the source of illumination is independent, and
2. The self-luminous type in which the source of illumination is within the instrument itself.

The essential portion of this instrument is a perforated mirror. This is mounted upon a convenient handle and supplemented behind by a disc containing convex and concave lenses.

The mirror serves to reflect light into the interior of the eye, while the aperture allows a portion of this light, after returning from the patient's eye, to pass into that of the observer. The mirrors commonly employed in the ordinary ophthalmoscope are concave and circular, a larger one for use in indirect examination, and a smaller, tilted one (which can be rotated either to right or left to reflect the light according to its position) for use in the direct examination. Both large and small plane mirrors may be added for special examinations, which require less intense illumination.

The lens disc is placed behind the mirror and provided

with a series of lenses, which follow each other in regular order from the weaker to the stronger. The disc can be rotated by means of the finger applied to its milled edge so that any lens can be placed behind the perforation in the mirror. Opposite each lens is a number indicating its strength in dioptries.



FIG. 23.—MORTON'S OPHTHALMOSCOPE.

FIG. 24.—OPHTHALMOSCOPE OF MORTON TYPE, SELF-LUMINOUS.

FIG. 25.—MAY'S ELECTRIC OPHTHALMOSCOPE.

nating, the lighting current being usually supplied by a dry-cell battery contained in the handle. It is now in very general use and is especially useful to students and practitioners, as with this instrument a satisfactory view of the fundus is obtained even in daylight, with great ease, without much special training, and with less necessity for dilating the pupil; for bedside use it is indispensable. It does not entirely replace

the ordinary ophthalmoscope, however, and it is wise for the student to perfect himself in the technique of the latter. To obtain satisfactory results from any electric ophthalmoscope it is necessary to observe important precautions in adjustment to ensure perfect illumination of the fundus. Such directions are, or should be, supplied by the manufacturer of the instrument, and vary with different types. Once the illumination has been correctly adjusted for the various forms of examination, direct, indirect, etc., the procedure is the same as when a non-luminous ophthalmoscope is used.

The Ophthalmoscope Examination.

Before examining the fundus one should explore the media. This preliminary step is important, since it will explain modifications in the picture obtained by subsequent methods, or failure to see the fundus in cases in which changes in the media exist. One mode of obtaining such information, oblique illumination, has already been described; it is particularly applicable to the anterior media. A second method is:

Examination with the Ophthalmoscope at a Distance.

This method explores all the media—cornea, aqueous, lens, and vitreous. The light is reflected from the mirror into the eye, and, returning from the background, traverses the media before reaching the eye of the examiner through the aperture in the mirror.

With the reflecting ophthalmoscope the source of illumination is placed just above the patient's head, so that the light strikes the patient's temple, leaving his face in darkness. The patient faces the examiner, the latter standing or sitting directly in front. The ophthalmoscope is held in front of either eye of the observer, so that he can look through the perforation, and is steadied against the side of the nose and supra-orbital margin. The distance between patient and examiner is about 15 inches.

From the mirror the light is reflected into the eye of the patient. Reaching the background, it is coloured orange-red by contact with the choroidal vessels and retinal and choroidal pigment. This tinted light returns through the patient's eye and enters the eye of the examiner by means of the aperture in the mirror. The tint varies with the colour of the background of the individual, depending upon the abundance of choroidal and retinal pigment; hence it is brighter in persons of light complexion, and darker in others. It is also influenced by the amount of illumination, and consequently the reflex is brighter when the pupil has been artificially dilated. The patient is told to move the eyes in various directions, and in this manner different parts of the media are explored.

In the normal eye a homogeneous orange-red reflex (fundus reflex) is obtained (Fig. 26, Plate II.). If any details of the vessels of the fundus are seen, the eye is *ametropic* (Fig. 27, Plate II.). If, when the observer moves his head from side to side, these vessels appear to move in the same direction, the eye is *hypermetropic*; if in the opposite direction, it is *myopic*.

If fairly large and dense opacities exist in any of the media, they will appear as dark or black spots upon the coloured background of the pupil. They are dark because they intercept a certain part of the light (Figs. 29, 31, 33, 35, Plate II.).

Opacities of the media may be either fixed, in which case they move only with the eye, or movable (floating), when they float about after the eye has been rapidly moved and then suddenly stopped; the latter occur only in the vitreous. The situation of opacities of the media can often be estimated by oblique illumination. Another method consists in noting the displacement of the opacity with regard to the pupil, when the observer's head is moved slowly from side to side. When there is no apparent motion of the opacity, it is in the plane of the iris; when it appears to move in the opposite direction, it is in front; and when in the same direction, it is behind this plane. A third method is based upon the relationship of the motion of the opacity to that of the eyeball. If, when the patient moves his eye, the opacity moves




FIG. 26.—NORMAL FUNDUS REFLEX.
Ophthalmoscope at a distance.




FIG. 27.—FUNDUS REFLEX IN AMETROPIA.
Ophthalmoscope at a distance.




FIG. 28.—OPACITY OF THE CORNEA.
Oblique illumination.




FIG. 29.—OPACITY OF THE CORNEA.
Ophthalmoscope at a distance.




FIG. 30.—SENILE CATARACT (CORTICAL).
Oblique illumination.




FIG. 31.—SENILE CATARACT (CORTICAL).
Ophthalmoscope at a distance.




FIG. 32.—SENILE CATARACT (NUCLEAR).
Oblique illumination.




FIG. 33.—SENILE CATARACT (NUCLEAR).
Ophthalmoscope at a distance.




FIG. 34.—LAMELLAR CATARACT.
Oblique illumination.




FIG. 35.—LAMELLAR CATARACT.
Ophthalmoscope at a distance.

FIGS. 26 TO 35.—EXAMINATION OF THE MEDIA WITH OBLIQUE ILLUMINATION, AND WITH THE OPHTHALMOSCOPE AT A DISTANCE, THE PUPIL BEING MODERATELY DILATED.

with (in the same direction as) the eye, it must be in front of the centre of rotation of the globe (which corresponds to the anterior portion of the vitreous, about 10 mm. in front of the retina); if it moves in the opposite direction, it must be behind this point; if it has no motion, it must be exactly at the centre. In both of these tests the greater the apparent motion the more removed is the opacity from the plane of the iris and the centre of rotation of the globe respectively.

The presence or absence of squint should be determined at this stage by the method described on p. 412.

Additional detail of changes in the media and iris may be obtained by placing strong convex lenses (from 5 to 20 D.) in the sight-hole of the ophthalmoscope, gradually approaching the eye as the strength of the lenses is increased.

The examination may usually be made through a pupil of natural size. It may be necessary to dilate the pupil, however. Moderate dilatation is secured by instilling one drop of a 4 per cent. solution of cocaine; after fifteen minutes the pupil will be of sufficient size, and the effects will pass off in half an hour, thus causing no discomfort to the patient. A 5 per cent. solution of euphthalmin acts more energetically, and the effects pass off in a few hours. Greater dilatation follows the instillation of one drop of a 1 per cent. solution of homatropine, or of a mixture of 1 per cent. homatropine and 1 per cent. cocaine; these cause mydriasis in from twenty to thirty minutes, and the effects last from twenty-four to forty-eight hours. Atropine should not be used to dilate the pupil for ophthalmoscopic examination. Before using a mydriatic one should be quite certain that none of the symptoms of glaucoma are present. After examination of the eyes of an elderly person under a mydriatic, it is a wise precaution to instil a miotic.

The Indirect Method of Ophthalmoscopic Examination.

With the indirect method we obtain an inverted image of the fundus, magnified about four diameters. The source of illumination, when external, is just above the patient's head

and the examiner is about $\frac{1}{2}$ metre in front of the patient. In the aperture of the ophthalmoscope we place a 2 D. convex lens, which enables the examiner to obtain a clear image with his accommodation at rest. Placing the ophthalmoscope before either eye, we obtain the fundus reflex. A strong




FIG. 36.—INDIRECT METHOD OF OPHTHALMOSCOPIC EXAMINATION.

convex lens (14 D.) of about 3 inches focus (called the objective lens) is now held at about its focal distance in front of the eye to be examined. This lens is grasped at its edges, or by its handle, by the thumb and index finger of the left hand, and steadied by placing one of the other fingers against the forehead of the patient (Fig. 36). If a clear view of some part of the

background is not obtained, we vary the distance from the patient by slowly moving the head backward or forward, until there appears a distinct aerial, inverted image of the fundus at a short distance in front of the lens, corresponding to its focus.

After having seen the right fundus, we proceed to the examination of the left, without making any changes in the position of the light, ophthalmoscope, patient, or examiner. We merely move the lens so as to cover the patient's left eye, now steadying it with the middle finger placed upon the forehead; the little and ring fingers are flexed into the palm of the hand, so that they will not obstruct the right or free eye of the patient, and thus prevent him from gazing in any direction which we indicate. In the examination of the left eye we may, if we prefer, hold the ophthalmoscope in the left hand and the lens in the right.

We always begin the examination by looking for the entrance of the optic nerve (the disc or papilla), this being the most prominent feature of the background. The optic nerve entrance is a little to the inner or nasal side of the visual axis; hence, in order to bring it into view, it is necessary to direct the patient to move the eye in somewhat, which will rotate the posterior pole of the eyeball outward. When we are directly in front of the patient, this is accomplished by causing him to look at our little finger, held virtually at the end of the handle of the ophthalmoscope, held horizontally, when we examine the right eye, and at our left ear when we examine the left eye.

To see the parts surrounding the disc, we move the lens or the head slightly in various directions, always remembering that the image is inverted, and that it moves with the lens, but in the opposite direction to that taken by the head. More peripheral parts are brought into view when the patient moves his eye up, down, to the right, and to the left.

When the patient looks directly at the ophthalmoscope, it brings the macula into view; but since he must accommodate when fixing so near an object, the pupil will contract. On this account it is well to dilate the pupil when we wish to get a view of the macular region with the indirect method.

The beginner may encounter a number of difficulties in using the indirect method. He may have trouble in bringing the disc into view, because the patient persists in watching the ophthalmoscope instead of looking across the examiner's shoulder. Owing to defects in the manufacture of the instrument, there are often very confusing reflexes from the margins of the sight-hole and perforation of the mirror. There is frequently a very annoying reflection of the flame from the cornea or from the surfaces of the lens which we hold before the patient's eye. These reflexes may be obviated by a slight inclination of the lens, a change in the angle of the mirror, or a little variation in the position of the examiner or source of illumination, which experience alone will teach us.

The Direct Method of Ophthalmoscopic Examination.

With the direct method we obtain an erect picture of the fundus magnified about fourteen diameters.

The examiner sits or stands to the side of and facing the patient (Fig. 37). The ophthalmoscope is supported as in previous methods, and brought directly in front of the patient's eye as close as possible. There should not be a greater distance than an inch between the eye of the patient and that of the observer. With external illumination the light is placed to the side of the patient slightly above the level of his ear.

In the examination of the right eye, when using the reflecting ophthalmoscope, the examiner and the light must be on the right side, and consequently the ophthalmoscope must be placed before the right eye of the observer. When the left eye is being examined, the light and examiner must be to the left, and the observer must use his left eye. The small tilted mirror must be so rotated that it reflects the light into the eye.

When both examiner and patient are emmetropic, and both relax their accommodation, the observer looks through the sight-hole and obtains a clear view of the fundus without any lens. The patient is told to look at the opposite wall, directly forward, over the shoulder of the examiner. This brings the disc into view. The parts around the disc are next

examined. The periphery of the fundus is brought into view when the patient looks in various directions. The macular region is found to the outer side of the disc, the distance corresponding to about twice the diameter of the papilla. When the pupil has been artificially dilated so that it cannot contract in accommodation, the macula can also be brought



FIG. 37.—DIRECT METHOD OF OPHTHALMOSCOPIC EXAMINATION WITH ELECTRIC OPHTHALMOSCOPE.

into view by directing the patient to look into the aperture of the mirror.

The size of any particular lesion is compared with that of the disc; a normal disc is $1\frac{1}{2}$ mm. in diameter. Changes in the level of the fundus (elevations or depressions) are measured in dioptres; an elevation of 1 mm. corresponds to 3 D.


The beginner is often annoyed by reflexes from the surface of the cornea. These can be obviated by a slight change in the angle of the mirror, the position of the examiner or that of the light.

If the observer be ametropic, he must either wear his correcting distance glasses or have a special correcting lens fitted behind the aperture, or he may rotate his correcting lens before the aperture from one of those contained in the disc of the instrument. When the patient is ametropic, a suitable lens must be rotated into place behind the aperture; if he is myopic, this will be the weakest concave lens, and if hypermetropic, the strongest convex lens, which will give a distinct picture. This gives an indication of the manner in which the direct method is employed for the estimation of errors of refraction.

The emmetropic observer will be unable to obtain a distinct view of the fundus of a myopic eye by the direct method without putting behind the sight-hole in the ophthalmoscope mirror a concave lens of a strength at least sufficient to neutralize the myopia. He can examine a hypermetropic eye either by putting up a convex lens or by using his accommodation. But in the direct method the observer must learn to relax his accommodation. The beginner often finds this difficult, since he cannot forget that he is looking at a very near object, and he accommodates accordingly. He is very apt to place a concave lens in the sight-hole to neutralize the effects of such efforts, even though the patient has no myopia. Relaxation of accommodation is absolutely indispensable in using the direct method for the purpose of estimating errors of refraction. It is encouraged by keeping both eyes open and looking in the distance with the uncovered eye. The sensation of relaxing the accommodation is experienced when we hold a book at the usual reading distance and stare at the print until it becomes blurred and fades. Such an exercise may be used to cultivate the habit.

The Indirect and Direct Methods Contrasted.—The *indirect method* gives us a larger field, though a smaller magnification, and hence presents a general view of the background, which is inverted (Fig. 38). It can be used successfully independent of errors of refraction in the patient's eye. On account of greater illumination we are often able to get details of the fundus, even when slight opacities of the vitreous exist.

The *direct method*, on the other hand, gives us an erect picture, which is more highly magnified (Fig. 39), though a smaller portion of the field is seen at a time; hence it permits



FIGS. 38 AND 39.—THE INDIRECT AND DIRECT METHODS OF OPHTHALMOSCOPY CONTRASTED.

of more minute exploration of particular parts to which our attention has been directed by the indirect method. It is also




FIG. 40.—OPHTHALMOSCOPIC EXAMINATION AT A DISTANCE.

the method of using the ophthalmoscope for the estimation of errors of refraction.

Theory of the Ophthalmoscope.—As ordinarily seen, the

pupil appears black because the light which leaves it is necessarily reflected in the direction from which it came. If the eye of the observer be placed so as to intercept the returning rays, the interior of the observed eye will appear illuminated. With the ophthalmoscope light is reflected into an eye under examination, and the observer's eye is placed in the path of the returning rays and receives some of these through the perforation in the mirror.

Fig. 40 illustrates the illumination of the interior of the eye with the ophthalmoscope at a distance. E represents the eye of the examiner and P that of the patient. Divergent rays of light, proceeding from the lamp L, strike the ophthalmoscopic mirror O, are reflected and made convergent, passing into the eye P, crossing in the vitreous, and illuminating the fundus between A and B. From any point of this illuminated area, C for instance, rays are reflected, pass out of the eye, being made parallel by its refracting apparatus, and proceeding, pass through the aperture of the mirror O into the eye of the examiner E. The dioptric apparatus of E brings these rays to a focus on the retina, and they form at C' an image of C.

Fig. 41 explains the indirect method. From L divergent rays proceed to the mirror O, are reflected and made convergent, passing into the examined eye P, crossing in the vitreous. They illuminate the fundus between A and B. From any portion of this illuminated area, C D for instance, rays are reflected, and passing out of the eye, are rendered parallel by its refracting apparatus. They fall upon the convex lens (L), and are brought to a focus at C' D', forming an enlarged inverted image in the air at the focus of the lens (L), which image can be seen by the eye of the examiner E.

Fig. 42 illustrates the direct method. Divergent rays proceeding from L to the mirror O are reflected and made convergent, passing into the examined eye P, crossing in the vitreous. The fundus from A to B is lighted up. From any portion of this illuminated area, C D for instance, rays are reflected, pass out of the eye P, being made parallel by its dioptric apparatus, through the perforation of the mirror O,

into the eye of the examiner E. Here they are brought to a focus on the retina. They are convergent, and being prolonged backward, form a magnified and erect image of C D behind the eye of the patient P, at C' D'.




FIG. 41.—INDIRECT METHOD OF OPHTHALMOSCOPIC EXAMINATION.

Ophthalmoscopy with Red-Free Light.—If the fundus be examined by light from which the red rays have been excluded, by a suitable filter, the background will appear of a yellowish-green colour, the macula standing out as a lemon-yellow area, the disc white, the fundus reflexes very marked, and the




FIG. 42.—DIRECT METHOD OF OPHTHALMOSCOPIC EXAMINATION.

vessels almost black with sharply cut outlines; the nerve fibres become clearly visible, those going to the macula running in a straight line, those more peripheral forming elliptical arches; thus we are able to note any disappearance of such fibres,

for instance, in optic nerve atrophy. Slight alterations in the vessels, minute retinal hæmorrhages, ill-defined exudates, and obscure changes at the macula become more easily detected than with ordinary light. A disc, having an aperture fitted with green glass, is attached to some ophthalmoscopes, to be slid over the sight-hole; this is very imperfect compared with large and special apparatus; the light must be more intense than when ordinary light is employed; the method is used

only for the detection of minute changes in the background when doubt exists with the ordinary examination, and has not been found as useful as was once expected.

The Normal Fundus.

The normal fundus exhibits a great many variations in detail. It presents an orange-red surface, upon which we distinguish the disc, the bloodvessels, and the macula (see Plates III., IV., V., and VI.).

The *Disc* or *Papilla* represents the entrance of the optic nerve; it is usually circular, but sometimes oval. Its colour is light pink, more pronounced over the inner half, the outer portion being paler. The disc is much lighter in colour than




FIG. 43. — OPHTHALMOSCOPIC VIEW AND LONGITUDINAL SECTION OF THE DISC.

a, Central artery; *v*, central vein; *E*, physiological excavation; *s*, scleral ring; *c*, choroidal ring; *r*, retina; *ch*, choroid; *scl*, sclera.

the rest of the fundus, and is separated from adjacent portions by a sharply defined margin, especially at the outer side. This margin often presents two rings: an inner, the scleral (*s*, Fig. 43), of white colour, formed by exposure of the sclera when the opening in the choroid is larger than that in the sclera, and an external ring, the choroidal (*c*, Fig. 43) of dark colour, formed by an accumulation of pigment at

PLATE III.




FIG. 44.—NORMAL FUNDUS OF AVERAGE TINT.




FIG. 45.—NORMAL FUNDUS IN A PERSON
OF LIGHT COMPLEXION.

To face page 38.

PLATE IV.



FIG. 46.—NORMAL FUNDUS IN A PERSON
OF DARK COMPLEXION.




FIG. 47.—OPAQUE NERVE FIBRES.

PLATE V.



FIG. 48.—NORMAL FUNDUS OCULI IN A PERSON OF
VERY FAIR COMPLEXION.




FIG. 49.—NORMAL FUNDUS IN A NEGRO.

To face page 38.

the margin of the aperture through which the optic nerve passes. This pigmented ring may be complete or incomplete. In the latter case it is generally found at the outer border. The margins of the normal disc are occasionally slightly indistinct, especially above and below. This appearance is sometimes seen in hypermetropic eyes of young subjects, and must not be mistaken for neuritis.

The centre of the papilla presents a funnel-shaped depression (E, Fig. 43), formed by the separation of the nerve fibres. This appears whiter than the rest of the disc. It is known as the 'physiological depression' or 'cup.' It may be comparatively large and occupy one-half or more of the disc, but never the entire papilla, in which respect it differs from the pathological excavations of glaucoma and of optic nerve atrophy (Figs. 177, 178, 179). At the bottom of this physiological excavation, when marked, we frequently see grayish spots. These represent the openings in the lamina cribrosa, the connective-tissue layer through which the fibres of the optic nerve pass (Fig. 50, Plate VI.).

The *Central Artery and Vein* of the optic nerve (*a* and *v*, Fig. 43) pass along the inner wall of the excavation, and upon reaching the surface of the disc usually divide into superior and inferior branches; each of these soon divides and subdivides, giving off nasal and temporal branches; from these smaller twigs are derived, which become terminal and do not anastomose. Small branches are often given off from the main trunks and pass across the disc. The macular region is devoid of larger vessels, though finer branches are seen to approach this area. The arteries are readily distinguished from the veins by their smaller calibre, bright red colour, and straighter course. They present a bright reflex running along the centre. The veins are of greater thickness, of a darker red colour, more tortuous, and the light streak is fainter. Arteries and veins usually follow the same course. The veins sometimes present a distinct pulsation, most marked where the central trunk appears on the disc, and increased by pressure upon the eyeball; this is physiological. Pulsation in the retinal arteries, on the other hand, is pathological,

and occurs in glaucoma, cardiac disease, and in profound anæmia.

The *Retina* itself is transparent. The colour of the background is derived from the choroidal vessels, and modified by the pigment-epithelium layer of the retina and the pigment of the choroid. It is bright orange-red in persons of fair complexion, while in darker individuals it has a deeper, brick-red colour. The fundus presents a granular or stippled appearance, caused by the pigment cells. When the pigment-epithelium layer of the retina is well developed, the choroidal vessels cannot be seen. More often considerable detail of the vessels of the choroid will be visible. This occurs under two conditions: In some cases there is no obscuration by the pigment layer of the retina, and the choroidal pigment is very abundant and collected into the intervacular spaces; then these stand out as dark islands separating bright red lines and bands, which anastomose freely, the choroidal vessels (Fig. 48, Plate V.). In other instances there is very little pigmentation in either retina or choroid, allowing the choroidal vessels to be seen plainly, now presenting the picture of bright-red anastomosing channels with brighter interspaces (Fig. 45, Plate III.). The choroidal vessels are most markedly visible in the periphery, and are readily distinguished from retinal vessels by being less sharply defined, flat, having no light-streak, by their free anastomosis, and by the fact that they obviously lie in a plane posterior to the latter.

The region of the *Macula Lutea* (Fig. 51, Plate VI.), physiologically the most important part of the fundus, is situated rather less than two disc-diameters to the temporal side of the entrance of the optic nerve, in the line of direct vision. Very often this region presents scarcely any distinctive feature. It is always devoid of visible vessels, and is somewhat darker than the rest of the fundus. Frequently a bright spot is seen in its centre corresponding to the position of the fovea centralis, or there may be two or three of these bright spots. Sometimes the macular region is represented by a bright spot surrounded by an area of dark red colour, about the size of the disc, oval horizontally, and this again encircled by a bright halo;

PLATE VI.




FIG. 50.—"PHYSIOLOGICAL CUPPING" OF THE
OPTIC DISC.




FIG. 51.—THE NORMAL MACULA LUTEA.

this reflex is best seen in the indirect method, and is most marked in children of dark complexion, especially if they be hypermetropic.

Physiological Variations.—In children of dark complexion the fundus not infrequently presents a bright lustre, which changes its position with movements of the mirror. It is most marked along the bloodvessels; it resembles the shimmer of watered silk. Another peculiar but physiological appearance is sometimes occasioned by opaque nerve fibres. In such cases the axis cylinders of some of the optic-nerve fibres regain their medullary sheath at the disc, and continue in this condition for some distance beyond the papilla, presenting one or more whitish areas extending for a variable distance from the disc and terminating in brush-like extremities (Fig. 47, Plate IV.).

Transillumination.

An ocular transilluminator is an instrument shaped like a pencil and having a tiny electric lamp and system of lenses which project a beam of light through a little hole at the end. If a solid tumour is suspected to underlie an accessible part of the sclerotic, but cannot be seen ophthalmoscopically, the eye is lightly anæsthetized and the instrument is pressed against the eye in this situation, the light being, if possible, directed slightly forward. A solid tumour may cast a shadow noticeably darker than the red illumination of the pupil obtained by placing the instrument on another meridian in the same latitude.

Most electric ophthalmoscopes can be adapted for this purpose by using a special attachment; with the May ophthalmoscope it is sufficient to apply the exposed electric bulb to the external surface of the eyelids and press firmly against the eyeball, causing the light to shine through lid and sclera from various directions. .

The Corneal Microscope and Slit-Lamp (Gullstrand's) furnish a combination which, though elaborate and expensive, is coming into use for the study of minute changes in the anterior

portion of the eye. The slit-lamp furnishes an extremely brilliant light which is condensed into a very sharp and narrow band which, traversing the parts to be examined, shows them in optical section, the remainder of the eye being in darkness. This illuminated section is then examined with the binocular microscope. With this intense, concentrated illumination and added magnification we are enabled to study microscopic changes in the conjunctiva, cornea, iris, lens, circumlental space, ciliary body, and even the anterior portion of the vitreous. The apparatus is expensive, and its use needs considerable practice. But, as this method of examination is of great importance and now forms part of the curriculum of students of ophthalmology, a separate chapter is devoted to it.

CHAPTER IV

THE SLIT-LAMP AND CORNEAL MICROSCOPE

By T. HARRISON BUTLER

OBLIQUE illumination of the eye by the light of a lamp or ordinary electric bulb of the 'Osram' type gives information about the cornea, the iris, and the anterior part of the lens, but its usefulness is limited because the diffuse light has little penetration and cannot be brought to a sharp focus, and by the low magnification of the loupe. The last difficulty can be overcome by the binocular corneal microscope, which has a useful magnification up to about 35 diameters, and gives stereoscopic vision. If instead of the usual frosted globe we use a 'half-watt' bulb of 60 candle-power we can focus the image of its circular filament as a bright line upon any object, or alternatively by throwing the condensing lens out of focus we obtain diffuse illumination superior to that given by the orthodox bulb. We can project a somewhat ill-defined beam into the depths of the eye, gaining the necessary penetration to view objects deep in the globe.

The slit-lamp, invented by Allvar Gullstrand of Upsala, is in effect a small searchlight which projects a sharply defined beam into the eye, which can be focussed as desired. The latest model of the Zeiss slit-lamp is shown in Fig. 52. The microscope seen in Fig. 53 slides upon an adjustable table, and the slit-lamp is fitted to it in such wise that the beam can be projected at any angle and can rapidly be transferred from one eye to the other. The light passes through an adjustable slit varying the width of the beam from 1 to $\frac{1}{40}$ mm. The adjustable chin and forehead rests are seen in the illustration.

Having at command a beam of bright light—the nitra-lamp used has 50 candle-power—which can be directed at any angle and focussed upon any spot, we possess selective methods of examination. We can focus directly upon an object—a foreign body in the cornea or the surface of the iris—*direct focal illumination*. Alternatively we can focus the beam on to a reflecting surface behind the object under examination, and observe it by transmitted light—*retro-illumination*. We can form a bright secondary source of light on the iris and examine the cornea




FIG. 52.—THE CORNEAL MICROSCOPE AND GULLSTRAND'S SLIT-LAMP.

by transmitted light, or by focussing the beam upon the posterior capsule of the lens we can illuminate the iris from behind, showing up atrophied areas or fissures in its pigmented layer. If the cornea is vascularized the actual circulation of the blood in the tenuous vessels is clearly visible. A third method makes use of the *zone of specular reflection*. We can so adjust the angle of the beam and the line of gaze of the patient to the axis of the microscope that the light from a reflecting surface is received by the eye of the examiner. On

the edge of the bright area the endothelium cells lining the back of the cornea can be distinguished, and the *shagreen* surface of the anterior and posterior lens capsules can be studied. The use of this *mirror light* gives valuable information regarding changes in the cornea and lens.

An examination with the slit-lamp can be made with either the broad or the narrow beam. The *broad beam* is chosen mainly to gain information about the reflecting surfaces of the eye. These are: the anterior surface of the cornea; Descemet's membrane; the posterior corneal surface; the anterior capsule of the lens; the surfaces of the various *nuclei* of the lens; the posterior capsule; and, finally, the surface of the retina. The optical conditions of the eye limit the range of the apparatus to the anterior third of the vitreous, but if the refraction of the

cornea is abolished by using a *contact glass*, the posterior parts of the vitreous and the retina are brought within the scope of the instrument. The *narrow beam* is used for examining the eye in *optical section*, and is of supreme value. The ordinary methods are limited to two dimensions; the slit-lamp gives the third dimension. We can estimate the depth of structures in the eye, mainly by inspection, taking advantage of the stereoscopic properties of the microscope, but also, if necessary, by actual measurement with the micrometer eye-piece, and by using the micrometer drum, which is fitted to the microscope and registers the travel of the tubes in tenths of a millimetre. We focus the




FIG. 53. — MICROSCOPE HEAD WITH DR. ULBRICH'S GRADUATED DRUM ATTACHED.

endothelial cells of the cornea and then the surface of the iris; the travel of the microscope read on the drum will give a fairly

accurate measurement of the depth of the anterior chamber, which may be useful in glaucoma.



FIG. 54.

We can view the cornea in optical section, and estimate its thickness and the nature of its curvatures. The depth of an ulcer can be gauged by inspection, and its extent mapped out by moving the beam across its surface. In Fig. 54 a corneal ulcer is seen on the left illuminated by the broad beam, and on the right in optical section by the narrow beam. The

dark colouration shows the depth of penetration of fluorescein. The changes in conical cornea become obvious to the eye, and the site of a foreign body or the location of an inflammatory focus is determined with accuracy.

The Architecture of the Lens.

Using the narrow beam the lens can be divided into a series of concentric *nuclei*. The *capsule* is seen as a thin bright line, and under it is a clear area limited by the *subcapsular line*. In glassblowers' cataract a thin film may be seen peeling off from the anterior surface of the capsule. This is the *zonular lamella*. The fibres of the suspensory ligament are inserted into this structure, and can be seen with the aid of the slit-lamp in such conditions as congenital coloboma of the iris and subluxation of the lens. They are arranged in bundles which lie in the valleys between the ciliary processes, one on each side, and run back to an insertion near the ora serrata. Their arrangement is seen in Fig. 55.




FIG. 55.—A CONGENITAL COLOBOMA OF THE IRIS, SHOWING THE FIBRES OF THE SUSPENSORY LIGAMENT ARRANGED IN BANDS.

The first lens nucleus is called the *cortex*, and within it is the *adult nucleus*. The next nucleus is the *infantile nucleus*,




FIG. 56.—AN OPTICAL SECTION OF A TYPICAL NORMAL LENS.

and then we have the *fœtal* nucleus, consisting of two plano-convex bodies placed plane surface to plane surface. Finally,

in the centre of the lens there is the *embryonic nucleus*, or central lucid interval. This is the oldest part of the lens. The appearance of a typical adult lens in optical section is shown in Fig. 56.

With the broad beam we can examine the surfaces of some of these nuclei with their characteristic *sutures*. The student must refer to a manual on the slit-lamp for a full description of




FIG. 57.—THE POSTERIOR Y SUTURE AND THE ARC LINE WITH A SECOND ARC ABOVE.

The hyaloid artery is very distinct.

these sutures, which are intimately concerned with the growth of the lens. The most interesting, and one well known before the advent of the slit-lamp, is the *Y suture* on the surface of the foetal nucleus. This suture can often be seen with an ordinary loupe in some examples of lamellar cataract. The anterior *Y* is erect, the posterior inverted. In all textbooks of anatomy and in most on the eye the reverse position is described. Simple inspection with the slit-lamp will show the correct

posture of the Υ s. From the arms of the Υ the lens fibres can be seen radiating out to the equator of the nucleus, where they bend over to join the opposite Υ suture.

In most children and in some adults the anterior remnant of the hyaloid artery can be seen hanging down into the retrolental space like a piece of twisted string. It is attached to the posterior lens capsule somewhat nasally from the posterior pole. The posterior Υ suture with the lens fibres and the hyaloid remnant are shown in Fig. 57.

The Clinical Value of the Slit-Lamp.

It is obvious that the exact localization given by the apparatus is of great value. In industrial areas it is constantly used to decide whether a foreign body has perforated the cornea, and often this examination shows that there is a fragment of metal within the globe. The age of a perforating corneal scar can be estimated, information sometimes of critical value in a medico-legal case. The nature and position of opacities in the lens are determined with exactitude, and their progress can be watched. Many new kinds of cataract have been discovered with the slit-lamp, and the nature of well-known types is now better understood. In old persons the central part of the lens sclerosis and the surface of the adult nucleus may now reflect almost as much light as the capsule itself. If this sclerosis is accompanied by increase in the index of refraction, the central nucleus, having a stronger curvature than the outer surface of the lens, refracts light more than the peripheral part of the lens; thus originates the *lens with a double focus*. In old persons the pupil is generally small, so the change in refraction leads only to lental myopia; but if the pupil be wide there is much blurring of the retinal image, and without the slit-lamp it would be difficult to account for the defective acuity, because the lens is often quite transparent.

In inflammatory conditions of the eye the slit-lamp is indispensable. With its aid we are able to see particles, perhaps actual cells or clumps of cells, floating in the aqueous in the

anterior chamber and retrolental space, and in the vitreous. Whereas the cornea is colder than the iris, there are convection currents in the anterior chamber; therefore, these particles circulate, rising on the iris side, falling behind the cornea. If the inflammation becomes more intense, the aqueous becomes albuminous and the convection currents cease; the particles are at rest. The beam of light now becomes visible as it traverses the anterior chamber, the so-called *flare*. The detection of flare and the presence of particles in the aqueous are often the earliest signs of such serious inflammations as *sympathetic ophthalmitis*, and the use of the slit-lamp enables us to detect this formidable complication some days earlier than we could by the ordinary methods of examination. Immediate treatment with N.A.B. can be initiated, and in most cases of sympathetic ophthalmitis both eyes can be saved. Generally the presence of cells in the anterior chamber is an indication of choroiditis, and careful search with the ophthalmoscope will often detect a focus in the fundus oculi.

The nature of corneal precipitates elucidated with the slit-lamp is of diagnostic value. Pigmented deposits are not of serious import. Clumps of lens cortex after an extraction may adhere to the posterior surface of the cornea, and cause anxiety till the slit-lamp reveals their true nature. Pellucid keratic precipitates, on the other hand, indicate serious inflammation of the ciliary body and choroid. Gelatinous precipitates upon the surface of the iris and at the pupil margin have the same significance as pellucid keratic precipitates. They come and go, and may be mistaken for tubercles. We are often in doubt whether the globe contains a malignant growth or an inflammatory exudate. This is especially the case in children; the diagnosis rests between glioma and pseudoglioma.' The detection with the slit-lamp of definite signs of inflammation suggests that the condition is inflammatory in nature.

It is obvious that not all can possess a slit-lamp, which nowadays costs from £40 to £80, but one who has mastered slit-lamp technique can apply his knowledge to the simple examination with the loupe and half-watt lamp, or he can project

a parallel beam from an electric ophthalmoscope into the eye. He can now apply the varied methods of illumination, and, knowing what he is looking for, he will gain far more from oblique illumination than one who uses the ordinary frosted bulb and has no knowledge of the slit-lamp.

Further knowledge of the slit-lamp can be gained from 'The Microscopy of the Living Eye,' by Koby (translated into English by Goulden and Harris); and 'An Illustrated Guide to the Slit-Lamp,' by T. Harrison Butler.

CHAPTER V

AFFECTIONS OF THE EYELIDS

Anatomy and Physiology.—The eyelids consist of movable folds formed, from before backward, of skin, loose connective tissue, muscular tissue, tarsus and fascia, and conjunctiva. In addition, they present eyelashes, numerous glands, bloodvessels, lymphatics, and nerves.




FIG. 58. — LONGITUDINAL SECTION OF THE UPPER LID.

S, Skin; O, orbicularis muscle; C, conjunctiva; T, tarsus; M, opening of Meibomian gland; L, lashes.

The integument is thin and delicate, and joined to the subjacent muscles by loose areolar tissue, free from fat. These characteristics explain the readiness with which extravasations of blood and œdematous swellings occur in this region.

The margin of each lid presents in front a rounded, anterior lip, from which the eyelashes (cilia) spring. These form two or three rows of short, thick, curved hairs, their roots deeply embedded in the connective tissue and muscle. They are provided with sebaceous follicles, known here as Zeis's glands. In this situation are also found modified sweat glands, known as the glands of Moll, which open into the hair follicles of the cilia, into the duct of a Zeis gland, or on to the surface of the lid margin. Behind, the lid margin presents a sharp posterior lip; directly in front of this are the openings of the Meibomian glands. The surface between these two lips is known as the intermarginal space. The margins of the lids unite at an acute

angle externally (external canthus). At the internal canthus the junction presents a rounded space, which is occupied by a small, reddish elevation of modified skin, the caruncle.

In and behind the subcutaneous connective tissue we find the muscles of the eyelids. The levator palpebræ superioris arises from the apex of the orbit, and is attached to the upper border and anterior surface of the tarsus and to the skin of the middle of the upper lid. The orbicularis muscle lies between tarsus and integument, being

attached to the latter, but gliding loosely over the former. It forms a flattened ring surrounding the palpebral aperture. We also find a layer of unstriped muscular tissue inserted into the peripheral margins of the tarsal plates and known as the palpebral involuntary muscles.

The tarsus consists of a thin plate of dense fibrous tissue, giving to each lid its firmness; it is larger in the upper than in the lower lid. The tarsi are connected with the lateral walls of the orbit by means of the internal and external tarsal ligaments, and to the upper and lower margins by an aponeurotic layer of fibrous tissue known as the palpebral fascia or ligament. In the substance of the tarsus, occurring in parallel rows, are found the Meibomian glands, thirty to forty in the upper and twenty to thirty in the lower lid. These are elongated sebaceous glands, with blind extremities and numerous cæcal appendages, filled with fatty secretion and opening on the free margin of the lid.

The palpebral conjunctiva is thin, vascular, and closely adherent to the tarsus.

The arterial supply of the lids is derived from the ophthalmic and the facial arteries; in the upper lid they form a superior arch running along the upper tarsal margin, and an inferior arch placed near the free border of the lid; in the lower lid there is merely one arch near the free edge. The veins empty into the ophthalmic, temporal, and facial. The lymphatics pass to the pre-auricular, submaxillary, and parotid lymphatic glands. The third nerve supplies the levator, the facial the orbicularis, and the sympathetic the palpebral involuntary muscles. The sensory nerve supply is derived from the fifth.

The lids protect the eyes from external injury, foreign bodies, undue exposure, and excessive light. They serve to distribute the tears and the secretions from the various glands, thus lubricating the eyeball, keeping the surface of the cornea moist and transparent, and washing away any dust which may have found its way into the eye.

The Common Affections of the Eyelids are blepharitis, hordeolum, chalazion, trichiasis, entropion, ectropion, ptosis, tumours, and injuries.

Blepharitis.

A very common chronic inflammatory condition of the margin of the lids, in which reddening and thickening is usually associated with the formation of scales and crusts (Fig. 61, Plate VII.). It occurs under two forms: (1) *non-ulcerative*; (2) *ulcerative*.

Symptoms.—In the superficial or *non-ulcerative* form the margins of the lids are swollen and reddened, and usually

present numerous whitish scales at the bases of the lashes. The latter fall out readily, but are replaced, since there is no destruction of the hair follicles. Persons with very fair complexions are especially liable to this affection.

In the deep or *ulcerative form* the edges of the lids are reddened and swollen, and present yellowish crusts, which glue the lashes together. On removing these crusts, small ulcers are seen about the attachments of the lashes; these ulcers bleed readily. The lashes become distorted, fall out, and grow scarce, since they are not replaced on account of destruction of the hair follicles. In both forms there will be itching, soreness, epiphora, and sensitiveness to light.

Sequelæ occur especially in the ulcerative form. There may be permanent loss of a greater or lesser number of lashes, hypertrophy of the lid margin (tylosis), trichiasis, and ectropion. Sometimes, in very chronic cases, all lashes, save a few scattered, fine ones, are lost, and to this condition the term *madarosis* is applied.

Etiology.—Poor hygienic surroundings; debilitated conditions of the system; following the exanthemata, especially measles; exposure to irritating atmosphere—smoke, wind, dust; late hours; insufficient sleep; uncorrected errors of refraction, especially hypermetropia and astigmatism; chronic conjunctivitis; nasal affections; lacrymal disorders; pediculi; lack of cleanliness. The disease occurs at all ages, but is very common in children.

Treatment.—The disease is apt to be obstinate. Removal of the cause, if possible, is of the greatest importance. Cleanliness, change of faulty habits, and correction of errors of refraction are great aids to local treatment. The edges of the lids must be washed gently but thoroughly with soap and water, so as to dissolve away all scales and crusts, dried and then covered with the ointment of the yellow oxide of mercury, ammoniated mercury, or ichthyol. To remove the crusts soap and water, or water to which a little borax or bicarbonate of soda has been added, should be used and applied upon cotton-wool. In the ulcerative form an occasional application of a 1 or 2 per cent. solution of silver nitrate to the raw spots

PLATE VII.




FIG. 59.—HORDEOLUM OR STYE.




FIG. 60.—CHALAZION OR MEIBOMIAN CYST.




FIG. 61.—CILIARY BLEPHARITIS.






FIG. 62.—ECTROPION.



will prove useful. In severe and long-standing cases it will be necessary to pull out all the lashes, and then to apply the treatment given above. Expression of infected Meibomian ducts is often very useful. This is done by rolling a glass rod along the inside of the lid, and exerting counter pressure with the thumb or finger on the outside. Staphylococcal vaccines have been found beneficial in some obstinate cases.

Œdema of the Lids is a very common symptom, being favoured by the structure of these parts. It may be (1) *inflammatory*, accompanying affections of the lids and adjacent parts, such as styes, dacryocystitis, and affections of the nasal accessory sinuses, or existing as a symptom of violent inflammations of the interior of the eye, such as iridocyclitis, acute glaucoma, panophthalmitis, and orbital cellulitis; (2) *traumatic*, when due to injuries, including the sting of insects; (3) *systemic*, in renal and cardiac disease; and (4) *non-inflammatory*, of which a rather frequent type is *angioneurotic œdema*, a recurrent variety which comes on rapidly, is often marked enough to close the lids, is unaccompanied by any change in the eyes, causes much alarm to the patient, and disappears about as quickly as it came on. This form is most frequently seen in women, especially at the menstrual period; it is allied to urticaria, and is most promptly relieved by a brisk saline cathartic and large doses of sodium bicarbonate.

Herpes Febrilis sometimes forms on the lids as well as on the lips and alæ of the nose, in the course of febrile affections, especially when associated with inflammation of the respiratory tract. Small vesicles, occurring in groups, are found in a slightly reddened base; they dry up in a few days and leave no scars; they may be associated with an eruption of vesicles on the cornea (p. 147).

Herpes Zoster Ophthalmicus.

Herpes zoster ophthalmicus is characterized by a unilateral herpetic eruption following the distribution of the ophthalmic division of the fifth nerve. The affection begins with severe neuralgic pain of one side of the head and face, and constitutional disturbance. The eruption presents vesicles situated

upon inflamed bases. The vesicles are at first filled with clear fluid, but this soon becomes cloudy; subsequently discoloured crusts form and drop off, leaving permanent and disfiguring scars. In some cases the nasal branch is also attacked, and




FIG. 65.—HERPES ZOSTER
OPHTHALMICUS.

then the eyeball becomes implicated; the cornea presents one or more vesicles changing to ulcers, and the iris and ciliary body and even the entire globe (panophthalmitis) may be involved.

The affection is due to disease of the Gasserian ganglion. It is most frequently observed in elderly patients of feeble constitution. Its duration is from several weeks to several months. The prognosis is usually good, but is serious when the cornea is involved.

Treatment.—Bland dusting-powders (bismuth subcarbonate, zinc oxide, or talcum) or

10 per cent. ichthyol ointment. Internally quinine and iron and the salicylates are most useful. The severe pain may need to be controlled by aspirin or morphine. The eyeball may become involved, necessitating treatment for corneal ulcer or possibly iridocyclitis.

Syphilis of the Eyelids is occasionally seen as a primary sore, in the secondary stage or in the form of gumma. Chancre having the same characteristics as when found elsewhere occurs upon the lid margin, usually near the inner canthus, accompanied by enlargement of the pre-auricular and sub-maxillary lymph glands; it might be mistaken for sty, suppurating chalazion, dacryocystitis, vaccinia, or rodent ulcer.

Vaccinia of the Eyelids is now and then met with as the result of the careless inoculation with the secretion from vaccine pustule elsewhere. It presents an ulcer covered with

greyish exudate or crust, situated at the margin of the lids—usually the lower, sometimes upon both; it is accompanied by marked swelling and redness of the lids, and by enlargement of the pre-auricular and submaxillary lymph glands.

Hordeolum or Stye.

A circumscribed acute inflammation of the tissues about the follicle of an eyelash, generally ending in suppuration.

Symptoms.—A red swelling (Fig. 99, Plate VII.) appears at the margin of the lid, accompanied by pain, tenderness, and often by considerable œdema. Very soon a yellowish point will be seen, indicating suppuration.

Etiology.—Styes occur at all ages. They are very common in young adults. They often appear in crops. They are frequently associated with a deranged condition of the system, constipation, and uncorrected errors of refraction.

Treatment.—It is sometimes possible to abort a stye by the use of cold compresses. As a rule, however, this is unsuccessful. Hot compresses are then indicated to hasten suppuration. As soon as a yellow spot is seen, the pus should be evacuated either by pulling out one or more lashes or by a horizontal incision. To prevent the formation of others, the general health should be looked after, constipation relieved, and errors of refraction corrected. Cod-liver oil and various preparations containing Vitamins A and D in concentrated form may be usefully employed. Calcium sulphide, gr. $\frac{1}{8}$ t.i.d., or syrup of the hypophosphites with iron, $\mathfrak{z}$ i. t.i.d., may be of service. In obstinate cases the use of an autogenous vaccine may be of value. The tendency to recur is often checked by treatment of the blepharitis which is frequently present.

Chalazion.

Chalazion (tarsal tumour, tarsal cyst, Meibomian cyst) is an enlargement of one of the Meibomian glands in consequence of stoppage of its duct, accompanied by a chronic inflammation in the surrounding tarsus. It occurs most frequently in adults. Uncorrected refractive error is believed to be a pre-

disposing cause. Very often several are found at the same time.

Symptoms.—The process develops slowly with insignificant or no symptoms until, after weeks or months, it has reached the size of a small or large pea. Then it presents a noticeable swelling (Fig. 60, Plate VII.), which feels hard, and is adherent to the tarsus, but not to the skin. On everting the lid, its situation is shown by discoloration of the conjunctiva, and sometimes by a small mass of granulation tissue. Sometimes chalazia disappear spontaneously; they sometimes remain for years without alteration in size and without inflammatory symptoms; occasionally they suppurate, this change being accompanied by inflammatory symptoms. Occasionally they form in the duct of the Meibomian gland and then project as a reddish-gray nodule from the edge of the lid (marginal chalazion). Rarely, a large swelling may cause a temporary astigmatism, which disappears after the contents are evacuated.

Pathology.—A chronic inflammatory enlargement of a Meibomian gland, accompanied by involvement of the surrounding tissues through infiltration with lymphocytes and fixed connective-tissue cells. The duct becomes obstructed through proliferation of its epithelium. Much of the gland tissue is replaced by small round cells (granulation tissue) containing giant cells (non-tuberculous); the blood-supply is cut off by the surrounding fibrous tissue and the contents then undergo hyaline degeneration, forming a jelly-like mass, and after longer duration they become liquefied, the chalazion then forming a pseudo-cyst with thick fibrous walls.

Treatment.—When small, they need not be interfered with. Occasionally we can cause their disappearance by the frequent application of ointments of the yellow oxide of mercury, ammoniated mercury, or boric acid, associated with massage and hot compresses. When larger, we remove them by operation, usually through the conjunctiva. The eye is cocainized; the lid is everted. A vertical incision is made through the conjunctiva and wall of the chalazion with a small scalpel (Fig. 66) or Beer's knife (Fig. 67); the contents (Meibomian secretion, granulation tissue, and mucilaginous

fluid) are removed and the walls thoroughly scraped with the chalazion scoop (Fig. 68). When a suppurating chalazion points externally it may be necessary to evacuate it through the skin. In this case the incision should be horizontal, so that the scar may be hidden in the folds of the integument.

After the operation the cyst will be filled with a blood-clot; this causes a continuation of the disfigurement for several days. Absorption may be hastened by gentle massage for



FIG. 66.—SMALL SCALPEL.




FIG. 67.—BEER'S KNIFE.



FIG. 68.—CHALAZION SCOOP.

a few minutes several times a day. Daily massage also tends to prevent recurrence of chalazia.

Marginal chalazia are often difficult to get rid of, but electro-coagulation (diathermy) has been found useful.

Infection of Meibomian ducts, causing chronic and swollen lid margins after a frothy secretion on the conjunctiva, and the oozing of glairy fluid from the ducts, is a common complication of blepharitis and is best treated by expression of the ducts at frequent intervals.

Trichiasis.

Trichiasis is an inversion of a varying number of lashes, so that they rub against the cornea (Fig. 70).

Distichiasis is an infrequent condition, usually congenital, in which the lashes can be separated into two rows, the posterior of which is directed backward so as to rub against the eyeball (Fig. 71).

In both of these conditions the margins of the lids have a normal position, the displacement affecting the lashes only.

Symptoms.—The misdirected lashes cause mechanical irritation and injury to the cornea, with irritation, pain, lachrymation, photophobia, opacities, and ulceration.




FIG. 69. FIG. 70. FIG. 71. FIG. 72. FIG. 73.

Figs. 69-73—Section of the Upper Lid, showing Normal and Abnormal Position of Tarsus and Lashes. Fig. 69, Normal lid; Fig. 70, trichiasis; Fig. 71, distichiasis; Fig. 72, entropion; Fig. 73, ectropion.

Etiology.—The most frequent cause is cicatricial contraction of the conjunctiva and tarsus in old cases of trachoma. Other causes are blepharitis, burns, injuries to the lids, and operations upon the lids.

Treatment.—1. *Epilation.*—When the misdirected lashes are few in number, we may epilate with the cilia forceps (Fig. 74), repeating this every few weeks, since the lashes grow again.

2. *Electrolysis.*—A sponge electrode corresponding to the positive pole is applied to the temple, and a fine platinum needle forming the negative pole is introduced into the hair follicle, destroying the latter; a very weak galvanic current (2 milliampères) is employed. This method results in a permanent cure, but is often rather painful, in spite of 2 per cent. novocain solution injected into the lid margin.

Diathermy has been used with good results, and is less painful.

3. *Operation*.—When a great number or all of the lashes are misdirected operations must be performed. These have



FIG. 74.—CILIA FORCEPS.

for their object correction of the faulty position or transplantation of the lashes. Since trichiasis is frequently associated with entropion, these operations will be considered in connexion with the latter disease.

Entropion.

A rolling in of the margin of the lid (and with it the lashes (Figs. 72 and 75).

Varieties.—There are two forms: (1) Cicatricial, due to cicatricial changes in the conjunctiva and tarsus, most commonly affecting the upper lid.

(2) Spasmodic, due to spasm of the palpebral portion of the orbicularis muscle, almost always occurring in the lower lid. The second variety is generally found in old persons (senile entropion) who are predisposed through relaxation of the palpebral skin and the deep position of the eyeball resulting from the absence of fat.




FIG. 75. — TRICHIASIS OF THE UPPER LID; ENTROPION OF THE LOWER LID.

Symptoms.—Those due to mechanical irritation and injury to the cornea: irritation, pain, lachrymation, photophobia, opacities, and ulceration of the cornea.

Etiology.—Cicatricial form: principal cause, the cicatricial

changes in old cases of trachoma, also burns and other injuries to the lids, and operations upon the lids. Spasmodic form: atrophy or absence of eyeball, blepharospasm, inflammatory conditions of the lids and conjunctiva, and the prolonged wearing of a bandage.

Treatment.—Non-operative treatment may be of service in the spasmodic variety. Wearing an artificial eye over the stump left after enucleation, or over an atrophic globe (if tolerated), often relieves the symptoms. If a bandage causes the entropion, we must either leave this off or apply a small roll of lint to the orbital margin beneath the bandage, exerting




FIG. 76.—HORN PLATE.




FIG. 77.—KNAPP'S ENTROPION FORCEPS.

pressure in such a manner as to neutralize the inversion. In other cases we try to remove the cause. The lid may be kept everted for a few days by collodion painted on the external surface, or by adhesive plaster passing from the margin of the lid to the cheek. If these simple means do not answer, an operation is indicated. In the cicatricial form operation is always necessary.

Operations for Trichiasis (Distichiasis) and Entropion.—

The choice of an operation (there are a great many) is influenced by the peculiarities existing in the individual case. The object of these operations is to remove the displaced lashes

from contact with the eyeball either (1) by changing the direction of the lashes from a faulty to a correct one; (2) by transplanting the offending zone; or (3) by straightening the curved tarsus.

In these operations we use either a horn plate (Fig. 76) or Knapp's entropion forceps (Fig. 77), to protect the eyeball, check hæmorrhage, and give proper support to the lid. The horn plate is passed beneath the lid and pressed forward. If the lid-clamp be used, its solid blade is passed beneath the lid, and the latter secured by tightening the screw of the instrument.

The *Jaesche-Arlt Operation* attaches the zone of hair fol-




FIG. 78.—THE JÆSCHE-ARLT OPERATION AND ENTROPION INCISIONS.

FIG. 79.—THE JÆSCHE-ARLT OPERATION COMPLETED.

licles at a higher level by shortening the skin of the lid. The lid is split through its entire length in the intermarginal space (Fig. 78), so that the anterior lip contains the hair follicles. A second incision, dividing the skin down to the tarsus, is made 4 mm. from and parallel to the margin of the lid. A third incision extends upwards in a curve between the two ends of the second incision. The elliptical piece of skin bounded by the second and third incisions is dissected away without injury to the orbicularis, and the margins of the defect are united by fine silk sutures. In this manner the strip of integument containing the cilia is drawn upward and the lashes are tilted forward, away from the cornea. The

area from which the skin and lashes have been displaced may be allowed to cicatrize, or may be covered by the excised strip of integument properly trimmed, or by a strip of mucous membrane from the mouth; these inserts become attached in a few days.

The *Streatfeild-Snellen Operation* aims at straightening the inverted lid by the removal of a wedge-shaped piece from the tarsus. A transverse incision is made through the skin, 2 mm. above and parallel to the margin of the lid along its entire length. A strip of orbicularis is excised, thus




FIG. 80.—THE STREATFEILD-SNELLEN OPERATION FOR ENTROPION.

One suture has been tied.

FIG. 81.—THE STREATFEILD-SNELLEN OPERATION FOR ENTROPION.

View in section.

exposing the tarsus. A wedge-shaped piece, the apex of which is directed toward the conjunctiva, is removed from the tarsus along its entire length. The cut surfaces of the tarsus are brought into contact by three sutures, provided with needles at both ends, in the following manner: One needle is passed through the tarsus above the groove; both needles are then carried down in front of the wound in the tarsus, and then between tarsus and skin, and brought out just above the free margin of the lid (Fig. 81) about 4 mm. apart. More pronounced eversion is produced if the threads are brought out behind the cilia, at the posterior lip of the lid

margin (Herbert's modification). The two threads are tied upon a bead and then turned up over the forehead and secured by plaster. The cutaneous wound closes of itself.

Operations for Spastic (Senile) Entropion include: (1) *Excision of a horizontal strip of skin* with the underlying orbicularis, the width being gauged so that when pinched up it shall cause the disappearance of entropion without producing ectropion; the margins of the wound are then united by silk sutures. (2) *Injection of alcohol*: a few drops of 4 per cent. novocain are injected into the outer fibres of the orbicularis after puncture of the outer extremity of the lower lid near its margin; leaving




FIG. 82.—CANTHOPLASTY.

the needle in place, 0.2 to 0.3 c.c. of 95 per cent. alcohol are then injected into the same site for a distance of 5 mm.; the procedure is usually painless, successful for some time, often permanently so. (3) *Canthoplasty*.

Canthoplasty consists in an enlargement of the palpebral fissure by division of the external canthus. The lids being separated and stretched at the external canthus with the fingers, one blade of blunt-pointed, straight scissors is introduced behind the external commissure as far as possible, and the entire thickness divided, the wound in the skin being made a little longer than that in the conjunctiva. This

leaves a rhomboidal wound. The conjunctiva at the apex of the wound is loosened from underlying tissue and stitched to the centre of the incision in the skin. A second suture is passed through the upper, and a third through the lower part of the wound, uniting conjunctiva to skin (Fig. 82).

The sutures are inserted so as to prevent reunion, thus making the effect permanent. If a temporary enlargement is desired, we omit the sutures; the operation is then known as canthotomy or temporary canthoplasty.

The indications are blepharospasm associated with spastic entropion, and certain cases of trachomatous pannus. Temporary canthoplasty is indicated in acute purulent conjunctivitis, when swelling of the lids exerts injurious pressure upon the eyeball, in blepharospasm, and in the removal of an enlarged eyeball or an orbital tumour.

Ectropion.

An eversion of the lid with exposure of more or less conjunctival surface (Fig. 73, and Fig. 62, Plate VII.). It may affect the upper or the lower lid, or both.

Symptoms.—Epiphora (from eversion of punctum) causing excoriations and eczema of the lower lid, which, in turn, through contraction, increase the deformity. The exposed conjunctiva becomes reddened and hypertrophied. In marked cases the cornea may suffer, as a result of imperfect closure of the lids.

Etiology.—(1) Cicatricial contraction from wounds, operations, burns, ulcers, and caries of the orbital margin or surrounding surfaces (cicatricial ectropion). (2) Chronic conjunctivitis and blepharitis associated with considerable hypertrophy. (3) Relaxation of the skin and orbicularis in old people (senile ectropion), affecting only the lower lid. (4) Affections of the facial nerve, causing paralysis of the orbicularis (paralytic ectropion), affecting only the lower lid. (5) Spasmodic contraction of the marginal portion of the orbicularis (spasmodic ectropion), seen especially in children with acute conjunctivitis associated with considerable blepharospasm.

Treatment.—*Non-operative*: The spasmodic form is frequently relieved by a suitable retaining bandage applied after the lid has been properly placed. In the paralytic form we employ a bandage, at the same time attempting to cure the facial paralysis. In the senile form, the patient should be instructed, when wiping away the tears, to press upward and inward and not downward and outward. In slight cases of ectropion associated with much conjunctival hypertrophy, painting the exposed surface with a 2 per cent. solution of silver nitrate may be of service. Careful massage of a cicatrix may give some relief. When these simple pro-




FIG. 83.—ADAM'S OPERATION FOR ECTROPION.

A, Incisions; B, completed.

cedures do not answer, and in cicatricial ectropion, we must resort to operative intervention.

Operations for Ectropion.—In some cases the lower lid margin appears to be everted by a chronic hypertrophy of the conjunctiva and subconjunctival tissue. Two horizontal incisions, removing a wedge-shaped piece of the whole width of the hypertrophied mass, will effect a cure. No sutures are needed. In senile and paralytic forms of ectropion the lid may be replaced by: (1) Galvano-puncture; (2) reduction of the length of the lid-border; (3) tarsorrhaphy.

Galvano-puncture is indicated only in slight cases, and effects may be only temporary. The conjunctiva of the lower lid is punctured by a blunt-pointed electrode, heated to a dull heat,

along a line 4 mm. from lid margin; the punctures are 4 mm. apart; the cautery penetrates conjunctiva and tarsus but not the skin.

Shortening the Margin of the Lid (Adam's Operation) is applicable when there is considerable elongation. A wedge-shaped piece is excised from the whole thickness of the lid, the base corresponding to the margin of the lid, and varying from 5 to 10 mm. in width, according to the amount of shortening required (Fig. 83, A); the edges are brought together by a hare-lip pin and the cutaneous margins by silk sutures. The piece may be excised from the centre of the lid; but, to prevent notching, it is better to operate at the external canthus (Fig. 83, B).

For cicatricial ectropion a great many operative procedures have been advocated. An essential condition for success is the thorough division of all cicatricial adhesions, so that the lid assumes a natural position, the object of any operation being to prevent recicatrization. If the ectropion is slight and but little skin has been lost, it may be sufficient to divide the cicatricial bands subcutaneously, or to cut out the scar portion and bring the margins of the wound together by sutures. A procedure very frequently used is—

The V Y Operation (Wharton Jones).—A V-shaped incision is made with the apex directed away from the palpebral margin, the incision including the cicatrix. The skin is freed from underlying parts, not only in the V-shaped area, but also to either side. The V-shaped area is then slid upwards until slight inversion of the lid-margin is produced. The margins of the incisions are then brought together by sutures in such a manner that the figure Y results (Fig. 84).

In more extensive cicatricial ectropion a plastic operation is usually required (blepharoplasty).

Blepharoplasty consists in covering the defect formed by the excision of a cicatrix, new growth, or extensive ulceration, with skin-flaps with a pedicle, taken from some adjacent part, or by means of skin-grafts. In such operations it is customary to close the lids temporarily by several sutures so as to

prevent the contraction of the cicatricial tissue from undoing the result accomplished by the operation. Of the innumerable blepharoplastic operations with pedunculate skin - flaps, Knapp's, Dieffenbach's, and Fricke's methods are the ones most commonly employed.

Knapp's Method (lower lid) consists in detaching a lateral flap on each side of the defect in the lid, freeing it from adjacent tissue, drawing the two flaps over the defect, and uniting them by a vertical row of sutures.

Dieffenbach's Method (lower lid) makes use of an adjacent




FIG. 84.—THE V Y OPERATION FOR ECTROPION.

A, Incisions: sutures in place; B, completed.

quadrangular flap taken from the cheek and slid inward so as to cover the defect of the lid.

Fricke's Method (upper or lower lid) consists in taking a tongue-shaped flap somewhat larger than, and having the shape of, the defect in the lid from the temple or cheek; the base of the flap adjoins one end of the lid-wound, and is the part which becomes twisted when the flap is transplanted into the defect.

Skin-Grafting.—The defect is filled in by one large piece of skin or by a number of smaller ones after the lid has been fastened in its proper position by temporarily suturing the two lids together. The grafts are taken from some part of the body in which the skin is thin and delicate, such as the

inner side of the arm or thigh. The area of the graft or grafts must be one-third larger than the defect to be covered, to allow for shrinkage. The graft may consist of the entire thickness of the skin (Wolfe's method), or comprise only the epidermis (Thiersch's). The area to be covered must be clean and free from blood. When in place, the graft is covered with a layer of rubber or silk protective, and then with an anti-septic dressing. The dressing is not disturbed for three days, and the original protective layer over the graft is often left in place still longer.

Skin-grafting is now used very extensively and with very good results. If a portion of the graft sloughs, the corresponding defect can be freshened and another graft applied. This method causes less disfigurement than when pedunculate flaps are used. Thiersch's grafts, being thinner and softer than Wolfe's, produce better results cosmetically, and the lid is not so heavy.

Tarsorrhaphy.—The object of this operation is to reduce the width of the palpebral fissure by uniting the edges of the lids at the outer commissure. The edges of the lids are approximated at the outer canthus to the required extent, so as to give the operator exact knowledge as to how much




FIG. 85.—TARSORRHAPHY.

union is desired. A horn spatula is passed behind the outer commissure, and the desired length of the border of each lid is excised, including the hair follicles. The length of the flap varies according to the effect desired (about 3 to 6 mm.); its breadth is about 1 mm. To obtain firmer adhesion, the border of the lid, excluding the

cilia, is denuded for 2 or 3 mm. beyond the point at which the first incision stops. The denuded edges are then brought together by silk sutures (Fig. 76). This operation is indicated in lagophthalmos, especially in Basedow's disease, in some cases of senile and paralytic ectropion, and in connexion with

blepharoplasty. Median tarsorrhaphy, a union of the central portion of the lids, is performed when the latter are to be kept closed for some time but ultimately separated; this operation is indicated in blepharoplasty, to keep the lids in position during healing, and in neuromyopathic keratitis when the eye is to be kept covered for a lengthy period.

Ankyloblepharon is the adhesion of the margins of the two lids; it may be partial or complete, congenital or acquired; it is often associated with symblepharon.

Blepharophimosis is an apparent contraction of the palpebral fissure at its outer canthus, due to this angle being covered and hidden by a vertical fold of skin. It may be seen in lengthy cases of chronic conjunctivitis in which, as a result of epiphora, dermatitis develops and draws the adjoining skin over the canthus.

Blepharospasm, spasm of the orbicularis, closing the lids, is a symptom of ocular disease or of a neurosis. The *tonic form* (persistent) is present with foreign bodies, fissure at the outer canthus, corneal and uveal disease; it is due to irritation of the terminal filaments of the *trigeminus*; rarely it is hysterical; treatment consists in removing the cause. The *clonic variety* (intermittent), often consisting merely of *fibrillar twitchings*, especially of the lower lid, unimportant though annoying, may depend upon errors of refraction, conjunctivitis, or overwork. Intermittent spasm may be a form of habit chorea. Another variety, very obstinate, is sometimes seen in elderly persons in whom the lid spasms may be accompanied by similar movements of the neighbouring facial muscles, constituting a form of *tic*; this condition may be relieved for a variable period or even permanently by the injection of alcohol into the orbicularis.

Lagophthalmos is an incomplete closure of the palpebral fissure when the lids are shut, as a result of which there is exposure and consequent injury to the bulbar conjunctiva and the cornea. The condition may be due to congenital or acquired shortening of the lids, ectropion, paralysis of the orbicularis and protrusion or enlargement of the eyeball; it is seen also in unconscious and moribund individuals.

Ptosis.

A drooping of the upper lid due to paralysis or deficient development of the levator. All degrees of ptosis occur. When marked, it interferes with vision by covering the pupil. Patients attempt to raise the lid by forced action of the occipito-frontalis muscle, wrinkling the skin of the forehead (Fig. 86); they favour exposure of the pupil by throwing the head backward; this attitude is characteristic.



FIG. 86.—EPICANTHUS AND BILATERAL PTOSIS.

Etiology.—Ptosis may be *congenital* or *acquired*. When congenital it is usually bilateral, due to deficient development of the levator, and often associated with other congenital defects. Acquired ptosis is usually unilateral; it is caused by paralysis of the branch of the third nerve which supplies the levator, and is usually associated with paralysis of other ocular muscles supplied by the oculomotorius. When not associated in this way it is not infrequently the result of cerebral disease.


Mechanical ptosis is a variety due to (1) increased weight of the lid (trachoma, tumours, etc.); (2) lack of support (atrophy of globe and after-enucleation); and (3) lack of connective-tissue connexion between skin and levator and upper border of orbit, in which the relaxed skin forms a fold which falls over the margin of the lid.

Treatment.—In the ordinary variety of the acquired form we seek the cause of the paralysis of the third nerve and treat this; syphilitic cases respond well to treatment. Electricity

is used. If this treatment fails to remedy the deformity after a lengthy trial, in the congenital variety and in some mechanical cases, operation is indicated.

Operations for Ptosis.—Operations for ptosis usually produce some improvement, but a really satisfactory result is rarely obtained. Care should be taken that closure of the lids be not made difficult by the operation.

Of the operations which aim at replacing the action of a defective levator palpebræ by attaching the lid to the occipito-frontalis muscle, that of Hess (or one of its modifications) is generally employed.



FIGS. 87 AND 88.—HESS'S OPERATION FOR PTOSIS.

Hess's Operation.—A 3 cm. incision through the skin of the eyebrow permits undermining down to the lid-margin. Three double sutures are then introduced, so as to form loops about 7 mm. from the lid-border, and passed upward beneath the brow, emerging 1 cm. above the incision, where they are tied upon small rolls of gauze (Figs. 87 and 88). The skin wound is closed with sutures. The double threads are allowed to remain for two weeks. In this operation the skin of the lid is displaced upwards, adheres, and gives the occipito-frontalis greater purchase; the effects are therefore better than when simple sutures are used.

Motais' Operation endeavours to assist or replace the action of the levator by exposing and freeing the middle third of the

superior rectus muscle and stitching this to the upper border and anterior surface of the tarsus. It can only be performed when there is good upward movement of the eyeball. There is some risk of temporary diplopia and depression of the globe from weakening of the superior rectus.

Advancement of the Levator may be done when the loss of action of this muscle is incomplete, more especially in older people with gradually drooping lids. The tendon, with its expansions to the tarsus, is exposed through a skin or conjunctival incision, and is either folded on itself or a portion resected and reattached to the tarsus. In suitable cases this operation may prove very satisfactory. To determine whether or no the levator palpebræ has any action, the upper lid is closed, and firm pressure made by the surgeon's thumb on the upper margin of the orbit. This pressure stops any assistance from the frontalis, and if the lid can be raised there is some power in the levator.

Epicanthus.

Epicanthus is a congenital condition, sometimes associated with ptosis, usually bilateral, in which a perpendicular fold of the skin extends from the root of the nose to the inner end of the brow, concealing the inner canthus and caruncle. In Mongolians it is a racial characteristic. In slight degree it is often seen in young children associated with a flattened bridge of the nose, and often disappears with the development of the face. When sufficiently marked to constitute a deformity, it can be relieved by excising an elliptical piece of skin from the root of the nose, long axis vertical, and suturing together the free margins.

Tumours of the Eyelids.

Benign Tumours include xanthelasma, molluscum, verruca (wart), fibroma, cyst, nævus, and milium.

Xanthelasma is a flat or slightly raised yellowish discoloration beneath the skin, found most frequently near the inner canthus in elderly women. It is due to degeneration of the muscle fibres.

Molluscum is a small, white, rounded tumour about the size of a small pea, presenting a depression at its apex; several

usually occur at the same time. They are considered contagious by some authorities. They represent a diseased condition of the sebaceous glands. They should be incised, the contents forced out, and the base touched with silver nitrate.

Milium is a small, yellowish-white elevation about the size of a pin's head, due to retention in a sebaceous gland.

Small cysts with transparent contents, due to obstruction in the outlet of sweat glands, are often seen on the lid-border; they give rise at times to irritation and disfigurement, and should be punctured with the point of a Graefe knife.

The others resemble tumours of the same class occurring in other parts of the body. Benign tumours of the lids may be excised, providing too great a loss of skin is not occasioned by the operation.

Malignant Tumours.—*Sarcoma* is rare. *Carcinoma*, when it attacks the lids, usually assumes that form of epithelioma known as '*rodent ulcer*.' This occurs in elderly persons, especially at the inner end of the lower lid-margin. It begins as a small pimple or wart, covered by a crust, soon changes to an ulcer with indurated walls, and spreads, if unchecked, to neighbouring parts. Its growth is, however, slow, and many years may elapse before it assumes any considerable size.

Treatment of Rodent Ulcer.—Exposure to radium, which is effective in early stages and is indicated in every case; the hypertrophic type is the most favourable; results are bad in cases of deep ulceration, infection, or previous X-ray or radium treatment. When radium fails excision is necessary, the defect in the lid being covered by blepharoplasty. When radium treatment is not available in early stages, carbon dioxide snow may be used—applied for forty to sixty seconds on alternate days at first, and then once a week.

Injuries of the Eyelids.

These are quite common, and include contusions, wounds, burns, and insect bites. Ecchymosis and œdema are often marked symptoms, on account of the looseness of the subcutaneous connective tissue.

Ecchymosis ('black eye') is usually of no importance,

merely causing disfigurement, which lasts one or two weeks. If seen immediately, cold compresses may be of service. After a day or two hot compresses and gentle massage are indicated to promote absorption of the extravasated blood. Occasionally in debilitated individuals, especially if associated with abrasion, abscess of the lid results, and may require horizontal incision. In fracture of the base of the skull, blood may travel along the floor of the orbit, and after a day or two appear in the lower lid and bulbar conjunctiva.

Insect bites give rise to a great deal of swelling, which is best controlled by cold compresses.

Incised wounds cause considerable gaping, if vertical, on account of division of the orbicularis, and then the scar is apt to be noticeable; if horizontal, the lips of the wound do not tend to separate, and usually heal without deformity. Incised wounds should be cleansed and stitched at once, using fine silk and delicate needles. A vertical wound of the margin must be carefully sewed so that no indentation will remain.

Lacerated and contused wounds, if extensive and accompanied by much swelling, should not be closed at once. The wound should be thoroughly cleansed, and cold compresses applied to reduce the swelling, after which the edges may be brought together. Injured parts, however slenderly attached, should not be removed. Care must be taken not to produce deformity or shortening by too tight sutures. It may be advisable to use skin-grafts.

Burns should be irrigated with solution of boric acid, dried, and covered with a bland oil or ointment. When granulating, skin-grafts should be supplied if the defect is extensive.

Emphysema associated with injury to the lids denotes a solution of continuity of the walls of the orbit, permitting communication with the neighbouring air cavities. The lids will present a soft swelling of considerable size, often closing the palpebral aperture; bubbles of air, becoming displaced in palpation, give rise to the sensation of crepitation. A firm bandage will hasten the disappearance of the air. The patient must be instructed to avoid any straining efforts, such as blowing the nose, which will increase the emphysema.

CHAPTER VI

DISEASES OF THE LACRYMAL APPARATUS

Anatomy and Physiology.—The lacrymal apparatus consists of a secretory portion, the lacrymal gland, and an excretory portion, which collects the tears and conducts them into the inferior meatus of the nose.

The *lacrymal gland* is a small, oblong body, placed in the upper and outer part of the orbit, and divided into two portions. The upper part, the larger, about the size of a small almond, is situated in a depression in the orbital plate of the frontal bone, the lacrymal fossa, to which it is fixed by connective tissue; the lower division, the smaller, is known as the accessory lacrymal gland, and is placed just beneath the outer part of the conjunctiva of the fornix. In structure the lacrymal resembles the salivary glands, consisting of acini containing cuboidal cells. The excretory ducts of both portions of the gland, the lacrymal ducts, six to twelve in number, pass downward and empty into the external half of the superior fornix conjunctivæ by separate orifices.

The excretory portion of the lacrymal apparatus (Fig. 89) consists of the puncta, the canaliculi, the sac, and the duct. The *puncta* are two minute openings, one of which is seen upon an elevation on each lid about 6 mm. from the inner canthus; they are the orifices of the *canaliculi*. The latter extend vertically for a short distance, and then continuing at right angles, pass horizontally inward in a curved course, and empty separately or together into the lacrymal sac.

The *lacrymal sac*, situated at the inner side of the internal canthus, is the upper, dilated portion of the lacrymo-nasal duct, and is placed in a groove formed by the lacrymal bone and the nasal process of the superior maxillary bone. It measures 12 mm. in the vertical




FIG. 89.—DIAGRAMMATIC SKETCH OF THE EXCRETORY PORTION OF THE LACRYMAL APPARATUS.

and 6 mm. in the horizontal and transverse diameters: its walls are thin; it is covered in front by the internal tarsal ligaments and some fibres of the orbicularis muscle.

The *nasal duct* passes downward and slightly outward and backward in a canal formed by the superior maxillary, lacrymal, and inferior turbinated bones, and terminates below in the fore-part of the inferior meatus of the nose; its length varies from 18 to 24 mm., and its diameter from 4 to 6 mm.; it is somewhat contracted where it joins the sac and again at its lower extremity. Both sac and duct are formed of fibrous and elastic tissues, and mucous membrane lined with columnar epithelium which may be ciliated. The lower part of the duct is surrounded by a dense plexus of veins.

The *lacrymal secretion* is a slightly alkaline liquid containing a comparatively large amount of sodium chloride. Ordinarily the lacrymal gland secretes just enough to moisten the eyeball, and this is lost by evaporation. As the result of psychical stimulation or of irritation of the eye or the nose, there is increased secretion. The conveyance of tears from the conjunctiva to the lacrymal sac is effected by the act of winking, the lubrication of the margins of the lids by fatty material ordinarily preventing the tears from flowing over.

Epiphora.

Epiphora ('watery eye'), an overflow of tears upon the cheeks, is a very pronounced symptom in all affections of the tear-conducting apparatus. It may also be dependent upon increased secretion (foreign bodies, inflammations, exposure to bright light and smoke, affections of the nose, irritation affecting the terminal twigs of the trigeminus). The two forms may be combined. It is very common in old people, especially in the open air or in cold weather, often without manifest lesion of conjunctiva or tear passages.

Anomalies of Puncta and Canaliculi.

Normally, the lower punctum is directed backward and upward toward the eyeball. In *eversion of the punctum* (lower) it looks forward and away from the depression in which the tears accumulate, and the result is epiphora. The condition may be due to a relaxed state of the lids in old age, to conjunctivitis, blepharitis, and ectropion.

Contraction and Obliteration of the Puncta and Canaliculi may be congenital, or acquired as a result of wounds and

chronic inflammations of this region. Foreign bodies (eyelash or concretion) may obstruct the canaliculi.

A leptothrix or streptothrix infection of the canaliculus may occur, characterized by dilatation of the canaliculus with discharge oozing from the punctum, and, as a rule, with no obstruction to the passage of tears. On opening the canaliculus a large cavity with pockets is found containing little roundish masses of leptothrix or streptothrix in mucoid secretion.

Diseases of the Lacrymal Apparatus may be divided into those of the gland and those of the conducting portion. The former (acute and chronic dacryoadenitis) are very rare; the latter (acute and chronic dacryocystitis) are very common.

Chronic Dacryocystitis.

A chronic inflammation of the lacrymal sac frequently due to an obstruction in the nasal duct. It is also known as blenorrhœa or catarrh of the lacrymal sac, and as mucocele.

Symptoms.—The constant symptom is epiphora, increased by exposure to cold, wind, dust, smoke, etc. There is fulness in the region of the lacrymal sac. By pressing upon the distended sac a viscid fluid of whitish, yellowish, or greenish colour (depending upon the amount of pus) escapes from the puncta; but sometimes the sac is emptied in the reverse direction, and the accumulation is pressed into the nose.

The course is chronic, and extends over years. A long period may elapse before the patient seeks relief. After the mucopurulent material has filled the sac for a long time there is atrophy of its mucous membrane and distension of its walls (Fig. 63, Plate VII.). A form of chronic conjunctivitis affecting chiefly the inner canthus (lacrymal conjunctivitis) and blepharitis are frequently present; eczema occurs sometimes and there may be more or less ectropion. As a result of contamination by micro-organisms from the conjunctiva (especially streptococci and pneumococci, Figs. 113 and 110, Plate IX.), a purulent inflammation of the lining of the sac may be set up.

The infectious character of the accumulation in the lacrymal sac is shown, when any abrasion or ulcer of the cornea exists,

by the readiness with which the wound or ulcer becomes infected. In operations upon the eye such a condition is a very frequent cause of infection, and must be radically treated before these are performed.

Etiology.—In most cases there is stricture of the nasal duct. This usually arises from some affection of the nasal mucous membrane, in the first place. The duct is predisposed to obstruction by the existence of a plexus of veins encircling its lower end. As a result of rhinitis, there is swelling or cicatricial contraction of the mucous membrane of the duct. More rarely pressure from tumours (polyps and hypertrophies), ulcerations, caries, and periostitis are responsible.

The affection is not very uncommon in the new-born and in infants, in whom it is due to imperfect canalization of the nasal



FIG. 90.—WEBER'S PROBE-POINTED CANALICULUS KNIFE.



FIG. 90a.—NETTLESHIP'S DILATOR.

duct. It is usually cured by the simplest measures, but may, after two or three months, require syringing under anæsthesia, or very occasionally the passage of a probe.

Treatment.—In recent and slight cases we may relieve the epiphora by curing the nasal affection which produces the obstruction. Locally, stimulating and astringent remedies, such as zinc sulphate and the ointment of the yellow oxide of mercury, may be applied to the inner part of the eyeball, followed by gentle massage over the sac so as to favour entrance of some of the remedy. The patient is instructed to empty the sac by pressure several times a day. The sac should be washed out with warm and weak solutions of salt, boric acid, or zinc sulphate, using a small syringe with delicate nozzle (Fig. 91). It is often necessary to dilate the lower punctum with the fine conical sound (Fig. 90a) before

the nozzle can be introduced. If the nasal duct is pervious, the solutions will enter the nose and escape from the anterior nares when the patient inclines the head forward. In this case, continued syringing will cure the trouble. And even if the fluid cannot at first be got to flow through the nasal duct, one usually succeeds if one keeps up the treatment long enough. Daily washing out of the sac gradually brings it




FIG. 91.—LACRYMAL SYRINGE.

into a more healthy condition, and, as the congestion of the mucous membrane of the nasal duct subsides, the duct again becomes pervious; a drop or two of 2 per cent. novocain in 1:10,000 adrenalin syringed into the sac is often of great help in reducing congestion prior to syringing.

When conservative treatment fails one may try probing, but this is not done so frequently as formerly; it is painful,



FIG. 92.—COWPER'S PROBES.



FIG. 93.—ONE OF TEALE'S PROBES.

requires tedious repetitions, and is usually only of temporary benefit. Even with the greatest care cicatricial stricture is likely to follow injury of the delicate mucous lining. But when a case has been probed before, continued probing will probably be required. Though it is possible to pass the smallest probes after dilatation of the canal, for larger sizes it may be necessary to slit up the canaliculus at its outer end, after

anæsthetizing with a few drops of 2 per cent. novocain in 1:10,000 adrenalin syringed along the duct.

To Slit Open the Canaliculus.—The surgeon stands behind and supports the patient's head against his body. Weber's




FIG. 94.—SLITTING OPEN THE LOWER CANALICULUS.

probe-pointed canaliculus knife (Fig. 90) is most frequently used. The lower lid is pulled outward by the thumb of one hand, and with the other the knife is introduced vertically




FIG. 95.—PASSING A PROBE INTO THE NASAL DUCT.

until it passes the punctum, and then horizontally. Its edge is upward, and looks toward the eyeball so as to cut into the conjunctiva and not into integument (Fig. 94). It is pushed

horizontally inward until its extremity meets with the firm resistance of the inner bony wall of the sac; then the knife is raised into a vertical position, slitting the outer third of the inner wall of the canaliculus.

To Pass Probes into the Nasal Duct.—Commencing with a small size—say a No. 3—we pass this horizontally inward exactly as the knife is passed, the surgeon standing behind the patient. When the probe reaches the inner wall of the sac, which we can be certain of when in lifting the probe there is no wrinkling of the skin of the lower lid, it is raised so that its lower end points toward the furrow between nose and cheek. It is then pushed downward gently until it reaches the floor of the nasal fossa (Fig. 95). If the probe does not pass readily, we must not use force, but withdraw it slightly and try again, or try a smaller or larger size. The probe is left in from fifteen to thirty minutes, and the proceeding is repeated every other day, gradually using larger probes; then the intervals between probing are increased.




FIG. 96.—LACRYMAL STYLE.

Sometimes the stricture is cut, a strong narrow knife being passed in the same manner as a probe, and the obstruction divided in two or three directions; this is immediately followed by probing. The treatment is painful, and the benefit, if any, is only temporary. In former days leaden or silver styles (Fig. 96) were passed into the nose after dealing with the stricture in an endeavour to make the opening permanent. They were left in for days or weeks; nowadays they are very rarely employed.

In all obstinate and long-standing cases, when there is much discharge, when a lacrymal fistula is present, when any operation on the eyeball is in prospect, or when the patient is exposed by his occupation to the risk of corneal injuries, extirpation of the sac is indicated.

Excision of the Lacrymal Sac.—Instruments required are: Strong fixation forceps, dissection forceps, small scalpel, re-

tractor (Fig. 97), scissors, blunt dissector and spoon (Fig. 98). A general anæsthetic may be used in the case of a very timid patient. Local anæsthesia (see Chapter XXXI.) is preferable,




FIG. 97.—RETRACTOR.



FIG. 98.—BLUNT DISSECTOR AND SPOON.

as there is almost complete absence of hæmorrhage. About 30 minims of novocain (2 per cent.) in adrenalin chloride (1:10,000) are injected at the site of the operation with a hypodermic syringe. The skin of the eyelids is pulled outwards with the finger so as to make the internal palpebral ligament prominent. A curved incision is made from just above the upper border of the internal palpebral ligament downwards and slightly outwards for about $\frac{1}{2}$ inch along the margin of the orbit. The incision is only about $\frac{1}{8}$ inch internal to the inner canthus. A more median position would involve risk of wounding the angular vein, with troublesome hæmorrhage. The incision enters into the sac above, but is shallower below. The crest of the lacrymal bone, which can be felt by the finger, forms the guide to the sac. The retractor is inserted. The mucous membrane of the sac is grasped with




FIG. 99.—EXTIRPATION OF THE LACRYMAL SAC.

forceps, and is separated from the bone on the inner side. Hæmorrhage is stopped by keeping the incision stretched by retractors, and by pressing into the wound for two or three minutes a plug of wool soaked in adrenalin. The sac is then freed all round, especially remembering the top of the sac. The sac is pulled up and the duct cut as low down as possible. The duct is then curetted with a small Volkmann's spoon. When all hæmorrhage is stopped, the wound is closed with sutures. A firm pressure pad is applied over the sac for two or three days. The sutures are removed on the fourth or fifth day. When the sac has been completely removed results are very satisfactory, but a certain amount of lachrymation must be expected to occur in cold weather, or in strong winds. Care must be taken to keep the conjunctiva healthy, and weak astringent drops (zinc. sulph. gr. $\frac{1}{2}$ ad $\mathfrak{z}$ i.) for occasional use will be found beneficial. Under ordinary conditions there should be little or no epiphora.

When there is slight, or no, persistent purulent discharge, epiphora is very troublesome to the patient, no operation on the eyeball is in prospect, and the sac is shown by X-rays, after injection of lipiodol, to be distended, good results may be obtained by drainage of the sac into the nose. This may be done intra-nasally (West's operation), when the services of a rhinologist are required, or externally (Toti's operation or one of its modifications). In the latter type of operation the lowermost part of the inner wall of the sac is cut away after complete exposure, a portion of the underlying bone and periosteum is removed by punch or dental drill, and a corresponding opening made in the subjacent nasal mucous membrane.

Acute Dacryocystitis.

An acute inflammation of the region of the lacrymal sac occurring in the course of a chronic dacryocystitis, as a result of an acute exacerbation. It is also known as abscess of the lacrymal sac (Fig. 64, Plate VII.).

Symptoms.—The skin over the lacrymal sac becomes red and swollen. This condition extends to the lids and conjunctiva, and is often sufficiently pronounced to lead to

a suspicion of erysipelas. There are great pain and tenderness, some fever and constitutional disturbance. After two or three days a yellow discoloration appears at one point. Here pus will be present, and should be evacuated. This will be followed by relief and a subsidence of symptoms.

The opening may heal completely, and the case again have the symptoms and slow course of chronic dacryocystitis. In other cases the opening persists, and the escaping fluid changes its character and becomes watery. This constitutes lacrymal fistula. As long as this remains open the patient is safe; as soon as it closes, he is liable to have a recurrence of abscess. Sometimes merely a minute passage is left, insufficient to admit a probe, from which a drop of fluid escapes from time to time.

Etiology.—Lacrymal abscess involves not only the sac, but the surrounding connective tissue as well. The decomposed contents of the sac find a small defect in the lining, through which micro-organisms (especially streptococci, Fig. 113, Plate IX.) reach the neighbouring tissues and excite inflammation and suppuration.

Treatment.—If the case is seen early, we try to prevent the formation of abscess, by pressing out the accumulation and syringing with mild antiseptic solutions (boric acid 4 per cent., or bichloride 1 in 10,000). If this cannot be done, or is not effective, as is often the case, hot fomentations should be used.

As soon as fluctuation occurs, we make a free incision through the skin and the anterior wall of the sac. After evacuation the incision is packed with a strip of gauze, which is changed daily, until all inflammatory signs have disappeared, and the fluid is no longer purulent. We try to restore permeability of the duct, after which the fistula closes spontaneously. If this does not happen after the duct becomes pervious, we freshen and unite the edges of the opening, or cauterize them with the actual cautery, or scrape out the track with a sharp spoon. In most cases it will be advisable to extirpate the sac, but not until all acute symptoms have subsided.

CHAPTER VII

DISEASES OF THE ORBIT

Anatomy.—The orbit is formed of bony walls having the shape of a quadrilateral pyramid; the apex corresponds to the optic foramen; the base is directed forward and corresponds to the strong, thick, projecting, anterior margin. The nasal wall, the thinnest, is formed largely by the lacrymal bone and the os planum of the ethmoid, also by the frontal and sphenoidal bones; it presents in front the groove for the lacrymal sac. The inner walls of the orbits are almost parallel, but the outer diverge considerably from each other from behind forward.

The posterior portion of the orbit presents three *openings* leading to adjacent cavities: (1) the optic foramen, transmitting the optic nerve and the ophthalmic artery; (2) the superior orbital fissure (sphenoidal fissure), transmitting the ophthalmic vein, the nerves for the ocular muscles, and the first branch of the trigeminus; (3) the inferior orbital fissure (spheno-maxillary fissure), transmitting the maxillary nerve and infra-orbital artery.

Besides communicating with the cavity of the skull by means of the openings at the apex, the orbit is *surrounded by a number of other cavities*. These are the nasal fossæ and accessory cavities—the ethmoidal and sphenoidal sinuses, the frontal sinus, and the maxillary sinus; these relations are important.

The *contents* of the orbit consist of the eyeball and optic nerve, the ocular muscles, the lacrymal gland, bloodvessels, and nerves; the spaces between these are filled with fat and fasciæ.

The *eyeball* is composed of the segments of two spheres; the anterior (cornea), about 12 mm. in diameter, is the smaller and more prominent; the larger, posterior, corresponds to the sclera. The eyeball measures about an inch in diameter (24.5 mm. from side to side, 24 mm. from before backward, and 23.5 mm. from above downward).

The *orbital fascia* is extensive and presents numerous subdivisions. It serves as *periosteum* to the walls of the orbit (*periorbita*). A portion closes in the opening of the orbit, forming an anterior wall and extending from the margin of the orbit to both tarsi, and to the external and internal tarsal ligaments, thus constituting the *septum orbitale*. Prolongations of the orbital fasciæ surround the muscles and connect them with one another, the lids, and the margins of the orbit.

In addition, a layer of fascia surrounds the globe from the cornea to the posterior part, separating the organ from the orbital fat and attached to the sclera by fine trabeculae. This investment is known as *Tenon's capsule*. Anteriorly it merges into the subconjunctival connective tissue; while posteriorly it disappears around the optic nerve, and is pierced by all the structures which are attached to the globe. In wide movements of the eyeball, both globe and its capsule move together as a whole upon the surrounding fat. Where the tendons of the ocular muscles pierce Tenon's capsule, the latter is reflected upon them, becoming continuous with their fasciae.




FIG. 100.—CORONAL SECTION SHOWING THE ORBIT AND ADJACENT CAVITIES.

The *arteries* of the orbit are derived from the ophthalmic. The *veins* empty into the ophthalmic veins, which pass through the superior orbital fissure to the cavernous sinus. The *nerves* of the orbit are motor and sensory; the motor nerves, the third, fourth, and sixth, supply the ocular muscles; the sensory nerves are the first and second branches of the trigeminus. The ciliary ganglion lies to the outer side of the optic nerve. It receives motor fibres from the third, sensory fibres from the fifth, and sympathetic filaments from the carotid plexus. It gives off the short ciliary nerves which enter the eye at its posterior part. The orbit contains no lymph vessels or lymphatic glands.

Affections of the Orbit frequently present a characteristic symptom, exophthalmos.

Exophthalmos (proptosis) is a protrusion of the eyeball from the orbit (Fig. 101). It is caused by inflammations, tumours, and injuries of the orbit, dilatation of adjoining cavities, Basedow's disease, pulsating exophthalmos, thrombosis of the cavernous sinus, and in some cases by paralysis of the recti muscles and tenotomies of these muscles. When of high degree, it may cause interference with the mobility of the eye, imperfect closure of the lids (lagophthalmos), with resulting keratitis from exposure, ectropion of the lower lid, diplopia (if lateral displacement is added), and interference with vision from inflammation and atrophy of the optic nerve.



FIG. 101.—EXOPHTHALMOS.

Enophthalmos is the recession of the eyeball into the orbit. With the exception of the cases seen in the aged and in extreme emaciation, due to the decrease of orbital fat, it is rare. Other causes are paralysis of the sympathetic, injuries to the orbit causing cicatricial contraction, and fracture of the orbital wall.

Orbital Periostitis.

An inflammation of the orbital periosteum, either acute or chronic in its course, and either limited to a portion of the margin of the orbit or spreading more deeply.

The products of inflammation often consist merely of a thickening of the membrane. Sometimes there is a deposit of bone or gumma (syphilis); there may be the formation of an abscess, with or without caries or necrosis of a part of the wall of the orbit.

Symptoms.—These depend upon whether the affection runs an acute or a chronic course, the part of the orbit involved, and the nature of the products of inflammation.

The most common variety is that attacking the margin of the orbit. In such a case there may be no other symptoms

than pain, tenderness on pressure at the orbital margin, hard, immovable swelling in this situation, and some swelling of the lids and conjunctiva. Such a case frequently results in complete absorption of the products of inflammation; less commonly, periosteal thickening or bony deposit remains. If, on the other hand, there is pus, a periosteal abscess is developed at the margin of the orbit, which perforates through the skin, leaving a fistula through which the probe detects either bare or necrosed bone. Such a fistula remains open for months until all the dead bone has been extruded, and after it heals there is a depressed scar and sometimes ectropion and lagophthalmos.

If the periostitis is situated more posteriorly, there will be more pain of a deep-seated character, tenderness upon pressure upon the globe, swelling and redness of the lids and conjunctivæ, perhaps exophthalmos, and marked constitutional disturbance. Such a case may result in absorption of the products of inflammation, or in periosteal thickening or bony deposit. But if such a deep-seated process goes on to the formation of an abscess, it becomes much more serious, and presents the symptoms of orbital cellulitis (*q.v.*), from which it frequently cannot be differentiated. The pus tends to find its way to the surface, but this may take some time. Meantime there may be danger to life through extension to the cranial cavity and the occurrence of meningitis or cerebral abscess.

Etiology.—Injuries, tuberculosis (in children); tertiary syphilis (in adults); chronic “rheumatism”; extension from neighbouring cavities or bones. Rheumatic and syphilitic cases usually run a chronic course, and produce periosteal thickening without any tendency to suppuration.

Treatment.—That of syphilis, rheumatism, or tuberculosis. Locally, moist, warm compresses. Incision as soon as there are any signs of suppuration. Early incision by means of a narrow knife is indicated as soon as we suspect the existence of pus, so as to prevent the extension of suppuration to the brain. The opening is drained by means of a strip of iodoform gauze, until pus no longer escapes. Caries and necrosis may require subsequent operative intervention.

Orbital Cellulitis.

Orbital cellulitis is an inflammation of the cellular tissue of the orbit, terminating in suppuration. It runs an acute course, accompanied by marked constitutional symptoms.

Symptoms.—Great swelling of the lids, chemosis, exophthalmos, impairment of mobility of eyeball, violent pain in the orbit and side of head, increased by pressure against eyeball. These local signs are accompanied by marked constitutional symptoms, with high fever; cerebral symptoms may be added. Vision may not be affected, or it may be reduced or abolished, owing to the occurrence of optic neuritis. After these symptoms have lasted about a week, pus appears at a certain part of the skin of the lids and perforates, or it may empty into the fornix. After early evacuation of pus in an uncomplicated case, the symptoms rapidly subside, and the opening heals.

Complications.—Optic neuritis; less frequently, thrombosis of the retinal vessels; occasionally panophthalmitis. Cases following influenza are especially apt to be complicated with corneal ulcer. Extension of the process to the brain may be fatal.

Etiology.—(1) Injuries and operations followed by infection; (2) extension of inflammation from neighbouring parts; (3) influenza; (4) facial erysipelas; (5) metastasis (pyæmia, puerperal septicæmia, etc.); (6) sometimes no cause can be discovered.

Treatment.—Hot fomentations. Early and deep incision in the spot where we suspect the abscess to be situated, either through the conjunctiva or through the skin. Even when we do not strike pus, we relieve tension, and prepare a route for the subsequent evacuation. The wound should be irrigated frequently with warm hydrogen peroxide lotion (3 volumes) and kept freely open until it heals from the bottom.

Thrombosis of the Cavernous Sinus (almost always infective) may be due to extension of a thrombus in the orbital veins occurring in orbital abscess, or may be caused by other neighbouring pus foci situated in the pharynx, tonsils, teeth, and the accessory nasal sinuses, or may follow erysipelas, caries of the petrous bone, and metastasis in pyæmia and the

infective diseases. The signs and symptoms are similar to those of orbital abscess, from which it can be differentiated by the existence of severe cerebral symptoms, œdema over the mastoid region (through venous stasis in the emissary veins), extension to the opposite side, marked distension of the retinal veins and the more constant existence of neuro-retinitis. The disease is always fatal.

Tenonitis.

Tenonitis is rare as a primary affection and consists of serous inflammation of Tenon's capsule, resulting in cure in a few weeks. Its symptoms are moderate swelling of the upper lid, localized swelling over the insertion of one of the recti muscles or diffuse chemosis, slight exophthalmos, limitation of movements of the eye, and some pain on motion of eyeball. It may follow an injury or a tenotomy of one of the recti muscles, in which the wound has become infected; it has also been ascribed to influenza, gout, and rheumatism. It may be difficult to differentiate between this affection and a mild case of orbital cellulitis. As a secondary affection it is found to accompany severe forms of iridocyclitis, and also panophthalmitis. Treatment consists of cold or warm fomentations and the treatment of the etiological factor.

Pulsating Exophthalmos.

Pulsating exophthalmos presents the following symptoms: exophthalmos, pulsation of the eyeball and surrounding parts, bruit heard over the eye and forehead, noises in the head, pain, marked distension of the bloodvessels of the retina, conjunctiva, and lids, and occasionally optic neuritis. Compression of the carotid of the same side causes a diminution or disappearance of the pulsation and bruit. It is most frequently produced by an arterio-venous aneurism due to rupture of the carotid into the cavernous sinus, generally caused by traumatism. Some cases are due to aneurism of the ophthalmic artery or one of its branches, or of the internal carotid, or to a vascular tumour. The condition may be fatal from hæmor-

rhage. Treatment consists in the ligation of the common carotid of the same side; it may also be necessary to tie the common carotid of the opposite side after a sufficient interval has elapsed to allow the cerebral circulation to recover, or, alternatively, to ligature the ophthalmic veins or the angular veins. Many cases are cured in this manner.

Tumours of the Orbit.

These are of infrequent occurrence. They may arise from the walls or the contents of the orbit. The symptoms will depend upon the size, position, and nature of the tumour. Exophthalmos is usually present. The direction of the protrusion and the impairment of motion of the eyeball will be determined by the exact situation of the tumour. Pressure upon the optic nerve may cause optic neuritis and atrophy. The tumour may perhaps be felt by the tip of the finger passed between the margin of the orbit and the eyeball. Benign tumours usually grow slowly, and frequently give few symptoms; malignant tumours increase in size very rapidly, and cause much pain. Benign tumours of the orbit include dermoid cyst, aneurism, angioma, pulsating exophthalmos, meningocele, osteoma, and distension of neighbouring cavities. Malignant tumours, sarcoma and carcinoma, usually necessitate complete exenteration of the orbit. After removal a course of radium or X-ray therapy is indicated. In the case of a benign tumour requiring removal, an attempt should be made to preserve the eyeball, especially if the sight be good. When such tumours are not situated too far posteriorly they can often be removed by a free external canthotomy down to the bone, with an incision into the upper and lower fornix at right angles to the canthotomy opening; additional approach may be provided by temporary tenotomy of the external rectus muscle. If this does not procure sufficient space, Kronlein's operation (temporary loosening and displacement of the external orbital wall (p. 100) may be resorted to.

Good results have sometimes been obtained by the implantation of radon seeds into the growth.

The Ocular Manifestations of Disease of the Nasal Accessory Sinuses.

The accessory sinuses of the nose (frontal sinus, anterior and posterior ethmoidal cells, sphenoidal sinus, and maxillary antrum) surround the orbit, being separated by bony walls which are very thin in places. They are lined by an extension of the nasal mucous membrane, and, as a result of such relationship, often become infected. If the natural outlet of the sinus becomes blocked, there will be an accumulation of secretion and consequent distension of the walls of the sinus, often with encroachment upon the orbit and exophthalmos. If this retention is of a mucoid character, the condition is known as mucocele; if of a purulent character, as empyema. Such a sinusitis may run an acute or a chronic course.

Frontal sinus disease may be accompanied by a bulging at the upper and inner angle of the orbit, with tenderness on pressure over this area, and sometimes redness of the overlying skin, severe frontal headache, and dizziness on stooping. There may be protrusion of the eyeball downward and outward, diplopia, oedema of the lids, conjunctival and episcleral congestion, and lacrymation. Orbital periostitis and cellulitis may result.

Ethmoidal disease may present a tumefaction at the upper and inner part of the orbit, with swelling of the integument of the adjacent lids, displacement of the globe downward and outward, diplopia, marked pain, conjunctival and episcleral congestion, and lacrymation. The process may involve the orbit, causing periostitis or cellulitis.

Disease of the sphenoidal sinus is usually associated with ethmoidal disease. The optic nerve is in close relationship to the walls of this cavity. A case of disease of this sinus may present no external evidence of inflammation, and yet may give rise to optic neuritis, neuro-retinitis, and retrobulbar neuritis, leading to optic nerve atrophy if the cause is not removed. There may be a central, paracentral, or annular colour scotoma, which later may become absolute, usually

with but little contraction of the visual field. Another fairly constant symptom is enlargement of the blind spot. There may be deep-seated pain, and tenderness when the eyeball is pressed backwards.

Antrum disease is not so often accompanied by ocular symptoms. There may be pain, swelling of the lids, conjunctival congestion, and lachrymation, but involvement of the orbit is rare.

Any chronic sinus disease may be the underlying cause of acute, or relapsing, intra-ocular inflammations (iritis, iridocyclitis, etc.).

Injuries of the Orbit.

These include contusions, incised and penetrating wounds, foreign bodies, and fracture of the orbital wall. A prominent sign is hæmorrhage into the orbit, causing exophthalmos and perhaps ecchymosis of the lids and conjunctiva. A blow or stab with a blunt body may rupture the globe or dislocate it in front of the eyelids.

Penetrating wounds may destroy the eyeball, injure the optic nerve, causing blindness, or sever some of the muscles, resulting in paralysis and diplopia; if infected, such wounds are followed by orbital abscess. *Foreign bodies* may be tolerated if aseptic; if infected, suppuration ensues.

Fracture may involve the anterior wall, or the inner wall causing emphysema, or the apex involving the optic canal and injuring the optic nerve; the last may result from direct injury or indirectly (*contrecoup*) and produce blindness with-




FIG. 102.—ENUCLEATION SCISSORS.

out ophthalmoscopic evidence, followed in a few weeks by atrophy of the optic nerve.

Treatment consists in cleansing and disinfecting wounds and endeavouring to extract foreign bodies (with the aid of an X-ray examination). If the situation is such that extensive manipulation would be necessary for its removal, and we have reason to believe that the substance is aseptic (such as shot), it is often better to allow the foreign body to remain. Free exit for secretions must be maintained. A bandage aids in the absorption of blood and air.

Congenital Anomalies of the Eyeball.

These are usually bilateral. Anophthalmos is a small solid or cystic mass occupying the place of the eyeball. Microphthalmos consists of an eyeball of diminished size in all diameters. Buphthalmos (congenital glaucoma) is an increase in size of the eyeball, with symptoms of glaucoma, usually resulting in blindness (p. 235).

Operations upon the Eyeball.

Enucleation of the Eyeball.—The instruments required are: Eye speculum (Fig. 185), fixation forceps (Fig. 186), blunt-pointed strabismus scissors (Fig. 336), squint hook (Fig. 337), strong, curved enucleation scissors (Fig. 102), needle-holder (Fig. 338), fine, curved needle, and thin black silk.

Operation.—A general anæsthetic is necessary in children and nervous patients. In others local anæsthesia may be used if desired, with an injection of 3 per cent. novocain into the orbit, as described on p. 467. After introduction of the speculum, the conjunctiva is divided all around the cornea, *as close to its border as possible*, and dissected back as far as the insertions of the recti muscles. A squint hook is passed beneath the tendon of the internal rectus, and the latter is divided with the strabismus scissors close to its insertion; then the other straight muscles are cut in the same way, together with the subconjunctival connective tissue, for some distance beyond the equator. The points of the scissors

must always be directed toward the eyeball, and the latter stripped as clean as possible to avoid any unnecessary removal of orbital tissue. Instead of commencing with a circum-corneal division of the conjunctiva, we may begin with a tenotomy of the internal rectus, and then divide the conjunctiva as we pass from tendon to tendon. The hook is passed around the globe to make sure that the attachments of the muscles have been completely divided. The eyeball is then dislocated forward by pressing the speculum backward, and thus the optic nerve is put on the stretch. The enucleation scissors (Fig. 102), closed, are passed between sclera and conjunctiva, feeling for the optic nerve; they are withdrawn a little, slightly opened (Fig. 103), and the nerve is divided close to the sclera. The eyeball is held between the thumb and index finger of the left hand, and the oblique muscles and other unsevered attachments are divided. The socket is irrigated with a large quantity of perchloride of mercury lotion, 1 in 5,000. The orbit is plugged for a few minutes to control hæmorrhage, and the conjunctiva is usually closed either with a single suture, which is passed through its edge at intervals and tied like the string of a pouch, or by interrupted sutures. The eye is bandaged and the patient kept in bed for a day.




FIG. 103.—ENUCLEATION OF THE EYEBALL.

Care should be taken to avoid rupturing the globe. A collapsed globe makes the operation more difficult. Troublesome hæmorrhage may occur; it can be controlled by hot water or by a firm bandage. When an eyeball containing a malignant growth is enucleated, as much of the optic nerve as possible should be removed. In rare cases infection of

the wound has led to abscess, thrombosis, and even fatal meningitis. The danger of meningitis is somewhat increased in enucleation of suppurating eyeballs; hence many oculists consider panophthalmitis a contra-indication to enucleation, and postpone operation until after the suppurative process has ceased.

The indications for enucleation are: (1) Injuries of the ciliary region, when the eye is completely blind, or the traumatism so extensive that the form of the eyeball cannot be preserved; (2) to prevent or cure sympathetic ophthalmia; (3) severe pain in a blind eye; (4) iridocyclitis, phthisis bulbi, and glaucoma, when accompanied by severe pain or inflammatory symptoms, and when the eye is blind or is certain to become so; (5) malignant tumours, either intra-ocular or epibulbar (excepting small tumours of the iris, which can be entirely removed by iridectomy); (6) anterior staphyloma, if the eye is blind, troublesome, and disfiguring; (7) panophthalmitis; (8) foreign bodies in the eye when they cannot be removed and cause irritation, or the eye is blind.

To give a larger and more movable stump in cases where there is no sympathetic ophthalmia, new growth or panophthalmitis, an insert may be placed in Tenon's capsule; this insert may be a hollow sphere of glass, gold or platinum, or a ball of fat, rather smaller than the globe.

The tendons of the recti muscles and the edges of Tenon's capsule are drawn together by a purse-string catgut suture, and the edges of the conjunctiva then united by a horizontal row of fine silk sutures.

Evisceration of the Eyeball.—In this operation the cornea and entire contents of the eyeball are removed, the sclera alone remaining.

The instruments required are: Eye speculum (Fig. 185), fixation forceps (Fig. 186), curved strabismus scissors; Graefe knife (Fig. 254) or Beer's knife (Fig. 67), sharp spoon, needle-holder (Fig. 338), small curved needles, catgut, and silk sutures.

Operation.—After insertion of the speculum, the eye is transfixated just behind the cornea with a Graefe or Beer's knife, which is made to cut its way out at the upper sclero-

corneal junction; the other half of the cornea is removed with the scissors. The contents of the eyeball are then removed thoroughly with a sharp spoon, care being taken that nothing but sclera is left. The cavity is irrigated and allowed to fill with blood, or else dried. The scleral edges are brought together in a vertical line with catgut sutures, and the conjunctiva is united horizontally with silk sutures.

Recovery is less rapid than after enucleation, and the pain and reaction are greater; the support for an artificial eye is usually better. The operation may be substituted for enucleation in all cases excepting malignant tumours, foreign bodies, and sympathetic ophthalmitis.




FIG. 104.—ARTIFICIAL EYES.

a, Front view; *b*, rear view; *c*, section of shell eye; *d*, section of Snellen 'reform' eye.

Evisceration with Insertion of an Artificial Vitreous (Mule's Operation).—Following evisceration, after the scleral cavity has been cleansed and hæmorrhage checked, a hollow sphere of glass or silver (or some other substance) is introduced. This ball must not be too large. Its introduction is facilitated by slitting the sclera and by the use of a special inserting instrument. The wound is then closed and dressed as after the ordinary evisceration. There is considerable reaction after this operation, and the patient is confined to his room for a week. The stump is undoubtedly superior to that furnished by any other method, but it frequently happens that the ball is extruded.

Artificial Eyes (Fig. 104) are worn after enucleation and evisceration, for cosmetic purposes, and to fill out the cavity left between the lids. They can be worn as soon as the

socket is free from inflammation, usually after several weeks. The artificial eye should be washed frequently, and must be removed every night. After a year its surfaces and edges usually become roughened, and it must be replaced by a new one. When there is a stump of good size, a shell-shaped artificial eye may be indicated; but with a small stump or after enucleation, the more modern Snellen 'reform' artificial eye gives better cosmetic effect; the latter has a certain thickness, and is hollow (Fig. 104, *d*).

Contracted socket may result from cicatricial bands or scar tissue due to injuries; or from irritation of a roughened glass eye, the lower fornix may become shallow; in either case an artificial eye will no longer stay in place. After thorough division of cicatricial bands, the conjunctival sac may be relined by a large skin-graft kept in place by a guttapercha mould, or by a pedunculated graft from adjacent skin. Sometimes an 'epithelial inlay' can be satisfactorily employed to enlarge the lower fornix. A dental wax mould, covered with a Thiersch graft, is buried from without in a cavity made in the lower lid beneath the fornix; when the graft has taken the cavity is opened from the conjunctival surface and the mould removed.

Kronlein's Operation is employed for the removal of innocent retro-ocular growths without sacrifice of the eye. It consists in incision down to the bone along the temporal margin of the orbit, separation of the periosteum from the temporal wall of the orbit, temporary resection and bending outwards of the bony wall, and replacement of the parts after removal of the tumour.

Exenteration of the Orbit is practised in early malignant diseases of the orbital contents. It consists in stripping the periosteum from the walls of the orbit and removing the whole contents up to the apex.

CHAPTER VIII

DISEASES OF THE CONJUNCTIVA

Anatomy.—The conjunctiva is a thin layer of mucous membrane which lines the eyelids, and is reflected on to the eyeball, forming a sac, the conjunctival sac. We distinguish three divisions: (1) The palpebral or tarsal conjunctiva, covering the under surface of the lids; (2) the ocular or bulbar conjunctiva, coating the anterior portion of the eyeball; and (3) the fornix, the transition portion, forming a fold between lid and globe. The conjunctiva differs somewhat in structure in each of these portions.

The *palpebral conjunctiva* is thicker than the other portions. In the greater part of its extent it is closely adherent to the subjacent tarsus, allowing the Meibomian glands to be seen through it. Its surface is smooth, but presents a number of minute projections, or papillæ. It is covered with cylindrical epithelium. Its stroma is of an adenoid character, containing a large number of lymph corpuscles, which sometimes project as small rounded nodules (lymphoid follicles) considered by some as normal and by others as pathological; the tubular depressions between these constitute Henle's glands. Numerous epithelial crypts are also found.

The *conjunctiva of the fornix* is similar in structure to that of the lids. It constitutes a very loose fold, ensuring great freedom of movement to the eyeball. It is richly supplied with bloodvessels. This and its lax condition explain its liability to marked swelling in inflammations of the conjunctiva. It has opening into it the lacrymal ducts and the accessory lacrymal glands.

The *bulbar portion of the conjunctiva*, thin and transparent, covers the anterior surface of the eyeball, being loosely attached to the sclera by connective tissue (episcleral tissue), with the exception of the margin representing the boundary between cornea and sclera (limbus), where it is firmly adherent. In structure it resembles the rest of the conjunctiva, but contains no glands. It is covered with laminated pavement epithelium, which is continued uninterruptedly over the cornea, and constitutes its outer layer. Near the inner canthus it forms a crescentic fold (*plica semilunaris*), the rudiment of the nictitating membrane or third eyelid of the lower animals.

The *vascular supply* of the conjunctiva is derived from the blood-vessels of the fornix—the posterior conjunctival (derived from the palpebral) and from the anterior ciliary. The latter pass forward along the recti muscles and pierce the sclera near the limbus to reach the interior of the eye, giving off one set of branches which form vascular loops surrounding the cornea, and supplying it with nourishment, and another set (anterior conjunctival) which pass backward in the conjunctiva and anastomose with the posterior conjunctival. This arrangement, together with the posterior ciliary arteries and the retinal system of vessels, constitutes the entire vascular system of the eye. Thus the bulbar conjunctiva presents two vascular systems—the posterior conjunctival and the anterior ciliary. The nature of the injection in any given case is of some value in locating the seat of the congestion.

The *nerves* of the conjunctiva, branches of the fifth, terminate in free ends or tactile corpuscles, especially in the palpebral portion. *Lymphatic vessels* are found in considerable numbers in the conjunctiva, forming a superficial and a deep layer.

Conjunctival and Ciliary Injection.

The differences between conjunctival and ciliary or circumcorneal injection (Figs. 105 to 107, Plate VIII.) are as follows:

Conjunctival Injection.

1. Derived from posterior conjunctival vessels.
2. Accompanies diseases of the conjunctiva.
3. More or less muco-purulent or purulent discharge.
4. Most marked in fornix conjunctivæ.
5. Fades as we approach the cornea.
6. Bright brick-red colour.
7. Composed of a network of coarse, tortuous vessels, anastomosing freely, and placed superficially, so that the meshes are easily recognized.
8. Can be moved with the conjunctiva by pressure on lower lid.

Ciliary Injection.

1. Derived from anterior ciliary vessels.
2. Accompanies disease of the cornea, iris, and ciliary body.
3. Often lacrymation, but no conjunctival discharge.
4. Most marked immediately around the cornea; hence called 'circumcorneal.'
5. Fades toward the fornix.
6. Pink or lilac colour.
7. Composed of small, straight vessels, placed deeply, so that the individual vessels cannot be recognized easily, but are seen indistinctly as fine, straight lines radiating from the cornea.
8. Cannot be displaced by movement of the conjunctiva.

PLATE VIII.




FIG. 105.—CONJUNCTIVAL INJECTION.




FIG. 106.—CILIARY INJECTION.




FIG. 107.—IRITIS WITH INTENSE CILIARY INJECTION.

The pupil has partially dilated under atropine.




FIG. 108.—SUBCONJUNCTIVAL HÆMORRHAGE.

In severe forms of diseases of the anterior part of the eye these two types of congestion are often found associated, as one would expect when one remembers that the two systems of vessels anastomose freely.

When very pronounced, particularly when there is much venous congestion, ciliary injection assumes a violet colour. A form of injection of this sort involves the episcleral tissue between the equator of the eyeball and the cornea, presenting a deeply placed, violet-coloured patch seen in scleritis and glaucoma.

CONJUNCTIVITIS.

Inflammations of the conjunctiva are known as conjunctivitis, or ophthalmia. The varieties are:

1. Catarrhal: (a) acute, (b) chronic, (c) follicular.
2. Purulent: (a) ophthalmia neonatorum, (b) gonorrhœal.
3. Membranous: (a) non-diphtheritic or croupous, (b) diphtheritic.
4. Granular or trachoma.
5. Phlyctenular.

Acute Catarrhal Conjunctivitis.

An acute catarrhal inflammation of the conjunctiva, accompanied by mucoid or muco-purulent discharge. It is also known as acute muco-purulent and acute simple conjunctivitis.

Objective Symptoms.—The palpebral conjunctiva and that of the fornix are of a brilliant red colour and swollen. There is slight congestion of the bulbar conjunctiva; but in severe cases this may become marked, and there may be added œdema of the bulbar conjunctiva (chemosis), small conjunctival hæmorrhages, and œdema of the lids. The secretion, increased in amount and altered in character, varies according to the severity. In mild cases, it is mucoid; in severer forms, it is muco-purulent. In very-marked examples the amount of pus may be so considerable that for a day or two we may be in doubt whether the disease is not the beginning of a

purulent inflammation. The secretion accumulates during the night, and dries upon the edges of the lids during sleep.

Subjective Symptoms.—Itching and smarting sensations referred to the lids, which feel hot, heavy, and as though sand or a foreign body were underneath. There is more or less photophobia. There may be some blurring of sight when the altered secretion lies upon the cornea. The symptoms are usually worse toward evening. They vary in severity with the degree of inflammation.

Course.—The patient may perhaps get well in two or three weeks, even without treatment. Sometimes the acute symptoms subside, and a chronic catarrhal conjunctivitis remains. Blepharitis may be present. In severe cases small, grayish infiltrations may form at the corneal margin. The coalescence of a number of these may cause a marginal ulcer, which is usually unimportant, superficial, and heals readily, but occasionally becomes deep and serious. Rarely iritis occurs as a complication.

Etiology.—The disease occurs at all ages and at all times during the year, but is most common in the spring and autumn. The causes may be divided into: (1) Mechanical—foreign bodies, exposure to wind, dust, smoke, etc. (2) Epidemic—in spring and autumn, and depending upon some atmospheric condition (the presence of certain micro-organisms). (3) Infection—through contact with fingers, towels, handkerchiefs, etc., of patients suffering from the disease. The discharge is contagious, especially when free and containing much pus; hence the affection often presents a number of examples in the same household or school. (4) Exanthemata, accompanying or following measles, scarlatina, and small-pox. (5) Associated with coryza, hay fever, and influenza.

Clinical Varieties.—Certain forms of this disease are distinguished by qualifying adjectives, indicating the etiology.

Traumatic Conjunctivitis is the name often given to acute catarrhal conjunctivitis when excited by the presence of a

PLATE IX.




FIG. 109.—GONOCOCCUS.




FIG. 110.—PNEUMOCOCCUS.




FIG. 111.—KOCH-WEEKS BACILLUS.




FIG. 112.—MORAX-AXENFELD
DIPLO-BACILLUS.




FIG. 113.—STREPTOCOCCUS.




FIG. 114.—DIPHTHERIA BACILLUS.

FIGS. 109 TO 114.—MICRO-ORGANISMS FOUND IN VARIOUS FORMS OF
CONJUNCTIVAL, LACHRYMAL, AND CORNEAL DISEASES.

foreign body or by traumatism. Under this head are included the forms of conjunctivitis due to intense light; for example, the electric arc light or that used in electric welding, and that produced by reflection from snow (snow blindness). In such cases there are symptoms of conjunctivitis, and, in addition, marked photophobia, lacrymation, intense smarting of the lids, contraction of the pupil, and sometimes œdema of the lids and superficial ulceration of the cornea.

Lacrymal Conjunctivitis (Fig. 63, Plate VII.) accompanies dacryocystitis. It is most frequently limited to the inner third of the palpebral and ocular conjunctiva. It is caused by the purulent secretion from the inflamed lacrymal sac, which contains streptococci (Fig. 113, Plate IX.).

Exanthematous Conjunctivitis, when associated with the exanthemata; most commonly seen in measles.

Acute Epidemic Conjunctivitis, popularly known as 'blight' or 'pink eye,' is a very contagious form of acute catarrhal conjunctivitis occurring most frequently in the spring and autumn in which the symptoms are apt to be marked and the discharge is profuse. The contagious element is the Koch-Weeks bacillus (Fig. 111, Plate IX.), or the pneumococcus (Fig. 110, Plate IX.). This is the variety which not infrequently gives rise to such severe objective symptoms, including swelling and redness of the lids and copious discharge, that one may be in doubt whether the affection is an example of the catarrhal or of the purulent form of conjunctivitis. It is advisable, when first called to treat severe cases of this sort, to examine a smear of the conjunctival secretion under the microscope, and to decide definitely upon the nature of the infection only after the responsible organism has been identified.

The *Morax-Axenfeld Diplo-bacillus* (Fig. 112, Plate IX.) causes a subacute conjunctivitis (angular conjunctivitis), which is more common in adults. It soon passes into the chronic form.

Other Clinical Varieties of acute catarrhal conjunctivitis have been classified according to the micro-organism which seems responsible for the inflammation. One form, which

clinically resembles acute epidemic conjunctivitis, presents an abundance of pneumococci (Fig. 110, Plate IX.) in the discharge. In another variety, streptococci (Fig. 113, Plate IX.) are found in considerable numbers. Staphylococci, xerosis bacilli, and diplococci, resembling in appearance the pneumococci, are usually present in the normal conjunctival sac, so that it is difficult to classify clinical varieties according to bacteriological findings.

Follicular Conjunctivitis is considered by some authorities as a variety of acute catarrhal conjunctivitis. It will be described separately.

Treatment.—Though the disease tends eventually to get well without interference, treatment reduces the duration and prevents the change into chronic conjunctivitis. The conjunctival sac should be irrigated several times a day with salt solution (3i. to Oi.) or saturated solution of boric acid, or with perchloride of mercury lotion (1 in 20,000, gradually increased to 1 in 10,000). When there is abundant tenacious discharge frequent irrigation with warm hydrogen peroxide lotion (1 to 2 volumes) may be used with advantage. A bland ointment (vaseline or boric-acid ointment) is applied to the edges of the lids at night, to prevent them from becoming glued together during sleep. A 10 per cent. solution of protargol or argyrol may be applied to the everted conjunctiva daily, as long as the discharge is abundant. The patient should be cautioned concerning the contagiousness of the discharge.

If the disease shows a tendency to become obstinate or chronic, weak astringent solutions are indicated (alum, zinc chloride or sulphate $\frac{1}{2}$ grain in 3i.) accompanied with an occasional application of a 1 per cent. silver nitrate solution or the alum-stick to the everted lids.

Chronic Catarrhal Conjunctivitis.

A chronic catarrhal inflammation of the conjunctiva, presenting similar symptoms to those found in the acute form, but associated with only slight changes in the quantity and quality of the normal secretion. It is also known as chronic simple conjunctivitis.

Objective Symptoms.—The conjunctiva of the lids is reddened and smooth. In old cases it may be hypertrophied and velvety. The secretion is usually but slightly altered, and there is very little increase; there may be enough to make the eyelids stick together in the morning or to present some dried secretion at the inner canthus. There is apt to be some excoriation at the outer angle (angular catarrh). In some cases there appears to be less than the normal amount of secretion (dry catarrh).

Subjective Symptoms are the same in kind as in the acute form: Itching, burning, dryness, feeling of sand or foreign body, heavy feeling in lids, some sensitiveness to light, and the eyes tire easily. These symptoms are worse at night.

Course.—The disease is probably the most common of ocular affections. It usually occurs in adults and frequently in old persons. It is apt to be of lengthy duration.

Complications.—Blepharitis is frequently present. Eczema of the lower lid and eversion of the inferior punctum occur not infrequently, and sometimes ectropion and corneal ulceration.

Etiology.—It may be the sequel of an acute catarrh. It may be caused by improper hygienic surroundings, vitiated atmosphere (overcrowding), irritating atmosphere (smoke dust), insufficient sleep, late hours, alcoholic excesses, exposure of the conjunctiva in ectropion, eye-strain, overuse, local irritation, such as trichiasis, chronic dacryocystitis, etc. It is usually bilateral, but when due to local irritants it may be unilateral.

Treatment.—We must endeavour to remove the cause of the inflammation. Locally: Astringent solutions (zinc, borax, alum, tannic acid, silver nitrate $\frac{1}{10}$ grain, or $\frac{1}{8}$ to $\frac{1}{2}$ i.); ointments of the yellow oxide and ammoniated mercury; silver nitrate, 1 per cent., brushed on the everted lids once a week, or the occasional use of the alum or sulphate of copper stick; bland ointments to the edges of the lids at night to prevent gluing together and excoriations. As in all chronic catarrhal affections, the remedies must be changed from time to time.

Conjunctival Concretions are sometimes seen, especially in elderly persons, appearing as small, bright yellowish, or whitish points beneath the conjunctiva; they consist of accumulations in the tubular depressions in the conjunctiva (Heale's glands) and are composed of amyloid material and crystals; the condition is sometimes called *lithiasis conjunctivæ* but the concretions do not contain calcium; sometimes they project through the surface, acting like foreign bodies and causing discomfort; under such circumstances they should be picked out with the point of a needle or knife after surface anæsthesia.

Angular Conjunctivitis is a subacute or chronic inflammation caused by the Morax-Axenfeld diplo-bacillus. It occurs usually in adults. There is redness and perhaps swelling of the palpebral conjunctiva and lid margins, especially at the angles (hence the name). The patient complains of smarting and itching and a feeling as though there were sand in the eye. On waking in the morning the lashes may be glued together with a scanty tenacious discharge. Collyria of zinc sulphate (gr. i. to ii. in 1 ounce) or chloride ($\frac{1}{2}$ grain in 1 ounce) usually acts almost as a specific. In obstinate chronic cases a useful prescription is hydrarg. oxycyanid. gr. i.ss., zinci sulph. gr. x., aq. distil. $\mathfrak{Z}$ xx., to be used in an eye-bath three or four times a day.

Collyria.—For the proper methods of using collyria see p. 455. This is of great importance, as the want of success in treating conjunctival affections is often due to these remedies not being properly applied.

Follicular Conjunctivitis.

This disease, also known as follicular catarrh, may be regarded as an obstinate form of catarrhal conjunctivitis, with the occurrence of 'follicles' upon the lower lid (Fig. 119, Plate X.).

Objective Symptoms.—In addition to the appearances found in catarrhal conjunctivitis, the conjunctiva of the lower lid presents a variable number of small, pale, round granules, about the size of the head of a pin. If many are present, they may be arranged in rows. They are most

abundant in or near the fornix. Occasionally some are found on the upper lid. These follicles consist of small masses of adenoid tissue, in which respect the microscopic appearance resembles that of trachoma.

Subjective Symptoms are identical with those of catarrhal conjunctivitis. In many cases patients do not complain of any symptoms, and the existence of the granulations is discovered accidentally.

Course.—The disease may be either acute or chronic. In either case the course is obstinate; in chronic cases the follicles may persist for months, and even years. It is sometimes difficult, especially in acute cases, to differentiate between follicular catarrh and granular conjunctivitis, and we may have to await the results of several weeks' treatment in order to decide definitely. The follicles, in this disease, disappear after a time, leaving the conjunctiva in a natural condition, and they affect principally the lower lid. In trachoma, on the other hand, there are permanent changes in the conjunctiva, and when the granulations are confined to one lid it is usually the upper which is involved.

Etiology.—It occurs most frequently in children and in young persons; it is often found in schools and asylums. The exact cause is not definitely known. Contagion seems responsible in some cases. Poor hygienic surroundings especially indoor life, seem to predispose. The various causes of catarrhal conjunctivitis act as exciting factors, as does the prolonged use of atropine and eserine.

Treatment.—The same as that given for acute and chronic catarrhal conjunctivitis. It is of especial importance to correct any interference with the general health, and to place such children under the best hygienic surroundings. Locally, the ointment of the yellow oxide of mercury, applied within the conjunctival sac, is a favourite remedy. The occasional use of a 1 per cent. solution of nitrate of silver may be of service. When the patient no longer complains of any symptoms and the follicles persist, they may be allowed to remain, and treatment discontinued.

Purulent Conjunctivitis.

An acute purulent inflammation, due to contagion from gonorrhœal virus. The contagious elements are the gonococci (Neisser). They occur in gonorrhœal secretion, are found in the pus-cells and conjunctival epithelium, and are arranged in pairs (diplococci) and generally in colonies (Fig. 109, Plate IX.). The disease is also known as acute blennorrhœa of the conjunctiva.

Clinical Varieties.—(1) Adult purulent conjunctivitis or gonorrhœal ophthalmia or conjunctivitis; (2) infantile purulent conjunctivitis or ophthalmia neonatorum (occurring in the new-born).

Gonorrhœal Ophthalmia or Adult Purulent Conjunctivitis.

Symptoms.—*First Stage, Infiltration.*—After a period of incubation varying from a few hours to two or three days (short in severe cases), there occur great swelling and redness of the lids, so that the latter cannot be opened voluntarily, and can be separated only with difficulty. The conjunctiva of the lids and fornix is intensely swollen and reddened, and is uneven. There is chemosis (œdema of the ocular conjunctiva, causing it to swell up around the cornea). The secretion is at first serous, somewhat coloured with blood, and containing a little pus. The eye is tender to touch. The patient complains of a hot, smarting pain in the eye, and a dull aching in the brow and temple. There is some constitutional disturbance, slight fever, and some swelling of the pre-auricular gland. This stage lasts about two days, and is followed by the

Second Stage, the Stage of Purulent Discharge.—The swelling of the lids and conjunctiva and the chemosis diminish, and the eye becomes less tender. A very profuse, purulent discharge appears, and escapes continually from between the lids. This condition continues for two or three weeks, all symptoms gradually diminishing.

Third Stage, Convalescence, or Papillary Swelling.—The eye may return to a normal condition in two or three weeks. More frequently, however, there is a stage of papillary swelling,

a chronic inflammation of the lids, the palpebral and retro-tarsal conjunctiva remaining thickened and red, and presenting, especially over the tarsus, an uneven granular or velvety appearance, with hyperæmia of the ocular conjunctiva.

Course.—The disease occurs in various degrees of severity. Cases in which there is slight infection, or in which the disease is acquired from a chronic gonorrhœa (gleet), are the mildest. The very intense cases have probably been acquired through contagion from the secretion of a very virulent gonorrhœa, and especially from contamination during the early stages.

Etiology.—The disease is always acquired through infection from gonorrhœal secretion, either directly, the fingers of the patient transferring the virus from the genitals, or indirectly by means of contaminated towels, etc. It is not often seen nowadays in Britain.

Complications.—A very frequent and important complication is corneal ulceration. This begins with a circumscribed grayish infiltration, becoming yellow and breaking down, so that ulcers are formed. The ulcers vary in situation, size, and course. They may be central or marginal; the latter may be confluent, so as to form an annular ulcer. The ulcers may perforate, and this be followed by cicatrization with or without incarceration of the iris, staphyloma, and other sequelæ of corneal ulceration. Panophthalmitis may result. Severe and early involvement of the cornea is most common in intense attacks. In such cases serious and permanent damage to the eye is very common.

Prognosis depends upon the severity of the case, and upon the behaviour of the cornea. It is always grave.

Treatment—*Prophylactic.*—Great precautions must be observed to prevent infection of the eyes of the physician, nurse, and attendants through spurting of the discharge




FIG. 115.—CHEMOSIS OF THE CONJUNCTIVA.

during examination or treatment. Protecting glasses should be worn whenever exposed to this risk. Contaminated fingers must be carefully disinfected. Materials which have been used for cleansing the eye must be burned.

The non-affected eye should be protected from infection by the application of Buller's shield (Fig. 116). This consists of a watch-glass, securely held in place by adhesive plaster applied to the side of the nose, the cheek, and forehead. The junction of skin and plaster is sealed by a layer of collodion. The centre of the glass is left uncovered by plaster, to permit inspection of the eye, and a small part of the outer margin of the watch-glass is usually left free in order to allow air to enter and contribute to the comfort of the shielded eye.




FIG. 116.—BULLER'S SHIELD.

Treatment of the First Stage.—An injection of gonococcal vaccine or a foreign protein (e.g., 5 c.c. sterilized milk) into the glutei should be given as soon as possible. Locally iced compresses are used continuously day and night. The eye must be carefully

cleansed, and the secretion removed as rapidly as it forms. When very abundant, this will be necessary every quarter or half an hour. For this purpose a saturated solution of boric acid is most frequently employed, being allowed to trickle in between the lids from a piece of absorbent cotton dripping with the solution. Then the secretion which has been washed out is gently wiped off the margins of the lids.

The iced compresses may be used continuously at first. But when the tense, reddened, and swollen condition of the lids becomes less marked, the application of cold must be reduced to every other hour, or every third hour. Too much refrigeration interferes with the nutrition of the cornea. When the cornea is involved, we must carefully gauge the amount of cold so as not to use an excess. In the *later stages*,

when there is little swelling, and corneal infiltration or ulceration exists, hot applications may be used, in order to improve nutrition by stimulating the flow of blood to the part.

Other cleansing and antiseptic solutions may be tried: mercury perchloride (1 : 20,000 to 1 : 10,000), hydrogen peroxide (1 volume), permanganate of potassium (1 : 500), etc. The instillation of a few drops of a 25 per cent. solution of argyrol or protargol every three hours is of value.

In the initial stage, if the disease be very severe, from three to six leeches, applied to the corresponding temple, may be of service. Occasionally there is so much tension that the eye cannot be cleansed, on account of the difficulty in separating the lids, and in addition harmful pressure is exerted upon the eyeball. In such cases it may become necessary to widen the palpebral fissure by a division of the external canthus (temporary canthotomy, p. 66).

Treatment of the Later Stages.—When the lids have lost their swollen and angry appearance, and the discharge begins to diminish, a 1 or 2 per cent. solution of nitrate of silver is brushed upon the everted conjunctiva once a day. This may be done even though the cornea be implicated. It is continued until the patient is well, or until the papillary swelling has persisted for some time. After each application the eye should be irrigated with warm water to remove any excess of caustic, lest the cornea be attacked. Then, if silver no longer exerts a favourable influence, we may apply glycerole of tannin (5 to 10 per cent.), the alum-stick, or sulphate of copper pencil once a day.

The treatment of corneal complications resembles that of infected corneal ulcers, and is described in the next chapter.

Ophthalmia Neonatorum or Infantile Purulent Conjunctivitis.

An acute purulent conjunctivitis occurring in the newborn within the first three weeks of life, presenting similar symptoms, complications, and course, and requiring the same treatment, as in the gonorrhœal ophthalmia of adults. It is notifiable in Great Britain (Fig. 118, Plate X).

Symptoms.—The period of incubation being the same as in adults, the first symptoms are usually noticed on the second or third day after birth. When the onset is later than the third or fourth day, infection has taken place subsequent to the birth of the child.

The symptoms are the same in kind as those of gonorrhœal ophthalmia, but are often less severe, and more apt to be limited to the palpebral and retrotarsal conjunctiva. Both eyes are usually involved. The cornea is implicated in a much smaller proportion of cases, especially if the affection is treated from the start. If the case is seen early, before the cornea is affected, and properly managed, this part very often escapes destruction or serious damage.

The prognosis, therefore, with early and proper treatment, is generally favourable.

Etiology.—Infection by gonorrhœal secretion from the genitals of the mother during parturition. In rare cases infection occurs before birth. Sometimes it occurs subsequent to the birth of the child, through infection from sponges, napkins, towels, or the fingers of the nurse, which have been in contact with the genitals of the mother.

Most cases are undoubtedly of gonorrhœal origin. Some cases, however, result through infection from simple catarrhal (non-gonorrhœal) secretion. Such cases run a mild course, and are not usually complicated by corneal ulcers.

Treatment.—The conjunctival sac should be irrigated every hour with warm normal saline solution or with warm solution of potassium chlorate (gr. x. ad ʒi.). Vaseline should be applied to protect the corneal epithelium. Applications of 1 per cent. solution of nitrate of silver may be used after the swelling and redness have diminished and the discharge begins to be less profuse, continued throughout the stage of papillary swelling. Once a day will be sufficient, followed immediately by warm-water irrigation. Other useful remedies are, in the acute stage irrigation every hour with warm H_2O_2 solution ($\frac{1}{2}$ volume), and in the later stage zinc chloride drops ($\frac{1}{4}$ grain ad ʒi.) twice a day.

If there is much swelling of the lids, iced compresses may

be applied, but we must be careful not to use them too continuously, and they must be stopped as soon as the redness and swelling begin to diminish. Occasionally canthotomy is necessary. In adults, the sensations of the patient guide us to a certain extent, and we use the pads less often when they no longer feel grateful, as happens when the redness and swelling subside. In infants we cannot receive this information; hence great care must be used not to injure the cornea by excessive cooling, especially if there be corneal infiltration. In such cases hot compresses are often substituted for the cold.

Every day the cornea should be examined. If any ulcer be observed, warm lotions should be used every hour and atropine ointment ($\frac{1}{4}$ per cent.) thrice daily. Over-zealous treatment with too strong antiseptic lotions (especially perchloride of mercury) or silver nitrate solution, may cause serious injury to the cornea.

The general health of the infant must be looked after, since enfeebled conditions render treatment unsatisfactory, and favour corneal complications. Gonococcal vaccines are of no benefit.

Credé's Method of Prophylaxis.—Ophthalmia neonatorum is practically preventable. Through the adoption of Credé's method, its occurrence has been made infrequent in lying-in hospitals and in private practice. The method consists in cleansing the eyes of the child with water immediately after birth, and instilling one drop of a 2 per cent. solution of nitrate of silver into each eye. This often causes a slight redness of the conjunctiva for a day or two. It acts by destroying any gonococci which may have entered the conjunctival sac. Antiseptic irrigation of the vagina of the mother before delivery is also useful as a prophylactic measure.

Catarrhal Conjunctivitis in the New-Born.—Sometimes we meet with a slight catarrhal conjunctivitis in the new-born, lasting a few days, and presenting merely hyperæmia, slight swelling, and a little mucoid discharge. These are not examples of ophthalmia neonatorum. But at the start we may be in doubt whether they are not purulent cases, and it will be safer to treat them as such until the character of

the inflammation becomes certain. In such cases bacteriological examination of the discharge may furnish valuable information.

Membranous Conjunctivitis.

This variety is uncommon, attacks children, and occurs in two forms: (1) diphtheritic and (2) croupous. The subdivision is based upon anatomical differences since, in both varieties, the Klebs Loeffler bacillus may be present or absent in the secretion, and this has no relationship to the severity of the local symptoms; both may run a mild or a severe course.

In diphtheritic conjunctivitis the Klebs Loeffler bacillus is usually present, but not always, there are severe constitutional symptoms, and the membranous secretion which forms on the surface also penetrates the substance of the palpebral conjunctiva.

In croupous conjunctivitis there is commonly an absence of the Klebs-Loeffler bacillus, the constitutional symptoms are much less severe, and the secretion forms a membranous *layer upon the surface* of the palpebral conjunctiva, but does not invade its substance.

Diphtheritic Conjunctivitis.

An acute inflammation of the conjunctiva, associated with exudation and infiltration, purulent discharge, with tendency to necrosis of the involved tissues. The disease is rather rare and occurs in children. It spreads by infection. The secretion usually contains the Loeffler bacillus (Fig. 110, Plate IX.) and is contagious.

Symptoms.—The lids are very much swollen, reddened, hot, and tender. The conjunctiva of the lids and fornix is intensely inflamed and is covered by a grayish-yellow exudation, which also infiltrates the underlying tissues. In this way the lids become hard and cannot be everted. The exudation causes compression, and, as a result, there is a tendency to sloughing of the infiltrated parts. Besides this fibrinous exudation, there is a discharge of a thin, cloudy fluid. With these local signs there are the prostration and

other constitutional symptoms of diphtheria, and often local evidences of the disease in other parts of the body.

At the end of a week the exudation disappears, partly through absorption, partly through necrosis and sloughing, causing a loss of substance covered by granulations. The secretion now becomes more abundant and purulent.

The defects in the lining of the lid gradually cicatrize, this process causing various deformities: symblepharon, trichiasis, and entropion. There is frequently corneal ulceration. When the diphtheritic process is severe, the cornea is seriously involved and sight is always lost.

The *prognosis* in regard to sight is always serious; in regard to life, it depends upon the constitutional effects and general condition of the child.

Treatment.—*Prophylactic.* The precautions described under gonorrhœal ophthalmia must be employed in this disease to protect physician, nurse, and attendants. Besides being contagious, the disease is infectious; hence the patient should be isolated; other children must be removed. The second eye must be shielded.

Treatment of the Affected Eye.—Careful cleansing with weak antiseptic solutions (boric acid, corrosive sublimate). Cold compresses may be applied, but must be used cautiously on account of the enfeebled circulation. After a short period hot compresses are used. When the exudation has separated, we apply a 1 per cent. solution of nitrate of silver. We endeavour to prevent sequelæ due to cicatrization by frequent separation of the lids from the globe, and by keeping the two surfaces apart by a piece of fine linen smeared with some bland ointment. Corneal ulceration must be treated as described in the next chapter.

Constitutional.—We must remember that the eye affection is merely the local manifestation of a constitutional disease. Hence the general treatment of diphtheria must be carried out. Injections of antitoxin are of great value, producing a rapid improvement in the local as well as the general condition.

Croupous Conjunctivitis or Non-diphtheritic Membranous Conjunctivitis.

A form of inflammation in which there is the deposit of an exudation upon the surface of the conjunctiva, upon which it hardens into a membrane. There is no infiltration into the tissues; this constitutes the essential difference between croupous and diphtheritic conjunctivitis. Bacteriological examination may reveal the Loeffler bacillus and other micro-organisms, and these may be identical with those found in diphtheritic membrane. There are, however, no constitutional symptoms such as accompany diphtheria.

Symptoms.—Those of catarrhal conjunctivitis. A fibrinous membrane forms upon the palpebral conjunctiva; when this exudation is pulled off, a raw surface presenting a few bleeding points is seen; under such circumstances the membrane re-forms.

Etiology.—It may be caused by irritants (mechanical, chemical, or thermic), or by the action of various micro-organisms. Examples of the former are nitrate of silver, acids, lime, molten lead, burns, and injuries in general.

Treatment.—That of catarrhal conjunctivitis. As soon as the membrane shows no tendency to re-form, applications of a 1 per cent. solution of nitrate of silver are useful.

Granular Conjunctivitis, Trachoma, or Granular Lids.

An inflammation, generally of lengthy duration, accompanied by hypertrophy of the conjunctiva and the formation of 'granules,' with subsequent cicatricial changes. It occurs at all ages. It is not common in England except among immigrants from Eastern Europe. There is more or less secretion, which is contagious. It is a very important affection on account of its disastrous complications and sequelæ, which are responsible for many cases of partial or total blindness.

Subjective Symptoms.—More or less photophobia, lachrymation, itching, and burning sensations, feeling of foreign




FIG. 117.—CATARRHAL CONJUNCTIVITIS.



FIG. 118.—OPHTHALMIA NEONATORUM.




FIG. 119.—FOLLICULAR CONJUNCTIVITIS.




FIG. 120.—TRACHOMA.




FIG. 121.—PHLYCTENULAR CONJUNCTIVITIS.




FIG. 122.—EPISCLERITIS.

body, pain, and visual disturbance. In a good many cases there are no subjective symptoms.

Objective Symptoms.—Swelling of the lids, narrowing of the palpebral aperture, and drooping of the upper lid (from weight and swelling). There is a variable amount of mucopurulent discharge, marked in recent cases, scanty in chronic forms. The conjunctiva of the tarsus and fornix is reddened, thickened, and uneven, on account of hypertrophy and the occurrence of granules. The ocular conjunctiva is often somewhat injected.

Forms.—Various forms of trachoma have been described; these represent either (1) the various stages of simple trachoma or (2) secondary infections of trachoma. So we distinguish three forms of simple trachoma: (1) papillary, (2) granular, and (3) mixed.

1. *Papillary Form.*—A large number of small elevations (*papillæ*) are seen upon the greatly thickened conjunctiva, giving the latter a velvety appearance, or, if the *papillæ* are larger, a granular aspect. This form affects only the tarsal conjunctiva, and usually only the upper lid. The *papillæ* are caused by the hypertrophied conjunctiva being thrown into folds, covered by an increase in epithelium, the connective-tissue interior being infiltrated with cells. This is the *first stage* of simple trachoma.

2. *The Granular Form* presents a preponderance of trachoma granules (Fig. 120, Plate X.). These are grayish or yellowish, rounded, translucent bodies showing through the conjunctiva. They may be small and rounded, larger and warty, or flattened and succulent. They are present principally in the fornix, and when numerous are arranged in rows. In the tarsal conjunctiva they are less numerous, smaller, and less distinct, being hidden by the *papillæ*. Occasionally trachoma granules are formed upon the semilunar folds and the bulbar conjunctiva. The granules are rounded collections of lymph corpuscles in a connective-tissue reticulum, resembling what we see in Peyer's patches in the intestines; they may present an incomplete capsule in old cases. This is the *second stage* of simple trachoma.

3. These two stages are commonly seen together in the 'Mixed Form,' but may occur separately; papillæ are more prominent in palpebral conjunctiva, granules in the fornices.

The *third stage* is that of cicatrization, which is also often combined with the two previous stages. This cures the trachoma, and the papillæ and granules disappear; but the conjunctiva does not return to a normal condition, the cicatricial changes and contraction leading to certain sequelæ; the seriousness of the latter depends upon the severity of the process and the amount of hypertrophy and subsequent cicatrization. In the tarsal conjunctiva the cicatricial process causes narrow whitish bands and scars (Fig. 135, Plate XI.), sometimes a network; in advanced and severe cases the entire surface may be replaced by a pale, smooth cicatricial membrane. In the fornix cicatrization changes the conjunctiva into a pale, bluish-white membrane, and as a result of contraction the transition fold is shortened or disappears.

The *fourth stage* is that of complications due either to invasion of the cornea by the trachoma or to contraction of the scar tissue in the lids and conjunctiva (see below).

Clinical Varieties.—Clinically, trachoma presents a number of varieties due to secondary infections. Occasionally the invasion is acute and accompanied by marked inflammatory symptoms and profuse purulent discharge; such cases resemble purulent conjunctivitis. The finding of gonococci in the secretion and the presence of the trachoma granules serve to differentiate, but frequently the swelling hides the latter; we may have to wait several days, until the swelling subsides somewhat, before we can decide.

Simple trachoma begins insidiously; it may exist unknown for months, before the subjective symptoms become annoying. Most cases of trachoma are chronic in their course, and the duration is months or years.

So, clinically, two forms of trachoma are met with: (1) inflammatory trachoma, with inflammatory symptoms, very contagious, leading to cicatrization of the conjunctiva and various sequelæ when unchecked; and (2) simple or non-inflammatory trachoma, in which, with moderate or marked

deposition of granulations of the granular form in the palpebral and retrotarsal portions of the conjunctiva of both lids, there are but slight or no symptoms of irritation or discomfort.

Besides these differences in the intensity of the inflammatory symptoms there are great variations in the amount of change in conjunctiva and cornea. There are mild cases, in which there are but little hypertrophy and insignificant cicatricial changes in the conjunctiva, so that afterward we can scarcely be sure that trachoma has existed; such mild cases usually remain free from corneal complications.

In moderate and severe cases there always remain permanent cicatricial changes, which enable us to diagnose the previous existence of trachoma. When the cornea is implicated the case is always a serious one.

Trachoma does not always progress uninterruptedly; there are often intermissions and exacerbations. Relapses are quite frequent, especially when treatment has been discontinued too soon.

Complications.—The most frequent are pannus and corneal ulceration, both causing disturbance of sight.

Pannus consists of a newly formed vascular tissue, which usually covers the upper part of the cornea (Fig. 135, Plate XI.). The affected portion of the cornea presents a cloudy appearance, and is grayish and translucent; its surface is uneven and vascularized, the bloodvessels springing from the conjunctival vessels at the limbus. The process advances until it covers the upper half of the cornea. Finally, the entire cornea may be covered, in which case vision is reduced to perception of light. Unless subsequent changes occur, complete retrogression is possible, so that the cornea can become transparent again. In marked cases iritis is apt to develop. Pannus is not merely due to mechanical irritation, but to a change similar to that which occurs in the conjunctiva.

Ulcers of the cornea occur with or without pannus. They leave opacities, which interfere with vision according to their seat and density.

Sequelæ.—Complete cure is usually effected in the mildest

cases alone, or in more severe forms only when they are subjected to early treatment. Sequelæ are very common, affect the conjunctiva, cornea, and lids, and produce permanent disability of the eye.

1. Trichiasis and entropion occur as a result of cicatricial contraction of the conjunctiva with curving of the tarsus; they are more pronounced in the upper lid. As a result of this distortion of the lid with consequent changes in the position of the cilia, there is mechanical interference with the cornea, causing ulceration.

2. Ectropion (usually of the lower lid) follows in some cases, as a result of hypertrophy of the conjunctiva and contraction of the orbicularis.

3. Symblepharon results from cicatricial contraction of the conjunctiva; when considerable, there is obliteration of the fornix. This condition restricts the movements of the eyeball.

4. Corneal opacities result from pannus and corneal ulcers. After lasting some time, pannus changes into a thin, permanent layer of connective tissue.

5. Staphyloma of the cornea follows in some cases.

6. Xerosis, a contracted, dry, and scaly condition of the conjunctiva, with changes in the cornea, may occur in very severe cases.

Etiology.—Trachoma is caused by contagion from another eye, being transferred through the secretion. The danger of contagion depends upon the amount of secretion in any given case. The transfer from one eye to another may take place by the finger, but usually by towels, handkerchiefs, and the like, which are used in common by many persons. Hence the disease spreads most extensively in schools, asylums, and barracks, and among people who live crowded closely together, and who are careless in regard to cleanliness. It is scarcely ever seen in private practice in England. It is common in Russia, Poland, Hungary, Japan, and China; also Ireland. It occurs with especial frequency in certain countries—Arabia, Egypt; it is endemic in the latter country. During the Napoleonic wars the affection was carried to Europe by soldiers (hence often called Egyptian

ophthalmia). In Europe it occurs much more extensively in the east than the west, and much more frequently in low lands (Belgium, Holland, Hungary) than in elevated countries (Switzerland). Negroes are said to be comparatively immune to the disease. The causation is still unknown, but the contagious principle in the secretion is thought to be a micro-organism. Noguchi isolated an organism, *bacterium granulosis*, from cases of trachoma with which he was able to produce trachoma in monkeys, but the evidence that this disease can be transferred to the human subject by tissue substance of experimental trachoma of monkeys is contradictory and inconclusive.

Pathology.—There is nothing characteristic. The adenoid stroma of the conjunctiva is infiltrated with lymphocytes. The papillæ are caused by hypertrophy of the conjunctiva thrown into folds with intervening depressions, covered by an increase in epithelium. The granules are round collections of lymphocytes, with a few phagocytes in the interior, in a delicate connective-tissue reticulum which merges with the surrounding tissues; the structure resembles that of Peyer's patches in the intestines, adenoids present in the throat, and is practically the same as that of the follicles in follicular conjunctivitis. In later stages stroma and cells tend to undergo hyaline degeneration, and there is the formation of fibrous tissue which forms an incomplete capsule for the granules and also gives rise to cicatricial bands.

Treatment consists in an attempt to reduce the inflammatory symptoms and secretion, and to check and remove hypertrophy of the conjunctiva, thus shortening the duration and diminishing the liability to conjunctival cicatrization and to sequelæ. This is accomplished either by the use of certain irritating applications, or by mechanical (surgical) means.

Irritating Applications.—The conjunctiva may be rubbed thoroughly with a solution of mercuric chloride (1 : 500) upon cotton-wool, wound on a glass rod. Sulphate of copper in the form of a crystal or pencil is the favourite local application. Nitrate of silver (1 or 2 per cent. solution), glycerole of tannin (5 to 25 per cent.), and the alum stick are also employed.

Surgical treatment includes expression, grattage, excision, curetting, electrolysis, X rays, radium, and galvano-cautery. Expression is the most popular of these mechanical methods, and has the widest range of usefulness. The kind of treatment best suited for trachoma depends upon the nature of the affection, the presence or absence of inflammatory symptoms, and the stage of the disease. Mechanical treatment is indicated in the granular and mixed forms of trachoma, with well-marked translucent granulations, when there is an absence of severe inflammatory symptoms; it is particularly useful in the form which Knapp calls simple or non-inflammatory. Irritating applications are indicated as supplementary treatment to surgical procedures, and for cases of chronic trachoma, in which the granulations are of small size, or of the papillary variety, particularly when there is considerable thickening of the conjunctiva.

In acute forms and in acute exacerbations of chronic cases, when there is much discharge, solution of nitrate of silver, 1 or 2 per cent., is applied to the conjunctiva, the excess being washed away with water or salt solution. In



FIG. 123.—SULPHATE OF COPPER STICK.

many cases of this sort, however, it is often advisable to suspend temporarily all irritative treatment, and to prescribe cold compresses, instillations of a 25 per cent. solution of argyrol, and mild cleansing and antiseptic washes.

During the cicatricial stage copper is no longer indicated; the ointment of the yellow oxide of mercury is then of service.

If treatment is not continued until every trace of hypertrophy has disappeared, relapses are very common.

Sulphate of Copper.—The pencil is applied to the everted lids once a day, or every other day; it is drawn lightly across the conjunctiva two or three times, but applied only to the hypertrophied portions. The application should include the palpebral portion of the transition fold of the upper lid; in

passing the copper stick under the tarsus the cornea is protected by the lower lid (Fig. 126). The stick of copper sulphate should have a flat, blunt end, as shown in Fig. 123, and not be pointed or conical. After each application, the excess of copper sulphate is washed off with water or solution of boric acid; subsequently iced compresses are applied for half an hour or longer. This treatment is continued for months until every trace of hypertrophy has disappeared. After a time the applications are made more lightly and less frequently.

Expression is best performed, in severe cases with many follicles, with Knapp's roller-forceps,




FIG. 124.—KNAPP'S ROLLER-FORCEPS FOR TRACHOMA (FULL SIZE).

by means of which the granulations are squeezed out between two fluted rollers at the end of the shafts (Fig. 124). The operation is painful, and a general anæsthetic is required. The upper lid is everted, and the trachoma follicles are squeezed out between the two extremities of the forceps. One extremity is passed back into the fornix and the other over the tarsus. Using moderate compression, the forceps is drawn forward, pressing out the contents




FIG. 125.—TYRREL'S FORCEPS.

of the granules (Fig. 118). This procedure is repeated until the lid is free from granulations and presents a dark-red surface with small red points. The lower lid is then operated upon in the same manner. After expression the conjunctiva is often brushed vigorously with a solution of mercuric bichloride, 1 : 500. Care must be taken not to cause abrasions of the cornea and not to tear the conjunctiva. If the granulations are hard and horny, it may be well to scarify them

before using the roller-forceps. There are swelling and ecchymosis for a day or two after the operation, but no other evidences of reaction. Cold compresses and irrigations with solution of boric acid are indicated for several days; then any remaining roughness is treated with gentle applications of the sulphate of copper crystal every other day for a few weeks, or until the lids are normal.

When granulations are scanty, or when they are situated on




FIG. 126.—METHOD OF APPLYING THE SULPHATE OF COPPER STICK TO THE CONJUNCTIVA OF THE UPPER LID.

FIG. 127.—THE OPERATION OF EXPRESSION FOR TRACHOMA AS APPLIED TO THE UPPER LID.

the caruncle or plica where roller-forceps cannot reach them, they are best expressed with Tyrrel's forceps under local anaesthesia. Single granulations may thus be dealt with, and the process readily repeated, even in children, as required.

The other mechanical or surgical means of treating trachoma are used much less frequently than expression. *Grattage* consists in scrubbing the granulations, with or without previous scarification, with a stiff tooth-brush until all the granules

are removed, and then thoroughly rubbing in a solution of mercuric bichloride, 1 : 500. *Excision* consists in the removal of the fold of conjunctiva, about 10 mm. broad, containing the granules. Both of these methods cause more injury to the conjunctiva than is the case with expression. Some surgeons recommend exposure of the everted conjunctiva to the influence of X rays or of radium. Either of these agents may cause much damage in unskilful hands. Carbon dioxide snow is useful in dealing with many follicles in upper tarsus—applied for thirty seconds, and repeated two or three times at a week or ten days' interval.

Treatment of Complications.—Recent pannus is best relieved by the treatment of the conjunctiva. In addition, we may use atropine occasionally, so as to keep the pupil dilated and prevent posterior synechiæ, since iritis is frequently present in these cases. If the pannus is very dense, we may apply the copper directly to the cornea. The practice of rubbing jequiritol, or a 3 per cent. infusion of the jequirity seed, into the everted conjunctiva, which was at one time advocated for such cases, has now been abandoned, as it is difficult to control the reaction, and destruction of the cornea has occurred in some instances. The operation of peritomy, the excision of a narrow strip of conjunctiva surrounding the cornea with a view of cutting off the vascular supply, is occasionally performed for the relief of severe cases of pannus. For active ulceration, nitrate of silver is often used, and atropine if iritis is suspected.

General treatment must not be neglected. The eye should be kept cleansed by the frequent use of solution of salt, boric acid, or bichloride of mercury (1 : 10,000). The hygienic surroundings of the patient should be made as perfect as possible, with proper ventilation, plenty of outdoor exercise, and good food.

Prophylaxis is very important. The patient and his family must be warned of the contagiousness of the secretion, and impressed with the necessity for keeping the patient's handkerchiefs, towels, wash-basin, etc., apart from those of other persons. In schools, asylums, institutions, and barracks,

the prevention of epidemics of trachoma is a very serious matter, requiring constant vigilance, careful inspection of every new addition or inmate, and the isolation of trachoma cases so long as the latter are capable of conveying the disease.

Parinaud's Conjunctivitis is a disease of the conjunctiva due to infection with a leptothrix; the source of infection is said to be the mouth of cats and possibly other domestic animals. It is usually limited to one eye. The local symptoms comprise gray or yellowish areas beneath the conjunctival epithelium, representing infiltration of the tissues with large mononuclear cells with surrounding tissue necrosis; with this there is a general enlargement of the lymph follicles; there is no ulceration. Accompanying these conjunctival changes is marked inflammatory enlargement of the preauricular gland with no tendency to suppurate. There is but slight constitutional disturbance, no pain and but little discomfort. Prognosis is favourable and cure results within several weeks. Treatment is that of conjunctivitis in general. The disease may have to be differentiated from tuberculous conjunctivitis.

Phlyctenular Conjunctivitis.

This disease, also known as pustular conjunctivitis and as scrofulous ophthalmia, is a circumscribed inflammation of the conjunctiva, accompanied by the formation of one or more small reddened projections called phlyctenulæ. The latter consists of accumulation of lymphoid cells, which soften at their apices, forming small ulcers. The phlyctenulæ may occur upon the ocular conjunctiva, and then the disease is called phlyctenular conjunctivitis. They may occur upon the cornea, when the affection constitutes phlyctenular keratitis; or they may occur, and most frequently do occur, at the limbus, and then we might speak of the disease either as a keratitis or as a conjunctivitis. Very frequently they occur in all three situations in the same individual. The pathology, symptoms, and treatment being the same in all cases, it is convenient to describe the three varieties collectively under the title of phlyctenular ophthalmia.

Objective Symptoms.—The essential sign is the occurrence of one or more small, grayish elevations or nodules, about the size of a millet-seed, at some part of the conjunctiva or cornea, frequently at the limbus. The phlyctenule is surrounded by an area of conjunctival hyperæmia (Fig. 121, Plate X.). The non-affected parts of the ocular conjunctiva are but slightly changed from the normal. The phlyctenule soon presents a small ulceration at its apex, which then occupies the level of the surrounding conjunctiva. It heals without leaving behind any changes in the conjunctiva. The entire process lasts from a few days to two weeks.

Generally a number of phlyctenulæ appear at the same time. In this manner the entire ocular conjunctiva may be reddened. In such cases the palpebral conjunctiva will be congested. The nodules may become absorbed without going through the stage of ulceration.

When the phlyctenule appears upon the cornea, the infiltrations and subsequent ulcers are usually superficial, and heal without the production of lasting changes in the cornea. But sometimes they spread into the corneal substance, and then leave a permanent opacity. In rare cases the ulcer perforates the cornea. Sometimes a number of them may, by confluence, spread along its surface.

Fascicular Keratitis.—The ulcer resulting from the phlyctenule may advance from the margin to the centre of the cornea, drawing after it a fascicle of bloodvessels. In this manner there is formed a narrow, red band of vessels, extending some distance over the cornea (Fig. 136, Plate XI.). At the apex of this fascicle is seen a small, gray crescent, corresponding to the advancing margin of the ulcer, which has healed in the peripheral parts. This form of ulceration always remains superficial. When the process terminates, the bloodvessels gradually disappear and a superficial linear opacity remains.

The phlyctenule may, in severe cases, involve the deep layers of the cornea, forming deep infiltration. This either becomes absorbed completely, or leaves an opacity of the cornea. It rarely becomes purulent, and a deep ulcer results.

There is usually considerable lacrymation. If there is any discharge, it is mucous or muco-purulent, and not abundant.

Subjective Symptoms.—Photophobia is marked when the cornea is involved, slight or absent in conjunctival cases. When prominent, there is considerable blepharospasm, and the eyes can be examined only with difficulty.

Course.—The phlyctenules usually occur in crops. Before one is completely cured another is apt to appear. In this way the course may become protracted, and may extend over weeks. Each phlyctenule lasts from a few days to a week or two. Relapses are very common. The affection occurs most frequently in children and in young persons, but is also seen in adults. In adults a single large phlyctenule often gives the local appearances of episcleritis.

As a result of constant lacrymation, there are frequently added blepharitis, excoriations, and eczema of the lids.

The prognosis is favourable. Serious results are rare. The phlyctenulæ usually leave no traces. In some cases corneal opacities of greater or lesser density remain, and if these are central, sight will be interfered with. Frequent recurrences may result in a number of cloud-like opacities of the cornea with superficial bloodvessels.

Etiology.—The disease is very common. It seems dependent upon some constitutional error. It occurs frequently in children who suffer from the tuberculous or so-called scrofulous diathesis. It is especially frequent among the lower classes, in whom dirt, poor food, and improper hygienic surroundings are contributory factors; also in children debilitated from disease. One frequently sees other manifestations of the pre-disposing diathesis, such as swelling of the cervical lymphatic glands, adenoid vegetations, eczema, coryza, blepharitis, chronic otorrhœa, etc. Sometimes, however, the affection occurs in children of the better classes, apparently in good health.

Pathology.—Phlyctenules begin with an infiltration of the sub-epithelial layer of the conjunctiva and of the cornea respectively with many mononuclear leucocytes, to which polymorphonuclear leucocytes are added in some cases, with

no changes in the epithelial covering at first, except infiltration with leucocytes; this forms a nodule somewhat raised above the surface. When the hyperplasia of cells is pronounced, necrosis takes place, the nodule bursts and an ulcer is formed. There is an abundance of staphylococci and, with these, the micro-organisms present with catarrhal conjunctivitis. During reparation, in many cases when the infiltration is superficial, the defect is replaced with epithelium; in cases of deeper involvement, with scar tissue.

Treatment.—*Local.*—Finely powdered washed calomel dusted upon the eyeball once a day. This is believed to be slowly changed to corrosive sublimate by the action of the tears, and in this way to keep the eye bathed in an antiseptic fluid. The calomel is apt to cause irritation if the patient is taking iodides internally, probably owing to local formation of mercuric iodide. A favourite remedy is the ointment of the yellow oxide of mercury (1 or 2 per cent.); a piece about the size of a hemp-seed is deposited in the conjunctival sac and rubbed about with the lids. When there is a great deal of irritation, it is wise to withhold the yellow oxide ointment until less inflammation exists. If the symptoms of irritation are very prominent, it is better to irrigate with solution of boric acid, and to apply cold pads if the phlyctenulæ involve the conjunctiva, and hot compresses if they form upon the cornea.

If there is infiltration or ulceration of the cornea, atropine, hot compresses, and mild antiseptic washes are indicated. If there is fascicular keratitis, the ointment of the yellow oxide of mercury is employed. In such cases we can often cut short the progress of the disease by cauterizing the advancing edge of the ulcer with a fine galvano-cautery point (Fig. 138), or with tincture of iodine (p. 149). Bandages should not be applied. It is only in extreme cases of very deep ulceration that a bandage is indicated.

In corneal cases the photophobia and blepharospasm are often very annoying symptoms. Instillation of solution of holocain will give temporary relief. Douching the eye with cold water several times a day may be effective. If a fissure of the outer canthus is present, touching this with a 2 per

cent. solution of silver nitrate or the stick of copper sulphate is of value.

General treatment is of great importance. Suitable and nourishing diet, improved hygienic surroundings, and cold sponging and bathing are useful. The nose and nasopharynx should receive proper treatment. These patients should not be allowed to remain in the house and in the dark; as they are inclined to do on account of the photophobia. Smoked glasses are prescribed to relieve this symptom. Many cases benefit markedly from exposure to ultra-violet light. Preparations of iron (syrup of the iodide), quinine, and arsenic are useful for internal administration, and cod-liver oil is of great benefit.

Spring Catarrh.

A rather uncommon disease of the conjunctiva, of chronic course, lasting for years, recurring during warm weather and disappearing entirely or to a great extent in winter. It affects both eyes and is not contagious. It is also known as vernal catarrh. The disease occurs chiefly in children, most frequently in boys. It attacks either the tarsal or the bulbar conjunctiva, or occasionally both.

Objective Symptoms.—In the palpebral form, the upper palpebral conjunctiva presents hard, flattened papillæ, separated by furrows, giving a cobblestone appearance. Both upper and lower palpebral conjunctivæ are bluish-white in colour, as though covered with a thin layer of milk. In the bulbar form, the conjunctiva adjacent to the inner and outer portions of the limbus presents gelatinous hypertrophies, sometimes slightly pigmented; these may involve the cornea for a short distance; occasionally the hypertrophy surrounds the cornea. There are conjunctival congestion and some mucoid secretion. During winter these changes become less marked or almost disappear; they return with warm weather.

Subjective Symptoms include a feeling of heat, lacrymation, intense itching, and photophobia; these become worse in warm weather and disappear in the winter.

Course.—The disease continues in this intermittent way for several years or longer, finally becoming extinct and leaving

no traces or very slight changes behind. Its etiology is unknown. Occasionally a grayish line concentric with the limbus but with a clear area between the linear opacity and the limbus, thus resembling the arcus senilis, is left, and there may also be slight thickening, discoloration, and irregularity in the palpebral conjunctiva. The diagnosis is evident; it can scarcely be mistaken for trachoma, since the type of individual, the milky appearance, recurrence in hot weather, freedom of fornix from involvement, and absence of complications will prevent error.

Pathology.—The papillæ consist of dense fibrous material in the subconjunctival tissue, later undergoing hyaline degeneration; the epithelium is much thickened and sends short prolongations into the fibrous deposits; the hypertrophied epithelium together with the underlying hyaline tissue account for the milky appearance. Eosinophile leucocytes are abundant in the nodules and also in the secretion. The bulbar hypertrophies consist of translucent infiltrates, similar in structure to that of the papillæ, with hypertrophy of the epithelium, which is prolonged slightly into the border of the cornea.

Treatment.—The subjective symptoms can be made less annoying by the use of boric acid solution, corrosive sublimate (1 : 5,000), the alkaline wash (p. 457), acetic acid (2 drops of the dilute acid to an ounce of water), and sodium carbonate 1 per cent.; a few drops of the last are instilled four times a day, preceded by irrigation with cold boric acid solution. Another remedy often used is 3 per cent. ointment of the yellow oxide of mercury, a small portion being placed in the conjunctival sac followed by massage with the upper lid. Cold compresses and the wearing of smoked glasses add to the patient's comfort. Temporary relief from the distressing subjective symptoms follows the instillation of 1 per cent. holocaine in 1 : 10,000 adrenalin. Exposure to radium, when the granulations are of large size, has resulted in some cures, but results are not quite conclusive in a disease so subject to fluctuations. This remedy must be used cautiously, with dosage and protection to the globe regulated by an expert, since cataract has followed such applications.

Symblepharon.

A cicatricial attachment between the conjunctiva of the lid and eyeball (Fig. 128). It may affect both lids, but usually the lower. Sometimes it includes part of the cornea. It is called anterior or partial, when extending bridge-like from lid to globe, leaving a free portion of conjunctiva corresponding to the fornix; posterior, when it involves only the fornix; and complete when it affects all the conjunctiva. It is caused by




FIG. 128.—SYMBLEPHARON.

occurs after injuries, especially burns from lime, acids, and molten metal; sometimes it follows trachoma, and occasionally diphtheritic conjunctivitis.

Symblepharon often interferes with the movements of the eyeball, and this may cause diplopia. Traction upon the adherent part excites irritation. In severe cases the cornea is included and sight interfered with; or,

if there is inability to close the lids, lagophthalmos and its sequelæ may be present.

Treatment.—If anterior and not extensive, we divide the band and keep the two raw surfaces from uniting by separating them daily with a probe until they have cicatrized separately. The interposition of a small roll of absorbent cotton saturated with some bland oil or ointment may aid in this purpose. Or the band may be ligated and the ligature allowed to slough through.

In more severe forms, and in all cases of posterior and complete symblepharon, the separated raw surfaces must be covered with conjunctiva or with grafts of skin or mucous

membrane to keep them from uniting. This may be done (1) by loosening the adjacent bulbar conjunctiva and sewing it over the defect, (2) by transplanting pieces of mucous membrane from the lip or from the rabbit's conjunctiva, (3) by skin-flaps, passed from adjacent surfaces, and (4) by Thiersch skin-grafts, taken from other parts of the body, and supported on an artificial eye or piece of sheet-lead until adhesion has taken place.

Pinguecula.

A small, slightly raised spot of yellowish colour, situated at the inner and outer sides of the cornea, where the conjunctiva is most exposed to wind and dust; it is especially marked in old people, and most conspicuous when the conjunctiva is injected. It is not formed of fat, as its name implies, but is a thickening of the conjunctiva, due to excessive development of yellow elastic tissue and the deposit of hyaline substance. It never calls for interference.

Pterygium.

A triangular fold of membrane occupying the interpalpebral fissure, extending from the inner or outer part of the ocular conjunctiva to the cornea (Fig. 129). The apex is immovably united to the cornea; the base spreads out and merges with the conjunctiva.

Symptoms. — When recent, pterygium is rich in blood-vessels, and hence of a red colour. Later it changes into a white, tendinous membrane. The central portion is firmly attached to the sclera, but above and below its margin is represented by a fold of conjunctiva. It grows slowly towards the centre of the cornea, giving rise to moderate symptoms of conjunctival irritation, and it may eventually cover a considerable part




FIG. 129.—PTERYGIUM.

of the cornea; finally it becomes stationary. Besides more or less irritation, it causes disfigurement, and it spreads over the cornea, interfering with vision. It is generally situated to the inner side of the cornea, less frequently to the outer side, or in both situations. It may occur in one or both eyes.

Etiology.—It occurs usually in elderly persons who are exposed to wind or dust (farmers, coachmen, masons, sailors).

Pathology.—The view that pterygium originates from ulceration of the pinguecula extending into the cornea and drawing the conjunctiva after it is no longer held. While the pathology is not definitely known, the affection is thought to consist of a degenerative process due to long-continued irritation; this causes a thinning of the conjunctiva and later hypertrophy; with this there is destruction of Bowman's membrane and the epithelium corresponding to the invaded portion of the cornea, and a deposit of fibrous tissue.

Treatment consist in removal by one of a number of different operative methods. The pterygium may be dissected away with a sharp scalpel or Beer's knife (Fig. 67), and cut off, the conjunctival defect being closed by uniting the upper and lower borders, undermining the conjunctiva if necessary to bring the edges together. The apex of the pterygium must be thoroughly excised from the cornea, and its attachment in this situation scraped or cauterized with the actual cautery, to prevent recurrence. Instead of cutting off the pterygium, it may be dissected up and stitched underneath the detached conjunctiva, either above or below; or it may be divided into two halves, of which one is transplanted above and the other below, being held in the conjunctival pocket by a stitch. Some cases show a tendency to recur, even after repeated removal.

Pseudo-ptyerygium is an attachment of a fold of conjunctiva to the cornea as the result of ulceration of the latter; a fine probe can pass under it at the limbus to distinguish it from a true pterygium. It may be separated from its corneal attachment, if causing much disfigurement.

Subconjunctival Hæmorrhage.

Bright or dark red patches involving more or less of the bulbar conjunctiva (Fig. 108, Plate VIII.), unaccompanied by inflammatory symptoms. Ecchymosis is seen after injuries, operations, and inflammations of the eyeball. It is frequently observed in old persons with brittle bloodvessels, being excited by various straining efforts, and in children after whooping-cough. It is of no importance, and becomes absorbed within a week.

Injuries of the Conjunctiva.

These are very common, and include:

1. *Foreign bodies* in the conjunctival sac, consisting of dust, iron, coal, or ashes. They usually adhere to the inner surface of the upper lid, causing severe pain and irritation, and are readily removed after eversion of the lid.

2. *Wounds*.—Extensive wounds of the conjunctiva should be closed with a stitch.

3. *Burns* are quite common, being due to boiling water, steam, lime, mortar, molten metals, and acids. Following the accident a grayish eschar forms. This separates and leaves a granulating surface, which heals by cicatrization. In this way symblepharon often results.

The treatment consists in the complete removal of the caustic substance as soon as possible. Solid particles are removed with absorbent cotton or forceps. Then the conjunctival sac is washed out with solutions which tend to neutralize the corrosive substance, or render it insoluble. In the case of lime, mortar, or caustic alkalies, we flush out with a solution of boric acid; or we may wash out the eye with oil. If the corrosive agent consisted of an acid, the eye is irrigated with a weak solution of sodium bicarbonate. But since it is of the utmost importance to remove the irritating and destructive agent without any delay it is unwise to wait for neutralizing solutions, but to irrigate with great quantities of water immediately, whether the foreign material is acid or alkaline. Solid particles are removed with absorbent cotton

or forceps. Subsequently we use cold compresses, atropine, keep the conjunctival sac filled with a bland ointment, such as 1 per cent. boric acid, and apply a loose gauze and cotton dressing. Unless the burn is very superficial, there will be eschars; when these loosen and come away, raw and granulating surfaces will be exposed; these have a great tendency to form adhesions; the latter must be separated frequently. Symblepharon is, however, very apt to occur notwithstanding the greatest care in separating these adhesions.

CHAPTER IX

DISEASES OF THE CORNEA

Anatomy.—The cornea is the clear, transparent, anterior portion of the external coat of the eyeball. It is nearly circular, but is slightly wider in the transverse than in the vertical direction; its radius of curvature is somewhat shorter than that of the sclerotic; the junction of the two is known as the limbus, but their tissues are in complete continuity. The cornea is composed of five layers, from without inward: (1) Layer of epithelial cells; (2) Bowman's membrane; (3) the proper substance of the cornea; (4) Descemet's membrane; and (5) a layer of endothelium.

The *epithelium* covering the front of the cornea is of the stratified variety, formed of flattened, scaly epithelial cells superficially, of polygonal cells beneath these, and of columnar cells most deeply. Practically it is part of the bulbar conjunctiva.

Bowman's membrane is a thin, homogeneous membrane which separates the corneal epithelium from the proper substance of the cornea. Although usually described as a separate membrane, it is really a part of the corneal substance, and when highly magnified is seen to be composed of fine fibres which are intimately connected with the subjacent layer.

The *proper substance of the cornea*, the thickest layer, is formed of connective tissue arranged in lamellæ, the planes of which are parallel to the surface of the cornea; these are




FIG. 130.—SECTION OF THE CORNEA, SHOWING MINUTE ANATOMY.

A, Layer of epithelial cells, etc.

connected with one another, and cross at right angles in alternating layers. The ultimate fibrils of which the lamellæ are composed, as well as the different bundles of fibrils forming the lamellæ, are held together by means of a transparent cement substance. The corneal substance is traversed by a system of spaces or lacunæ, situated in the cement substance separating the laminae, and sending off prolongations in every direction. These form small canals, by means of which the lacunæ of the same plane and those placed above and below communicate. The spaces are partly filled with branching cells (corneal corpuscles), the branches of the cells passing into the small canals, and communicating with adjoining cells. The cells do not completely fill the lacunæ, but leave room for the passage of lymph and lymph corpuscles. The proper substance of the cornea passes uninterruptedly into the sclera.

Descemet's membrane (the posterior elastic lamina) is a thin, firm, structureless, transparent, and highly elastic layer, placed posterior to the proper substance of the cornea. At the periphery of the cornea

it passes over into radiating bundles of elastic fibres which form the *ligamentum pectinatum*.

Posteriorly, next to the anterior chamber is a single layer of flattened, hexagonal cells, the *endothelium*.

The cornea is not provided with bloodvessels. The capillary loops from the anterior ciliary vessels form a ring around the circumference of the cornea. Its nutrition is provided for by the system of lymph canals just described. It is richly supplied with *nerves* derived from the ciliary nerves.




FIG. 131.—ARCUS SENILIS.

The line between cornea and sclera is known as the *limbus*. Near the margin of the cornea, just within the sclero-corneal junction, we frequently find an opaque, whitish ring, or part of a ring. This is known as the *arcus senilis* (Fig. 131). It is due to a deposit of fatty granules, and most frequently occurs in advanced age, though occasionally it is found in younger persons.

Inflammations of the Cornea.—Keratitis.

Keratitis in general presents the following symptoms:

Objective Symptoms.—(1) Infiltration, with dulness of surface and diminution of transparency. This may be followed

by (a) complete absorption of the infiltration; (b) incomplete absorption, leaving opacities; and (c) suppuration, with formation of an ulcer. (2) Limited or general vascularization, the bloodvessels being derived from the conjunctival loops at the limbus. (3) Circumcorneal injection. (4) There is often a complicating conjunctivitis. (5) Neighbouring deep parts are frequently involved (iris and ciliary body), as a result of which there may be pus in the anterior chamber (hypopyon).

Subjective Symptoms.—Pain, photophobia, blepharospasm, lacrymation, and interference with vision.

Varieties.—Keratitis may be divided into suppurative and non-suppurative.

Suppurative Keratitis.—The common forms are (1) phlyctenular keratitis, and (2) ulcers of the cornea. The uncommon forms are (3) keratitis from imperfect closure of the eyelids (lagophthalmos), (4) neuro-paralytic keratitis, and (5) xerotic keratitis.

Non-suppurative Keratitis.—The common forms are (1) interstitial keratitis, and (2) vasculo-nebulous keratitis (pannus). The uncommon forms are (3) vesicular keratitis, (4) keratitis profunda, (5) sclerosing keratitis, and (6) ribbon-shaped keratitis.

Phlyctenular Keratitis has been described under the title Phlyctenular Conjunctivitis (p. 128), and the special symptoms arising when the cornea is involved have been pointed out.

Ulcer of the Cornea.

An infiltration, followed by suppuration and loss of substance of the cornea. The affection is of very common occurrence.

Subjective Symptoms.—Pain, photophobia, lacrymation, and blepharospasm. Sometimes all these symptoms are slight, or even absent, and yet the ulcer may be a very extensive and serious one.

Objective Symptoms.—An ulcer begins with a dull, grayish, or grayish-yellow infiltration of a circumscribed portion of

the cornea. Suppuration takes place in this area, the superficial layers are cast off, and thus there is loss of substance. The process may progress in two directions; it may either travel over the cornea so as to involve a greater area, or it may become deeper; it may extend both in area and in depth. Very often the advance takes place in one direction across the cornea; sometimes there is at the same time a tendency to heal at the opposite side, so that the ulcer merely changes its situation (creeping or serpiginous ulcer). There is nearly always more or less grayish infiltration of the cornea immediately surrounding the loss of substance, and considerable ciliary injection (Fig. 132, Plate XI.).

If the ulcer is small and superficial it may cleanse itself in the course of a few days. The destroyed portion of the cornea will be cast off, the infiltrated border will become clear, and repair set in; this is accompanied by the appearance of bloodvessels which spring from the limbus; the process terminates in cicatrization. When the ulcer extends no deeper than the epithelium, the cornea may remain perfectly transparent. But when some of the proper substance of the cornea has been destroyed, new connective tissue takes its place, and such a scar is always more or less opaque. The seat of the ulcer may also be marked by a slight depression (corneal facet).

The detection of the extent of infiltration and ulceration is facilitated by the instillation of a few drops of a 2 per cent. solution of fluorescein, which stains green all such ulcerated or infiltrated parts.

When the ulcer is deeper both subjective and objective symptoms are more pronounced, and the complications and sequelæ are more serious. Neighbouring structures give evidences of inflammation—conjunctivitis, congestion of the iris, even iritis with its symptoms, including hypopyon. Hypopyon is a collection of pus in the anterior chamber. The pus is not derived from the ulcer, but is an exudation from the inflamed iris and ciliary body. It collects at the bottom of the anterior chamber (Fig. 133, Plate XI.), or it may partially or completely fill this space. It remains sterile as




FIG. 132.—CORNEAL ULCER.




FIG. 133.—CORNEAL ULCER WITH
HYPOPYON.




FIG. 134.—LEUCOMA ADHERENS.




FIG. 135.—PANNUS, AND TRACHOMATOUS
SCARRING OF EYELID.




FIG. 136.—PHLYCTENULAR ULCERS OF
CORNEA. FASCICULAR KERATITIS.




FIG. 137.—INTERSTITIAL KERATITIS.

long as there is no perforation of the cornea, and may disappear and reappear in the course of the affection. It may either remain fluid, or when mixed with fibrin it may form a semi-solid, globular mass. Such an ulcer may heal with no other permanent injury except marked corneal opacity (Figs. 145, 146), or there may be bulging (anterior staphyloma) (Fig. 141). But deep and spreading ulcers frequently have their course modified by the occurrence of perforation of the cornea, which, in healing, affects the usefulness and safety of the eye in various ways.

The course of a simple ulcer is influenced greatly by the state of health and cleanliness of the patient. Even the smallest ulcer or abrasion in the cornea of a workman should not be regarded lightly.

Perforation of the Cornea is often preceded by a protrusion of Descemet's membrane through the floor of the ulcer, forming a small transparent vesicle. Perforation may be spontaneous, or it may be caused by increased pressure resulting from the blepharospasm, various straining efforts, such as crying, sneezing, or coughing, or occasionally by force exerted in examining the eye. The aqueous humour escapes, often carrying the iris into the wound; the eye feels soft; the anterior chamber is obliterated, and iris and lens are in apposition with the cornea. Perforation of the cornea has a favourable effect upon the course of the affection; the subjective symptoms are relieved, and the ulcer begins to heal as a result of diminished tension.

When the opening closes by cicatrization the iris may regain its normal position. But frequently it continues adherent to the walls of the perforation, or remains prolapsed, and becomes incorporated with the scar. Such a condition is called anterior synechia, and since the cicatrix forms a dense white opacity of the cornea it is known as adherent leucoma. Most frequently only a portion of the iris is drawn into the scar; the pupil is then more or less pear-shaped. Occasionally the entire pupillary margin may be adherent, causing both exclusion and occlusion of the pupil.

At the time of perforation the lens may become dislocated,

and occasionally it escapes. When it is pushed forward and lies in apposition with the margins of the opening, and then recedes after the anterior chamber is re-established, it frequently presents a proliferation of the subcapsular epithelium which has become irritated by the pressure of the lens upon the cornea, forming a white spot upon its anterior surface (Fig. 219), known as anterior capsular or anterior polar cataract.

Occasionally the perforation fails to close and a fistula of the cornea results; this condition exposes the eye to subsequent serous inflammation and jeopardizes its safety. Iridocyclitis and even panophthalmitis may follow perforation, especially if the suppurative process be a virulent one.

Etiology.—Ulcers of the cornea are usually found in adult and aged individuals; phlyctenular ulcers are the only ones which are common in children. Ulcers are much more frequent among the lower than among the better classes, and occur often in individuals in whom the general health is poor.

The process is essentially an infection by various micro-organisms (pneumococci, streptococci, staphylococci, Plate IX.), frequently introduced by the secretion of chronic conjunctivitis, and especially by that of dacryocystitis.

The exciting causes are: (1) Traumatism (foreign bodies, injuries)—this is one of the most frequent causes; (2) conjunctival inflammations (gonorrhœal ophthalmia, ophthalmia neonatorum, trachoma, diphtheritic conjunctivitis); (3) phlyctenular keratitis; (4) disturbances in the nutrition of the cornea (paralysis of trigeminus, keratomalacia, glaucoma); (5) infection during operations; (6) variola; (7) herpes.

Pathology.—The changes accompanying corneal ulcer comprise infection, infiltration, necrosis and repair. The infection is exogenous and is caused by the entrance of pyogenic organisms through some defect in the corneal epithelium. These organisms vary somewhat with the type of ulcer; in virulent forms the pneumococcus is often responsible either alone or mixed with other organisms (staphylococci, streptococci, gonococci, diphtheria bacilli). There is at first a gray zone due to *infiltration* of the epithelium, and if deeper, also

Bowman's membrane and corneal lamellæ with leucocytes; the leucocytes next invade more deeply and pass between the lamellæ. Then follows necrosis of the affected parts; the slough is cast off in part, leaving an excavation represented by the floor and walls of the ulcer covered by remains of the slough and by necrotic lamellæ separated by polymorphonuclear leucocytes. Finally, repair sets in, assisted by superficial bloodvessels which extend from the limbus and which supply the material which replaces the loss of substance; this replacement may consist entirely of epithelial cells growing in from the edges of the ulcer, providing the latter has been superficial. But if Bowman's membrane and the lamellæ have been destroyed, the defect is filled in with fibrous tissue; in such case the corneal corpuscles divide and form spindle-shaped cells which later are converted into fibrous tissue devoid of the transparency of the cornea, and a more or less opaque scar results.

Clinical Forms.—Certain variations in the course of corneal ulcers have already been considered. The nomenclature of ulcers of the cornea is quite extensive, and is founded upon peculiarities in the symptoms or course. The following warrant special mention:

Simple Ulcer is the name often given to a small and superficial ulcer, with symptoms of slight or severe irritation, no tendency to perforation, terminating in uncomplicated healing; phlyctenulæ and slight injuries often cause such ulcers.

Catarrhal Ulcer complicates catarrhal conjunctivitis in adults. With increase of subjective symptoms, peripheral punctate infiltrations appear, and coalesce into a crescent, changing to a superficial ulcer, concentric with, and just within, the limbus. The course is usually favourable and healing prompt; if opacities remain they do not reduce vision, being outside the pupillary area.

Deep Ulcer is one which shows a tendency to involve the deeper layers and to perforate rather than to spread over the cornea. The symptoms are apt to be marked, the iris is usually involved, and hypopyon is often present; hence the results are often serious.

Serpent Ulcer (Ulcer serpens, Infected Ulcer, Hypopyon Keratitis) is a very virulent form, in which the process tends to spread over a considerable portion of the cornea, and at the same time extends into its depth. The subjective symptoms are usually severe, though occasionally they are slight. The process begins as a grayish-yellow infiltration near the centre of the cornea, rapidly changing to an ulcer with sloughing margins, especially at the advancing side, where there is frequently a yellowish crescent (Fig. 133, Plate XI.). The rest of the cornea is often dull, gray, and infiltrated. The process advances very rapidly, much of the cornea becomes destroyed, and perforation takes place. There is early and intense iritis, and hypopyon is almost always present. Owing to the virulence of the ulcer and the accompanying iritis, much damage results to the eye. Adhesion and prolapse of the iris are frequent, the pupil is often occluded, and iridocyclitis and panophthalmitis are not uncommon. Considerable opacity of the cornea always ensues, and often staphyloma. There is, therefore, considerable impairment of vision, which not infrequently is reduced to perception of light. Pneumococci are found in the discharge.

Diplobacillary Ulcer is a milder type of hypopyon ulcer due to the diplobacillus, occurring in children, sometimes in adults, which has a more favourable course; it has a tendency to spread laterally, but is less apt to perforate.

Mooren's Ulcer begins near the upper margin of the cornea, and gradually spreads over it. The advancing edge is undermined; meantime another part of the ulcer may be healing, though this usually breaks down again. The ulcer involves perhaps half the thickness of the cornea, and does not perforate. Both eyes may be affected. The disease is very rare. Microscopically the floor of the ulcer consists of a collection of small round cells like granulation tissue. The cause is unknown, though the clinical course leads one to think that it may be of neuropathic origin—analogueous to the perforating ulcer of tabes. Treatment seems almost hopeless, but repeated applications of the electric cautery to the advancing edge of the ulcer should be tried.

Marginal Ring Ulcer is rare, of slow and intermittent course, successively involving different parts of the margin, so that it tends to extend all around the cornea.

Transparent Ulcer is small, superficial, central, shows no tendency to spread or to perforate, occurs chiefly in children, and is followed by little or no opacity, but often by a small pit (facet).

Herpetic Ulcers result from ruptured herpetic vesicles; they are seen in the course of acute febrile diseases, especially pneumonia and influenza, and are accompanied by marked symptoms of irritation. They tend to occur in groups or rows, and the affected spot is anæsthetic. They usually heal promptly unless secondarily infected.

Dendriiform Ulcer (Dendriiform Keratitis) is a chronic form of superficial ulcer, which commences with a grayish line and spreads by sending out branches which present small knob-like extremities. It is usually associated with febrile conditions, and was very commonly seen during the Great War after trench fever.

Abscess of the Cornea is a purulent infiltration in the substance of the cornea, covered both superficially and posteriorly by sound tissue. The subjective symptoms are those of infected ulcer; it is usually accompanied by iritis and hypopyon. The term *Ring Abscess* of the cornea refers to an infection, following perforating wounds (rarely operations), in which a yellow ring develops in the central part of the cornea, soon followed by necrosis of this part, and usually succeeded by panophthalmitis.

Treatment may be divided into (1) constitutional, (2) treatment of pre-existing local conditions, (3) local treatment of the ulcerative process.

Constitutional.—Since ulcers usually occur in persons in whom the general condition is below par, it is necessary to improve the tone of the system by attention to diet, fresh air, hygienic surroundings, condition of the bowels, etc., and often to administer tonics.

Treatment of Pre-existing Local Conditions.—Foreign bodies are to be removed and other local irritating conditions

remedied. The various forms of conjunctivitis and dacryocystitis must receive careful attention.

Local Treatment includes atropine (sometimes eserine), bandage, hot compresses, antiseptic lotions, scraping, cauterization, and paracentesis of the cornea, and division of the ulcer by Saemisch's method.

Atropine anæsthetizes the sensory nerve-endings in the floor of the ulcer, dilates the pupil, and inhibits lacrymation. One drop of a 1 per cent. solution or a 1 per cent. ointment made with vaseline may be used three times a day or oftener. The substitution of eserine for atropine when the ulcer is peripheral and likely to perforate is sometimes recommended, but, in view of the liability to iritis in such cases, eserine may prove a dangerous remedy as it contracts the pupil, which should be kept dilated when the iris is inflamed. It is also more irritating than atropine.

If there is intense pain, unrelieved by atropine, it may be necessary to use cocaine or holocaine for a day or two. These remedies, especially cocaine, should not be used for longer than is absolutely necessary, owing to their tendency to cause exfoliation of corneal epithelium.

Protection is afforded by smoked glass or by a bandage. A lightly applied bandage is not only comfortable, but by keeping the lids closed and immobile it prevents irritation of the ulcer; it also supplies beneficial warmth. When there is much discharge the bandage is contra-indicated in superficial ulcers. But in any case in which perforation of the cornea is liable to occur a firm (pressure) bandage is applied; this must be removed and replaced several times a day to permit cleansing of the eye and local applications.

Hot compresses should be applied for half an hour at a time, several times a day; they favour healing of the ulcer.

Antiseptic lotions, such as solutions of boric acid, sodium chloride, bichloride of mercury (1:6,000), act as cleansing agents, and are especially useful when there is much discharge. They must be used from irrigator or syringe, never from an eyebath when ulcers are present.

Other measures are sometimes resorted to: Iodoform

sprinkled upon the cornea or applied in the form of an ointment, subconjunctival injections of corrosive sublimate (1:5,000), protargol (10 per cent.), formalin (1:2,000).

Foreign Proteins.—In virulent forms of ulcer, polyvalent serum may be given advantageously, or boiled milk (5 to 10 c.c.) may be injected into the gluteal muscles.

To Limit Spreading.—If these remedies are insufficient and the ulcer spreads, we must destroy the infective focus either by scraping the floor and margins of the ulcer with a small sharp spoon or curette, or better, by cauterizing this area.

Cauterization is effected by pure liquid carbolic acid, applied with a sharpened match stick after drying the cornea with blotting-paper, and by the actual cautery or the galvano-cautery.

Tincture of iodine, or absolute alcohol, offers a very efficient mode of disinfecting and cauterizing corneal ulcers. A piece of absorbent cotton is wound firmly upon an applicator, dipped into tincture of iodine (or alcohol), and then exposed to the air for a few seconds so that there is no excess of liquid. It is now brushed upon the ulcer and its infiltrated margins. It is usually necessary to repeat the cauterization a number of times on successive days.

Galvano-Cautery.—After thoroughly anæsthetizing the eye and staining the ulcer with fluorescein, the electrode shown in Fig. 138 is placed cold upon the part to be cauterized,




FIG. 138.—GALVANO-CAUTERY.

the connexion made so that the burner assumes a deep red colour, and then the connexion quickly broken. Or the electrode, already heated, may be rapidly applied to the points to be cauterized. Successive points (especially the margins of the ulcer) are cauterized in this manner, each for a very short period, so as to prevent perforation and the conduction

of heat to deeper parts. An even better instrument in skilled hands is a little *platinum bulb* set in an ivory handle (Fig. 130). The bulb is heated in a spirit-lamp. It is held ready until it just ceases to glow in daylight; it is then quickly applied to the surface of the ulcer. The shape of the bulb is more suitable for the purpose than that of the wire electrode; the small size of the instrument admits of more delicate handling.



FIG. 139.—PLATINUM CAUTERY.

The *metri-cautery* in which the degree of electrically generated heat can be controlled, is very useful in infected and hypopyon ulcers. A thermometer registers the temperature at the point of the instrument, which is maintained at 75° to 85° C. by regulating the current. The head of the instrument is applied for one minute to the ulcer, after the cornea has been anæsthetized. Organisms are killed, but tissue not destroyed.

Paracentesis of the cornea is another valuable measure. This puncture is frequently made with a paracentesis trocar, which is provided with a thick shoulder to prevent the instrument from penetrating too far; it may be made with the lance-shaped knife (Fig. 192), broad needle (Fig. 217), or the Graefe knife (Fig. 207). After local anæsthesia and fixation of the eyeball with the fixation forceps (Fig. 186), the instrument is passed perpendicularly through the cornea, near its lower margin, unless the situation of the ulcer requires another site. As soon as its point reaches the anterior chamber, the handle of the instrument is depressed and the knife or trocar is pushed on horizontally, avoiding injury to the iris or lens, until the incision is about 3 mm. long. Then it is withdrawn slowly with pressure upon the posterior lips of the wound, so as to evacuate the contents of the aqueous chamber gradually. It may be necessary to repeat the paracentesis or to reopen the wound with a probe daily until the ulcer cleanses itself.

Threatened perforation should always be anticipated by paracentesis. Even when there is no danger of spontaneous perforation, the flushing out of lymphatic spaces due to the temporary lowering of pressure after paracentesis often starts the healing process in a previously intractable ulcer.

Saemisch's operation of splitting open the ulcer has been replaced to a great degree in favour of paracentesis and the thermo-cautery or metri-cautery, but it is useful in some severe forms of serpiginous ulcers. If contemplated, it should be done early before the cornea has been too extensively destroyed. A Graefe knife is thrust through clear corneal tissue 1 or 2 mm. to the outer side of the ulcer, made to traverse the anterior chamber, and brought out 1 to 2 mm. to the inner side of the ulcer. The edge of the knife is directed forward and the ulcer is split through its centre and the hypopyon removed.

After spontaneous perforation of an ulcer atropine is instilled, a pressure bandage applied, and perfect rest insisted upon. If there is a recent prolapse of the iris, the latter is freed from adhesion to the margins of the opening and then excised; this has the effect of an iridectomy. But if the prolapse has existed for some days it must be allowed to remain; subsequent operative interference may then be indicated.

After the healing process has become fairly initiated, certain mildly stimulating remedies, such as the ointment of the yellow oxide of mercury, or, better still, dionin drops (1 to 2 per cent.) once a day, are used to hasten cicatrization, and to clear up the cornea as much as possible.

Keratitis from Defective Closure of the Lids.—This form of keratitis is due to exposure of the cornea when it remains uncovered by the lids (lagophthalmos). Under such circumstances the cornea becomes desiccated, the conjunctival secretion and atmospheric dust settle upon it, infiltration and ulceration take place, with the subsequent course of ulcer of the cornea. The causes are paralysis of the orbicularis (facial paralysis), marked exophthalmos, and various deformities of the lids. Treatment consists in relieving the lagophthalmos, if possible, frequent irrigation of the conjunctival

sac with cleansing solutions, and closure of the lids by bandage or plaster. Unless the process has gone beyond certain limits, it can be controlled by this plan of treatment.

Neuro-paralytic Keratitis is a form of infiltration and ulceration of the cornea observed after paralysis of the trigeminus. The changes are considered by some to be trophic, by others to be due to exposure and lodgment of foreign substances upon the insensitive cornea. It begins with a large central infiltration of the cornea with characteristic exfoliation of the epithelium, followed by ulceration; in severe cases hypopyon and perforation may result. There is no pain or lacrymation, the course is chronic, and the result is considerable opacity of the cornea. Treatment consists in applying a bandage to the eye, or keeping the lids closed with plaster or median tarsorrhaphy. Atropine and hot compresses are of service.

Xerotic Keratitis (keratomalacia) is the result of lack of nutrition of the cornea. It is an uncommon disease, which occurs in greatly debilitated children. The process begins with dryness of the conjunctiva in both eyes; soon the cornea corresponding to the palpebral aperture becomes cloudy, desiccated, covered with scales, and ulcerates and often perforates. Night blindness is present. Most of the cases occur during the first year of life and the great majority of such patients die from the disease which is responsible for the corneal condition. In older children the affection may be less severe, but marked corneal opacities persist. Treatment consists in measures to increase the general strength (vitamin-rich diet, cod-liver oil butter); locally, hot compresses, anti-septic lotions, bandaging, and atropine are indicated.

Interstitial or Parenchymatous Keratitis.

A cellular infiltration of the middle and posterior layers of the cornea, of frequent occurrence in childhood, chronic in its course, not leading to ulceration, but accompanied by more or less inflammation of the uveal tract.

Objective Symptoms.—The affection begins either in the centre or at the margin of the cornea. If it starts in the

centre, this part will present a grayish infiltration, the superficial layers at first retaining their normal lustre. This central patch soon spreads, so that the whole cornea becomes implicated. If it commences at the periphery, one or more grayish crescents are seen, which soon spread toward the centre, and involve all the cornea. After the infiltration has become general, the cornea will become softened, of a dense grayish or sometimes yellowish-gray colour, so that the iris can no longer be seen, and vision is reduced to little more than perception of light. The surface of the cornea is now steamy, and resembles ground-glass. At this period, or even before, deep-seated bloodvessels (derived from the anterior ciliary) make their appearance, and pervade more or less of the cornea (Fig. 137, Plate XI.). They cover either the periphery, circumscribed sectors, or the whole cornea. This vascularization gives rise to a dirty-red or yellowish-red discoloration, which is known as the salmon-coloured patch. The progress thus far is accompanied by irritative symptoms, and lasts one or two months.

The inflammation then begins to subside. The periphery of the cornea clears up, the bloodvessels become fewer, the irritative symptoms disappear, and vision improves. This process may take several months, or even a year, the centre of the cornea being the last portion to clear. In favourable cases, after a year or more, nothing but a very faint central opacity and evidences of a few minute peripheral vessels can be found.

Not all cases will, however, run such a benign course. The anterior portion of the uveal tract is regularly involved. In mild cases this will consist merely in congestion of the iris. But in more pronounced types there will be iritis, choroiditis, cyclitis, and changes in the vitreous. In such cases, after the cornea has become less opaque, we may find evidences of these inflammations, in the form of adhesions of the iris to the lens (posterior synechiæ), changes in the iris and choroid, opacities of the vitreous, and even exclusion of the pupil and iridocyclitis. Staphyloma of the cornea may also follow. So that more or less serious impairment of

sight may ensue as a result of these inflammatory processes. Furthermore, the clearing-up process in the cornea may come to a standstill, leaving a dense opacity, which also causes loss of useful vision.


Subjective Symptoms.—During the period of infiltration and vascularization there will be photophobia, lacrymation, pain, and interference with vision, the intensity usually depending upon the severity of the process. These symptoms gradually subside during the progress of absorption.

Both eyes are usually involved. Frequently the inflammation in the second eye commences after that in the first has existed for some weeks or months.

Etiology.—The disease usually occurs between the fifth and fifteenth years, less commonly after this period, and rarely

after thirty; but it has been seen at all ages up to sixty. The great majority of cases are due to inherited syphilis. In few instances it depends upon acquired syphilis, tuberculosis, or occurs without known cause.

In many cases there will be other *Signs of Inherited Syphilis*, such as characteristic physiognomy, peculiar conformation of the skull (square forehead, prominent frontal eminences, depressed bridge of nose), radiating scars at angles of mouth (Fig. 140),



SHOWING THE SIGNS OF INHERITED SYPHILIS, INCLUDING HUTCHINSONIAN TEETH

scars in the mouth and pharynx, ozæna, enlarged cervical lymphatic glands, nodes on the bones, and more or less

impairment of hearing. The permanent teeth are developed, their angles rounded off, and there is often a centic notch in the free margin. These changes are especially marked in the upper central incisor teeth (Hutchinsonian teeth, Fig. 140).

Pathology.—The disease is essentially a deep keratitis with involvement of the uvea. Between the laminae of the substantia propria, especially those placed most posteriorly, there is cellular infiltration of fixed corneal cells, lymphocytes and eosinophil, gathered chiefly around capillary vascularization and causing thickening of the cornea. There is irregular thickening and haziness of the corneal epithelium. Bowman's membrane is wavy and irregular, and beneath it is a layer of proliferated fixed corneal cells in which are found minute capillaries. Descemet's membrane is wrinkled. The epithelium is usually missing from the posterior surface of the cornea and is replaced by an accumulation of lymphocytes. During resolution there is thinning of the cornea. The changes in the uvea are described under Uveitis (p. 189).

Treatment—Local.—Atropine, protection from light by smoked coquilles, or by a shade, hot compresses. When the cornea begins to clear, we employ mild, stimulating ointments, such as yellow oxide of mercury and calomel, often combined with gentle massage, or instil dionin in 1 per cent. solution. We must be careful not to apply stimulating treatment too early.

Constitutional.—Calomel, $\frac{1}{10}$ grain four times a day, gray powder, 1 grain twice a day, or potassic iodide, 5 grains, combined with corrosive sublimate, $\frac{1}{10}$ grain t.i.d. Syrup of the iodide of iron or other preparation of iodine, cod-liver oil, iron and quinine, and attention to the general health. In the uncommon cases occurring in adults we prescribe mercury by inunction or by hypodermic injection, also iodide of potassium.

Though salvarsan is usually less successful in the treatment of inherited syphilis than in the acquired disease, it has produced excellent results in some intractable cases, especially when combined with paracentesis of the anterior chamber. Since tuberculosis is occasionally the cause and sometimes

associated, injections of tuberculin are indicated when we suspect this factor.

Pannus (vasculo-nebulous, or vascular keratitis) has been described in connexion with trachoma (p. 121).

Uncommon Forms of Non-suppurative Keratitis.

Vesicular Keratitis comprises a number of uncommon inflammations of the cornea, in which the distinguishing feature is the occurrence of groups of small, clear vesicles, or of a single large transparent blister, with marked irritative symptoms. Vesicles occur in herpetic keratitis (herpes corneæ) and in the keratitis accompanying zoster corresponding to the distribution of the fifth nerve, bullæ in keratitis bullosa. Vesicles and bullæ are often found in blind eyes with increased tension, and damaged eyes with opaque and insensitive corneæ.

Superficial Punctate Keratitis complicates acute affections of the respiratory tract, and begins with the symptoms of acute conjunctivitis. Numerous small gray spots appear in the superficial layers of the cornea, beneath Bowman's membrane. These are accompanied by gray radiating lines, and by some general clouding. The disease resembles herpes, but there are no vesicles, no ulceration, and deeper parts are not involved. It occurs in young persons, is usually bilateral, and lasts several months, after which there is complete absorption.

Treatment comprises attention to the conjunctivitis and the bronchial affection, the use of atropine, hot compresses, and smoked glasses, and later the ointment of the yellow oxide of mercury.

Keratitis Profunda is a form of deep-seated inflammation of the cornea occurring in adults, in which a gray, central opacity of the cornea develops, accompanied by irritative symptoms. It becomes entirely or almost perfectly absorbed in a few weeks, and requires treatment similar to that of interstitial keratitis.

Sclerosing Keratitis is the name given to the corneal complication of scleritis (p. 168). The portion of the cornea

adjacent to the sclerotic nodule participates in the process, and a triangular opacity remains. The symptoms and treatment correspond to those of scleritis.

Ribbon-shaped Keratitis (*Transverse Calcareous Film of the Cornea*) is a whitish or grayish band, which extends across the cornea opposite the palpebral aperture, and often contains lime. It occurs usually in old persons and in eyes which have been seriously injured or lost by a previous intra-ocular affection. Treatment consists in gently scraping away the band and using solutions of sodium carbonate (gr. i. to ʒi.).

Keratitis Disciformis is an uncommon type, occurring in adults, in which a gray disc-shaped opacity develops in the middle layers of the cornea with a denser spot in the centre and sometimes concentric lines at the circumference, accompanied by symptoms of irritation. The cause is not definitely known, but it is thought to be due to infection through a tiny defect in the corneal epithelium. Duration is several months. A permanent central opacity remains.

A number of other forms of keratitis are described, but are of rare occurrence.

The so-called **Keratitis Punctata** (*K. P. descemetitis*) is a misnomer and not an inflammation of the cornea. It is described on p. 189.

Protrusions of the Cornea.

These may be either of inflammatory origin, when they are known as staphyloma, or of non-inflammatory origin, when they are called keratoconus.

Staphyloma of the Cornea.—A protrusion of a previously inflamed cornea (Fig. 141), formed of more or less corneal tissue, iris, and cicatrix. Very often the iris forms the chief part. It develops after corneal softening, ulceration, and perforation. It may be total, when it occupies the situation of the entire cornea, or partial, when it occupies only a portion of this area. It may be globular, conical, or lobulated. It is whitish with bluish areas representing spots where pigment shows through the thin cicatrix. It may be all white or all

bluish. Some bloodvessels are frequently seen on the surface. It varies in size, being small in some cases, and so large in others that the lids cannot close.

Symptoms.—Besides the objective signs just mentioned, there are changes in the eyeball, in the staphyloma, and in the lids. There is almost always increased tension, often




FIG. 141.—STAPHYLOMA OF THE CORNEA.

due to obliteration of the pupil. This condition causes pain, produces changes in the interior of the eye (atrophy of the optic nerve, retina, and choroid) which lead to blindness, results in an increase in the size of the bulging, and is responsible for staphyloma of the sclera. The conjunctiva becomes the seat of inflammation from mechanical irritation. The summit of the protrusion becomes dry and

ulcerated, and there is frequently rupture, followed by closure of the opening. This process may be repeated a number of times, until the eye is lost and a shrunken globe remains.

Even before these secondary changes have taken place, there is considerable deformity, and sight is very much reduced. In total staphyloma there will be merely perception of light; in the partial form the amount of sight will depend upon the condition of the cornea which is preserved, the position of the pupil, and the extent to which the curvature of the cornea has become altered.

Treatment.—(1) In partial staphyloma an iridectomy should be performed (p. 228) for the purpose of reducing tension, flattening the protrusion and preventing its increase, and to serve for optical purposes. We select the part of the iris corresponding to the most clear portion of the cornea. If there is no anterior chamber, and the iris lies against the posterior surface of the cornea, this operation is impossible on account of the risk of injuring the lens. In such cases we may excise a portion of the staphyloma, and unite with sutures,

followed by a pressure-bandage for a considerable period of time.

(2) In total staphyloma we resort to incision, abscission, or enucleation. Incision is followed by the extraction of the lens if it still be present. Abscission is performed by cutting through one-half of the protrusion with the knife, and separating the other half from the ring of corneal tissue forming its base with forceps and scissors. The lens is removed. The edges of the corneal gap are then brought together with sutures drawn through the corneal tissue, or, better, through the conjunctiva, which has previously been freed around the limbus. Enucleation, or one of its substitutes, is practised in certain cases in which the staphyloma is very large, or in which, for various reasons, abscission is not advisable.

Keratoconus, or Conical Cornea.—A non-inflammatory conical protrusion of the centre of the cornea (Fig. 142), due to a gradual atrophic thinning, in consequence of which the cornea is unable to resist the normal intra-ocular pressure. The condition is not of frequent occurrence, and is usually observed in young women. It is easily seen when well marked; when less developed, it is recognized by the annular shadow produced when the eye is examined with the ophthalmoscope at a distance, by the pear-like alteration in the shape of the image when Placido's disc is used, and by distortion of the picture of the fundus as seen by the ophthalmoscope. The condition tends to progress for many years before it comes to a standstill. When pronounced, it often presents a slight opacity at its apex. It never ulcerates. Conical cornea causes myopia and astigmatism, and seriously interferes with sight, especially peripheral vision, even after the best possible correction with glasses. 10 D. to 20 D. of myopic astigmatism are not uncommon. Treatment consists in providing concave sphero-cylinders; sometimes these improve vision considerably;




FIG. 142.—KERATOCONUS.

occasionally a disc with a stenopæic hole or slit placed in meridian of greatest visual acuity may be useful, but this limits the field of vision very greatly.

Contact Glasses may be of service. These are devices which, in the usual Zeiss form, are thin shells of glass fitting on to the sclera adjacent to the cornea by suction, and separated from the cornea by a tiny space which is filled with normal saline. Some experience is necessary before it is easy for the surgeon to select the correct glass for any individual case, and often two or three have to be tried. A second variety, the Mueller form, is like a transparent artificial shell eye, which fits over the whole front of the sclera and has to be specially made in Vienna for each case. With both varieties the refractive error, due to corneal irregularity, is banished by the provision of the artificial cornea. Many patients can apply these glasses for themselves after some practice, but they require special care in fitting in the first instance, and the eye has to be gradually accustomed to them. They can seldom be worn for longer than a few hours at a time. Their chief use is in cases of conical cornea, but in any case of irregular astigmatism



FIG. 143.—THERMO-CAUTERY FOR CONICAL CORNEA.

which glasses do not sufficiently benefit, or when, for any reason, cosmetic or otherwise (*e.g.*, for actresses), the use of spectacles is not considered advisable, they may be of service. They are costly (£5 to £6), and, in practice, their uses are very limited.

Repeated paracentesis, followed by the long-continued application of a pressure-bandage, pilocarpine or eserine to diminish tension, and abscission or cauterization of the apex of the cone to cause flattening by subsequent cicatrization (followed by optical iridectomy to enable the patient to see past the central leucoma) have all been tried, with some degree of success.

Worth's method of cauterization consists in making a graduated oval or crescentic excavation, with a thermo-cautery, midway between the apex and periphery of


the cornea, the shape and position of the excavation being so regulated as to reduce the refraction in the meridian of greatest curvature only. Daily paracentesis of the anterior chamber and pressure-bandages for about ten days are an essential part of the after-treatment. No iridectomy is needed.

A Graefe knife section of the cornea through the apex of the cone, at an angle of 45 degrees from the vertical meridian, followed after two weeks, when the scar has healed, by a second section at right angles to it, has proved beneficial in some cases, from contraction of the scars.

Keratoglobus is a globular enlargement of the cornea, which, although occasionally met with under other circumstances, is usually one of the manifestations of congenital glaucoma, and is described under this title (p. 235).

Opacities of the Cornea.

This term refers to a lack of transparency of the cornea resulting from inflammation, ulceration, or injury. According to density, the corneal opacity is called *nebula* when faint and cloud-like (Fig. 144), often overlooked until examined by oblique illumination; *macula* when more pronounced and



OPACITIES OF THE CORNEA.

Fig. 144, Nebula. Fig. 145, Macula. Fig. 146, Leucoma.

appreciable as a gray spot in daylight (Fig. 145); *leucoma* when dense and white (Fig. 146). When the iris is attached to the scar tissue, the condition is spoken of as *adherent leucoma* (Fig. 134, Plate XI.).

Opacities of the cornea interfere with perfect vision when they involve or encroach upon the pupillary area, the degree depending upon their density. Even slight opacities cause

considerable visual disturbance on account of the resulting diffusion of light. Denser opacities cause disfigurement.

Treatment.—Various measures are used to reduce the density of corneal opacities, or, if faint, to cause their disappearance. Such remedies are of value only when the opacity is recent. They act most successfully in children and when the change is superficial. Most commonly the ointment of the yellow oxide of mercury is placed in the conjunctival sac, after which the cornea is massaged for a few minutes, and then hot compresses are applied. Dionin may be of service.

When such measures are unsuccessful, and the leucoma entirely occludes the pupillary area, an iridectomy for artificial pupil may be performed, the coloboma being made opposite a clear part of the cornea.

One should not be in a hurry to make an artificial pupil in the eye of a child, especially if the opacity is the result of interstitial keratitis. One occasionally sees a patient whose cornea has only the faintest opacity, but on whom an optical iridectomy was performed years ago when the opacity was presumably quite dense.

To remove the disfigurement in cases of leucoma, tattooing and 'colouring' are often resorted to. The eye is anæsthetized, and the leucoma covered with a thick paste of India ink. The



FIG. 147.—MULTIPLE TATTOOING-NEEDLES.



FIG. 148.—GROOVED TATTOOING-NEEDLE.

pigment is then introduced obliquely into the corneal substance, either by means of an instrument consisting of a row

or bundle of round needles (Fig. 147) or with a grooved needle (Fig. 148). The colour fades in the course of a few years, and then the operation may be repeated. When the opacity covers only a part of the pupillary area, tattooing is useful in preventing the diffusion of light which is so annoying to the patient. When an area is to be 'coloured,' the superficial epithelium is scraped off and a neutral 4 per cent. solution of

gold chloride is applied freely on cotton-wool and allowed to remain for four minutes before being washed away. A very dark brown stain remains.

These operations are contra-indicated when the eye is at all irritable, and at any time are not without risk.

Injuries of the Cornea.

These comprise foreign bodies, burns, and wounds.

Foreign bodies, consisting of iron, coal, ashes, dust, etc., frequently adhere to or become embedded in the cornea, causing much pain, lacrymation, and photophobia. When the foreign body is small, it may be difficult to detect, unless we make use of oblique illumination. The mischief which a foreign body provokes depends upon the depth to which it penetrates, and whether or not it is infected. If present for a number of days, a surrounding area of infiltration appears, resulting in a small ulcer, and in this manner the foreign body may become dislodged. If it consists of iron or steel, this ring will become stained by rust. Foreign bodies are frequently the cause of ulcers of the cornea.

To Remove a Foreign Body.—The eye is cocainized. The patient is seated, facing a good light, with the surgeon standing behind and supporting the head. The lids are separated, and the eye-ball is steadied by the fingers of the left hand. The index




FIG. 149.—METHOD OF REMOVING A FOREIGN BODY FROM THE CORNEA, THE SURGEON STANDING BEHIND THE PATIENT.

finger is applied to the margin of the upper lid, and the middle finger to the lower lid, and the two fingers are separated, at the same time gently pressing backward (Fig. 149). If the patient is seated in a chair provided with a

suitable head-rest, the surgeon may stand in front and to the left. In separating the lids, the index finger is then applied to the lower lid and




FIG. 150.—METHOD OF REMOVING A FOREIGN BODY FROM THE CORNEA, THE SURGEON STANDING IN FRONT AND TO THE LEFT OF THE PATIENT.

the middle finger to the upper (Fig. 150). The instruments used are either the blunt spud, the gouge, or the foreign-body needle; these should be sterilized before use. When the foreign body is superficial, the blunt spud will answer (Fig. 151). Very often it can be removed with a little absorbent cotton wound around the end of the spud. When the foreign body has penetrated into the corneal substance, it must be picked or dug out with the needle (Fig. 153). The gouge (Fig. 152) is nowadays rarely used. In such

cases the instrument is passed behind the foreign body. The wound which results must be kept clean by frequent irrigation with solution of boric acid; frequently a protective bandage is indicated. If a ring of rust is present, this also should be removed. Care must be taken to inflict as little injury as possible, and, when the foreign body is deep, not to perforate the cornea. If it penetrates into the anterior chamber,

a keratome (Fig. 178) should first be passed through the cornea and behind the foreign body, so that the latter



FIG. 151.—FOREIGN-BODY SPUD.



FIG. 152.—FOREIGN-BODY GOUGE.




FIG. 153.—FOREIGN-BODY NEEDLE.

will not be pushed into this space during efforts at removal.

Burns of the cornea are treated like similar conditions of the conjunctiva.

Wounds may be non-penetrating or penetrating. Non-penetrating wounds are most commonly abrasions due to scratching with the finger-nail, twig of a tree, or the like. Such injuries, though very painful, heal readily unless infected. They should be kept clean by frequent irrigation with solution of boric acid. A bandage may be indicated, and sometimes atropine.

Perforating wounds are more serious, owing to the danger of prolapse of the iris and injury to the deeper parts. They should be treated by thorough cleansing, atropine, or eserine, according as they are central or peripheral, and a pressure-bandage. If extensive and irregular, but probably not infected, it may be advisable to make two curved concentric incisions in the conjunctiva and slide a bridge of the membrane over the wound.

In penetrating wounds of the cornea the question of the entrance of a foreign body into the globe presents itself, and then an X-ray examination is indicated, or the use of the giant magnet when iron or steel fragments are possibilities.

CHAPTER X

DISEASES OF THE SCLERA

Anatomy.—The sclerotic coat (sclera) is the tunic which with the cornea forms the external fibrous layer of the eyeball. It is strong, opaque, and inelastic, and serves to maintain the form of the globe. Its thickness is about 1 mm., but varies at different points. Its structure resembles that of the cornea, being composed of bundles of connective tissue with some elastic fibres, disposed in both longitudinal and transverse layers. These are connected by cement substance containing cell spaces, in which are lodged connective-tissue corpuscles. These parts are, however, much less regularly arranged than in the cornea. In the child, the sclera often has a bluish-white colour, owing to its being thinner, and allowing the dark pigment of the choroid to show through. The sclera is pierced about 2.5 mm. internal to the posterior pole of the eye by the optic nerve. Here it has blended with it the external fibrous sheath of the nerve. The part through which the nerve passes is known as the lamina cribrosa.

The outer surface of the sclera is white and smooth, covered by Tenon's capsule and the conjunctiva, to which it is enjoined by loose connective tissue (episcleral). In front it presents the insertions of the extrinsic muscles of the eyeball. Its inner surface, brown and rough (lamina fusca), is united by filaments of pigmented connective tissue to the choroid and lined by endothelium, forming the outer wall of the supra-choroidal lymph-space. Where it is pierced by vessels and nerves, a communication between the capsule of Tenon and the supra-choroidea is established. Though traversed by many bloodvessels, the sclera itself has a very scant vascular supply; but the episcleral tissue contains numerous vessels.

Affections of the sclera include superficial inflammation (episcleritis), deep inflammation (scleritis), staphyloma, and injuries.

Inflammation of the sclera (scleritis) may be either superficial or deep. The superficial form, called episcleritis, is limited to the tissues superficial to the sclera, and is relatively

harmless. The deep form, known as scleritis, involves the sclera itself, and extends to subjacent and contiguous parts, causing serious consequences. There is often an absence of a very sharp line of division between the two forms.

Episcleritis.

A not uncommon inflammation of the episcleral tissue, including the superficial layers of the sclera.

Symptoms.—There are some discomfort, lachrymation, slight pain, and photophobia. Occasionally these symptoms are more marked. A flat or somewhat raised inflammatory patch of a purplish colour is seen, usually on the temporal side, adjacent to the cornea or a short distance from the limbus (Fig. 122, Plate X.). There is no tendency to suppuration or ulceration. After a few weeks the purple spot will disappear; but others are apt to take its place. In this way the process may encircle the cornea. Owing to this tendency to relapses, the disease often lasts many months. Sometimes some discoloration of the sclera remains, but there is no interference with vision. Occasionally the adjacent portion of the cornea is implicated. The disease may resemble a marked case of phlyctenular conjunctivitis. It may merge gradually into scleritis.

Etiology.—It is usually observed in adults, especially in women. It is often found in 'rheumatic' and gouty individuals. Syphilis, tuberculosis, and menstrual disorders are predisposing factors.

Treatment should be of a sedative nature: warm fomentations; if photophobia be pronounced, smoked coquilles; if the cornea be implicated, atropine. Instillations of a 1 per cent. solution of holocain in 1:10,000 adrenalin will relieve the discomfort. Sub-conjunctival injections of bichloride of mercury (1:5,000) have been recommended, but injections of normal salt solution appear to be just as efficient and much less painful. The ointment of the yellow oxide of mercury, applied with gentle massage, is often of value, especially when the disease shows a tendency to become chronic. In chronic cases dionin may do good. When there is a rheumatic history,

sodium salicylate or aspirin in large doses (10 to 15 grains every two hours) should be given. In some cases tuberculin injections are indicated after a focal or general reaction to a diagnostic tuberculin test (p. 473). Other constitutional disorders should be prescribed for. Iodide of potassium may be ordered, also hypodermic injections of pilocarpine. In the chronic cases eserine often does good.

Transient Periodic Episcleritis is a variety of episcleritis which appears in sudden attacks lasting several days, reappears at intervals of several weeks or months, and may recur for years. It is seen in gouty and rheumatic adults. The treatment is that recommended for episcleritis.

Scleritis.

An infrequent inflammation of the sclera, in which the symptoms are acute, the course is prolonged, and the consequences are serious. In this disease the entire thickness of the sclera is involved, and after softening it tends to become thin and staphylomatous. Both eyes are frequently involved. Relapses are very common.

Symptoms.—Pain, usually severe, and frequently radiating to neighbouring regions, tenderness over ciliary region, lacrymation, and photophobia. The tension of the eyeball is frequently increased; secondary glaucoma often ensues.

There are well-marked dusky or violet patches adjacent to the cornea, often extending to the equator and frequently surrounding the limbus.

Complications.—The cornea is frequently implicated, and sclerosing keratitis (p. 156) ensues. Secondary glaucoma often results. Not uncommonly there are iritis, cyclitis, choroiditis, opacities of the vitreous, and secondary glaucoma; the combination of such complications is known as anterior uveitis. As a result of these changes, vision is often seriously interfered with, and sometimes lost. The thinning of the sclera results in staphyloma of the anterior portion of the globe, which presents a violet appearance and causes myopia.

Etiology.—The disease is most common in adults, and especially in women. Tuberculosis, syphilis, disorders of

menstruation, focal infections, or sometimes gout, are the predisposing causes. Exposure to cold is sometimes the exciting cause.

Pathology.—The middle layers of the sclera become œdematous, and there is infiltration of lymphocytes, polymorphonuclear and red cells; the swollen lamellæ break down and become necrotic; they may be replaced by fibrous tissue. The infiltration extends into the cornea and uveal tract. Later there is thinning of the sclera.

Treatment comprises the measures advocated in episcleritis energetically applied. In addition, the complicating keratitis and uveitis require appropriate treatment. Tuberculin injections are effective in some cases. After the acute symptoms have subsided, an iridectomy is sometimes advisable for diminishing glaucomatous tension or reducing the staphyloma, or for establishing an artificial pupil.

Staphyloma of the Sclera.

A thinning and bulging of the sclerotic occurring either at the anterior portion (ciliary region), the equator, or the posterior portion of the eyeball.

Ciliary and equatorial staphylomata are usually secondary to inflammations of the uveal tract and sclera which have caused softening of the sclera with or without increased tension. The former present a bluish-gray bulging, which may be limited, or may extend all around the cornea (Fig. 154). This bulging shows a tendency to increase; occasionally it bursts. Histologically the distinction is made between true ciliary staphyloma in which the ciliary body itself gives way, and intercalary staphyloma in which the bulging part is between the limbus and the ciliary region. Equatorial staphylomata are mostly seen in globes excised as being blind and painful from chronic glaucoma. Iridectomy, if feasible,




FIG. 154.—STAPHYLOMATA OF THE SCLERA.

is the only treatment, and is done for the purpose of arresting the process. In some cases, when the enlarged eyeball causes much discomfort and is sightless, enucleation or evisceration is advisable.

Posterior staphyloma is of common occurrence, and is associated with high myopia and choroiditis (p. 197). It is seen with the ophthalmoscope, presenting a white area of variable width which embraces the optic disc (Fig. 172, Plate XV.).

Injuries of the Sclera.

The important injuries include rupture and perforating wounds. These are serious on account of the danger of escape of the contents of the eyeball and infection of the interior.

Small, clean, perforating wounds often heal without reaction, and require no other treatment than cleansing and the application of a bandage.

Large, gaping wounds and ruptures frequently allow escape of the vitreous. There will be blood in the vitreous, diminished tension, and some of the underlying tissues (choroid, ciliary body, or iris), varying with the position, will be found in the wound. Such wounds should be cleansed, the prolapsed parts returned when not too seriously injured, and the opening closed by sutures in the sclera, or preferably through the conjunctiva. The patient must remain absolutely quiet, and the eye should be bandaged.

Sometimes such wounds fail to excite much inflammatory reaction; then they heal quite readily, often with incarceration of the prolapsed parts in the scar. But frequently they give rise to panophthalmitis, with ultimate phthisis bulbi, or to plastic iridocyclitis, with loss of sight. When the wound involves the ciliary body, iridocyclitis is apt to be set up, and the injury becomes more dangerous on account of the liability of such wounds to excite sympathetic ophthalmitis.

Ruptures of the sclera are produced by blows and blunt instruments. They usually occur near the corneal margin, generally above. The conjunctiva may not be broken.

When injuries of the sclerotic are very extensive and cause considerable loss of contents of the eyeball, and when we believe that useful sight cannot be hoped for, the eyeball should be removed at once. This becomes still more urgent when the wound involves the dangerous zone, the ciliary region.

The presence of a foreign body in the eye is a serious complication. The attempt should be made to extract the foreign body, as described on p. 242.

In any case in which there is any doubt as to the presence of a foreign body within the globe, X-ray examination should be made.

CHAPTER XI

DISEASES OF THE IRIS

Anatomy and Physiology.—The second or vascular coat of the eye (uvea or uveal tract) lies immediately beneath the sclera. It provides for the nourishment of the eyeball, and it is formed of three parts, which from before backward are known as the iris, the ciliary body, and the choroid. These three portions are so intimately associated that when one part becomes diseased, the others frequently participate.

The *iris* is a coloured membrane, circular in form, hanging behind the cornea immediately in front of the lens, and perforated in its centre by an aperture of variable size, the pupil. It serves to regulate the amount of light admitted to the interior of the eye, and cuts off the marginal rays which would interfere with the sharpness of the retinal image. Its peripheral border springs from the head of the ciliary body and the ligamentum pectinatum. Its free inner edge, the boundary of the pupil, lies upon the anterior capsule of the lens when the pupil is contracted or moderately dilated; with maximum dilatation of the pupil it hangs free in the anterior chamber. The iris separates the anterior from the posterior chamber of the eyeball. Its anterior surface presents great variation in colour in different eyes, and is marked by radially directed, wavy lines, converging toward the circle of irregular elevations and small depressions (crypts) situated near the pupil; other finer lines are seen extending from this ring to the pupil. This appearance is produced by the subjacent blood-vessels.

In structure the iris consists of a delicate, spongy connective-tissue stroma, containing branched pigmented cells, muscular fibres, and an abundance of vessels and nerves. It is covered anteriorly by endothelium, and posteriorly by the posterior limiting membrane and the retinal pigment layers.

The colour of the iris depends partly on the pigment in the stroma cells, which is variable, and partly on that in the cells of the retinal layers, which is constant.

The muscular tissue, the *sphincter pupillæ*, is a narrow band, about 1 mm. wide, situated close to and encircling the pupil posteriorly,

and supplied by the *third nerve*. The posterior limiting membrane consists of fibres which extend from the ciliary to the pupillary margin, which are regarded by some authorities as consisting of unstriped muscle fibres, and as contributing to the active dilatation of the pupil. The chief factor in active *dilatation* of the pupil is, however, the contraction of the thick muscular coats of the arteries, under the control of the *sympathetic*.

The posterior surface of the iris is covered by two strata of pigmented cells, the uveal layer, which extends to the free border, around which it turns a little, forming the black fringe of the pupillary margin.

The *vessels* of the iris come from the two branches of the ophthalmic known as the long posterior ciliary arteries. Each artery divides into an upper and a lower branch. These anastomose with the corresponding vessels of the opposite side and with the anterior ciliary, and form a vascular ring just behind the attached margin of the iris, the greater vascular circle of the iris. This gives off branches to the ciliary body and iris; the iris branches converge towards the pupil, and here form by anastomosis a smaller vascular circle, the lesser vascular circle of the iris. The veins of the iris follow the arrangement of arteries just described; in addition they communicate with the canal of Schlemm. They chiefly pass backward to the *venæ vorticosæ*.

The *nerves* are given off from the plexus in the ciliary body, and are derived from the third, the nasal branch of the ophthalmic, and the sympathetic.

Pupillary Membrane.—In the *fœtus* the pupil is closed by a thin, transparent, delicate membrane—the pupillary membrane. The membrane and its vessels are gradually absorbed in the seventh or eighth month of *fœtal* life. A few shreds may remain at birth; occasionally part or all of the membrane persists (persistent pupillary membrane.)

Occasionally the iris does not develop and *congenital aniridia* results. Rather more frequently it develops imperfectly, leaving a gap, frequently symmetrical, and usually situated below and to the nasal side. This is known as a *congenital coloboma* of the iris. It may or may not be associated with a similar defective development of the choroid.

Iritis.

An inflammation of the iris, which may be divided into the following:

Varieties.—(1) According to its course, into acute, sub-acute, and chronic.

(2) According to the pathological products, into plastic, serous, purulent, and tuberculous.

(3) According to its etiology, into syphilitic, rheumatic, gouty, diabetic, gonorrhœal, tuberculous, scrofulous, traumatic, sympathetic, secondary, and idiopathic.

The classification according to the nature of the pathological products is unsatisfactory, because one type merges into the other. It is better, therefore, to consider iritis in general, and then to mention the peculiarities of the different forms which have been named according to their etiology.

Objective Symptoms.—The iris looks altered (Fig. 156, Plate XII.). It appears swollen, dull, loses its lustre, its markings become indistinct, its colour changes and becomes greenish in blue or gray irides, and muddy in darker varieties. These changes are due to congestion of the iris and exudation of cells and fibrin into its substance; also to exudation into the anterior chamber.

The pupil is contracted, grayish, sluggish in action, and irregular (Fig. 155); the last peculiarity is due to adhesions




FIG. 155.—POSTERIOR SYNECHIÆ
CAUSING IRREGULAR PUPIL IN
IRITIS.

between the posterior surface of the iris and the anterior capsule of the lens (posterior synechiæ), best seen after the instillation of atropine.

The contents of the aqueous chamber show changes: there is frequently turbidity; there may be more or less dust-like deposit on

the posterior surface of the cornea (keratic precipitates or 'k.p.'), which often involves the lower part (Fig. 158, Plate XII.), or may give a cloudy appearance to the entire cornea. In this exudation there may be pus, which then gravitates to the bottom (hypopyon), or fibrin, which coagulates into a grayish mass (spongy iritis), or there may be blood (hyphæma). The anterior chamber may be deeper than normal. The tension of the eyeball, though usually normal, may be increased.

The anterior capsule of the lens may present evidences of



FIG. 156.—IRITIS IN LEFT EYE.




FIG. 157.—SYPHILITIC NODULES IN IRIS.




FIG. 158.—KERATITIS PUNCTATA.




FIG. 159.—PANOPHTHALMITIS.




FIG. 160.—PANOPHTHALMITIS.

exudation, and also small spots of uveal pigment where posterior synechiæ have been torn away.

There is always marked circumcorneal injection, and with this pink zone there is more or less conjunctival congestion.

Subjective Symptoms consist of pain, photophobia, lacrymation, interference with vision, and sometimes general malaise.

The pain is often severe, neuralgic in character, radiating to the forehead and temple, and worse at night. It is sometimes accompanied by tenderness of the eyeball, a symptom pointing to involvement of the ciliary body.

The diminution in the acuteness of vision depends upon the cloudiness of the anterior chamber and the deposits in the pupil and upon the posterior surface of the cornea. When very marked it indicates extension of the inflammation to the deeper parts.

Differential Diagnosis.—Iritis is most frequently mistaken for acute catarrhal conjunctivitis. Sometimes acute glaucoma is mistaken for iritis. The differential points between iritis and conjunctivitis are given in the table on p. 176.

With the differential points given in these tables in mind, there ought to be little difficulty in making a differential diagnosis in these three affections.

The error of mistaking iritis for conjunctivitis would be unfortunate in delaying the immediate instillation of atropine. But the error of mistaking glaucoma for iritis would be most serious, since the use of atropine is disastrous in glaucoma, while of the utmost importance in iritis.

Course.—Iritis may be acute and run its course in several weeks, or it may be chronic and last a number of months. A great many cases terminate favourably, especially when subjected to proper treatment early; the exudation becomes absorbed, and the iris returns to a normal condition, with no evidences or mere traces of former inflammation. Chronic cases present very mild inflammatory symptoms, or the latter may be entirely absent. Certain forms of iritis have a tendency to recur. Iritis may involve one or both eyes; when both eyes are attacked the second usually is affected a short time after the first.

<i>Acute Iritis.</i>	<i>Acute Conjunctivitis.</i>	<i>Acute Glaucoma.</i>
1. Iris swollen, dull, and discoloured. 2. Pupil small, gray, sluggish, irregular after use of atropine. 3. Anterior chamber of normal depth (deeper in serous form) and presents exudation. 4. Cornea transparent (may present deposits on posterior surface) and sensitive. 5. Ciliary (circumcorneal) injection; pink zone of fine vessels surrounding cornea and fading towards fornix. 6. Conjunctiva usually transparent. 7. Lacrymation, but no discharge. 8. Tension usually normal (occasionally increased). 9. Some ciliary tenderness. 10. Pain radiating to forehead and temple; worse at night. 11. Dimness of vision.	1. No change in iris. 2. Pupil is normal. 3. Anterior chamber normal. 4. Cornea transparent. 5. Conjunctival injection; coarse meshes, most pronounced in fornix, and fading towards the cornea. 6. Conjunctiva reddened and opaque. 7. Mucous or muco-purulent discharge. 8. Tension normal. 9. No ciliary tenderness. 10. Discomfort, hot, gritty feeling, but no real pain. 11. No interference with vision, except blurring caused by the discharge smeared over the surface of the cornea.	1. Iris congested, discoloured, dull, periphery pushed forward. 2. Pupil dilated, oval, immobile. 3. Anterior chamber shallow and aqueous; sometimes turbid. 4. Cornea steamy and sensitive. 5. Ciliary and episcleral injection (also conjunctival congestion). 6. Conjunctiva congested and chemotic. 7. Lacrymation, but no discharge. 8. Tension increased. 9. Ciliary tenderness. 10. Severe pain in and about eye, with headache. 11. Marked dimness of vision.

Complications.—The neighbouring parts of the eye are sometimes involved in severe forms of iritis—conjunctiva, cornea, ciliary body, choroid, vitreous, optic nerve, and retina. The association of inflammation of the ciliary body (cyclitis) with iritis (iridocyclitis) is so common that some authors describe the two conditions together, and regard pure iritis as rare. The following symptoms, occurring in the course of an iritis, point to the existence of cyclitis: Violent inflammatory symptoms, marked diminution in vision, tenderness in the ciliary region, deposits upon the posterior surface of the cornea, and increase or decrease of normal tension.

Sequelæ.—There are often posterior synechiæ and deposits upon the anterior lens capsule; less frequently there are seclusion of the pupil, occlusion of the pupil, atrophy of the iris, opacities of the vitreous, deposits upon the posterior capsule of the lens, and cataract. In seclusion of the pupil (annular posterior synechia), the iris is bound down throughout its entire pupillary margin, the pupil remaining clear (Fig. 162). This causes a loss of communication between the anterior and the posterior chamber; the aqueous secreted by the ciliary processes is hemmed in, the iris stretched (iris bombé) and atrophied, glaucoma results, and if unrelieved blindness follows. Occlusion of the pupil is a filling in with opaque exudate (Fig. 163).

Etiology.—Iritis is most frequently dependent upon some systemic disorder; in many cases, syphilis. It often depends on a toxin derived from a septic focus in some part of the body; this class includes infection from diseased teeth (pyorrhœa and periapical infection), tonsils, nose and nasal accessory sinuses, prostate, female genital organs, or the intestinal tract. Not uncommonly it is caused by chronic gonorrhœa; the so-called 'rheumatic' cases are probably due to focal infection. Much less commonly it is dependent upon tuberculosis, acute infectious diseases, or diabetes. It may be secondary, traumatic, or sympathetic.

Pathology.—Inflammation of the iris presents similar changes to those occurring in other connective tissue, modified

by the great vascularity of this membrane and the looseness of its stroma. There are dilatation of the bloodvessels and exudation of lymph, lymphocytes, and fibrin into the stroma and anterior chamber. The cells which find their way into the anterior chamber cause cloudiness of the aqueous and adhere to the posterior surface of the cornea, forming keratic precipitates. All these products of inflammation may be completely absorbed, or they may become organized, the fibrinous material forming adhesions between the iris and the




FIG. 161.—SECTION OF THE ANTERIOR PORTION OF THE EYEBALL, SHOWING THE IRIS IN ITS NORMAL RELATIONS.




FIG. 162.—SECTION SHOWING ANNULAR POSTERIOR SYNECHIA (SECLUSION OF THE PUPIL). IRIS BOMBÉ.




FIG. 163.—SECTION SHOWING TOTAL POSTERIOR SYNECHIA AND OCCLUSION OF THE PUPIL.

anterior capsule of the lens, and also deposits on the lens capsule. In chronic iritis and after repeated attacks of acute iritis, the iris may atrophy and degenerate; it then presents a change in colour to a dirty gray, red lines on its surface representing new bloodvessels; the pupil is sluggish in reaction and its margin thin and ragged.

Treatment.—(1) Atropine, (2) dionin, (3) leeches, (4) hot fomentations, (5) rest, (6) protection from light, (7) treatment of etiological factor.

Atropine diminishes the congestion of the iris, puts this part at rest, causes mydriasis, and thus prevents adhesions and tends to break up those which have already formed. It should be instilled every two hours at first, and after the pupil is dilated three or four times a day. When the inflammation is pronounced, the pupil will not dilate rapidly. The action of atropine is often increased by the addition of cocaine. Sub-conjunctival injection of a drop of atropine solution, or better, adrenalin, or a mixture of both, made near the limbus corresponding to the situation of an obstinate adhesion, will often cause its release, if recent. In certain cases symptoms of atropine-poisoning occur, either local or constitutional, necessitating the substitution of some other mydriatic (duboisine, hyoscyamine, scopolamine—see Chapter XXXI.); a solution of the aqueous extract of belladonna (1:8) often acts well in such cases. Exceptionally atropine causes an increase in inflammatory symptoms, and must be stopped; then a miotic may be of service. This action is apt to occur when there is increased tension, and sometimes when there is a complicating cyclitis. The use of miotics in iritis requires constant supervision, lest adhesions form and secondary glaucoma supervenes.

Dionin relieves pain to a certain extent, and favours the absorption of exudates, by locally stimulating the circulation of blood and lymph.

Two or three *leeches* applied to the temple usually produce a favourable effect. The use of the "artificial leech" is nowadays practically abandoned.

Moist, warm *compresses* for several hours each day diminish the pain and the inflammation in acute cases. In traumatic iritis iced compresses may be used in an attempt to check the inflammation in the early stage.

Absolute *rest* in bed in the early stage of acute iritis is an important aid in treatment.

Protection from light by means of smoked glasses or a shaded room is essential.

The injection of foreign protein (*e.g.*, 5 to 10 c.c. of sterile milk into the buttock) is often of great service, especially in cases with fibrinous exudate.

Other important indications are light diet, abstinence from alcohol, a brisk purge, and avoidance of all use of the eyes for near work.

Constitutional treatment must meet the indications in the different forms. In syphilitic iritis mercury is given, usually by inunction, up to the point of salivation; after the acute symptoms have subsided, mixed treatment (mercury and iodide of potassium) is substituted. Injections of salvarsan often produce rapid improvement. In certain cases in which no cause can be discovered small doses of mercury have a favourable effect. In rheumatic cases large doses of salicylate of sodium combined with small doses of aspirin improve the condition and relieve pain. This remedy also has a quieting effect upon the pain in other forms. Tuberculous cases often respond to general treatment by ultra-violet light; in some cases tuberculin is indicated. Gonorrhœal cases benefit from vaccines, and, at a later stage, from prostatic massage and irrigation of the posterior urethra.

Paracentesis is occasionally resorted to for the purpose of relieving tension, and also in obstinate cases to produce a favourable effect upon the progress of the disease. Iridectomy was formerly performed for the same reasons. As a rule, however, operative procedures are useful only after the inflammatory symptoms have subsided, for the purpose of remedying sequelæ.

Clinical Varieties.—*Syphilitic iritis* is a very common form. It occurs in the secondary stage of acquired syphilis; it is usually acute; the second eye becomes involved soon after the first; there is more or less plastic exudation; there are broad synechiæ; in some cases there are yellowish vascular nodules, usually situated at the pupillary border of the iris (Fig. 157, Plate XII.); pain is not pronounced; if properly treated relapses are not common. There are often other manifestations of syphilis in the eye, choroido-retinitis, etc. Occasionally interstitial keratitis is complicated with iritis.

An acute form of iritis is often ascribed to *rheumatism*. Perhaps it would be more correct to say that rheumatism and iritis are often due to a common cause. It frequently attacks

only one eye, but more often both; it occurs especially in adults; the effusion is serous with a few cells; pain is pronounced; relapses are common. Probably many of these cases are of gonorrhœal origin.

Gonorrhœal iritis is not uncommonly seen in persons suffering from gonorrhœa or gleet; many cases called rheumatic are really gonorrhœal. The exudate is plastic, pain is pronounced and relapses are common. Injections of gonococcic vaccine are indicated and are often valuable, as are also massage of prostate and irrigation of the posterior urethra, which are employed after the iritis has quieted down, to cure the cause and prevent relapses.

Iritis from Septic Infection.—Decayed teeth (pyorrhœa, periapical abscess), septic tonsils, infection of the nasal accessory sinuses, digestive trouble, colitis, chronic constipation, a septic wound, or any other source of auto-intoxication may cause iritis. The cause should be sought for, and if possible removed.

Diabetic iritis is plastic in character and chronic in course.

Tuberculous iritis is an uncommon form which occurs chiefly in children and young adults, and is not necessarily associated with tuberculosis in other parts. It occurs under two forms: (1) the miliary, in which small grayish nodules develop near the ciliary or pupillary border; (2) a single yellowish-gray mass in the ciliary part of the iris. The tuberculous deposits may, under favourable conditions, be completely absorbed. On the other hand, the process may lead to the destruction of the eyeball. The treatment is that of iritis in general, and of tuberculosis.

In a case in which the eye has been destroyed, but in which there is no evidence of tuberculosis in any other part, the eyeball should be removed.

Tumours of the Iris.

These may be (1) *inflammatory*—(a) syphilitic, (b) tuberculous, both of which have just been described; and (2) *new growths*: cyst, melanoma, and sarcoma, all of which are rare.

Injuries of the Iris.

These may be (1) non-perforating and (2) perforating.

1. *Non-perforating injuries* (concussion, blows upon the eyeball) may cause (a) mydriasis; (b) a tear in the pupillary margin, in both of these cases eserine being indicated;




FIG. 164.—IRIDODIALYSIS.

(c) iridodialysis, a separation of the ciliary border of the iris (Fig. 164), for which atropine is required.

2. *Perforating injuries* are usually complicated by wounds of the lens and other parts of the eye. A perforating wound of the eyeball may lacerate the iris or merely allow the latter to project through a wound of the cornea or of the ciliary

region (*prolapse*). In such cases the wound must be irrigated with a mild cleansing lotion, the prolapse excised, the cut edges carefully separated from the wound by a repositor, atropine instilled and the eye bandaged. If the anterior chamber has re-formed prior to such operation, replacement of the iris pillars is facilitated by making a small keratome incision at the limbus at right angles to, and as far as possible from, the wound. The repositor can be inserted through this incision if found necessary after excision of the prolapsed iris. A foreign body may pass through the cornea and lodge on the iris; in such a case the particle should be removed by fine non-toothed forceps after a preliminary incision with the lance-knife at the limbus; if magnetic it can be removed by a magnet (as described on p. 241). If these efforts are not successful the piece of iris on which the foreign body lies should be drawn through the wound and excised.

Operations upon the Iris.

Iridectomy is the only frequent operation upon the iris. It is described with glaucoma (p. 224), which is one of its indications.

Iridotomy and *Iridocystectomy* are operations, fortunately required infrequently, which are performed when, after loss of the lens following injury or cataract-operation, the pupil has been closed by inflammation or been drawn towards the cicatrix. In iridotomy the iris fibres are cut transversely with a Ziegler's knife, a Graefe knife or with special forceps-scissors introduced through a small corneal incision. In iridocystectomy a keratome incision is made through the cornea, and the capsule and the edge of the iris are drawn out by means of a blunt hook or forceps, and a piece excised. The object of both operations is the formation of an artificial pupil. However perfect the operative effect may be, the visual result is usually marred by vitreous opacities caused by the previous inflammation.

The Pupil.

The normal pupil is circular and regular in outline. It is larger in the young than in advanced life. Its size should equal that of its fellow; both should respond alike when one is subjected to a change in intensity of illumination. The movements of the pupil are contraction and dilatation.

The contracting fibres of the iris, the sphincter pupillæ (muscle), are supplied by the third nerve. The dilating fibres in the posterior limiting membrane are supposed to be supplied by the sympathetic. The bloodvessels of the iris, also supplied by the sympathetic, constitute the principal agents in active dilatation of the pupil; contraction of these vessels causes narrowing of the iris with dilatation of the pupil.

Contraction of the pupil is affected by stimulation of the oculomotor nerve and by paralysis of the sympathetic. Dilatation follows paralysis of the third nerve or stimulation of the sympathetic.

The oculomotor nerve fibres are conveyed through the ciliary

ganglion and short ciliary nerves. The nucleus of origin of the third nerve concerned in the movements of the iris is in the floor of the aqueduct of Sylvius, and can be divided into three portions: (1) that giving rise to the sphincter fibres of the iris; (2) accommodation (ciliary muscle); and (3) convergence (internal rectus). The sympathetic or dilating fibres are given off from the cilio-spinal centre of the lower cervical spinal cord.




FIG. 165.—VISUAL AND PUPILLARY REFLEX PATHS.

L, Left eye; *R*, right eye; *ON*, optic nerve; *LT*, left optic tract; *RT*, right optic tract; *POG*, primary optic ganglia; *OMN*, oculomotor nuclei; *OR*, optic radiations; *OL*, occipital lobe; *CIL*, ciliary ganglion. Division of the fibres at 1 abolishes the reaction of the pupil to light upon illuminating the left half of either retina. At 2, the same result with right homonymous hemiopia. At 3, right homonymous hemiopia with preservation of the reaction of the pupil to light.

The pupil contracts upon exposure to light, with accommodation, and with convergence. The light contraction may be direct or consensual. The direct light reflex is obtained by exposing one eye to increased illumination and observing the contraction of the pupil of this eye. The consensual or indirect light reflex is obtained by throwing light into one eye, and observing the contraction of the pupil of the other eye. The accommodation and convergence reflex is obtained by

directing the patient to look at an object held several inches in front of the face in the middle line; the pupils will be seen to contract. These three actions are associated.

When the pupil diminishes in size as the result of exposure to light, its contraction is not perfectly continuous; there are a number of minute oscillations during the contracting period. When these oscillations are pronounced and noticeable, and independent of illumination, the condition is known as *hippus*; it is bilateral, and of central origin, but its cause and significance are unknown.

The dilatation reflexes of the pupil are seen upon shading the eye (both direct and consensual) and upon looking at a distant object. In addition there is a sensory reflex; when sensory nerves are stimulated, as by scratching or tickling the skin, both pupils dilate.

The consensual contraction is explained by the fact that the light stimulus in one eye is carried by the optic nerve and passes to both optic tracts, and in this way to the nucleus of the third nerve of each side (Fig. 165). Blindness in one eye abolishes the direct reflex in this eye, but its consensual reflex is preserved.

In certain pathological conditions there may be loss of light reflex, without interference with sight. The Argyll-Robertson pupil, so frequently a symptom of locomotor ataxia, contracts with accommodation and convergence, but does not respond to light. It is usually accompanied by miosis. It is explained by an interruption in the path from the optic nerve to the oculomotor nucleus, the connexions of the centres for accommodation and convergence remaining unaffected. It should not be confused with the 'tonic pupil,' which is often associated with absent knee-jerk, but is not an indication of serious disease of the nervous system. Such pupils are usually unilateral; though apparently inactive to light, they often react after a stay in a dark-room. They are larger than their unaffected fellow, and, on convergence, contract only slowly through a wide range. The characteristics of the pupils—size, equality, and reflexes—are of great value in the diagnosis of various affections of the nervous system

and in the localization of cerebral lesions. Hence it is important to be familiar with the afferent and efferent routes which control the movements of the pupil (Figs. 165 and 243A, and Plate XXV.). The course of the afferent impulse is retina, optic nerve, both optic tracts, corpora quadrigemina, nuclei of origin of the third nerve in the floor of aqueduct of Sylvius (there being a communication between the two sides).

The efferent impulse travels on either side from these nuclei to the third nerve, the ciliary ganglion, short ciliary nerves, to the iris.

Mydriatics and miotics are described in the chapter on Ocular Therapeutics, and on p. 383. The hemianopic pupillary reflex is explained on p. 320.

CHAPTER XII

DISEASES OF THE CILIARY BODY

Anatomy.—The ciliary body is that part of the tunica vasculosa which extends backward from the base of the iris to the anterior part of the choroid. It consists of the ciliary processes and of the ciliary muscle. A longitudinal section is of triangular shape, with a narrow base directed forward, giving origin to the iris. The outer side of the triangle is formed by the ciliary muscle; the inner side can be divided into two parts: an anterior, which bears the ciliary processes, and a posterior portion, which is smooth.

The *ciliary muscle* (the muscle of accommodation) consists of non-striated muscular fibres arranged in bundles, anastomosing with one another frequently so as to form a sort of plexus, and running in three different directions—meridional, radiating, and annular. The proportion between circular and longitudinal fibres varies according to the refractive condition of the eye. The circular set is well developed in hypermetropia (Fig. 275, B), but atrophied in myopia (Fig. 276). When the ciliary muscle contracts, it draws the ciliary processes and choroid forward and inward, thus relaxing the suspensory ligament and allowing the lens to become more convex.

The *ciliary processes* consist of about seventy folds or thickenings, arranged meridionally, so as to form a circle. They have the same structure as the rest of the choroid, but are even more vascular. They serve to secrete the nutrient fluids in the interior of the eye which nourish neighbouring parts, especially the cornea, lens, and part of the vitreous. The inner surface of the ciliary body is covered by three layers: externally, a homogeneous membrane continuous with the posterior limiting membrane of the iris; next, pigment epithelium; internally, next to the vitreous, a layer of cylindrical non-pigmented cells.

The ciliary body is supplied by branches from the greater circle of the iris and by the anterior ciliary *arteries*. The *veins*, constituting the greater part of the ciliary processes, pass backward to the *venæ vorticossæ* of the choroid. A part of the veins from the ciliary muscle pass backward, pierce the sclera, and run beneath the conjunctiva

with the anterior ciliary arteries. These constitute the violet sub-conjunctival vessels seen running backward in ciliary injection and in deeper congestion (glaucoma). They anastomose with the conjunctival veins, and communicate with Schlemm's canal.

Cyclitis and Uveitis.

As already pointed out, iritis is frequently associated with cyclitis (iridocyclitis). While unmixed cases of cyclitis do occur, they are uncommon; usually, in inflammation of the ciliary body, adjacent portions of the uveal tract are involved.

Practically, the term iridocyclitis is reserved for those cases in which, with the symptoms of iritis, they are decided evidences of participation of the ciliary body.

Varieties.—Cyclitis (iridocyclitis) may be divided into (1) acute; (2) chronic, or uveitis—(a) mild type, and (b) severe type; (3) purulent; and (4) sympathetic.

1. Acute Iridocyclitis.

Acute iridocyclitis presents the picture of an attack of acute iritis with the addition of the following symptoms pointing to involvement of the ciliary body; marked circumcorneal injection, tenderness in the ciliary region, swelling of the upper lid, turbidity of the aqueous (occasionally hypopyon or hyphæmia), increased depth of anterior chamber, deposits upon the posterior surface of the cornea, abnormal tension (increase or decrease), and greatly reduced vision (due to vitreous opacities and deposits in the pupillary space).

Prognosis varies with the severity of the symptoms and the general health of the patient, but is always serious. Although in many cases the disease may run a comparatively mild course and the eye recover with little injury, a severe attack may result in blindness with atrophy of the globe.

Treatment is that of iritis.

2. Chronic Iridocyclitis or Uveitis.

Chronic iridocyclitis or uveitis is an inflammation of the entire uveal tract (iris, ciliary body and choroid), occurs chiefly in young adults, is chronic in course, has a tendency to

relapse, may involve one eye, but often both. The severity of the inflammation varies, and in this sense the disease is divided into two types: (a) mild, and (b) severe.

Etiology.—Certain constitutional diseases: syphilis, gonorrhœa, tuberculosis, influenza, acute infective diseases, chronic rheumatism, gout, arthritis deformans, and diabetes; to auto-intoxication from the gastro-intestinal tract; not infrequently from the influence of toxins of bacterial origin (focal infections) derived from the gums and teeth (pyorrhœa, periapical abscess), tonsils, nose, nasal accessory sinuses, genito-urinary tract, and skin (furuncles); or the infective agent may arise from perforating wounds of the eyeball, including operations such as cataract extraction.

Pathology.—In addition to the pathological changes which have been described under the heading Iritis (p. 178), which apply, practically, to acute iridocyclitis and the mild form of uveitis, there are the following additions in the severe type: Infiltration of round cells and fibrin in the ciliary processes, in the anterior and posterior chambers, and in the vitreous. The exudate in the ciliary body becomes organized with the formation of new bloodvessels and cysts. The exudate in the posterior chamber organizes and becomes infiltrated with pigment. The iris atrophies and the lens becomes cataractous. The choroid is greatly congested and infiltrated and later undergoes degeneration occasionally with formation of bone or lime plaques.

Symptoms.—(a) *The Mild Type* begins very insidiously with few, if any, symptoms of irritation. The patient complains chiefly of diminution of vision; there may be some ciliary congestion, slight pain, ocular fatigue, and limited photophobia. There occurs an exudate of serum and cells with slight admixture of fibrin, sometimes slightly pigmented, precipitated upon the posterior surface of the cornea as dots forming a triangular area over the lower part, apex above (Fig. 158, Plate XII.). The terms 'keratitis punctata,' or less commonly 'descemetitis,' which are usually employed to indicate this condition, are incorrect, and based originally on a false conception of the pathology of the deposits. The abbrevia-

tion 'k.p.' so frequently used should be taken as indicating 'keratic precipitates,' and not 'keratitis punctata' as heretofore. Sometimes a number of spots coalesce and form small masses, called 'mutton-fat' deposits, on account of their appearance. The anterior chamber is deeper than normal, and the aqueous may be slightly turbid. The pupil is somewhat dilated. Tension may be increased at first and lowered later, or there may be alternations of increase and decrease. Secondary glaucoma is an important complication. There are numerous minute opacities in the vitreous, and the latter may become fluid; there may be involvement of the choroid.

(b) *The Severe Type.*—The subjective symptoms resemble those of the mild form. But the objective signs are much more pronounced: the exudate is considerable, and consists largely of plastic material. On account of the nature of the exudate, this type is known as plastic iridocyclitis, in contradistinction to the mild type, in which the exudate is chiefly serous, which is called serous iridocyclitis. The plastic exudate forms in the anterior chamber, pupillary space, behind the iris, and in the vitreous, forming dense membranes; its subsequent contraction causes detachment of the retina; the iris atrophies; the nutrition of the lens is interfered with, and secondary cataract results; choroid and retina atrophy. Finally, the blind and degenerated eyeball shrinks (atrophy of the eyeball). The eye may now remain quiet, or there may be attacks of pain and tenderness from time to time. The complications are secondary glaucoma, which is quite common, and scleritis.

Prognosis varies with the type of disease. The liability to secondary glaucoma always makes the prognosis uncertain. In many instances of the mild form the disease subsides after a lengthy course, with or without relapses, and the eye returns to an apparently normal condition, with little or no interference with vision; or there may be only small iritic adhesions or vitreous opacities. Such a favourable outcome is possible because the exudate is largely serous with little or no plastic addition, and is capable of absorption.

In the severe form the prognosis is always grave. Vigorous treatment in the early stages may save some useful vision, but very frequently the sight is lost, owing to damage through the deposit, organization, and contraction of the plastic exudate.

Treatment is that of iritis, but the indications are for more vigorous interference. It is especially important to remove every source of infection as promptly as possible. Thus the removal of an infected tooth, a diseased tonsil, or the accumulation in accessory sinuses, the injection of salvarsan in a syphilitic, of tuberculin in a tuberculous subject, and colonic irrigations when there is intestinal auto-intoxication, will often prevent the mild from changing into the severe type.

Locally, atropine, hot moist compresses, hot bathing, or continuous dry heat applied in the form of an electric eye-warmer, or, if this is not available, an electric light bulb in a pad of cotton-wool. Sub-conjunctival injections of normal saline or other solutions are often used, but it is doubtful whether they do any good. Increase of tension in these cases is not a contra-indication for atropine; on the contrary, atropine may cause a reduction in tension. Lævo-glaucosan, which dilates the pupil but reduces tension, may prove very useful. Increase of tension should be dealt with by paracentesis of the anterior chamber, and this may have to be repeated a number of times. Atropine ointment 1 per cent. with a soft paraffin basis is much less irritating than the B.P. preparation made with lard. In obstinate cases the addition of 2 per cent. of cocaine increases the effect of the atropine.

Diaphoresis is very useful. Mercury by inunction or by the mouth, even in the non-specific cases, is sometimes of value; also iodides and large doses of salicylates. The use of thyroid extract in these cases has been reported upon favourably. Injection of foreign protein (*e.g.*, 5 to 10 c.c. milk into the glutei) may be useful.

Where a source of focal infection has been discovered, such as a tooth or the tonsil, a culture may be made from the infective material and then a vaccine. The use of vaccines in these cases is apt to be disappointing, but if other measures

do not quickly cause improvement no possibly hopeful treatment should be neglected.

After the eye has become quite quiet, unless hopelessly lost, an iridectomy may be indicated for optical reasons, or to restore communication between the anterior and the posterior chambers, and thus prevent subsequent glaucoma.

3. **Purulent Iridocyclitis (Purulent Uveitis).**

Purulent iridocyclitis (purulent uveitis) is described under the heading of Purulent Choroiditis on p. 198, and of Panophthalmitis on p. 205.

4. **Sympathetic Iridocyclitis (Sympathetic Uveitis).**

Sympathetic iridocyclitis (sympathetic uveitis) is described under the heading Sympathetic Ophthalmitis on p. 201.

Injuries of the Ciliary Body.

The ciliary region, represented by a pericorneal ring about 6 mm. wide, is known as the 'dangerous zone,' because infected penetrating wounds in this situation are apt to set up plastic uveitis, which may be followed by sympathetic ophthalmitis. After thorough cleansing, extensive ciliary wounds are closed by one or more sutures passed through the superficial layers of the sclera or through the conjunctiva; such wounds are often covered by a conjunctival flap. A prolapse of the iris should be excised, but a prolapsed ciliary body should not be removed. The presence of a foreign body in the globe must be excluded. A bandage is then applied. If the wound is very extensive and sight is lost, enucleation is indicated. Additional details of treatment are given in the paragraphs on Injuries of the Sclera (p. 170) and on Sympathetic Ophthalmitis.

CHAPTER XIII

DISEASES OF THE CHOROID

Anatomy and Physiology.—The choroid is a dark brown membrane placed between the sclera and the retina, extending from the ora serrata to the aperture for the optic nerve. It consists mainly of bloodvessels, united by delicate connective tissue containing numerous pigmented cells; these vessels are arranged according to their calibre into three superimposed layers.

This vascular structure is bounded on either side by a non-vascular membrane; accordingly, the choroid can be divided into five layers: (1) Externally, the suprachoroid, connected with the sclera by loose connective tissues. (2) The layer of large vessels. The spaces between these are filled with connective tissue and cells. The arteries are the short ciliary. The veins are arranged in curves (*vasa vorticosae*) converging to four or five principal trunks, which pierce the sclera behind the equator of the eyeball. (3) The layer of medium-sized vessels. (4) The layer of capillaries (*chorio-capillaris*). (5) The lamina vitrea, or membrane of Bruch, a structureless, transparent membrane, which is placed next to the pigmentary layer of the retina.

The *function* of the choroid is chiefly to serve as a nutrient organ for the retina, vitreous, and lens. It forms the dark coating of the interior of the eyeball.

Inflammations of the Choroid (choroiditis) may be (1) exudative or non-purulent, and (2) purulent.

While in the suppurative form there is implication of the entire choroid together with the rest of the uveal tract, in the exudative type the involvement is limited to certain parts of the choroid; this forms the basis for the classification into a number of clinical forms according to the situation of the areas involved.

Exudative Choroiditis is of frequent occurrence, and is observed at all ages; it is usually chronic in its course. In most instances the disease involves the retina as well as the

choroid and is then, properly speaking, an example of choroido-retinitis or retino-choroiditis. The affection may extend to the anterior portion of the uvea, constituting irido-choroiditis. Although the terminology indicates an inflammation, some of these types of exudative choroiditis are examples of a degenerative process rather than an inflammation.

Varieties.—Exudative choroiditis (Plates XIII., XIV., XV.) occurs in the following principal forms: (1) Diffuse, (2) disseminated, (3) circumscribed, (4) anterior, (5) central (6) juxtapapillary, (7) syphilitic, and (8) myopic. It will be of advantage to describe exudative choroiditis in general before giving the distinctive features of the several varieties.




FIG. 166.—PERIPHERAL SCOTOMATA IN DISSEMINATED CHOROIDITIS.

Subjective Symptoms.—There are disturbances of vision, both diminution in acuteness, appearance of specks, and distortion of objects (metamorphosia). The latter symptom may be divided into micropsia, when objects appear too small, and macropsia, when they appear too large. There are often flashes of light, sparks or bright circles before the eyes. In the later stages there may be defects in the field of vision, both scotomata and peripheral contraction.

These defects may pass quite unnoticed if situated peripherally, but will be very noticeable to the patient if centrally situated. There is no pain.

Objective Symptoms.—There are no external signs, but the ophthalmoscope reveals a well-marked picture. There are patches of exudation of plastic material, varying in size and position. At first these areas are yellowish or yellowish-white in colour, with ill-defined margins. The retinal bloodvessels are seen to be lifted and to pass over them. In very mild cases the exudates may become absorbed, leaving little visible change in the choroid. But usually, after several weeks or months, the exudate becomes organized, and the choroid and retina become atrophied and adherent to each other, giving

PLATE XIII.



FIG. 167.—DIFFUSE CHOROIDITIS.



FIG. 168.—DISSEMINATED CHOROIDITIS.

rise to atrophic areas of various shapes and sizes, which appear whitish in colour due to the deposits of fibrin, and the sclera showing through the thinned or absent choroid; these patches often present visible choroidal vessels and they are generally marked with more or less black pigment, especially at their margins. Not infrequently the vitreous is involved, and then there are opacities of this medium. Very often the retina becomes atrophied opposite the patches just described. The optic disc may participate in the changes, and present a yellowish-white dirty colour, a condition often spoken of as 'choroidal atrophy.'

Complications.—From this description it will be seen that neighbouring structures are frequently implicated: iris, retina, vitreous, and sclera. Choroiditis may also cause posterior polar cataract.

Etiology.—Frequently some constitutional disease, especially acquired and hereditary syphilis, but also anæmia, scrofula, and tuberculosis; many examples are found in myopia. It may depend on septic infection from the teeth, oral and nasal cavities, or upon intestinal auto-intoxication; many cases of obscure origin are spoken of as idiopathic.

Pathology.—In the early stages there are congestion and infiltration of the affected part of the choroid and overlying retina with lymph. This explains the appearance of the soft, fluffy areas seen with the ophthalmoscope. Later the exudation is augmented by leucocytes and fibrin, the membrane of Bruch is broken through, and the leucocytes invade the retina and vitreous, causing opacities in the vitreous. The exudates become organized, fibrous tissue is formed, which destroys the choroid and retina and causes atrophy and adhesion between the two layers in the affected areas. The retinal pigment proliferates and becomes heaped up at the margins, and, to a lesser extent, within the atrophic areas.

Prognosis depends upon the position of the patches of exudation, with subsequent atrophy. A single patch involving the macular region will seriously impair vision. On the other hand, the process may extend over a considerable part of the fundus, and yet vision remain good, if the macula escapes.

Treatment.—Removal of the etiological factor; iodides and mercury in syphilitic cases; inunctions of mercury are often used with success even in non-syphilitic cases; tuberculin in suitable patients; attention to the general health; diaphoresis is often valuable. Rest of the eyes, avoidance of bright light by the use of smoked glasses, sometimes a darkened room; sub-conjunctival injections of salt solution are often useful.

Diffuse Choroiditis (Fig. 167, Plate XIII.).—In this variety the patches of exudation are of large size; later, the coalescence of the atrophic spots produces a large area of white or yellowish-white colour, more or less pigmented, and showing some choroidal vessels.

Disseminated Choroiditis (Fig. 168, Plate XIII.) presents numerous round or irregular spots scattered over the fundus. The entire fundus may be studded, and yet the vision remains good if the macular region escapes. This form of choroiditis runs a very chronic course. After existing a long time, atrophy of the retina and optic nerve may be added.

Anterior Choroiditis, presenting foci of exudation similar to those found in disseminated choroiditis but limited to the periphery of the fundus, is usually due to syphilis, and is not infrequently found accompanying interstitial keratitis. As a rule, the diagnosis is made only in the later stage, when the exudates have changed to white spots with pigmented margins, and then generally only in the course of an ophthalmoscopic examination made for other reasons. This form of choroiditis sometimes accompanies high myopia. Vision is not affected.

Exaggeration of normal fundus pigmentation in the periphery is not unusual in normal eyes; it is often found as a senile change in the aged.

Central Choroiditis is a term which includes a variety of atrophic and degenerative changes situated in the macular region; the designation is incorrect, because the changes are not inflammatory in origin, but it is convenient and in general use. There are many variations in the ophthalmoscopic appearance, from that of a small patch of definite choroidal atrophy with patches of pigment (as illustrated in Fig. 170, Plate XIV.), to a much less definite area of atrophy with numerous small

PLATE XIV.



FIG. 169.—FUNDUS OF MYOPIC EYE.




FIG. 170.—CENTRAL CHOROIDITIS.

To face page 196.

scattered pigment spots, superficially situated in the retina about the macula. The resulting central scotoma causes marked reduction, or loss, of central vision. Such changes occur most frequently in high myopia, syphilis, and old age. Sometimes there is a preceding hæmorrhage, but more often there is none. When the degeneration occurs in an old subject, it is known as senile central choroiditis; in such patients there may be associated arterio-sclerosis of retinal vessels. The condition is usually bilateral, the second eye becoming involved after a varying interval. There is no treatment which is of any avail.

It is not uncommon to find the macular region and the area immediately surrounding it dotted with numerous small, round, yellowish spots, with more or less stippled pigment between them; they are hyaline excrescences of the membrane of Bruch and are known as 'colloid bodies'; they are found in both eyes, usually occur in elderly individuals, and do not affect vision.

Syphilitic Choroido-Retinitis is the name given to a diffuse inflammation of the choroid, associated with retinitis and changes in the vitreous, which occurs in the secondary stage of syphilis. At first there are diffuse cloudiness of the retina, numerous exudations in the choroid, especially in the region of the macula, and fine, dust-like opacities of the vitreous. Later, the cloudiness of the retina is replaced by atrophy, there are atrophic patches of the choroid, spots of pigment, and opacities of the vitreous (Fig. 171, Plate XV).

Choroiditis of Myopia; Posterior Staphyloma, or Sclero-Choroiditis Posterior.—The fundus of near-sighted eyes, especially if the myopia be of high degree, very often presents characteristic changes (Fig. 169, Plate XIV., and Fig. 172, Plate XV.). There is frequently a whitish, atrophic, crescentic area embracing the temporal border of the disc; this is known as *myopic crescent* (Fig. 169, Plate XIV.); it may extend and include the upper and lower margins, or it may encircle the disc; it is sometimes called 'conus,' but the latter term is properly reserved for a congenital anomaly of similar shape, situated at the lower edge of the papilla, found most often in

astigmatic eyes, and due to a gap between the optic nerve head and the choroidal aperture.

Owing to the elongation of the eyeball in high myopia there is a bulging of the sclera at the posterior pole which results, in addition to the error of refraction, in degenerative and atrophic changes in the choroid and retina in this situation; this ectasia is known as *posterior staphyloma* (Fig. 172, Plate XV.). The terms myopic crescent and posterior staphyloma are *not* synonymous.

When the myopic crescent or annular patch is separated from healthy choroid by a sharply defined margin, often pigmented, it is a sign that the process has come to a standstill. But when the border is ill-defined, it indicates that the changes are advancing (progressive myopia). Such knowledge is of great importance in emphasizing the necessity for attention to ocular and general hygiene. The size of the crescent is often, but not always, proportionate to the degree of myopia. More or less superficial atrophy of the choroid is often observed in myopia of high degree, allowing the larger choroidal vessels to become plainly visible. Besides choroiditis in the macular region, there may be patches of choroidal atrophy in other parts of the fundus. These often coalesce with the posterior staphyloma, so that an extensive white area is seen, spotted or bordered with more or less pigment. The early changes in the macular region may be represented by fine lines or fissures. Hæmorrhages, especially in the macular region, fluidity, and opacities of the vitreous also occur in myopia of high degree.

The treatment consists in avoidance of near work, rest for the eyes, smoked glasses, outdoor life, and attention to the general health. During the progressive stage, the treatment recommended for choroiditis in general is indicated.

Purulent Choroiditis.

In this affection the choroid, ciliary body, and iris are involved in a purulent inflammation which leads to the destruction of the eyeball. In most cases all the structures of the eyeball are involved, and panophthalmitis results

PLATE XV.




FIG. 171.—SYPHILITIC RETINO-CHOROIDITIS, WITH
DUST-LIKE OPACITY OF VITREOUS.



FIG. 172.—POSTERIOR STAPHYLOMA.

To face page 198.

(p. 205). Rarely, the inflammation is limited to the choroid (suppurative choroiditis), the purulent exudate forming an abscess of the vitreous with no external evidences of inflammation, but with loss of vision.

The **Symptoms**, those of irido-cyclitis already given, are acute and severe. The pain is severe; the congestion of the conjunctiva is pronounced, with chemosis and swelling of the lids. The cornea becomes clouded, and pus appears in the aqueous and vitreous, so that no details of the fundus are visible. Sight is rapidly lost. There is usually much constitutional disturbance.

In those cases in which the process remains limited, after the acute symptoms subside, a sightless, degenerated, and atrophied eyeball remains. A yellowish or grayish-yellow reflex is then obtained from the interior of the eye, due to the purulent degenerated mass. This is known as pseudo-glioma, on account of its resemblance in colour to glioma of the retina.

Etiology.—Infection of the interior of the eye by pyogenic microbes, either from without or from within the body. Ectogenous infection occurs most frequently from penetrating wounds, including operations, perforating ulcers, thin corneal scars, and prolapse of iris. Endogenous infection results from septic embolism (metastatic ophthalmia), and is seen most frequently in puerperal pyæmia, also surgical pyæmia; extension from orbital cellulitis; also in meningitis and cerebro-spinal meningitis, especially in children, and sometimes infectious diseases.

Treatment.—It is impossible to save sight. Pain should be relieved by morphine and by the local applications of hot moist compresses. If the process has involved all the ocular structures, the treatment of panophthalmitis is indicated.

Coloboma of the Choroid is a congenital defect of the choroid and retina, showing itself in a large white patch, representing the exposed sclera. It is usually situated below the disc. The retinal vessels are seen passing across this patch. There is a scotoma corresponding to the defect. This condition is often associated with coloboma (a cleft) in

the iris, and a notch at the equator of the lens opposite the coloboma.

Rupture of the Choroid sometimes results from contusions of the eyeball. The immediate effect of such an injury is an extravasation of blood into the vitreous. After this is absorbed, a long, yellowish-white streak with pigmented edges is seen, usually in the neighbourhood of the disc and to its outer side.

Tubercle of the Choroid occurs in acute miliary tuberculosis, and in tuberculous meningitis. The tubercles appear as small, yellowish-white spots, surrounded by some retinal haze, vary in number, and are found near the disc, in the macular region, or scattered over the fundus. They resemble the spots seen in recent cases of disseminated choroiditis, but are much smaller. Microscopically they consist of giant cells and small round cells in which tubercle bacilli are found. No treatment is indicated, since the systemic disease is always rapidly fatal when these spots are present.

Tuberculosis occasionally assumes the form of a solitary, irregular mass at the macula, which may be mistaken for glioma in infants, or even for sarcoma in adults. The mass may involve the sclera and perforate; in such cases enucleation is demanded; in others, injections of tuberculin may be indicated.

Sarcoma of the Choroid (see Intra-ocular Tumours, Chapter XV.).

CHAPTER XIV

SYMPATHETIC OPHTHALMITIS—PANOPHTHALMITIS

As its name implies, uveitis is an inflammation of the whole uveal tract—iris, ciliary body, and choroid. There are two forms: (1) serous and plastic, and (2) purulent. Both forms have been described in connexion with cyclitis, which, as already explained, is generally merely part of a uveitis. There are, however, two special varieties: (1) sympathetic uveitis, generally known as sympathetic ophthalmitis; and (2) the form of purulent uveitis known as panophthalmitis.

Sympathetic Ophthalmitis.

Sympathetic ophthalmitis (sympathetic ophthalmia) is a serous or plastic inflammation of the uveal tract in one eye, due to the effects of a similar inflammation in the other.

Etiology and Occurrence.—This inflammation is usually due to traumatic iridocyclitis of the first eye as a result of perforating injury, such as accidental or operative wound in the ciliary region, the danger being increased if iris or ciliary body be entangled in the wound. Foreign bodies in the interior of the eyeball are also apt to excite this disease. Sometimes it results from the iridocyclitis following perforating corneal ulcers. Rarely, if ever, it occurs without any perforating lesion. Suppurative inflammations of an eyeball very rarely cause sympathetic ophthalmitis.

It is, fortunately, not of very frequent occurrence. Children are especially liable. Asepsis and antiseptics have lessened the danger of its occurrence after operations. It usually begins between five and eight weeks after injury in the exciting eye, rarely before three weeks. It may, however, occur after many

months, and it has been seen in a child as early as the ninth day after the injury.

The eye which has been originally affected is known as the exciting eye; the one secondarily involved as the sympathizing eye.

Symptoms.—In most cases, but not invariably, the disease presents a stage known as sympathetic irritation. It is very important to recognize this stage, since removal of the exciting eye at this period will almost certainly prevent the progression of the affection from irritation to actual inflammation.

The Symptoms of Sympathetic Irritation.—The sympathizing eye is 'irritable'; there are marked photophobia and lachrymation; neuralgic pain in the eye and neighbouring parts; dimness of vision occurs when the eyes are used for near work. There may be bright and coloured sensations.

The exciting eye usually presents an iridocyclitis or uveitis, which may be slight or severe. When the sympathizing eye becomes affected, there may be symptoms of irritation and marked tenderness over the ciliary region in the exciting eye.

These symptoms of irritation in the sympathizing eye may be intermittent. Each attack may last a number of days or weeks, then subside, and recur a number of times. They may finally disappear entirely. But, as a rule, if the exciting eye is not excised, sympathetic inflammation results. Examination with the slit-lamp may give valuable information in a very early stage.

The Symptoms of Sympathetic Inflammation.—These may follow directly upon those of irritation, or may occur after the sympathizing eye has been quiet for a time. They may begin acutely or insidiously. When once established, the inflammation is chronic, and its duration is months or even one or two years. In the majority of cases blindness results, though occasionally, if the inflammation be mild, useful vision may be preserved.

The symptoms are photophobia, lachrymation, dimness of vision, and tenderness in the ciliary region. There will be circumcorneal injection, punctate deposits upon the posterior

surface of the cornea, increased depth of the anterior chamber, contracted pupil, and increased tension.

In mild cases the symptoms may not pass beyond those of cyclitis or iridocyclitis; but usually they develop into a plastic uveitis, including iris, ciliary body, and choroid, and giving the following signs: The iris is thickened, its colour becomes changed, its markings are obliterated, and it presents new bloodvessels upon its surface; it is firmly bound down by numerous and extensive posterior synechiæ. The plastic exudation fills up the pupil and more or less of the anterior chamber, which becomes shallow. Tension is diminished. The choroid and retina participate in the plastic inflammation, the vitreous presents numerous opacities, and the lens becomes opaque. Finally, there is detachment of the retina, the eyeball shrinks and passes into the condition of atrophy.

Theories of Transmission.—A number of theories have been put forward to explain the occurrence of inflammation in the sympathizing eye; none has been generally accepted as satisfactory. The theories advanced are: (1) Infection spreading through the sheath of the optic nerve of one side to the chiasma and sheath of the nerve of the opposite eye; (2) the action of a toxin generated by bacteria which have entered the exciting eye; reaching the second eye by lymph channels; (3) a focal allergic reaction to uveal pigment which is set free in the blood stream by the injured eye; (4) metastasis through the blood current of some form of specific organism (filterable virus) which is supposed to be pathogenic for the eye only; since such organisms have not been isolated, they are supposed either to be too small to be seen with the microscope, or else the infective agent is a toxin generated by the specific micro-organisms. At present the last is regarded as the most probable explanation.

Pathology.—The changes in the exciting and in the sympathizing eye are identical; they resemble a tuberculous infiltration and consist of nodular collections of small round cells throughout the uveal tract; later this infiltration becomes more diffuse, and there are added epithelioid and giant cells. The retina suffers very little change. In addition there are

the pathological changes of ordinary uveitis (p. 189) with its destructive sequelæ.

Treatment.—Prophylactic treatment is of the greatest importance, and refers to the removal of the exciting eye under the following circumstances: We should enucleate the injured eye if it be sightless, or its condition such (especially when the ciliary region is involved) that we cannot hope to preserve useful vision. This is particularly imperative if it is irritable, has ciliary tenderness, presents the signs of iridocyclitis, or contains a foreign body which cannot be extracted.

When, however, there is useful vision in the injured eye, or a good chance of obtaining fair sight in this eye, the question of enucleation is often a difficult one to decide, since symptoms of sympathetic irritation may appear and then subside, and yet sympathetic inflammation never develop. In such cases we are often justified in waiting, if the injured eye remain quiet and free from inflammation, providing we can keep the patient under constant observation.

After sympathetic inflammation has made its appearance, enucleation of the injured eye has no effect upon the progress of the disease, and the exciting eye may ultimately possess better vision than its sympathizing fellow. Hence, under such circumstances, the exciting eye should not be removed if it possesses vision; if blind and exhibiting signs of inflammation, it should be enucleated, since its presence may aggravate the condition in the sympathizing eye.

The treatment of the sympathetic ophthalmia itself consists in the use of atropine (unless this seems to aggravate the symptoms), hot compresses, absolute rest, shaded room, and smoked coquilles. Leeches to the temples are sometimes of advantage. Injection of foreign protein is often of value. Large doses of sodium salicylate, even up to 200 gr. per day, are frequently given, with good effect in some cases. Salvarsan has been tried in many cases, and is believed to have done good. Since the disease is of lengthy duration the general health of the patient must be looked after.

Though the prognosis is unfavourable, and most cases end in blindness, the treatment must be carried out rigidly and

patiently. In some cases at least, especially if the inflammation be of the serous type, fair vision may ultimately be obtained. Operative intervention may be indicated for improving vision, but such interference must not be considered until the eye has been quiet for several months.

Panophthalmitis.

An intense purulent inflammation of the entire uveal tract, which fills the eyeball with pus, and ends in complete destruction of this organ. It is due to infection. It resembles purulent choroiditis (which term is often employed as a synonym), but the inflammatory process is more extensive.

Etiology.—It is almost always due to infected wounds of the eyeball, whether accidental or as a result of operation. It may also result from infective ulcers, metastasis in pyæmia and puerperal septicæmia, meningitis, and cerebro-spinal meningitis, especially in children.

Symptoms (already described in connexion with purulent choroiditis, p. 198) are apt to be acute and severe. The disease is usually ushered in by a rise of temperature, general febrile symptoms, headache, and sometimes vomiting. There are severe pain in the eyeball, rapid loss of sight, intense ciliary and conjunctival congestion, marked chemosis, and swelling and redness of the lids (Fig. 159, Plate XII). The iris soon becomes involved, the anterior chamber and vitreous become filled with pus, the cornea is clouded and yellow (Fig. 160, Plate XII.), and tension increased. There is infiltration of Tenon's capsule, followed by exophthalmos and limitation of the movements of the eyeball.

Pus usually breaks through the anterior portion of the sclera, after which the pain and other symptoms subside. In the course of several weeks the process has run its course, leaving a shrunken, sightless eyeball (phthisis bulbi, Fig. 173).

The **Prognosis** is always unfavourable. Sight is invariably lost. The condition does not cause sympathetic ophthalmia.

Treatment.—The indications are to alleviate pain by the use of hot, moist compresses, and to incise the sclera so as to

allow the escape of pus. If the case is seen early, thorough and repeated cauterization of the focus of infection with the




FIG. 173.—PHTHISIS BULBI.

electro-cautery or the introduction of small rods of iodoform into the anterior chamber, or both measures combined, may be of value, but evisceration is usually required. It is not considered advisable to enucleate in the inflammatory stage, on account of the danger of setting up purulent meningitis (p. 98), but, after evisceration, the sclera can be divided backwards towards the optic nerve, the tendons severed

as in enucleation and the anterior portion of the sclera removed, leaving a frill round the nerve head. This removal of the sclera accelerates healing of the wound.

CHAPTER XV

INTRA-OCULAR TUMOURS

INTRA-OCULAR tumours are rare. Their recognition is, however, important, since early enucleation of the eyeball may save life. There are two varieties: (1) Sarcoma of the choroid, and (2) glioma of the retina.

Sarcoma of the Choroid.

This malignant growth usually occurs in adults between the ages of forty and sixty. It is always primary, single, and involves one eye only. It may be formed of spindle or of round cells, and is usually pigmented (melano-sarcoma). It forms a rounded mass which springs from the choroid, most commonly near the posterior pole, and advances toward the centre of the eyeball; later it perforates the lamina vitrea, grows more rapidly and forms a globular mass in the sub-retinal space, pushing the retina forward at its summit and causing a detachment of the retina at its sides (Fig. 174).




FIG. 174.—SARCOMA OF THE CHOROID, WITH DETACHMENT OF THE RETINA.

Symptoms.—There are four stages.

In the *first or quiescent stage* there will be a defect in the field and diminution in sight depending upon the exact seat of the tumour. With the ophthalmoscope a yellowish, brown, or black mass may be seen, over which retinal vessels can be traced, and behind these other vessels belonging to the tumour itself. But very frequently the retina becomes detached, and thus obscures this picture. The anterior ciliary veins

may be found dilated near the seat of the growth. This stage usually lasts from eighteen months to two years.

In the *second or irritative stage* the tumour enlarges in size, and gives rise to pain and other symptoms of inflammatory glaucoma.

In the *third or extra-ocular stage* the tumour bursts through the globe, and then increases very rapidly in size, and ulcerates. In most cases it perforates anteriorly, and a dark mass is seen. If it perforates posteriorly, exophthalmos results. It soon implicates neighbouring structures, including the brain.

The *fourth stage* is distinguished by the occurrence of metastases, most frequently in the liver.

Pathology.—The tumour is composed of round or spindle-shaped cells, more or less pigmented, developed from the stroma cells of the outer and middle layers of the choroid; it is usually vascular, but the normal vessel wall is missing, often consisting only of epithelium, or there may be no wall at all, just vascular channels.

Differential Diagnosis.—Sarcoma of the choroid may be mistaken for detachment of the retina, glaucoma, or possibly glioma of the retina. The last, however, occurs only during the early years of life. Ordinary detachment of the retina usually occurs suddenly in a myopic eye, or after a blow, and tension is diminished. There are no premonitory symptoms such as usually precede glaucoma, nor remissions in symptoms; one eye only is involved, and the characteristic field of glaucoma (nasal limitation) is not present, and usually sight is impaired before inflammatory symptoms appear. Transillumination with one of the instruments invented for the purpose will usually reveal any solid swelling in the anterior hemisphere.

Treatment.—Enucleation as soon as the diagnosis is established, cutting the optic nerve far back. It may be necessary to clear out the entire contents of the orbit. After enucleation or exenteration a few prophylactic exposures to radium or X-rays are indicated. There is always danger of local recurrence and of metastases in internal organs. The affection is

invariably fatal when not removed early, death taking place within five years.

Glioma of the Retina.

A malignant growth, also known as neuro-epithelioma of the retina, consisting of small cells with some soft basement substance and bloodvessels, developing from the granular layers of the retina, and probably congenital. It occurs in children under five, usually in one eye, at times in both, and occasionally in successive children of the same family.

Symptoms.—We distinguish three stages.

In the *first or quiescent stage* there are no inflammatory symptoms. The ophthalmoscope shows small whitish or yellowish masses with metallic lustre, growing into the vitreous. The surface presents newly formed bloodvessels, and may also show hæmorrhages and white patches. The attention of the parents is attracted by the striking yellow reflex, easily seen through the pupil, which is usually dilated. In pre-ophthalmoscopic days this appearance gave rise to the term 'amaurotic cat's eye.'

In the *second or irritative stage* there are pain, increase of tension in about 50 per cent. of cases, and other symptoms of glaucoma. The tumour increases in size, and extends into the vitreous. Very soon the growth can no longer be seen on account of turbidity of the media.

In the *third or extra-ocular stage* there is bulging of the eyeball (both staphyloma and exophthalmos), and then perforation takes place. The growth passes backward along the optic nerve to the brain (in this way it becomes fatal), and forward through the cornea and sclera, increasing in size rapidly, and involving all tissues with which it comes in contact. Metastases are rather rare, but may occur in neighbouring glands and in the cranial bones.

Pathology.—The tumour does not resemble glioma of the brain. It is composed of small, round, densely packed cells with large nuclei arranged around bloodvessels or in the shape of tubes, or aggregated in the form of rosettes, in which case the extensions inward resemble the rods and cones of

the retina; a short distance from the bloodvessels, the cells show a tendency to necrosis. The mass originates either from the outer or the inner surface of the retina; in the former case it separates retina from choroid, causing detachment of the former; if the growth is inward, polypoid masses will invade the vitreous, increasing in size, and eventually filling the whole eyeball. The tumour is always multiple, there being scatterings of foci in various parts of the eyeball. Extension of the growth is usually along the optic nerve, and thence into the cranial cavity.

Differential Diagnosis.—We must distinguish glioma from pseudo-glioma (p. 199), the degenerated eyeball which is the outcome of purulent choroiditis following meningitis or cerebro-spinal meningitis in children. In the latter affection there will be the history of a previous acute febrile disease, with inflammation of the eyeball, and tension is diminished. Ophthalmoscopic examination is the chief guide. When in doubt, such eyes being always sightless, we should enucleate.

Treatment.—*Enucleation as soon as possible*, cutting the optic nerve far back. If microscopic examination of the cut end of the optic nerve shows that the growth has involved this part or if the growth has perforated, the entire orbit must be cleaned out. Afterwards local treatment with radium or X-rays is advisable. Even then there is danger of recurrence. When excision has been refused, or in cases in which the second eye is involved, an attempt should be made to destroy the growth by radium. For this purpose radon seeds have been inserted directly into the tumour through a small scleral incision; in a few instances the growth has been checked by this means. When excision is practised early there is a fair chance of cure. Unless this is done, death occurs within two or three years.

CHAPTER XVI

GLAUCOMA

Anatomy.—The aqueous chamber is bounded in front by the cornea, behind by the lens and its suspensory ligament, and laterally by the ligamentum pectinatum and anterior portion of the ciliary body (Fig. 175). Its depth varies; it is comparatively deep in the young, in myopic eyes, and when the eye is focussed for distant objects. The iris divides the aqueous cavity into an anterior and a posterior chamber. The former lies in front of the iris. The latter is the annular space between the iris and the lens. Since the iris is in contact with the lens only at its papillary margin, this space increases in depth from the pupil to the peripheral border of the iris, and is triangular in cross-section. The posterior communicates with the anterior chamber by means of the pupil.

The portion of the anterior chamber where the sclero-corneal margin, iris, and ligamentum pectinatum meet is known as the *angle or sinus of the anterior chamber* (often called the iris angles). This region is of great importance. Upon its integrity depends the proper circulation of the lymph which nourishes the anterior portion of the eyeball.

The *ligamentum pectinatum* is formed by the breaking up of Descemet's membrane at the margin of the cornea, into bundles which connect the sclera with the root of the iris. These elastic laminae are covered by endothelium continuous with that covering the posterior surface of the cornea. In this way spaces are formed which are continuous with the cavity of the aqueous, are lined with endothelium, and are known as the *spaces of Fontana*. To their outer side, at the sclero-corneal junction, is *Schlemm's canal*, communicating laterally with the anterior ciliary veins.




FIG. 175. — SECTION OF THE EYEBALL AT THE SCLERO-CORNEAL JUNCTION, SHOWING ANGLE OF ANTERIOR CHAMBER.

S, Sclera; C, cornea; I, iris; L, lens; cm, ciliary muscle; Sch, canal of Schlemm.

With the exception of the conjunctiva, no portion of the eyeball contains lymphatic vessels. In place of such vessels and serving the same function, there are lymph channels and lymph spaces. These may be divided into those of the anterior and those of the posterior portion of the eyeball.




FIG. 176. — DIAGRAM-MATIC SECTION OF THE ANTERIOR PORTION OF THE EYEBALL.

1, Upper conjunctival sac; 2, lower conjunctival sac; 3, anterior chamber; 4, posterior chamber.

The *anterior lymph spaces* and cavities consist of the aqueous chamber and the parts immediately around the iris angle. The anterior and posterior chambers represent two large lymph spaces which collect the lymph of the anterior portion of the eye. This lymph is known as the aqueous humour, and consists of a clear, watery fluid, secreted by the epithelium covering the ciliary processes and the posterior surface of the iris. It first passes into the posterior chamber, then through the pupil into the anterior chamber, and leaves the eye through the spaces of the ligamentum pectinatum (Fontana's spaces) and Schlemm's canal, passing into the anterior ciliary veins; a portion passes into the lymph spaces of the iris, and thence to the suprachoroidal lymph space.

The *posterior lymph passages* consist of the hyaloid canal of the vitreous, and of the suprachoroidal space (between choroid and sclera), communicating with Tenon's space along the *venæ vorticosæ*.

Glaucoma is an important and common disease of the eye, which has for its characteristic sign an increase of intra-ocular tension.

Varieties.—It is (1) *primary*, when occurring without antecedent ocular disease; and (2) *secondary*, when it follows as a result of some pre-existing disease of the eye.

Primary Glaucoma occurs under two forms: (1) *Inflammatory* or congestive, and (2) non-inflammatory or non-congestive, usually spoken of as *simple*.

The inflammatory variety is again divided into (1) acute and (2) chronic.

These variations in clinical types of primary glaucoma are explained by the rapidity with which the increase of

intra-ocular pressure shows itself and the height to which it rises. When the increase of tension is rapid, the inflammatory type results; when gradual, the eyeball accommodates itself to a certain extent to the altered conditions, and symptoms of inflammation or congestion are absent or only very slightly marked. The disease is then known as simple glaucoma (non-inflammatory or non-congestive glaucoma). This type is always chronic in its course.

All forms of glaucoma present very characteristic remissions or intermissions in the course of the disease.

Acute Congestive Glaucoma.

Symptoms.—Glaucoma presents a clinical picture which varies with the type of disease, depending upon the suddenness of onset, duration, and the presence or absence of congestive signs; hence the classification into acute and chronic congestive or inflammatory cases. It is not always possible to draw a sharp line between the different clinical forms, and simple glaucoma may at any time change to acute or chronic congestive.

The affection can be divided into three stages: (1) the prodromal stage, (2) the stage of active glaucoma, and (3) the stage of absolute glaucoma. To these we may add a fourth stage, the stage of degeneration.

The Prodromal Stage.—This stage is present in most instances; it may, however, be absent. There will be some diminution in the acuteness of vision—the sight appears to be obscured by fog. A ring of rainbow tints will be seen around lights; the cornea, especially at its centre, will, upon careful inspection, be found slightly clouded. This condition (oedema) is the cause of the preceding symptoms. There will be a feeling of dulness or slight pain in the eye and head. The anterior chamber is rather shallow, the pupil somewhat dilated, often oval, and sluggish in reaction. The tension of the globe is increased. There is often slight circumcorneal injection.

These symptoms last for a number of hours, and then dis-

appear entirely. The eye then returns to a normal condition, except that there may be a diminution in the power of accommodation, so that the patient requires stronger glasses than are natural at his age. Hence a rapid increase of presbyopia should always excite suspicion of glaucoma. The attack may be precipitated by insomnia, worry, dissipation, insufficient food, some condition which causes venous congestion, and sometimes by a hearty meal, indigestion, or the local use of atropine. It is perhaps relieved by sleep. At first the attacks may be separated by intervals of weeks or months, but they soon become more frequent.

This stage may be absent or it may last for weeks or months, sometimes several years. Then the disease suddenly passes into the second stage.

The Stage of Acute Glaucoma.—The sudden onset which characterizes this stage may be due to one of the exciting causes which bring on the prodromal attacks. There are rapid failure of sight, contraction of the visual field, especially on the nasal side (Fig. 182), and severe pain in the eye, radiating along the branches of the fifth nerve and causing violent headache. This pain is sometimes so severe that it occasions nausea, vomiting, general depression, and febrile disturbances, such attacks having been mistaken for bilious attacks.

Objective examination reveals marked increase in tension; the cornea is clouded or steamy (due to oedema), often presents punctate opacities, and is insensitive (from pressure upon nerve filaments). There is pronounced circumcorneal injection of a dark red colour; the episcleral veins are prominent (Fig. 180, Plate XVI.). The pupil is dilated, oval, immobile, and often presents a greenish reflex. The iris is congested, discoloured, and dull. The anterior chamber is shallow, the aqueous sometimes turbid. The lens and the periphery of the iris are pushed forward. The lids are swollen and oedematous. The ocular conjunctiva is markedly congested and chemotic. No details of the fundus can be seen with the ophthalmoscope, on account of the clouding of the media.

In many cases in the course of a few days or weeks a decided improvement takes place. The pain subsides, congestion and œdema of lids and conjunctiva disappear, the cornea clears up, and sight improves. But the eye does not return to a perfectly normal condition. It is left in a condition known as the *glaucomatous state*. Vision is not so acute as it was before the attack, and the visual field is somewhat contracted, especially on the nasal side. The pupil remains dilated, oval, and sluggish, the iris discoloured, the anterior chamber shallow, tension increased, and there is




FIG. 177.

FIG. 178.

FIG. 179.

OPHTHALMOSCOPIC APPEARANCES AND LONGITUDINAL SECTION OF THE OPTIC-NERVE DISC.

Fig. 177, Normal disc; Fig. 178, disc in optic-nerve atrophy
Fig. 179, glaucomatous excavation.

more or less circumcorneal injection. The power of accommodation is diminished.

After a period of quiescence of variable length another attack occurs, similar to the first, and this is succeeded by others. Each attack causes greater reduction in sight.

After a while the increased tension causes excavation of the optic-nerve disc (Fig. 179), recognizable with the ophthalmoscope in the intervals between attacks, when the media are clear. The lamina cribrosa, the portion of the sclera which is perforated by the optic-nerve fibres, is most yielding, and hence bulges backward with the fibres of the nerve as a result of increased intra-ocular pressure. With the ophthal-

moscope a deep depression with very steep or overhanging margins is seen. This is known as the glaucomatous cup or excavation (Fig. 181, Plate XVI.). The bloodvessels bend sharply over the margins of this excavation, and often appear interrupted in this situation, being again seen, more or less faintly, at the bottom of the depression. They are pushed over toward the nasal side. The veins are distended and the arteries contracted. There is pulsation in the veins and in the arteries at the disc. Pulsation in the veins is often seen in health, but arterial pulsation is always pathological, and is an important symptom of glaucoma (it is also seen in certain forms of heart disease). If not spontaneous, it can be produced by slight pressure upon the eyeball. The optic nerve becomes atrophied, and the disc appears pale, or in late stages greenish or bluish. The disc is often surrounded by a whitish-yellow ring (glaucomatous halo or ring), due to atrophy of the choroid in this situation.

The Stage of Absolute Glaucoma.—With each succeeding attack the diminution in vision becomes greater, until finally blindness ensues. The condition is then known as absolute glaucoma. There are now no inflammatory or congestive symptoms, except a dark red zone of circumcorneal injection and dilated episcleral veins. The cornea remains clear or slightly clouded, and often more or less insensitive. The pupil is widely dilated, immobile, and often presents a greenish reflex. The iris is atrophied, narrow, gray, with a border of dark pigment. The anterior chamber is shallow. Tension is markedly increased. The fundus presents a deep excavation of the disc, the glaucomatous ring, and atrophy of the optic nerve. Pain may disappear entirely, but frequently continues, and the patient suffers from severe attacks at intervals.

The Stage of Degeneration.—After absolute glaucoma has lasted a variable length of time, the eyeball is apt to degenerate. The cornea becomes more or less opaque, and frequently covered by deposits or vesicles. The sclera bulges, and bluish-black staphylomata appear between the cornea and the equator. Detachment of the retina often takes

PLATE XVI.




FIG. 180.—ACUTE CONGESTIVE GLAUCOMA.



FIG. 181.—THE FUNDUS IN CHRONIC GLAUCOMA.

To face page 216.

place. The lens is apt to become cataractous. The patient may experience subjective sensations of light. The final result is that the eyeball either softens, shrinks, and atrophies, or else there are ulceration and perforation of the cornea, followed by iridocyclitis, with subsequent atrophy of the eyeball, or panophthalmitis and phthisis bulbi.

Glaucoma Fulminans is the name given to a form, of rare occurrence, in which very violent symptoms of inflammation develop suddenly, and in which blindness may ensue in a few hours, unless proper treatment be instituted.

Chronic Congestive Glaucoma.

This form of glaucoma is much more common than the acute variety just described. Its symptoms resemble those of the acute variety, but are less intense and more gradual in their onset. Very often the prodromal stage passes uninterruptedly into the stage of active glaucoma, and there is no succession of attacks. The ocular conjunctiva is congested and dusky, the episcleral veins being very prominent. There is circumcorneal injection of a dark red colour; the cornea is steamy and more or less insensitive; the anterior chamber is shallow, and the lens and iris are pushed forward; the pupil is dilated, oval, and rigid, surrounded by the discoloured, narrow, and atrophic iris, and presents a greenish reflex. There is pain, but this is not so intense as in the acute form. There are gradual loss of sight and progressive limitation of the field, especially on the nasal side. After having lasted a sufficient length of time, the ophthalmoscope reveals the same changes in the fundus which are found in acute cases.

The chronic form has the same termination as the acute: absolute glaucoma and finally degeneration of the eyeball. In many cases no sharp line of differentiation can be drawn between the acute and the chronic forms of inflammatory glaucoma.

Simple Glaucoma.

In simple glaucoma (chronic non-inflammatory glaucoma) there is an absence of any marked external symptoms; there are no inflammatory attacks and no pain.

The diagnosis is made by noting the increase of tension, by the picture presented when the ophthalmoscope is used, and by charting the visual fields.

This form develops very gradually, and may have lasted some time before the patient becomes aware of the existence of any abnormal condition. The eye may appear perfectly normal externally, or there may be slight circumcorneal injection and moderate dilatation of the episcleral veins. The pupil is slightly or moderately dilated and is sluggish. The tension is elevated, often moderately; sometimes the increase is not constant. After the disease has lasted a certain length of time, the ophthalmoscope shows glaucomatous excavation (Figs. 179 and 181, Plate XVI.), atrophy of the optic nerve, and the circumpapillary ring of choroidal atrophy, the degree of change depending upon the duration of the process.

There may be periods when the patient complains of symptoms like those in the prodromal stage: Foggy vision, coloured halos around artificial lights, and diminished accommodation. There are gradual loss of sight, premature presbyopia, and progressive contraction of the visual field, especially on the nasal side. The earliest defect in the field of vision is usually an enlargement of the 'blind spot,' and this commonly progresses as a wing-like extension either above or below or both (Seidel's sign). As this scotoma extends, due to the progressive atrophy of the nerve fibres which curve outwards from the optic nerve above or below the macula, it becomes sickle-shaped, and is known as Bjerrum's sign. A complete ring scotoma may thus be formed. Isolated scotomata may also occur.

Later the nasal contraction of the field takes place and is often greater above than below the horizontal meridian, or *vice versa*. A sharp right-angled defect, one side of the angle

corresponding to the horizontal meridian, is thus caused; this is known as *Roenne's nasal step*. Central vision is the last portion to be lost; the patient may be able to read and yet have a very contracted visual field.

The course of simple glaucoma is very insidious, and its duration is years. If unchecked, it terminates in blindness. Sometimes this form gradually changes into the chronic inflammatory type, and then goes through the stages of the latter disease.

Etiology.—Glaucoma is a disease of advanced life, occurring generally after forty years of age, infrequently before this period. The congestive or inflammatory form attacks women more often than men; the simple type occurs equally in both sexes. It usually involves both eyes, the second eye generally becoming affected months or years after the first.

The cause of glaucoma is unknown. Jews appear to be more liable to the disease than others. There is not uncommonly a history of heredity. Arterio-sclerosis and cardiac disease, chronic constipation, and the gouty and rheumatic diatheses,




FIG. 182.—THE FIELD OF VISION IN GLAUCOMA.

The upper illustration shows peripheral contraction, especially on the nasal side; also Seidel's sign. The lower illustration shows peripheral contraction, Roenne's nasal step, and Bjerrum's sign.

are predisposing factors. A disposition toward glaucoma exists in hypermetropic eyes (myopic eyes are far less liable to the disease) as well as in small eyeballs, and in those in which the cornea is of small size. The exciting causes may be the following: Emotions, especially of a depressing character, insomnia, worry, injudicious use of atropine, over-use of ametropic eyes, insufficient food, indigestion, dissipation, various fevers, especially influenza, and any condition which produces venous congestion.




FIG. 183.—ANGLE OF THE ANTERIOR CHAMBER IN THE NORMAL EYE.




FIG. 184.—ANGLE OF THE ANTERIOR CHAMBER IN RECENT INFLAMMATION OF GLAUCOMA.

Pathology.—In the early stage there is œdema of the cornea, congestion and swelling of iris and ciliary body; the root of the iris and the anterior part of the ciliary body are drawn forward to lie in apposition with the sclero-cornea. In later stages every part of the eye shows degenerative changes: there is round-cell infiltration of the deeper layers of the cornea and around Schlemm's canal; proliferation of tissue cells forming organized exudate, resulting in peripheral synechiæ which causes firm adhesion between the root of the iris and the floor and outer wall of the anterior chamber; hyalini- zation of the pectinate ligament and degeneration of the cornea. Schlemm's canal is almost obliterated; the iris, ciliary body and choroid are atrophied; there are also generalized sclerosis of bloodvessels, and multiple hæmorrhages. The lens becomes cataractous and the sclera is thickened and undergoes fatty degeneration. Pressure upon the cribriform plate leads to excavation of the nerve head. The nerve fibre layer and the

ganglion cell layer of the retina atrophy almost completely and are replaced by hypertrophy of glial and connective tissue.

Pathogenesis.—All the symptoms of glaucoma can be explained by the increase in intra-ocular pressure. But there is no general agreement as to the exact cause of this increase. The intra-ocular fluids, after being secreted in the vitreous chamber of the eye, pass round the lens, through the pupil into the anterior chamber, to the iris angle, and are thence excreted into the canal of Schlemm. It is probable that the disease is due to some interference with excretion. The obstruction to the escape of the intra-ocular liquids is thought to be situated at the angle of the anterior chamber (iris angle). It is believed that this angle is obliterated by pressure of the peripheral portion of the iris against the sclero-corneal junction (ligamentum pectinatum) by the congested and swollen ciliary processes; later there is an *adhesive inflammation* between these opposed surfaces through proliferation of the endothelium of Descemet's membrane and of the iris. As already explained, this iris angle forms the principal exit for intra-ocular fluids, and when it is blocked retention takes place. An additional causative factor is supposed to be the *narrowing of the circumferential space* (between the margin of the lens and the ciliary body) in eyes predisposed to glaucoma. This area serves for the passage of the lymph which is secreted by the choroid and part of that produced by the ciliary body; it is encroached upon by the increased size of the lens with advancing age and by the comparatively large size of the ciliary body and the smaller size of the eyeball in general in *hypermetropes*. This obstruction in the communication between vitreous and aqueous chambers would cause venous congestion, subsequent swelling of the ciliary body, overdistension of the vitreous, with the result of pushing the periphery of the iris against the sclero-corneal junction, thus blocking up the iris angle. A primary swelling of the vitreous is probably the chief factor in some cases. According to Duke-Elder, this is probably due to faulty metabolism in the vitreous itself rather than directly to physical and chemical changes in the blood. But no explanation of the production of glaucoma satisfactorily

fits all types of the disease; probably they are not all developed in the same manner.

Differential Diagnosis.—The inflammatory form of glaucoma has been mistaken for iritis and conjunctivitis. The use of atropine in such cases has caused great mischief. The dilated pupil, increase in tension, turbidity of cornea, as well as the subjective symptoms, ought to be sufficient to differentiate (see tables, p. 176). The peculiar greenish pupillary reflex has been diagnosed as cataract, and thus valuable time has been lost in awaiting the ripening of this supposed lens change. In acute cases, the violent headache and general constitutional symptoms have misled the medical practitioner and been responsible for the diagnosis of some general febrile disease at a time when active ocular treatment was urgent.

Simple glaucoma is sometimes mistaken for primary optic-nerve atrophy. In the latter case there will be absence of increased tension, the excavation of the disc is shallow and gradual (Fig. 178, also Plates XVI. and XXIV.), there is apt to be greater diminution in central vision, the form-fields present more uniform contraction, and the colour-fields show greater peripheric loss. There are, however, instances in which the differential diagnosis between these two affections is not easy.

Prognosis is bad in every case if proper treatment is not carried out without delay. Vision becomes worse, more or less rapidly, but progressively, until complete blindness results. With correct treatment, the prognosis is more favourable. The disease in most cases can be arrested, so that the patient retains what sight he has.

Treatment of Glaucoma.—This may be (1) operative, (2) non-operative, (3) general. It varies with the type of case with which we are dealing; operation is usually necessary, but sometimes, in a mild chronic case in an old person, the use of miotics, carefully watched, may serve to keep the disease permanently in check, and is then preferable to operation.

In acute congestive glaucoma, if the non-operative measures described below fail to reduce tension and relieve congestion in a few hours, it is necessary to perform an iridectomy (excision of a portion of the iris).

If the eye responds to treatment, one of the operations to form a filtering scar can be performed at leisure.

In chronic congestive glaucoma the procedure is the same, but iridectomy is less frequently necessary, as the eye more often responds to treatment.

In chronic non-congestive glaucoma, and in those cases of congestive glaucoma which have responded to treatment, operations having for their object the production of a filtering scar in the sclera just beyond the limbus are now generally performed. Such operation is advisable in any case in which there is even the slightest progressive loss of visual field, and should not be delayed.

The most popular of these operations is Elliot's (trephining the corneo-scleral margin combined with a small iridectomy). This and other operations having the same object are described on p. 229. Late infections of the eyeball, made possible by the thin barrier between the conjunctival sac and the interior of the eye at the seat of operation, occasionally occur after these filtering operations, but in our experience are very rare and certainly do not counterbalance the advantage of this form of operative treatment.

In any case of unilateral glaucoma the patient should be warned that the second eye usually becomes affected and must be kept under observation, and have the field of vision charted at intervals. If he is going right away from skilled observation and treatment, it may even be necessary to consider the question of operating on the unaffected eye.

In absolute glaucoma enucleation is indicated for relief of pain; neither iridectomy nor trephining should be advised, since there is a risk that the cause of the glaucoma may be an intra-ocular malignant growth (sarcoma).

Non-Operative Treatment consists chiefly in the local use of miotics, eserine salicylate ($\frac{1}{4}$ to 1 per cent.), and pilocarpine nitrate ($\frac{1}{2}$ to 1 per cent.). Of these eserine has the stronger action, but produces more irritation, especially when used for a long time. The solutions are instilled two or three times daily, or even hourly in acute cases; they act by drawing the iris away from the angle of the anterior chamber; hence, they

are of no value after the iris has become atrophic and is incapable of contracting, or when the iris angle is permanently blocked, conditions observed in old cases of glaucoma. They are usually merely palliative measures of temporary advantage, but are used in the prodromal stage to cut short the attack, or at other times if, for any reason, surgical intervention is inadvisable or the patient refuses operation. In a mild chronic case in an old person their constant use may serve to check the disease permanently, and in such cases they are preferred to operation.

During an acute glaucomatous attack, in addition to frequent instillations of eserine or pilocarpine, hot fomentations are applied continuously, or the electric eye warmer, if available; two or more leeches are applied to the temple; the patient is put to bed and pain relieved by morphine; thorough purgation is indicated. Adrenalin often acts favourably. Lævo-glaucon has been found useful in cases of secondary glaucoma with iritis. An intravenous injection of 50-100 c.c. of 30 per cent. warm sterile saline reduces tension temporarily and is useful in preparing an eye with high tension for operation. Diathermy has recently been successfully tried for the same purpose and for relief of pain.

Massage of the eyeball, applied gently to the closed lids, may be used with advantage in simple and chronic forms, and also after operation.

General Treatment comprises rest, sweating, proper dietary, purgatives, correction of ametropia, avoidance of excess in eating, drinking, and late hours, the induction of sleep, and the relief of any other conditions which have been mentioned as predisposing to glaucoma.

Operations for Glaucoma.

Iridectomy.—The *instruments* required include an eye speculum (Fig. 185), a fixation forceps (Fig. 186), a Graefe cataract knife (Fig. 187), a curved iris forceps (Fig. 188), a curved iris scissors (Fig. 189) or de Wecker scissors (Fig. 210), and an iris repositor (Fig. 208).

Cocaine may be employed in cases of simple glaucoma, and

in some cases of congestive glaucoma. A drop of adrenalin (1:1,000) is instilled prior to the cocaine, to aid absorption of the latter and to reduce congestion; many congested cases can be operated on without pain if 2 c.c. of 3 per cent. solution of novocain in 1:10,000 adrenalin are injected deep into the




FIG. 185.—LANG'S EYE SPECULUM.



FIG. 186.—FIXATION FORCEPS.



FIG. 187.—NARROW GRAEFE KNIFE.



FIG. 188.—CURVED IRIS FORCEPS.

orbit to paralyze the ciliary nerves. General anaesthesia must be employed for nervous or unruly patients.

Eserine or pilocarpine is always instilled in the unoperated eye both before and after operation, since we know this eye is predisposed to glaucoma, and the anxiety and excitement of operation may otherwise induce an attack.

Operation.—Iridectomy for glaucoma is usually done upwards, so that the defect is covered by the upper lid, thus limiting troublesome optical effects of the coloboma. The




FIG. 189.—CURVED IRIS
SCISSORS.

operator, standing behind the patient's head, introduces the speculum. A narrow Graefe knife is held between the thumb and two fingers of the hand corresponding to the eye to be operated upon, the fixation forceps being held in the other hand. The patient, if not under general anæsthesia, is told to look downwards a little. A firm hold is taken with the forceps of the conjunctiva just below the corneal margin. A gentle tension is kept on the forceps so as not to press on the globe and so increase the internal pressure. The knife is made to enter 1 mm. behind the limbus, at about the junction of the lower five-sixths with the upper sixth, passes across the anterior chamber (great care being exercised not to wound the iris or lens capsule), and emerges at a corresponding point 1 mm. behind the limbus on the opposite side, the incision being completed by to-and-fro movements. Towards the end of this section it is well to turn the edge of the knife a little backwards so as to get a little tongue of

conjunctiva at the apex of the flap, as this helps the healing of the wound.

An assistant now takes the fixation forceps if a general

anæsthetic is being used. Otherwise the patient is asked to continue to look a little downwards and the fixation forceps is laid aside. The operator passes the closed iris forceps through the scleral incision to the pupillary margin, opens the instrument, seizes the pupillary border of the iris between its branches, and makes a radial cut in the iris at one end of the wound (Fig. 190). The iris is then torn away from its base




FIG. 190.—CUTTING THE IRIS IN IRIDECTOMY FOR GLAUCOMA.

as far as the other end of the wound, where it is cut off with the scissors. The piece of iris removed should comprise the entire width, including the ciliary attachment. It was formerly taught that at least one-fifth of the circle of the iris should be removed, but a moderate iridectomy appears to be just as efficient as a wide one.

The resulting coloboma must be cleanly cut, and the pupillary margin of the iris must return to its natural position, producing a keyhole-shaped pupil (Fig. 191). No iris tissue must be left on the wound, since this causes subsequent irritation and complications. Proper replacement of the iris is accomplished by stroking the wound with a spatula, or, if this is unsuccessful, by passing the spatula into the incision and freeing the angles.




FIG. 191.—IRIDECTOMY.

Hæmorrhage into the anterior chamber is common. The blood is usually absorbed in a few days. It is not wise to

make too great efforts to dislodge the blood, since undue pressure may cause the lens to become cataractous.

Both eyes are bandaged, atropine instilled in the operated eye and eserine in the other, and the patient is kept quiet in bed. After two days, the unoperated eye may be left uncovered. Recovery is uneventful in most cases. In some cases the anterior chamber is not re-formed for several days. Cystoid cicatrix sometimes results—a condition which, though it has dangers of its own, facilitates elimination of fluid, and is deliberately aimed at by some surgeons with that object.

Other Indications for Iridectomy.—Besides (1) glaucoma, the operation may be indicated in (2) some cases of chronic and recurrent iritis and iridocyclitis; (3) complete circular synechia; (4) partial corneal staphyloma; (5) tumours and foreign bodies



FIG. 192.—KERATOME.



FIG. 193.—BLUNT IRIS HOOK.

in the iris; (6) recent prolapse of the iris; (7) as a part of the operation of extraction of cataract—here the coloboma should be smaller than in glaucoma (Fig. 191); (8) as a means of improving sight (artificial pupil, optical iridectomy) in central opacities of the cornea and lens, occlusion of the pupil, and keratoconus. Except in glaucoma iridectomy, in which the root of the iris is better exposed by cutting from within outwards, a lance-knife (keratome) is used to make the incision at the limbus, the root of the iris is not torn away, and the fold of iris which is withdrawn is cut away radially with the iris scissors.

In performing *optical iridectomy* a small incision (4 to 5 mm.) is made in the cornea, 1 mm. from the limbus, either with a keratome (Fig. 192) or with a 'broad needle' (Fig. 216), the iris drawn out with curved iris forceps (Fig. 188) or the blunt

hook (Fig. 193), and its pupillary portion excised, making as small a coloboma as is practicable. The best position for the artificial pupil is downward and inward; but when there is a corneal opacity, the site must correspond to the most transparent portion of the cornea. The effects of optical iridectomy are often disappointing; hence, before operating, it is well to dilate the pupil and, by applying a stenopeic slit held in different positions, to ascertain whether there is an improvement in sight under these circumstances.



FIG. 194.—EXCISION OF IRIS IN NON-GLAUCOMA IRIDECTOMY.

Formation of a Leaking Cicatrix.—

When treatment of glaucoma has been delayed until the iridic angle has been permanently obliterated by inflammatory adhesion of the base of the iris, simple iridectomy seldom produces lasting benefit. But if prolapse of one of the pillars of the coloboma should prevent sound healing of the incision in the sclerotic, a leaking scar results through which the excess of intra-ocular fluid filters out under the conjunctiva. Operations have been devised to produce a similar result without inclusion of iris in the scar.

Corneo-scleral trephining is the operation which, in *chronic* glaucoma, is most commonly employed, and has given the best results. Lagrange's operation and Herbert's operation are alternatives. Iris inclusion operations, in which a small knuckle of iris is included in the scleral wound, have been more frequently employed recently, especially in America.

Trephining for glaucoma was tried by Argyll Robertson fifty years ago, and by others since, but the operation as now performed was perfected by R. H. Elliot.

Instruments required are speculum, fixation forceps, iris forceps, squint scissors, iris scissors, trephine 1.5 mm. to 2 mm. diameter (Fig. 197), Took's knife (Fig. 196), iris repositor, needle holder, small curved needle, and silk.

Everything having been prepared as described in Chapter XXXI., and the eye having been cocainized and the conjunctiva blanched with adrenalin, the speculum is inserted. The patient looks downwards. The conjunctiva is grasped with fixation forceps about one-third of an inch above the corneal margin. A large flap is cut, as shown in the diagram, and dissected to the corneal margin in the centre. The ends



FIG. 195.—TOOK'S KNIFE.



FIG. 196.—ELLIOT'S TREPHINE.

of the incision must not approach the corneal margin, or the percolation of aqueous under the conjunctiva may be interfered with, and there is more tendency to the formation of a 'bleb' over the trephine hole, from which the aqueous does not so readily escape.

The flap is turned over the cornea, and held with non-toothed forceps. With the scissors the tissue is dissected




FIG. 197.



FIG. 198. — ELLIOT'S OPERATION FOR GLAUCOMA.

from the sclerotic in the neighbourhood of the cornea, and the incision is carried horizontally into the cornea with the Took's knife, splitting it for about 2 mm. The superficial laminae are turned back with the flap. The trephine is applied with its centre over the apparent corneo-scleral junction, and the disc cut with a few semi-rotary movements with light pressure. One feels the trephine go through the coat of the eye, the patient usually experiences a sharp but not severe

pain, aqueous escapes, and the pupil is displaced towards the wound. Usually the disc remains attached by a hinge on one side, and a knuckle of iris presents in the opening. Disc and iris knuckle are grasped together with iris forceps, and cut off with iris scissors. The iridec-tomy is a small peripheral hole. The flap is replaced over the wound, and its apex secured with one fine stitch. The upper lid is lifted by the lashes over the wound as the speculum is removed. Throughout the operation the greatest care must be taken to avoid button-holing the flap.




FIG. 199.—BLEB AFTER ELLIOT'S TREPHINE OPERATION.

After the operation one drop of sterile liquor atropinæ should be instilled, and this should be repeated every day for from seven to ten days, until the anterior chamber has re-formed and all danger of iritis, a frequent complication, has passed. Not infrequently a detachment of the choroid occurs after trephining, in which case the anterior chamber is slow in re-forming, but with further rest in bed and longer use of atropine this complication subsides in two weeks or so, and results in no harm.

Lagrange's Operation (Sclerectomy combined with Iridec-tomy).—An incision is made in the sclera with a narrow Graefe knife, 1 mm. from the limbus, opposite the iris angle, and counter puncture made at a corresponding point 7 mm. removed. At the termination of the section the edge of the knife is directed backwards so as to bevel the sclera, then continuing so as to make a 5 mm. conjunctival flap. The apex of the triangular piece of sclera under the conjunctival flap is then snipped off with curved scissors.

Herbert's Operation and Flap-Sclerotomy.—A keratome is passed into the anterior chamber, either subconjunctivally (Herbert), or after reflecting a conjunctival flap. The in-

cision is 2 mm. behind the limbus. A trowel knife is passed inside the sclera at one end of the incision, and a cut made outwards up to the limbus. A similar cut is made at the other end of the keratome incision. Thus a small rectangular flap is cut in the sclera, the hinge being towards the cornea.

In **Iridotaxis** a small knuckle of the pupillary margin of the iris is drawn up into a small keratome incision just behind the limbus, after turning down a conjunctival flap. This operation is more frequently done in America than in Britain.

In **Cyclodialysis** the ciliary body is in part separated from the overlying sclera, and part of the iris periphery detached by breaking through the pectinate ligament. This is done




FIG. 200.—LAGRANGE'S OPERATION.

by a repositor, or special instrument passed through a small scleral incision 7 mm. behind the limbus. This procedure may be employed when other operations have failed, but is not usually regarded as a primary substitute for other operations.

Sclerotomy (incision through the sclera) is sometimes performed for the temporary relief of glaucoma. The incision in the sclera is made either in front of the iris (anterior sclerotomy) or behind the ciliary body (posterior sclerotomy).

In **Anterior Sclerotomy** an incision is made with a Graefe knife 1 mm. behind the limbus, similar to that made in iridectomy, but the middle third is left uncut, and forms a bridge connecting sclera and cornea.

In *Posterior Sclerotomy* an incision is made through the sclera into the vitreous with a Graefe knife or with a small trephine. The site usually selected is between the external and inferior recti muscles. The cut must not approach the cornea nearer than 7 mm., so as to avoid endangering the ciliary body. The opening becomes blocked with vitreous, so that the operation seldom produces even temporary relief of tension.

Results of Operation in Glaucoma.—How the filtering scar operations relieve tension in glaucoma is obvious. The manner in which iridectomy relieves glaucoma is not definitely known. The earlier the operation is performed, the more sight is preserved. Hence it is advisable to do the operation as soon as possible. The best time is during the prodromal stage, in the interval between attacks. In inflammatory cases, during the stage of acute glaucoma, the operation is very difficult, on account of the severe congestion and the shallowness of the anterior chamber. Under such circumstances, it is advisable to try hot bathing and eserine every half-hour for a few hours in order to reduce tension and increase the depth of the anterior chamber, and then to operate. But if these measures do not act promptly, the operation must be performed without further delay.

The most favourable results of iridectomy are seen in cases of acute inflammatory glaucoma. In such instances pain and inflammatory symptoms subside rapidly, and sight returns up to the degree possessed before the onset of the attack. Furthermore, the results are generally lasting. If the acute attack has been preceded by a chronic glaucoma, the effects of iridectomy may be only temporary. Corneo-scleral trephining should then be done, but if there is much congestion Lagrange's operation or a Herbert's sclerotomy is easier to perform.

In chronic inflammatory glaucoma the results of iridectomy may be favourable, but not so brilliant as in acute cases. The operation relieves the pain and inflammatory symptoms, and the media again become clear; but since the disease has already caused permanent changes in the disc and optic

nerve, the restoration of sight is limited. Hence, in such cases, we now aim at performing one of the operations to produce a filtering scar. Although operations very often check the progress of the disease, even in favourable cases existing impairment of function remains. After iridectomy the astigmatism produced by the incision may cause some reduction in vision. With all operations, but less with trephining than with iridectomy, tension sometimes increases after a variable period and a second operation is required. In a certain number of cases there is progressive diminution in sight, notwithstanding all operative intervention.

In simple glaucoma miotics, in some cases, hold the disease in check, but most cases require operation as soon as the visual field shows any progressive sign of contraction. The younger the patient the more certain it is that operation will be necessary. The results of operation are less marked and less permanent than in congestive cases, but in many cases there is arrest in the progress of the disease; in less fortunate patients results are temporary, and the procedure has to be repeated; in some, the disease progresses till blindness ensues.

Secondary Glaucoma is the name given to those cases in which increased tension and other symptoms of glaucoma are developed as a result of some other ocular disease or injury. The cause is obstruction of the filtration angle produced by viscid products of inflammation, by the crowding of a swollen and congested iris and ciliary body against the pectinate ligament, increase in pressure within the eye from tumour, exudation, etc., or deposit of plastic organized into fibrin resulting in adhesion between the periphery of the iris and the sclero-corneal space. The clinical picture varies with the disease which it complicates. The consequences, if the condition be not relieved, are the same as in primary glaucoma.

The ocular affections which are most frequently followed by secondary glaucoma are: Ulcers or wounds of the cornea with prolapse of iris, corneal cicatrices and staphylomata with incarceration of the iris, iridocyclitis, uveitis, choroiditis, and myopia of high degree, total posterior (ring) synechia,

dislocation of the lens, traumatic cataract (swelling of the lens), the operations of extraction, needling of the lens and dissection of secondary cataract, intra-ocular tumours, and foreign bodies in the eye. In old persons with arterio-sclerosis, a form of secondary glaucoma with retinal hæmorrhages is seen, and is known as hæmorrhagic glaucoma.

The treatment depends upon the primary disease. We endeavour, if possible, to remove the cause. Hæmorrhagic glaucoma does not respond favourably to treatment. Iridectomy is liable to be followed by an aggravation of symptoms. The other agents used in glaucoma may be tried, but are usually of no benefit.

Congenital Glaucoma (hydrophthalmos, buphthalmos) is a disease of early childhood, either congenital or developing in infancy, and usually involving both eyes. There is an increase of intra-ocular tension which, on account of the yielding character of the sclera at this period of life, causes marked enlargement of the eyeball. The cornea is enlarged and bulging, and either remains clear or becomes clouded; the anterior chamber is very deep; the pupil is dilated, and the iris atrophied and tremulous; the sclera is thinned and bluish, owing to the uveal pigment showing through; the disc is deeply excavated. The disease progresses slowly. Though in some cases it comes to a spontaneous stop with the preservation of moderately good vision, it generally leads to blindness. The prognosis is therefore unfavourable. Since some cases have benefited from corneo-scleral trephining together with the use of miotics, these measures should be tried.

CHAPTER XVII

DISEASES OF THE VITREOUS

Anatomy.—The vitreous is a transparent, colourless mass of soft, gelatinous consistence, which fills the posterior cavity of the eyeball behind the lens. Its outer surface presents a thin, structureless covering, the hyaloid membrane. The vitreous is traversed from the optic disc to the posterior capsule of the lens by a canal—the hyaloid canal—serving as a lymph channel in the developed eye, and containing the hyaloid artery during foetal life. In structure the vitreous consists of a transparent network, in the meshes of which are clear liquid and round and branching cells, probably emigrated white blood corpuscles. The vitreous has no bloodvessels, but receives its nourishment from the surrounding tissues, the choroid, ciliary body, and retina.

Persistent Hyaloid Artery.—The hyaloid artery usually disappears entirely during the later months of gestation. Occasionally a greater or lesser remnant persists during life. This can be seen with the ophthalmoscope, as a grayish cord or thread, which arises from the optic disc and stretches into the vitreous, with a free extremity or occasionally attached to the posterior pole of the lens. Rarely the hyaloid canal is abnormally dense, and is visible as a grayish, tubular cord extending from disc to lens.

Fluidity of the Vitreous (*Synchysis*) is a liquid alteration in consistency which, when limited in degree, may be merely a senile change; but when pronounced, it is due to degeneration of this structure dependent upon disease of neighbouring parts: choroid, ciliary body and retina, and found often in myopia of high degree. When opacities are present, these are observed to move freely in such fluid vitreous; there is often diminished tension of the eyeball, tremulousness of the iris, weakness of the suspensory ligament, and sometimes a pre-

disposition to detachment of the retina. These complications increase the risk of loss of vitreous in intra-ocular operations.

Occasionally small, glistening opacities are found in degenerated eyeballs and in some which are normal in other respects, especially in old persons. They fall in a silvery shower when the eyeball is moved. They are usually crystals of cholesterin in a fluid vitreous, and are known as *synchysis scintillans*.

Asteroid hyalitis is the name given to an anomaly occasionally found in vitreous of normal consistence; the vitreous is studded with dull white globules. The condition is seen most frequently in elderly persons; it has no significance and little if any effect on vision.

Muscae Voltantes is the term employed for the appearance of spots before the eyes, without appreciable structural change in the vitreous or other media. They are caused by the shadows cast upon the retina by the cells normally found in the vitreous, and are present in all eyes under certain circumstances, such as exposure to a uniform bright surface, or in looking through a microscope. They are found more frequently in errors of refraction (especially myopia), and temporarily during digestive derangements. They occur as grayish shadows, which move with changes in the position of the eyes, having the shape of dots or globules, frequently collected into strings. They may have any shape. They are annoying, and sometimes alarm the patient, but are of no importance, and do not affect the acuteness of vision. The treatment consists in correcting any error of refraction, or in relieving the disturbance of digestion. They often persist until the patient ceases to look for them, and thus forgets their existence.

Opacities of the Vitreous.—These are quite common. They may occur as a consequence of changes in the vitreous itself, but usually they are the result of disease or of hæmorrhages from the neighbouring structures—ciliary body, choroid, and retina. They vary in number, shape, and size:

1. A diffuse cloud or a dust-like haziness often accompanies cyclitis, choroiditis, irido-choroiditis, and retinitis. When

dust-like it is suggestive of syphilitic choroido-retinitis and iridocyclitis.

2. The opacities may occur in the form of dots, flakes, threads, or membranous masses, the result of exudations or hæmorrhages.

3. Sometimes extensive membranes are met with, which are attached to the retina and provided with bloodvessels. These are supposed to result from chronic retinal disease, called *retinitis proliferans*.

Symptoms.—There is more or less disturbance of vision, depending upon the situation, size, and density of the opacities. The opacities are most frequently movable, indicating a fluid vitreous (synchysis), the result of disease of surrounding parts. On this account, the visual disturbance may vary according to the part of the vitreous occupied by the opacity, and the patient may be able to move the eyeball in a certain way so as to throw the opacity out of the line of sight. Fluid vitreous gives rise to diminished tension, often a tremulous condition of the iris, and may predispose to detachment of the retina; but it is often discovered in an eye which in all other respects appears normal.

Diagnosis is made with the ophthalmoscope at a distance. The vitreous opacities appear as dark spots upon a red ground, when the eye is moved in various directions. When faint, the opacities are best seen with diminished illumination and with a plane mirror. They may also be examined by direct ophthalmoscopy, by interposing stronger and stronger convex lenses in the sight hole of the ophthalmoscope, thus focussing more and more of the anterior portions of the vitreous cavity.

Prognosis varies with the size, density, and nature of the opacity. Syphilitic opacities and small hæmorrhages frequently clear up when treated early. Others become smaller and less dense after a time. A great many are permanent.

Treatment.—Anti-syphilitic treatment is indicated in specific cases. In others small doses of potassium iodide and mercury may be of service. Diaphoretics and cathartics are sometimes employed. Sub-conjunctival injections of physiological salt solution (0·6 per cent.) may be useful.

Hæmorrhages into the Vitreous.

These usually come from the choroidal vessels, and produce opacities of small or large size, causing the symptoms of opacities in the vitreous. When small, they have a red colour as seen with the ophthalmoscope; when large, no red reflex can be obtained, and the pupil appears black. The smaller hæmorrhages are often absorbed; the larger ones frequently leave dense membranous masses. They occur after injuries, after operations upon the globe, in choroiditis, myopia of high degree, and retinitis. They are not uncommon in old persons with atheromatous arteries. The exciting cause is often a strain of some sort, such as a cough. Sometimes they are found in the young without discoverable cause, and in such cases they may recur repeatedly, and may lead to serious results, since the blood is imperfectly absorbed; connective-tissue bands and masses form and may cause detachment of the retina. Tuberculosis is thought to be a factor in such cases (Eales' disease).

Treatment consists in absolute rest, bandage to the eyes, treatment of any accompanying ocular affection or of the general condition. Tuberculin is advised for the recurrent hæmorrhages of young adults. Calcium chloride is useful in preventing recurrences. Later, to favour absorption, iodides and iodine preparations, mercury, or sub-conjunctival injections of normal saline may be tried.

Foreign Bodies in the Vitreous.

The entrance and lodgment of a foreign body (wood, glass, or metal) within the globe usually causes severe inflammation and destruction of the eyeball as a result of iridocyclitis or panophthalmitis, unless the substance be promptly extracted. The gravity of the accident depends upon the nature of the foreign body and the presence or absence of infection. Particles of iron which penetrate the eye during smiths' work, riveting, or stone breaking are usually so heated as to be sterile. Occasionally these substances remain quies-

cent and become encysted; but even in such cases there is danger of subsequent inflammation. The presence of a particle of iron for any length of time is apt to cause a rusty-brown or greenish discoloration of the iris and lens, and occasionally in the cornea, known as siderosis bulbi; in the lens there is a characteristic deposit in the form of a circle just beneath the anterior capsule in addition; such eyes are apt to suffer from degenerative changes in the retina. A fragment of copper remaining within the globe almost invariably leads to destruction of the eye. Lead pellets, gold, silver, porcelain or glass, unless septic, may become encapsuled, and for a long time may cause little or no trouble, but eventually an eye containing a foreign body anywhere posterior to the lens usually becomes inflamed and disorganized. Stone in the vitreous leads to destruction of the eye, owing to the impossibility of removing it; moreover, stone is usually septic.

A foreign body in the iris is seldom left, being accessible, but if not septic, it may become encapsuled, and cause no trouble.

Diagnosis.—If the patient comes under observation soon after the injury, before the media have become hazy, we may possibly be able to see the particle with the ophthalmoscope, or a careful examination of the field of vision, disclosing a scotoma, may locate the foreign body. The site of the wound of entrance and the probable direction which the foreign body took must be taken into account. In many instances an X-ray photograph will reveal its presence. By means of a special localizing apparatus, an expert in this branch of radiography may then be able to define its exact situation. But owing to the bones, and the minute size of the foreign body, and the fact that it may not be more opaque than the surrounding structures, a negative result is never conclusive. If it be of iron or steel, the giant magnet (Fig. 201) will frequently indicate the presence of the foreign body by the production of pain when the point is brought near the eyeball, or the bulging of the iris or forward movement of the lens when the particle is within these structures.

Treatment.—If the substance is a piece of iron or steel, an attempt to extract it with a magnet should be made at once.

We should also try to remove other foreign bodies (glass, wood, copper, lead) as soon as possible after they have been located, by means of delicate forceps introduced through the original wound, or through an opening into the vitreous cavity made at the point at which the foreign body has been located; but if this does not seem feasible, they should be left alone, especially if there be no symptoms of infection or irritation, and the patient can be kept under constant observation. In such cases, however, the question of enucleation must be considered.

Magnet Extraction.—A piece of iron or steel embedded in the iris is best removed by making an incision with a broad needle or keratome 2 or 3 mm. from the limbus, and introducing the pole of a small portable magnet. If this fails a small iridectomy embracing the foreign body should be performed. If the piece of steel is free in the anterior chamber a similar incision should be made if possible without losing aqueous. The foreign body is drawn forward to the back of the cornea and drawn along towards the incision. As the pole of the magnet passes over the incision the posterior lip of the wound is depressed, and the foreign body may usually be drawn out through the wound.

If a magnetizable foreign body is in the lens, an attempt should be made to draw it forward into the anterior chamber by means of the giant magnet. If this proves successful, the patient is then put on the table and the foreign body removed as already described.

If the magnet extraction fails, or if the foreign body in the lens is non-magnetizable, in a young subject it is better to perform a curette evacuation of the lens at once. There is no need to needle the lens and wait. Such a course would increase the risk of sepsis, and the swollen lens pushing the inflamed iris against the cornea would certainly lead to extensive




FIG. 201.—HAAB'S
GIANT MAGNET.

anterior synechiæ. In an older person in whom there would be a definite firm nucleus the lens may usually be extracted as in a case of senile cataract. Unless the character of the wound is such as to introduce complications, a non-cataractous lens may be extracted more easily and completely than a lens whose cortex is only partially cataractous.

It is very seldom possible to remove a non-magnetizable foreign body from the vitreous. If it is of iron or steel, an attempt should be made to remove it by means of the giant magnet. Though it is a great advantage to know the exact position of a foreign body, the magnet should always be used at the earliest possible moment without waiting for X-ray




FIG. 202.—PORTABLE MAGNET.

localization, because a few hours later it may have become so firmly fixed by inflammatory exudate as to make removal impossible.

Haab's magnet is used as follows: The pupil is widely dilated, and cocaine instilled; the patient is seated before the magnet with his face supported and steadied by the surgeon's hands. The pole of the magnet is brought into contact with the cornea. The current is gradually turned on. The patient usually feels moderately severe pain. In a successful case the foreign body comes forward to the back of the lens as a rule, occasionally it passes through the lens, especially if this has been wounded. Perhaps it rests behind the iris, and is seen dragging the iris forward. By a little manipulation it

may then be worked into the anterior chamber, whence it may be extracted as already described. If the magnet fails to draw the foreign body forward, the direction may be altered a little, and the current switched on and off again several times. It sometimes happens that after several minutes of manipulation the foreign body suddenly frees itself.

Or Mellinger's ring magnet may be used if it is available. This has the advantage that the patient lies on the table, and the solenoid is lowered over his head. Soft iron rods in the axis of the solenoid become powerful magnets when the




FIG. 203.—MELLINGER'S RING MAGNET.

current is turned on. If efforts at extraction have failed, the foreign body should, if possible, be accurately localized by X-rays, and then an incision should be made and an attempt made to remove it with a small magnet.

Even after successful removal the prognosis is always serious; a small number of cases recover permanently useful vision, but frequently a detachment of the retina ultimately follows; in many cases infection and severe inflammation of the eyeball occur. If the attempt at extraction fails, enucleation is usually required to prevent possibility of sympathetic inflammation of the other eye.

CHAPTER XVIII

DISEASES OF THE LENS

Anatomy and Physiology.—The *crystalline lens* is a transparent, colourless body, bi-convex in shape, suspended in the anterior portion of the eyeball, between the aqueous and the vitreous chambers. It presents an anterior and a posterior surface (the latter being the more curved), an anterior pole, a posterior pole, and a rounded circumference (the equator). It is enclosed in a transparent capsule, and held in position by its suspensory ligament. The adult lens consists of a peripheral portion, the cortex, and a central part, the nucleus. The cortex is semi-solid, softer than the nucleus, and colourless; the nucleus is harder, and has a yellowish tint. There is, however, no sharp limitation, the transition being gradual. The nucleus increases in size with advancing years, and the cortex diminishes in proportion; in old age the entire lens is of the consistence of the nucleus, and is hard and unyielding. This change is known as sclerosis.

In structure the lens consists of concentric laminae formed of long, hexagonal fibres, the edges of which are connected by a cement substance, leaving fine lymph channels. The fibres either start or end along Y-shaped or stellate figures, the lines of which radiate from the anterior and posterior pole to the equator, each fibre encircling the latter. The septa corresponding to the branches of the stellate figure divide the lens into sectors. These stellate and Y-shaped figures can often be recognized in the adult lens by oblique illumination.

The *capsule of the lens* is a thin, homogeneous, elastic membrane which covers the lens, being known as the anterior capsule in front and as the posterior capsule behind. The anterior capsule is the thicker, and its posterior surface is lined by a layer of cuboidal epithelium from which the lens fibres are formed.

The *suspensory ligament* of the lens is a delicate membrane, extending from the ciliary body to the lens capsule. It covers the inner surface of the ciliary body from the ora serrata to the apices of the ciliary processes, and then passes to the lens, dividing into three layers attached respectively to the anterior capsule, the equator, and the posterior capsule. Between these layers and the equator of the lens is a space, triangular on section, known as the canal of Petit. It is in com-

munication with the posterior chamber by means of slit-like apertures between the fibres of the anterior portion of the suspensory ligament.

The lens is devoid of bloodvessels except in foetal life, its nourishment being derived from the ciliary body.

The *function of the lens* is to focus rays so that they form a perfect image on the retina. To accomplish this, the refractive power of the lens must change with the distance of the object, according to whether the rays are parallel or divergent. This alteration in the refractive power of the lens is known as accommodation, and is produced by a change of shape mainly affecting its anterior curvature.

The lens presents variations in physical characteristics at different periods of life. In the foetus, it is nearly spherical, slightly reddish, and softer than at a later period. In the adult, its anterior surface is less convex than the posterior, and its substance is firmer. Sclerosis, which consists of a process of toughening, due chiefly to loss of water, begins in the centre of the lens in childhood, and advances slowly until adult life, after which its progress is more rapid, increasing the size of the nucleus at the expense of the cortex. In old age, the lens increases in size, is flattened, and assumes a yellow tinge, becoming tougher and less transparent. This process of sclerosis accounts for the gray reflex seen in the pupil of the aged, which may be mistaken for cataract (senile reflex). It also explains the inability on the part of the lens of advanced years to change its shape for the purposes of accommodation (presbyopia).

CATARACT.

A cataract is any opacity of the lens or of its capsule.

Varieties.—Cataracts may be divided into:

1. *Primary*, when independent of any other ocular disorder.
2. *Secondary, or Complicated*, when accompanying or following some other disease of the eye, such as glaucoma, uveitis, etc.

Cataracts are divided, according to the part of the lens involved, into:

1. *Lenticular*, when situated in the substance of the lens.
2. *Capsular*, when affecting the capsule.
3. *Capsulo-lenticular*, when involving both lens and capsule.

They are also known as:

1. *Stationary*, when they remain incomplete.
2. *Progressive*, when they spread and tend to affect the whole lens.

Stationary Cataracts may be divided into:

1. *Anterior polar.*
2. *Posterior polar.*
3. *Lamellar.*
4. *Various uncommon forms.*

Progressive Cataracts may be divided into:

1. *Senile* { Cortical, when the opacities lie immediately under the capsule.
Nuclear, when the opacity is in the part of the cortex immediately surrounding the nucleus.
2. *Congenital and Juvenile.*
3. *Traumatic.*

In patients under about thirty-five all cataracts are of soft consistence throughout, and usually of a grayish-white colour. After this period the nucleus becomes hard and of a yellowish tint.

Etiology.—According to etiology, cataract may be classified as:

1. *Congenital*, due to faulty development or intra-uterine inflammation of the eye. To this class belong the anterior and posterior polar, lamellar, and occasionally complete cataract.

2. *Senile*.—This is the most common form. It usually appears after the age of fifty. It is a degenerative change in most cases. The real cause is unknown. Heredity has some influence.

3. Due to *General Diseases*: diabetes, and less frequently nephritis, gout, and general arterial disease.

4. Due to *Ocular Diseases*, causing complicated or secondary cataract. The most common examples are severe forms of ulcerative keratitis, iridocyclitis, choroiditis, myopia of high degree, glaucoma, and detachment of the retina.

5. *Traumatic*, by the production of an opening in the capsule, thus allowing the lens to absorb aqueous humour; occasionally by mere concussion.

6. Caused by long-continued exposure to intense light or heat (as in the case of glass-blowers, iron-founders, etc.).

Symptoms.—There is (1) diminished acuteness of the vision,

depending upon the situation and the kind of cataract. It is greatest when the opacity is central and diffuse, and least when the cataract is peripheral. When central, the patient sees best in dim light—with dilatation of the pupil. The interference with vision increases with the progress of the cataract, until finally there is mere perception of light. (2) The patient complains of seeing spots which occupy a fixed position in the field. (3) Occasionally there is annoying diplopia or polyopia, due to irregular refraction of the lens. (4) Myopia often develops during the early stages, due to increased density and refractive power of the lens. For this reason the patient may be able to discard his reading-glasses for the time, but he may need concave glasses for distant vision.

Physical Signs.—There are no inflammatory symptoms. Examination by oblique illumination will show a grayish or whitish opacity on a black ground, and with the ophthalmoscope at a distance a black opacity upon a red field. (Plate II.). The pupil should be dilated, in order that the lens and the fundus may be examined. During the stage of swelling of complete cataract the anterior chamber is reduced in depth. This swelling of the lens may cause increase of tension in an eye predisposed to glaucoma.

PROGRESSIVE CATARACTS.

Senile Cataract.

Senile cataract is the most common form. It is a disease of advanced life, though occasionally it is seen as early as forty years of age. As a rule, both eyes are involved, but generally one in advance of the other. The opacity may begin either in the superficial part of the cortex (cortical, Fig. 204), or in the part immediately surrounding the nucleus (nuclear, Fig. 205). Senile cataract involves the cortex, the nucleus remaining transparent throughout the process. The time required for full development varies greatly. It may ripen completely in a few months, or may require many years. It may become stationary at any stage of its progress.

The Stages of Cataract.—Four stages are usually described, though it must be remembered that each stage passes imperceptibly into the next.

1. Incipient Stage.—The opacity most frequently begins as streaks, which extend from the periphery of the cortex,



FIG. 204.—SENILE CORTICAL CATARACT.

A, Seen with oblique illumination; *B*, section of the lens; *C*, seen with the ophthalmoscope.

where they are wider, to the centre of the lens, where they narrow like the spokes of a wheel (Fig. 204). The periphery is affected first. These streaks appear grayish by oblique illumination, and black when seen with the ophthalmoscope. Between these sectors the lens is transparent. Less frequently, senile cataract begins

with dot-like or cloud-like opacities situated in any portion of the lens. Sometimes the portion immediately surrounding the nucleus becomes opaque, constituting so-called nuclear cataract (Fig. 205). The last form causes relatively great visual disturbance. Cataracts sometimes remain stationary in the incipient stage, with little impairment of vision. Hence it is often wise not to alarm the patient by acquainting him with his condition, though it may be advisable to communicate the knowledge to a relative, for one's own protection.




FIG. 205.—SENILE NUCLEAR CATARACT.

A, Seen with oblique illumination; *B*, section of the lens; *C*, seen with the ophthalmoscope.

2. The Stage of Swelling (maturing stage).—The lens absorbs fluids, swells, and by pushing the iris forward reduces the depth of the anterior chamber. It appears bluish-white, shining, and presents distinctly the markings of the stellate figure. During this stage the iris casts a shadow upon the lens when the eye is illuminated from the side, since the superficial portion of the lens is still transparent, the opaque layer being some distance behind the iris.

3. Mature Stage.—The lens loses most of its fluid, shrinks

somewhat, and becomes opaque and of a dull gray or amber colour, the stellate markings still being recognizable. The anterior chamber regains its normal depth, and there is no shadow thrown by the iris on the lens with focal illumination. Occasionally the entire lens is changed into a hard, dark brown mass (black cataract). In this stage the cataract can easily be separated from the capsule of the lens. It is then said to be 'ripe' for operation, since it can be extracted whole, without leaving any portion of the cortex behind.

4. *Hyper mature Stage*.—The cataract may continue in the mature stage for a long time. If changes continue, the surface of the lens loses its radial markings and becomes homogeneous, or presents irregular spots. The cataract may continue to lose water, and thus a shrunken, dry, flattened mass results (shrunken cataract), with some deepening of the anterior chamber. Or, the cortex may become soft, liquid, and milky, and the nucleus sink to the bottom of this fluid (*Morgagnian cataract*), the cataract appearing white with a brownish colouring below. Very old hyper mature cataracts often present the deposit of cholesterin or of lime-salts. The latter change (chalky cataract) is found chiefly in complicated cataracts. The anterior capsule may become thickened and opaque (capsulo-lenticular cataract). The lens (and iris) may become tremulous through stretching of the suspensory ligament. For these reasons operation upon over-ripe cataract is often less favourable and more difficult than during the mature period.

Pathology.—Senile cataract results from shrinkage of the nucleus together with the lens fibres, and presence of fluid in the spaces thus created. The lens fibres then swell, become vacuolated (*Morgagnian spherules*), then cloudy, and disintegrate. Finally the lens substance is changed into a soft mass consisting of fat, spheres of Morgagni, remains of lens fibres, and albuminous liquid; this mass separates from the capsule. The nucleus usually remains transparent, but becomes harder and denser.

Treatment.—*Extraction* of the lens by operation is the only

means of relieving a patient of senile cataract. Discission is applicable only to cataracts of the young. No medicinal treatment, whether local or constitutional, has proved of curative value. The daily instillation of dionin drops (1 per cent.) has seemed to delay the progress of lens opacities, but, as these in many cases cease to progress without treatment, one cannot be sure. When an incipient cataract is detected, the eye should be carefully examined, and the result noted for future reference. The patient should be re-examined from time to time. If glasses assist vision, they should be ordered. There is no objection to the patient's using a magnifying-lens or making any use he can of his failing sight. In cases in which the opacity is central, sight may be improved temporarily by the instillation of a weak solution of atropine ($\frac{1}{2}$ grain to ℥i.), to cause mydriasis and enable the patient to see through the peripheral transparent portion of the lens. But the effect of the mydriatic must be watched, and the possibility of a glaucomatous rise of tension must be borne in mind.

The most favourable time for extraction of senile cataract is when the lens is completely opaque, and there is no shadow thrown by the iris—*i.e.*, when the cataract is ripe. If operated upon before this time, the lens is not always removed cleanly, and some transparent cortex is apt to adhere to the capsule and be left behind. This becomes opaque subsequently, and is absorbed slowly; besides, the remains of cortex after extraction tend to produce irritation, and interfere with smooth healing. However, the removal of cortical remnants by irrigation of the anterior chamber with sterile normal saline greatly lessens the disadvantages of operating upon immature cataracts.

As a general rule we operate when the cataract of one eye is complete and the other has progressed far enough to cause considerable interference with vision. But there are exceptions to this rule. For instance, when useful vision is abolished in both eyes before either cataract is ripe; to increase the field of vision on the side of the cataract for sake of safety; when the cataract shows signs of hypermaturity before the

second eye is much affected; or for cosmetic reasons. Removal of both cataracts should never be performed at one sitting. *Artificial ripening* is very occasionally resorted to. An incision is made through the periphery of the cornea, and the aqueous humour allowed to escape, so that the cornea falls against the lens. Then the cornea over the pupil is stroked with a smooth instrument, or the spatula or spoon may be introduced into the anterior chamber and applied directly to the lens capsule. Such direct or indirect massage may be done with or without an accompanying iridectomy. After this operation, the lens may become opaque within a few weeks, and can then be extracted. Ripening operations are, however, not reliable nor free from danger. It is certainly safer and better to remove the immature cataract than to resort to such artificial ripening.

Extraction may be performed with (*combined extraction*) or without (*simple extraction*) an iridectomy. The question as to which is the better operation has been much discussed. The chief advantages of the simple operation (without iridectomy) are the avoidance of tags of capsule healing in the wound, less reaction, and less likelihood of iris becoming adherent to capsular membrane; its disadvantages are the slight additional pressure on the eye which is required to press the upper edge of the lens through the intact sphincter and the danger of prolapse of the iris. This latter risk is reduced, with retention of all the advantages of a simple extraction, if we perform a small peripheral iridectomy by excising a very small part of the outer portion of the iris after delivery of the lens. The combined operation is undoubtedly the safer with a very large lens, in an eye which has suffered from iritis, or in cases in which the vitreous is known to be fluid. A successful simple extraction leaves a beautiful round pupil, and the action of the iris is not impaired. But in the combined operation the coloboma is covered by the upper lid. The patients are usually elderly people, few of whom would choose to incur any extra risk to gain a slight improvement in appearance. Some operators perform a preliminary iridectomy, and an extraction several weeks later, as a means of lessening the

dangers of extraction when complications are feared; but the disadvantages of an extra operation probably outweigh any advantage there may be.

Uncomplicated monocular cataract, with good vision in the other eye, should not as a rule be removed, since, owing to the difference in refraction, the eyes will not work together. Extraction may, however, be performed in such cases for cosmetic effect, to prevent hypermaturity, or to extend the field of vision on the affected side, but should not be attempted if the cataract is secondary to any disease in the eyeball.

Aphakia.—After the extraction of cataract the patient is compelled to wear strong convex glasses, since, as a result of loss of the lens (aphakia), there is a high degree of hypermetropia and absence of the power of accommodation. This hypermetropia amounts to about 10 D. With it there is usually astigmatism (2 to 3 D.), generally 'against the rule,' a result of the incision. In an average case, therefore, a convex spherical lens of about 10 D., combined with a convex cylinder of 2 to 3 D., must be worn for distant vision. To this sphero-cylinder an additional convex sphere of 3 or 4 D. must be added for reading. Any previous error of refraction will, of course, modify this correcting lens. Glasses should not be prescribed until all signs of irritation have disappeared—generally at the end of a month. Changes in refraction, usually a lessening of the post-operative astigmatism, may continue for several months. The aphakial eye presents, besides hypermetropia and loss of accommodation, a deep anterior chamber and usually a tremulous iris. The images normally seen on the anterior and posterior surfaces of the lens are absent.

Prognosis.—A favourable result and useful vision should follow cataract extraction in nearly all uncomplicated cases. There is generally good, and often perfect, vision. The success of the operation is dependent not only upon skilful operation, but upon extreme gentleness on the part of the surgeon, and upon the establishment of a sympathetic understanding between surgeon and patient. There should be no eyes lost through patients 'behaving badly' during the operation. Before deciding to operate it is important to investigate the

condition of the other ocular structures, and especially that of the retina. This is done by testing the field of vision with a light from an ophthalmoscope mirror for light perception and light projection. There should be a good field and good perception and projection of light. Lid margins, conjunctiva, and lacrymal sac must be examined by inspection and perhaps bacteriologically. If they are not clean and healthy they must be made so before operation is undertaken.

Projection is tested by throwing light from the mirror of the ophthalmoscope upon the upper, lower, inner, and outer portions of the retina. There is good projection if, with the eye directed forward, the patient is able to state correctly the direction from which the light comes. This test may also be applied with the lighted candle made to approach the eye from various directions, at a distance of 1 metre, and also at a greater distance—3 to 4 metres. Although the cataract be fully matured, there should be good perception of light, even with faint illumination. Fingers can frequently be counted at several inches.

Cataract Extraction.

Besides the division of cataract extraction into combined and simple, as explained above, there are two variations associated with either method. In one, the capsule of the lens is opened by means of the cystotome or capsule forceps and the operation is called *extra-capsular*; this is the most common method of extraction. In the other, no opening is made in the lens capsule, and the cataract is delivered enclosed in its capsule; this method is known as *intra-capsular* extraction.

There are many possible variations in technique, full discussion of which would take up much space. It seems, therefore, best to describe in detail the operation best suited to the beginner.

The combined operation is therefore described first, and the points of difference in the simple extraction afterwards.

Instruments Required.—An eye speculum—Lang's (Fig. 185) with solid blades has the advantage of keeping the lashes away from the knife. Fixation forceps (Fig. 186). A narrow Graefe.

knife (Fig. 207). Cystotome (Fig. 209). Two iris repositors (Fig. 208). Two curettes (Fig. 216). Curved iris forceps (Fig. 188). De Wecker's iris scissors (Fig. 211).

There should also be ready, although one hopes not to have to use them, a lens scoop (Fig. 212), or wire vectis (Fig. 213), and an irrigator for the anterior chamber. The irrigator consists of an undine (Fig. 214), to the nozzle of which is attached 18 inches of fine rubber tube with a fine flat canula (Fig. 214A) at the other end. After the irrigator has been sterilized some normal saline solution is run through it, it is filled with sterile normal saline solution, and the end with the canula is dropped into the large opening of the undine until required. The apparatus should stand in a bowl of warm water to keep it at about blood heat ready for use. The eye is well cocainized



FIG. 206.—THIN, BROAD CATARACT KNIFE.



FIG. 207.—NARROW GRAEFE KNIFE.

for fifteen minutes prior to operation; a drop of cocaine is instilled in the opposite eye while the patient is on the table; a general anæsthetic is seldom used. Sometimes a little 3 per cent. novocain is injected in the palpebro-temporal region to paralyze the orbicularis and to prevent squeezing. Very nervous patients may be quieted by a hypodermic injection of morphine and atropine given half an hour prior to operation, but this is seldom necessary in this country. The patient lies on the table with his head well up to the end, and the chin slightly raised. See that he is in a comfortable, easy position, and that the height and the light are exactly right for the operator. Tell the patient to keep looking downwards during the operation; explain to him that the lids will be held open by a prop to prevent blinking, but that he must be careful not

to squeeze the lids together voluntarily lest it should injure the eye.

Insert the speculum and cut short the lashes of the upper lid which protrude on the outer side. This is done with a pair of scissors greased with sterile vaseline, which prevents loose lashes falling into the eye. Suppose, for example, the left eye is to be operated upon. The patient looks slightly downwards, and the surgeon, standing behind the patient's head and holding the Graefe knife with the left hand, with the fixation forceps held in the right hand, takes a firm hold of the tissues near the lower margin of the cornea. The surgeon's left wrist is steadied against the patient's head. The Graefe knife is held between the thumb and two first fingers, the flat of the blade being parallel to the plane of the corneal margin with the edge directly upwards (anatomically).



FIG. 208.—SILVER IRIS REPOSITOR.



FIG. 209.—CYSTOTOME (SICKLE).

The point of the knife is entered just posterior to the plane of the transparent margin of the cornea about 2 mm. above the horizontal meridian. The knife is thrust across the anterior chamber and made to emerge at the corresponding point on the nasal side, so that the section comprises a little less than one-half the circumference of the cornea (Fig. 210 *a*). If there is room without touching the side of the nose, the thrust is continued for the whole length of the knife blade, cutting upwards at the same time. Then the knife, still cutting upwards, is withdrawn until the point has nearly reached the nasal side of the section. Other to-and-fro movements are made if necessary, and the corneo-scleral section is completed. Conjunctiva lies in front of the knife near the apex of the incision (Fig. 210 *b*). When the conjunctival flap is long enough, the edge of the knife is turned forwards and the apex

of the flap cut through. During the incision a slight pull should be maintained on the forceps so as to **keep up** a little negative pressure in the eyeball. During the whole section great care must be taken to keep the knife exactly in the same plane without any twisting or turning, lest aqueous escape and the iris fall forward in front of the knife. The conjunctival flap should be about $\frac{1}{4}$ inch wide and about the same length. Various larger conjunctival flaps are sometimes prepared in advance with sutures in place to be tied on completion of the operation. Sometimes an uncut conjunctival bridge is left after completing the section, but makes the expression of the lens more difficult. Such variations are useful in patients




FIG. 210.—EXTRACTION OF CATARACT.

a, Section of cornea; *b*, cutting conjunctival flap; *c*, expressing lens.

with a chronic cough, or in those known to be liable to 'squeeze.'

On completing the section the conjunctival flap is turned back on to the cornea as the knife is removed. An assistant, who knows beforehand exactly what he has to do, passes his hand low down at the side of the patient's face and takes the weight of the speculum from the eye. The fixation forceps is laid aside, the patient continues to look slightly downwards, the surgeon takes the curved iris forceps in his left hand and the de Wecker scissors in the right, and with the forceps grasps a small piece of the iris near the upper margin of the pupil, draws it slightly out of the wound, and snips it off with the

scissors, the blade of the scissors being held radially (Fig. 194, p. 229). The coloboma should be quite narrow.

The cystotome is then introduced flatwise, and its point turned to cut the capsule gently and without pressure; the incision may be various shapes -- **T**-shaped, **A**-shaped, or peripheral and concentric with corneal margin. One of the curettes or a Daviel's spoon is then laid flat on the sclerotic encircling the lower margin of the cornea. It is gently rocked to and fro, and at the same time moved slightly upwards. By this means the lens is loosened and separated from its capsule. When the expresser has travelled well within the corneal margin it is gradually tilted so that its convexity presses backwards in order to tilt the lens. At the same time the posterior lip of the incision is pressed backwards with the silver spatula. The rocking movement of the expressor is continued as the lens travels forwards and upwards, presenting in the wound and coming out through the wound (Fig. 210 c). Do not hurry with the expresser, and take care that it does not travel in advance of the lens. If any part of the periphery of the lens appears to have been left behind the iris, it may perhaps be coaxed out by the expresser applied to the cornea. If there is soft lens matter remaining in the anterior chamber after the delivery of the lens, take the clean curette in place of the spatula, insert it just within the lip of the wound, and continue manipulation with the expresser. All, or nearly all, the soft lens matter will then run out along the groove of the curette. Gently wipe away with a piece of soft linen any blood or lens matter which may be on the wound. Take a clean spatula, insert it within the wound, and gently stroke the




FIG. 211.—DE
WEECKER'S IRIS
SCISSORS.

pillars of the iris into position. Then with the iris forceps search along the wound for any invisible tags of capsule which might become incarcerated in the wound. If a tag of capsule is grasped in the forceps it should be snipped off. Replace the conjunctival flap carefully with the spatula, lift the upper lid



FIG. 212.—SPOON.




FIG. 213.—WIRE LENS VECTIS.

by means of the lashes or the loose skin of the lid grasped between the finger and thumb. Remove the speculum and carry the lid away from the eye and downwards over the eye, and tell the patient to keep it gently closed. Instil a drop of 1 per cent. atropine into the operated eye and lay a piece of sterile linen greased with parolin over the closed lids of each eye with a bit of absorbent gauze over it. This makes the first dressing very easy to remove. Secure the two pads with the Moorfields bandage (Figs. 349 and 350).




FIG. 214.—UNDINE IN STAND.

Accidents and Modifications.—

In making the incision, if aqueous should escape and the iris fall across the knife, the incision should




FIG. 214A.—CANULA.

be continued as if nothing had happened. The resulting coloboma will be large and ugly, but the eye will in all probability do quite well.

The object of the usual small conjunctival flap is not to cover the wound, but to assist in the nutrition of the cornea, and promote rapid healing. The larger flaps that are sometimes made when complications are anticipated may cause blood-clot to remain between the lips of the wound, and this may give rise to considerable post-operative astigmatism as the scar contracts.

If the lens does not present in the wound, it may be due to the capsule not having been sufficiently opened, in which case the capsulotomy should be repeated. It may be due to too great haste with the expresser. Try again, taking plenty of time over it, and see that the expresser does not over-run the lens. It may be due to the incision being too small. This is a bad mistake which should never be made if the surgeon thinks out the proposed procedure carefully beforehand. The only thing to do is to enlarge the incision with short-bladed, angled, blunt-pointed scissors.

Rarely, with a diseased eye and a weak suspensory ligament, the lens may become dislocated backwards. Attempts at expression should not be continued lest it disappear into the vitreous. The spoon or wire vectis should be pushed in well behind it, and it should be hooked upwards against the back of the cornea.

If the vitreous should present in the wound before the delivery of the lens, no attempt should be made to deliver it in the ordinary way, as this would certainly result in loss of vitreous. The scoop or vectis should be used at once. Vitreous may escape after the delivery of the lens and during manipulation for the removal of the cortical débris. Exceptionally this may be unavoidable, but it is usually due to want of gentleness in the manipulation or to violent squeezing on the part of the patient.

In operating upon an immature or hypermature cataract there may be tenacious lens matter which cannot easily be removed by manipulation from the anterior chamber. If this is quite small in quantity it may be left to absorb, otherwise an attempt should be made to remove it with the irrigator. The nozzle should be introduced just within the wound,

slightly depressing the posterior lip. Only just sufficient head of fluid should be employed to enable it to enter the anterior chamber freely.

Simple Extraction differs from the combined operation essentially in the omission of iridectomy. It avoids the risk of capsule being caught up in the wound. Before the operation the pupil should be fully dilated with atropine. This facilitates the delivery of the lens, avoids straining the sphincter, and makes post-operative prolapse less likely.

Extraction with Peripheral Iridectomy.—Many surgeons, nowadays, prefer to make a small 'button hole' in the iris, as peripherally as possible, after performing a simple extraction without preliminary atropinization. This allows the aqueous to circulate freely between the posterior and anterior chambers and thus diminishes the risk of prolapse. One prefers this operation in all suitable cases, reserving the combined extraction for cases complicated by disease in the eye when vitreous is fluid; rather more pressure is needed to free the lens than in the combined operation.

After-Treatment.—The patient is directed to lie quietly upon his back. An anodyne is often advisable. After twenty-four hours he may change to the side of the unoperated eye. His food should be fluid for the first day. The bowels need not be relieved artificially for three or four days. If the patient has a motion before this, he must be cautioned not to strain. The wound is inspected after twenty-four hours (or forty-eight hours if the patient is comfortable), extreme gentleness being especially essential at this first dressing. Atropine is instilled each day. On the fourth or fifth day the unoperated eye may be left free. After a week the patient may sit up in bed for an hour or two, and a day or two later may sit in an easy-chair the greater part of the day. After ten days nothing but smoked glasses need be worn.

Post-operative Complications.—If iridectomy has been performed one of the pillars may prolapse into the wound within a few hours. If it is not completely prolapsed it should be replaced with the spatula inserted into the wound. Any part of the iris which has remained for a few hours outside the eye

should be excised and the stump replaced with the spatula. In simple extraction a considerable prolapse of the iris is usually announced by severe pain. It should be dealt with as soon as possible. It may be possible to do what is required under local anæsthesia, or a general anæsthetic may be needed. It is generally inadvisable to attempt to return any part of the iris which has been completely prolapsed; as it may not be sterile; moreover, the prolapse may probably recur. An incomplete prolapse may perhaps be smoothed out, with the spatula introduced into the wound at one side.

Hæmorrhage into the anterior chamber is from new capillary vessels which are bridging across the wound. It is more likely to occur if an exceptionally wide conjunctival flap has been made. The time after operation is almost exactly 120 hours. It is especially important, therefore, that the patient be kept very quiet for the first six days. The hæmorrhage may completely fill the anterior chamber. Nothing should be done beyond keeping the patient quiet. The blood may absorb completely, or the part in the area of the pupil may partly organize and add to the thickness of the capsular membrane.

Rarely there may be iritis due to the traumatism. Suppuration of the wound and panophthalmitis were frequent in pre-aseptic days, but are now very rare. Iridocyclitis with k.p. is a late complication due to auto-intoxication which we are not always able to guard against.

After-Cataract.—When the eye has healed after the extraction of cataract, examination with a lens and focal illumination shows a membrane in the pupillary area. This consists of posterior capsule, perhaps fragments of anterior capsule, and in some cases organized remains of blood-clot or lens matter enclosed between the posterior and fragments of the anterior capsule. There may be bands due to iritis. Sometimes the capsule is perfectly transparent and smooth, and does not interfere with vision. If it is wrinkled it may interfere with sight even though it be transparent. It may be so dense as to interfere very greatly with sight. Within a few weeks of the extraction this membrane is usually quite easily cut.

The tendency is for it to become tougher and denser, so that a membrane which scarcely reduces vision at all at first may cause considerable diminution later. Usually discission with proper precautions is practically free from risk. It is therefore good practice to needle every case as soon as the eye is quiet, perhaps two or three weeks after operation. The eye having been cocainized and prepared and the pupil widely dilated with atropine, the speculum is inserted. The surgeon, standing behind the patient's head, grasps the eyeball with the fixation forceps a little below the corneal margin. A very sharp, short Ziegler's knife is introduced through the sclerotic just behind the periphery of the cornea, and below the original incision, and the point thrust through the membrane, the cutting edge upwards. The membrane should be cut from behind forwards with gentle sawing movements, to avoid



FIG. 215.—ZIEGLER'S KNIFE.

injury to the vitreous, carefully avoiding any dragging or tearing.

Extraction of Cataract in its Capsule (Intra-Capsular Extraction) was the method originally advocated by Smith and performed by him with success upon large numbers of cases in India. It is a combined extraction with the omission of capsulotomy. As performed by Smith, the lens was dislocated and expressed within its capsule, after an iridectomy, by strong pressure on the cornea with a squint hook, the lens being delivered lower pole uppermost. The risk of loss of vitreous prevented the operation from becoming popular in Britain, though the removal of the whole capsule with the lens and the avoidance of after-cataract is doubtless of great advantage, especially when patients come long distances and cannot readily return for a second operation. But it has now become usual to modify the technique by combining traction with pressure in the removal of the lens, thereby considerably reducing the risk of vitreous loss, and the operation has gained

some adherents in this country and a number abroad. It is contra-indicated in juvenile and traumatic cataracts, and also when there are posterior synechiæ or other complications.

Intra-Capsular Extraction after Subluxation with Capsule Forceps, as practised by Elshnig, Sinclair and others, consists in subluxating the lens by traction. After corneal section, with the pupil well dilated with atropine, the lens capsule near the lower pole is grasped firmly with special capsule forceps, and the lens rocked gently from side to side to rupture the suspensory ligament; backward pressure is then applied to the lower part of the cornea, while traction on the capsule is maintained, until the lens tumbles over and presents in the section lower pole uppermost; the upper part of the suspensory ligament is then separated. Prior to this operation it is usual to paralyze the orbicularis by an injection of novocain to block the facial nerve. When the operation is completed, after instillation of atropine, the upper lid is drawn down over the eyeball by a suture which is fastened to the cheek by a strip of plaster.

Other operations have been devised to withdraw the lens in its capsule without pressure. The best known is that of Barraquer, who attaches a special instrument to the lens capsule by means of suction, obtained by means of an electric motor. This operation is known as *Phakoerisis*, and the apparatus as the *Erisophake*. The most recent method is that of Lacarrère, who, by diathermy, fuses the lens to two small wires attached to a suitable insulated handle, and then withdraws it from the eye.

Curette Evacuation is performed for the removal of soft or traumatic cataracts, in patients between 20 and 40. It is rarely required in children, as soft lenses completely dissolve



FIG. 216.—CURETTE.

and drain away in the aqueous. Instruments required: speculum, fixation forceps, broad needle (Fig. 217), curette (Fig. 216), and silver spatula (Fig. 208). Local anæsthesia

is sufficient except in the case of very young children. The broad needle is entered through the cornea near its periphery on the temporal side, and it is thrust in, making a large rent in the anterior capsule of the lens. During the withdrawal of the instrument the incision is enlarged to a width of about 5 mm. The tip of the curette is then introduced just within the wound, slightly pressing back the posterior lip; the soft



FIG. 217.—BROAD NEEDLE.

lens matter runs out along the groove of the curette. If no more lens matter comes away while a considerable amount is left, close the eye for two or three minutes to allow a little aqueous to form and try again. The curette should not be introduced far within the wound in an attempt to remove the lens matter mechanically. Any which will not come away may be left to absorb, or washed out with an irrigator.

Congenital Complete and Juvenile Complete Cataract.

These forms of cataract are infrequent. The lens is uniformly white or bluish-white, or it may have a pearly lustre. It is always soft. Sometimes it is fluid and milky. These forms of cataract may occur in eyes which are otherwise perfectly healthy, or they may be complicated cataracts, with changes in the retina, choroid, or optic nerve. One or both eyes may be affected. The congenital complete cataract is due to a disturbance of development, or to some intra-uterine ocular inflammation. The complete cataract of young people (juvenile) may be due to heredity, or arise without known cause. In some cases there is a history of convulsions.

Treatment consists in discission (needling). This should be done as early as possible, so that disuse of the function of sight may not cause amblyopia. The needle operation must usually be repeated a number of times. Sometimes there

are remains of the lens which do not become absorbed, and must subsequently be removed by extraction. Semi-fluid cataracts are removed by linear extraction.

Discission of the Lens (Needling).

Indications.—In zonular, congenital complete, and juvenile complete cataracts (soft cataracts), previous to the fifteenth year.

Operation.—In young children a general anæsthetic is required; in others local anæsthesia is sufficient. The pupil must be dilated. The speculum is introduced, and the eye-ball steadied with the fixation forceps. A knife-needle (Fig. 218) is thrust through the sclerotic near the margin of



FIG. 218.—KNAPP'S KNIFE-NEEDLE.

the cornea, and then through the capsule of the lens, making two cross-cuts, each about 4 mm. in length. The cuts must be superficial, especially if this is the first operation, so that there will not be too rapid swelling of the lens. The lens substance may be broken up a little by rotating the needle. After some of the swollen lens matter has been absorbed (several weeks) the operation must be repeated. At the second operation the discission may be deeper and bolder. At the last of the several operations the incision must include the posterior capsule.

After-Treatment.—There is usually very little reaction. The pupil must be kept dilated with atropine. The lens substance swells, protrudes through the opening in the capsule, and pieces fall into the anterior chamber and become absorbed. Usually three operations are required. The entire duration of treatment is several months.

Complications.—Rapid and extensive swelling of the lens may cause a sudden rise of tension of the eye, requiring removal of the lens by linear extraction without delay. A bold

discission is sometimes done with a view of extracting the lens a few days afterwards, as soon as there is marked swelling. Iritis may occur after discission, occasionally iridocyclitis, and very rarely loss of the eye.

Traumatic Cataract.

This form of cataract is the result of a perforating wound of the lens capsule. Occasionally it ensues after contusions of the eyeball, without perforation (concussion cataract), though probably a rupture of the capsule occurs in such cases. Within a few hours after the injury the lens becomes cloudy at the seat of the wound from absorption of aqueous humour and swells. Opaque and swollen lens substance protrudes through the wound in the capsule, and often falls into the anterior chamber. The swelling and clouding continue until, after a few days, the entire lens has become opaque. Then the lens substance becomes absorbed. In favourable cases in young persons this process continues until there is spontaneous cure with a clear, black pupil. More frequently, however, part of the lens remains opaque in the capsule and requires subsequent operation. Occasionally the opacity of the lens remains limited to the injured portion, due, probably, to closure of the small capsule opening. The course described may be less favourable. Inflammation of other parts of the eye may result—iritis or iridocyclitis, or when infection occurs, panophthalmitis. The swelling of the lens may cause iritis or glaucoma.

Treatment.—Immediately after the injury absolute rest, iced compresses, and atropine are to be employed. If the rapid swelling of the lens causes inflammation or much increase of tension, the cataract should be removed by extraction. But if such complications do not arise, it is wiser to allow absorption to proceed, and to defer operative intervention until there is no irritation or inflammation, and spontaneous improvement has come to a standstill.

Stationary Cataracts.

Anterior Polar or Pyramidal Cataract.—This lenticular opacity occurs in the form of a small, round, white opacity, often pyramidal in shape, situated at the anterior pole of the lens, beneath the capsule (Fig. 219). It may be congenital or acquired. The acquired form originates from an ulcer of the cornea in early childhood. Such an ulcer perforates, and allows contact and pressure between lens and cornea, setting up an irritation in the anterior capsule which results in a proliferation of the subcapsular epithelium. Afterwards the anterior chamber is restored. Sometimes there is an accompanying corneal opacity. As a rule this form of cataract does not interfere with vision sufficiently to require treatment.



FIG. 219.—ANTERIOR POLAR CATARACT.

A, Seen with oblique illumination; B, section of lens; C, seen with the ophthalmoscope.

Posterior Polar Cataract.—This form may be congenital (capsular) or acquired (cortical).

The *congenital form* is a capsular opacity consisting of a small, round, white deposit, situated at the posterior pole.



FIG. 220.—ACQUIRED FORM OF POSTERIOR POLAR CATARACT.

A, Seen with oblique illumination; B, section of the lens; C, seen with the ophthalmoscope.

With the ophthalmoscope it appears as a black dot upon the red fundus-reflex. It represents the remains of the hyaloid artery at the point of attachment to the posterior capsule of the lens. It causes but trifling interference with vision, and requires no treatment.

The *acquired form* is a grayish, stellate opacity of larger size, situated in the cortical layer of the lens, at its posterior pole (Fig. 220). It is a form of secondary cataract, which develops in connexion with high myopia, choroiditis, disease of the vitreous, and retinitis pigmentosa. It remains stationary

for many years, but is apt finally to become complete. In this affection there is considerable impairment of vision, caused not only by the cataract, but also by the accompanying disease of the deep structures. It does not admit of treatment.

Lamellar or Zonular Cataract.—This variety of partial, stationary cataract is either congenital or forms in early childhood, and usually affects both eyes. It is the most common form of cataract seen in children. It is sometimes hereditary, and often associated with a history of convulsions and with the changes of rickets, especially in the teeth and bones. It consists of a gray, disc-like opacity of the layer surrounding the transparent nucleus, with clear cortex on




FIG. 221.—ZONULAR CATARACT.

A, Seen with oblique illumination; *B*, section of the lens; *C*, seen with the ophthalmoscope.

the outside (Fig. 221). When the pupil is dilated, examination by oblique illumination shows a grayish disc surrounded by clear lens substance. From the margin of the opacity short striæ are often seen projecting into the surrounding transparent cortex. The cataract is most dense at the margin of the disc. This distinguishes it

from nuclear cataract. By the use of the ophthalmoscope at a distance the cataract presents a dark disc surrounded by a zone of red fundus reflex. The disc is somewhat lighter in the centre than at the periphery, and in the former situation allows some light to pass.

Lamellar cataract may remain stationary, or the opacity may increase. It causes interference with vision. The amount may be slight or decided, depending upon the extent and density of the opacity.

Treatment.—When sight is considerably interfered with we can improve vision by iridectomy, by discission in the young, or by extraction in older persons. Iridectomy (small coloboma downward and inward) is indicated when the vision is very materially improved after the use of a mydriatic. Its advantages are that the patient does not require strong convex lenses, and often retains binocular vision. Its dis-

advantages are the elongated pupil, and some dazzling due to this. Removal of the lens by discission or extraction is indicated in those cases in which there is little or no improvement in sight after dilatation of the pupil, and when there are indications of progress of the cataract.

Various Uncommon Varieties of Stationary, Partial Cataract are met with. These include (1) central cataract, a small, white opacity in the centre of the lens; (2) fusiform cataract, a spindle-shaped opacity running from the anterior to the posterior pole; (3) punctate cataract, consisting of a number of very small white (very occasionally bluish) dots variously distributed through the lens; and (4) discoid cataract, an ill-defined disc situated between the nucleus and the posterior pole. These opacities are usually congenital, cause little interference with vision, but are often associated with other ocular defects.

Complicated or Secondary Cataracts.

These accompany or follow other diseases of the eye. The most frequent ocular affections which lead to cataract are iridocyclitis, choroiditis, severe forms of corneal ulcer, glaucoma, retinitis pigmentosa, and detachment of the retina. Such cataracts frequently begin in the posterior part of the lens, often have distinctive features, and tend to degenerate. It is important to establish the fact that a cataract is complicated when the question of operation presents itself. The treatment of complicated cataract is usually very unsatisfactory, and the prognosis is always less favourable than in uncomplicated cases. This is because the operation is rendered difficult and the effect on sight disappointing by the complicating ocular disease. Many cases cannot be operated upon.

Dislocation of the Lens.

Dislocation of the lens may be partial (subluxation) or complete (luxation), congenital or acquired.

Symptoms are disturbance of vision, interference with accommodation, a change in refraction, monocular diplopia, and tremulous iris. They differ according to whether the

displacement is partial or complete. In addition there are complications and sequelæ which are often serious.

Partial Dislocation (Subluxation) may consist of a tilting of one edge of the lens, or of a lateral displacement—upward, downward, inward, or outward. In such cases the anterior chamber will be of unequal depth, being increased where the



FIG. 222.—DISLOCATION OF THE LENS UPWARD AND OUTWARD.

lens is absent. The convex edge of the lens can usually be seen (Fig. 222) in some part of the pupil, the portion of the latter which is free from lens being particularly black. With the indirect method of ophthalmoscopy the optic disc appears double, one image being seen through the lens and the other through the free pupil. Movements of the eyeball disclose a tremulous condition of the lens and iris (iridodonesis).

There are considerable myopia and astigmatism in the area corresponding to the lens, the convexity of the lens being increased through relaxation of the suspensory ligament; also marked hypermetropia in the aphakial area. Monocular diplopia may be complained of, two images being formed on the retina. The subluxated lens may become opaque.

Complete Dislocation (Luxation) of the lens may occur *anteriorly* into the aqueous, or *posteriorly* into the vitreous cavity. In traumatic cases in which there is rupture of the sclera the lens may lie *beneath the conjunctiva*.

When dislocated anteriorly, the lens is easily recognized. If it is transparent, it looks like a large drop of oil with a curved, golden margin when seen by oblique illumination. The anterior chamber is increased in depth.

When displaced into the vitreous, the lens sinks into the lowest part, and either becomes attached to the fundus by exudation or moves about. When opaque, it can be seen with the ophthalmoscope and sometimes with the unaided eye.

The anterior chamber is deep, the iris tremulous, and the pupil very black. The eye is, as in aphakia, in a condition of extreme hypermetropia, and has lost its power of accommodation.

Complications and Sequelæ.—A partial dislocation often changes to a complete one. When subluxated, the lens may remain clear a long time, but completely dislocated lenses soon become opaque. Choroiditis and iridocyclitis, secondary glaucoma, and even sympathetic ophthalmia, sometimes follow. Displacement into the vitreous is tolerated better than anterior luxation.

Etiology.—Dislocation of the lens may be either congenital or acquired. In order that the lens can become dislocated there must be some defect in the suspensory ligament, such as rupture, stretching, or imperfect development.

The *congenital form* is partial, usually upward, often becomes complete in after-years, is generally bilateral, and often hereditary.

The *acquired forms* are either traumatic or spontaneous. Traumatic dislocation is generally the result of contusions. The predisposing cause of spontaneous dislocations is a change in the suspensory ligament seen in fluid vitreous, choroiditis, and myopia of high degree, detachment of the retina, and hypermature cataract. The exciting cause may be insignificant, such as various straining efforts.

Treatment.—In partial dislocation, if no symptoms of irritation are produced, treatment consists in prescribing suitable glasses, usually strong convex lenses to correct the refraction of the aphakial portion. When the lens is dislocated into the anterior chamber it should be removed, by discission in young persons, by extraction in older cases. The lens should first be pierced by a needle to prevent its dislocation into the vitreous, and be removed by a spoon or wire scoop after corneal incision. If dislocated into the vitreous, an attempt to extract it is almost certain to fail. Strong convex glasses are prescribed for the aphakia. If inflammatory symptoms arise in a case in which the dislocated lens cannot be removed, an iridectomy may be tried; if in such cases the eye is sightless, enucleation is indicated.

CHAPTER XIX

DISEASES OF THE RETINA

Anatomy.—The retina is a thin, delicate membrane which consists, among other parts, of an expansion of the optic nerve. It is placed between the hyaloid membrane of the vitreous internally and the choroid externally. It extends forward to the ciliary body, where its termination is called the ora serrata; devoid of nerve fibres, simpler and thinner, it is continued over the inner surface of the ciliary body and the posterior surface of the iris. In the living eye, it is transparent and of a purple-red colour. Under the influence of light it is quickly bleached; after death it soon becomes opaque and white. The retina is connected with the subjacent choroid at the entrance of the optic nerve and at the ora serrata. Elsewhere it simply lies upon this tunic, but is not attached to it. When we detach the retina, the pigment cells which form its outermost layer adhere to the choroid, and on this account were formerly described as part of the latter.

The inner surface of the retina presents in the axis of the eyeball the *yellow spot or macula lutea*, about 1 to 2 mm. in diameter, and in its centre a small depression, the fovea centralis. This is the region of most distinct vision, and the part of the retina which is made to receive the image when we wish to get an exact impression of an object. About 3 mm. to the inner side of the posterior pole of the eye is a pale, round area, the *head of the optic nerve (papilla, or disc)*, corresponding to the point where the optic nerve pierces the retina (Fig. 43). The circumference of the disc is slightly elevated above the surface of the retina, but the centre presents a depression, the physiological cup or excavation. Here the bloodvessels of the retina enter the eye. The ophthalmoscopic appearances of the background of the eye and the distribution of the retinal vessels are given in Chapter III.

The *central artery* of the retina, accompanied by the corresponding *vein*, pierces the optic nerve about 12 mm. from the globe, and passes between the bundles of fibres to the inner surface of the retina at or near the middle of the disc. Excepting at the papilla, where minute communications are sometimes found between retinal and


ciliary vessels, the retinal arteries have no anastomosis. They are terminal branches; hence in obstruction of the central artery there is no compensatory collateral circulation, and blindness results. The retinal vessels lie in the inner layers. The external layers are destitute of bloodvessels, and are nourished by the adjacent chorio-capillaris. The fovea has no bloodvessels. In this situation, the chorio-capillaris is thickened. The bloodvessels are surrounded by lymphatic sheaths forming the lymphatics of the retina.

The *minute anatomy* of the retina is very complicated. We distinguish two kinds of tissue: (1) nervous elements, of which there are eight layers; and (2) supporting tissue (Mueller's fibres). The supporting tissue comprises the internal and external limiting membranes and numerous fibres serving to keep the delicate nerve-tissue in proper position.

Microscopic examination shows the following layers of the retina, from within outward (Fig. 223): (1) the internal limiting membrane; (2) the layer of nerve fibres, consisting of the expansion of the fibres of the optic nerve, destitute of medullary layer after piercing the eyeball; (3) the layer of ganglion cells, a stratum of large branching nerve cells; (4) the inner plexiform layer; (5) the inner nuclear layer; (6) the outer plexiform layer; (7) the outer nuclear layer; (8) the external limiting membrane; (9) the layer of rods and cones, the light-perceiving layer; (10) the layer of pigment cells, which bounds the retina externally, and consists of a single stratum of hexagonal pigmented cells.

The rods are much more numerous than the cones, excepting at the macula, where the cones preponderate. At the fovea there are no rods, and the cones, longer and narrower than elsewhere, are found exclusively. In this spot, also, all the layers of the retina are much thinner, there is no nerve-fibre layer, and Mueller's fibres are arranged obliquely. The disc consists of optic-nerve fibres exclusively. It has no other retinal nerve elements, and has no power of sight; hence it is called the blind spot.

Choroidal Surface.



Vitreous Surface.

FIG. 223.—SECTION OF THE RETINA, SHOWING THE MINUTE ANATOMY. (Modified from Schultze.)

1, Internal limiting membrane; 2, layer of nerve fibres; 3, layer of ganglion cells; 4, inner plexiform layer; 5, inner nuclear layer; 6, outer plexiform layer; 7, outer nuclear layer; 8, external limiting layer; 9, layer of rods and cones; 10, layer of pigment cells.

Physiology.—The action of light changes the visual purple contained in the outer segments of the rods into a colourless substance. When the eye is in the dark, most of the pigment is stored in the body of the cell, and is withdrawn from between the rods. After exposure to light, the pigment granules push their way inward into the processes extending between the rods and cones, and the latter become contracted and shortened. The function of the pigment cells is the renewal of the visual purple of the outer segments of the rods after the bleaching produced by exposure to light.

The rods and cones, the terminal organs of the optic nerve, receive waves of light, which fall upon the retina and convert these vibrations into impulses, which are carried by the optic nerves and tracts to the brain. Here they produce the sensation of light. When the image of an object falls upon the macula, there is distinct vision; when it falls upon any other part of the retina, there is indistinct vision. Two points give rise to separate visual impressions when their images are at least 0.002 mm. apart, since this represents the diameter of the cones at the fovea. Images which are closer than this would only stimulate one cone, and consequently create but one visual impression. In other words, to be seen distinctly, two objects must subtend a visual angle of one minute or more (p. 12).

Images of an object give rise to a single visual impression when they fall upon corresponding retinal areas; otherwise there are double images. In binocular vision certain portions of the retina are associated thus, the upper halves of the retina correspond, as do also the lower halves; but the nasal side of one retina corresponds to the temporal half of the other, and *vice versa*.

Rays of light impinging upon the retina come from the opposite side of the field; thus, the upper part of the retina is used for seeing objects in the lower part of the field, the temporal portion of the retina for the nasal part of the field, etc. The image on the retina is always inverted.

Affections of the Retina may be divided into:

1. *Inflammation*, the various forms of retinitis: (1) Simple, (2) albuminuric, (3) diabetic, (4) leukæmic, (5) syphilitic, (6) hæmorrhagic, (7) purulent, (8) uncommon forms of retinal changes.

2. *Vascular Changes*: (1) Anæmia, (2) hyperæmia, (3) hæmorrhages, (4) arterio-sclerosis, (5) embolism, (6) thrombosis.

3. *Pigmentary Degeneration* (retinitis pigmentosa).

4. *Detachment*.

5. *Tumour*: glioma (see chapter on Intra-ocular Tumours).

Retinitis.

Inflammation of the retina presents varied clinical types. There are, however, certain signs and symptoms which are more or less common to all varieties. Retinitis may be (1) primary, or (2) secondary, when it is an extension of inflammation of neighbouring ocular structures. It usually extends to both the papilla and the choroid. When the involvement of the entrance of the optic nerve is marked, we speak of the affection as neuro-retinitis; when the choroid is prominently implicated, we call the condition choroido-retinitis. The disease may be confined to one eye; but since it is generally dependent upon a constitutional factor, it is almost always bilateral. It may be acute in course, but, as a rule, it lasts many weeks, or even several months.

Subjective Symptoms.—(1) Diminution in acuteness of vision, varying with the severity and extent of the retinitis, but generally considerable. It may be especially marked at night, constituting night-blindness. (2) Changes in the field of vision. There may be concentric or irregular contraction or scotomata. (3) Alterations in the shape of objects: micropsia, objects appearing smaller than they really are; macropsia, objects appearing larger than normal; metamorphosia, a distortion of the shape of objects, straight lines appearing wavy and bulging. (4) Diminution of the light sense. (5) Feeling of discomfort in the eyes. (6) Photophobia may be present, but pain is rare.

Objective Symptoms.—There are no external signs; the objective symptoms are all ophthalmoscopic: Diffuse clouding of retinal details, especially in the region of the papilla; congestion of the disc, with indistinctness of its edges; circumscribed exudations appearing as soft, white, or slightly yellow spots or patches, discrete or confluent, varying in size, and found principally along the retinal vessels and at the macula; tortuosity and distension of the vessels, which may be obscured in parts by swelling and exudation; hæmorrhages of various shapes and sizes, rounded when occurring in the deeper layers,

and feathery or flame-shaped when superficial. There may be opacities in the vitreous.

Course.—The inflammation may subside completely, and useful vision return; or certain changes may occur in the retina as a result of atrophy, causing considerable impairment or absolute loss of vision. These changes are: Atrophy of the retina, allowing the choroidal vessels to become visible; bright, white patches and bright dots replacing hæmorrhages or exudation, and frequently pigmented; contraction of the vessels, which are bordered by white lines; atrophy of the disc, which presents an indistinct outline and a pale, dirty colour (post-neuritic atrophy).

Prognosis depends upon the severity of the inflammation, the parts of the retina involved most, and the clinical form of the retinitis.

Pathology.—The changes consist in congestion, œdema, exudation of white blood corpuscles and of fibrin, fatty degeneration, and extravasation of blood. The white spots are due to exudation of leucocytes and of fibrin, swelling of nerve fibres and cells, and fatty degeneration of the retinal elements and of exudation. The walls of the bloodvessels become thickened, so that in extreme cases the lumen may be filled with connective tissue. When the retina becomes atrophied, the nerve elements disappear and the membrane is replaced by connective tissue more or less pigmented.

Etiology.—Retinitis occurs occasionally as a local lesion. But generally it is merely a manifestation of a constitutional disease, such as nephritis, diabetes, syphilis, affections of the vascular system, etc. It may be due to auto-intoxication, or to extension from the choroid or ciliary body.

Treatment.—The local treatment consists in absolute rest for the eyes, protection from light, either by smoked glasses or the darkened room, and the use of atropine. Internally we give small doses of mercury, also iodide of potassium, diaphoretics, and sometimes cathartics. In addition, it is of the greatest importance to treat the constitutional condition which is the cause of the retinal lesion.

Simple Retinitis.

This disease, also known as serous retinitis and œdema of the retina, is an inflammation of the superficial layers of the retina, slight in degree and simple or serous in type; the evidences of inflammation are limited to swelling, vascular distension, and occasionally hæmorrhages. It is regarded, not as a distinct disease, but as the first stage of the more common forms of retinitis.

There is limited impairment of vision (often merely a blurred sensation), some distortion of images, and moderate peripheral contraction of the field. The ophthalmoscope reveals a hazy fundus especially around the disc, the margins of which are indistinct, veins somewhat dilated, tortuous, and hidden in places by the œdema, and at times hæmorrhages.

The prognosis is good when the affection remains of this type and does not change to the deep form. Treatment consists in the removal of the cause and the observance of the directions given above.

The Deep or Parenchymatous Type is a more intense inflammation involving the deeper layers of the retina; the pathological changes are more extensive, including, besides those occurring in the serous form, exudation, changes in the vessel walls, and hæmorrhages, and consequently capable of causing greater destruction with atrophy and permanent visual damage.

There are often marked ocular discomfort, much disturbance of vision and distortion of objects, peripheral contraction of the field, and scotomata. The ophthalmoscope discloses, in addition to the picture presented by the serous type, scattered yellowish patches of exudate, especially in the macular region, changes in the walls of the bloodvessels, and hæmorrhages.

This variety depends upon constitutional causes or represents an extension of or association with neighbouring ocular diseases. The prognosis is always serious; though some cases recover with fair or even good vision, many are left with marked impairment of this function. Treatment has been given under retinitis in general, and is discussed again in the clinical forms of retinitis in the following pages.

Albuminuric Retinitis.

Retinitis of Bright's Disease (Renal Retinitis) presents ophthalmoscopic signs which are often pathognomonic. It is usually bilateral, rarely unilateral.

Symptoms.—The subjective symptoms are those of retinitis in general (p. 275). The degree of disturbance of vision depends upon the severity of the inflammation and especially upon the position of the exudations and hæmorrhages. Minute changes in the macular region will cause considerable reduction in acuteness of vision, while extensive involvement of the rest of the fundus may affect the sight comparatively little.

Ophthalmoscopic Signs (Plate XVII.) are those of retinitis in general: swelling and haziness of the retina and of the papilla, distension and tortuosity of the retinal vessels, especially veins, and hæmorrhages either in the form of flame-shaped or round spots, or larger extravasations. To these are added the distinctive feature: white spots found chiefly at the macula and surrounding the disc, less frequently elsewhere. At the macula there may be at first merely a few dots, but later there are more pronounced spots, and these are usually arranged in radiating lines which form a star-shaped figure with the fovea for a centre; or when less complete, the lines resemble the sticks of an open fan; they have a certain brilliancy due to fatty degeneration of retinal elements and of exudation. Near the disc, and often more or less surrounding it, are larger white spots; these may coalesce and form a ring around the disc.

Though this is the most frequent picture of albuminuric retinitis, there are other and less characteristic signs in nephritis; there may be simply retinal hæmorrhages, simple retinitis, hæmorrhagic retinitis, neuritis, or even a picture of choked disc such as we are in the habit of associating with cerebral tumour. On the other hand a brain tumour may present ophthalmoscopic appearances identical with those of a typical case of albuminuric retinitis.

Albuminuric retinitis occurs under two forms: (1) the inflammatory, when swelling, congestion, and hæmorrhages

PLATE XVII.



FIG. 224.—ALBUMINURIC NEURO-RETINITIS.




FIG. 225.—ALBUMINURIC NEURO-RETINITIS
(LATER STAGE).

To face page 278.

are the predominating features; and (2) the degenerative, when the white spots and hæmorrhages occur without swelling or congestion. The two forms are usually associated in varying proportions.

Etiology.—The affection is usually a complication of chronic interstitial nephritis; much less frequently of chronic parenchymatous nephritis; it may occur with any form of nephritis, including that of scarlatina and pregnancy. From one-quarter to one-half of all patients with nephritis present some form of retinal lesion.

Pathology.—The retina presents œdema, hypertrophy of its elements, deposits of fat and fibrin, and hæmorrhages. The retinal vessels are thickened and the seat of hyaline changes with proliferation of the lining epithelium; these changes are similar to those taking place in the vessels of the kidney. The spots in the macular region are caused by fatty degeneration of exudate and retinal elements; their arrangement in a star-shaped figure depends upon the disposition of Henle fibres in this situation.

Course and Prognosis.—Though the retinitis is often a late symptom of Bright's disease, the disturbance of vision may be the first symptom which calls attention to the nephritis. Occasionally ophthalmoscopic examination first reveals the presence of Bright's disease in a patient who was unaware of any defect of sight other than presbyopia. There is no fixed relationship between the course of the nephritis, the amount of albumin, and the degree of retinitis. There are cases in which the vision is but slightly affected even in the late stages, and others in which sight is seriously affected quite early. The condition is of great prognostic importance, and indicates, with but few exceptions, a fatal termination in from six months to two years. The exceptions usually are cases occurring during pregnancy and scarlatina.

Treatment should be directed to the nephritis. No local treatment is of any value.

Gravidic Retinitis is the name given to retinitis complicating the albuminuria of pregnancy. Its signs and symptoms are the same as in the other forms of albuminuric retinitis,

DISEASES OF THE RETINA

but they tend to clear up after delivery. It usually occurs during the final months of pregnancy, and the prognosis in regard to vision is often good, especially if labour be induced prematurely. When it occurs in the early months, the prognosis is less favourable, and the condition may warrant the induction of abortion in order to save sight.

Uræmic Amblyopia is the term used for loss of sight during an attack of uræmia, without any changes in the retina. It is cerebral, not retinal, being due to retention in the blood of waste products which should have been excreted by the kidneys. It occurs in pregnancy and during the late stages of scarlatina. Similar attacks may also occur in patients who have albuminuric retinitis. It appears suddenly, affects both eyes, and is associated with other symptoms of uræmia—headache, vomiting, convulsions, and coma. The pupils are dilated, but respond to light. After lasting a short time, or for a day or two, normal vision usually returns. Treatment is that of uræmia.

Diabetic Retinitis.

This form occurs as a late manifestation of glycosuria in elderly persons, but is not common. The ophthalmoscopic appearances (Fig. 226, Plate XVIII.) resemble those of albuminuric retinitis in some cases, but in others they are characteristic: small, bright, white spots in and around the macular region, grouped irregularly and not in the form of a stellate figure; sometimes larger spots; numerous punctate or larger hæmorrhages. There is no swelling of the optic nerve and retina. In rare instances in young diabetics nearing a fatal termination there is a striking picture of the fundus known as retinal lipæmia: the retinal vessels are full and have a very light red colour upon a background slightly paler than normal; this is due to excessive fat in the blood. The prognosis depends upon the systemic condition. The treatment is that of diabetes.

Leukæmic Retinitis.

This variety presents marked swelling of the retina and disc, and numerous hæmorrhages. The bloodvessels are greatly dilated and extremely tortuous, and the blood is very

PLATE XVIII.




FIG. 225.--DIABETIC RETINITIS.




FIG. 227.--THE FUNDUS IN AMAUROTIC FAMILY
IDIOCY.

To face page 280.

pale. The entire fundus is pale, and has a yellowish hue. There are white and yellow spots of exudation, and some of these present a pink border. They consist of white blood corpuscles surrounded by red blood cells.

Syphilitic Retinitis.

A common form of retinitis found in both hereditary and acquired syphilis (Fig. 171, Plate XV.). In acquired syphilis it occurs in the secondary stage, during the first or second year, and usually involves both eyes. It is generally associated with choroiditis and often with iritis.

Ophthalmoscopic Signs.—In acquired syphilis there may be indistinctness of the fundus due to swelling of the retina and disc, and to fine, dust-like opacities of the posterior portion of the vitreous. These opacities cause the disc to appear red and hazy; scattered grayish or white spots often fringed with pigment, especially in the macular region, but also peripherally; circumscribed white exudations along the large bloodvessels, forming white lines. Later, deposits of pigment may be so pronounced as to resemble somewhat the picture of retinitis pigmentosa. There may be one or more large white exudates, leaving later an atrophic area with pigmented edges.

In hereditary syphilis general gray discoloration, spots of black pigment and atrophied areas are common, especially towards the periphery.

Subjective Symptoms consist of more or less diminution in the acuteness of vision, diminution in the light sense, night-blindness, annoying flashes of light, distortion, and changes in size of objects, central and ring scotomata, and, later, contraction of the field of vision.

Course and Prognosis.—The progress is slow, and relapses are common. The prognosis depends upon the stage during which treatment is begun. If begun early and carried out vigorously, the prognosis is good, though some impairment of vision usually remains. Neglected cases are often followed by disseminated choroiditis, pigmentary degeneration of the retina, and optic-nerve atrophy.

Treatment consists in the thorough use of mercury by inunctions, and afterwards iodide of potassium, rest of the eyes, protection from light, and atropine. Salvarsan may be tried.

Hæmorrhagic Retinitis.

In this form of inflammation, numerous and recurrent hæmorrhages are added to the other signs of retinitis. The extravasations of blood, both flame-shaped (superficial) and round and irregular (deep), are usually scattered all over the fundus. This affection usually occurs in elderly individuals as a result of diseases of the heart and bloodvessels. It may be monocular or bilateral. The prognosis is unfavourable. New hæmorrhages are apt to be added to the residua of the old ones. Sometimes the affection terminates in hæmorrhagic glaucoma. It is often a forerunner of cerebral hæmorrhage. Treatment consists in rest for the eyes, smoked glasses, sometimes local abstraction of blood, and ergot. Constitutional treatment is of the greatest importance in the endeavour to guard against hæmorrhage in other parts of the body.

Septic Retinitis.

This affection, also known as metastatic retinitis, results from the lodgment of septic emboli in the retinal arteries in the course of puerperal and other forms of septicæmia and pyæmia, and also from infected wounds and foreign bodies. In the first stage there are small white spots and hæmorrhages around the disc and in the macular region. Very soon, however, the uveal tract is invaded and the signs of purulent choroiditis (p. 198) appear. The inflammation ends in panophthalmitis or in degeneration of the eyeball without perforation (pseudo-glioma). Non-infected embolus gives rise to characteristic retinal changes (p. 286).

Uncommon Forms of Retinal Changes.

A number of pathological conditions are found in the retina, which, though uncommon, have received names describing the clinical picture in each case. Though labelled 'retinitis,'

they are not inflammatory in origin, but due to circulatory changes. Among these are retinitis circinata (presenting a crescentic or annular figure formed of white patches surrounding the macula; probably caused by previous deep hæmorrhages and seen mostly in elderly women); angoid streaks (pigmented striæ resembling a system of obliterated bloodvessels); striate retinitis (yellowish or grayish streaks radiating from the disc to the periphery; representing either former hæmorrhages or cured retinal detachment); punctate retinitis (numerous small, white, or yellowish scattered spots); proliferating retinitis (dense vascularized masses of connective tissue which project from the retina into the vitreous and due to organization of a vitreous hæmorrhage; one form, Eales' disease, occurs in young people as the result of recurrent hæmorrhages and is thought to be tuberculous in origin); exudative retinitis (Coats' Disease) with large raised yellowish-white area from cicatrization of old hæmorrhages into deeper layers of the retina.

Retinal Changes due to Excessive Light are seen after injurious exposure of the eye (1) to the sun, especially in watching an eclipse with insufficient protection; (2) to the electric light, as in electric welding; and (3) to sunlight reflected from snow (snow blindness). There are pigment changes at the macula, and, corresponding to this, a central, positive scotoma which may become less marked, but does not disappear entirely. The conjunctivitis which results from exposure to excessive light is described on p. 105.

Symmetrical Changes at the Macula in Infancy (amaurotic family idiocy).—This condition presents a clinical picture (Fig. 227, Plate XVIII.) which somewhat resembles that of embolism of the central artery: a red spot at the macula surrounded by a grayish-white zone about twice the size of the disc. This is followed by optic-nerve atrophy. The disease occurs in infants who present general muscular and mental weakness. There is gradual loss of sight. Death results in a year or two. It attacks several children of the same parents. Nearly all recorded cases have been of Jewish parentage.

Contusion of the Retina (œdema of the retina) is a tran-

sitory clouding of the retina resulting from contusion of the eyeball. It causes some diminution in acuteness of vision, which frequently disappears, with the retinal change, in a few days. Sometimes, however, later degenerative changes occur in the macular area or elsewhere leading to some permanent defect in vision. Such changes may be evident to the ophthalmoscope either as a fine pigment stippling or as a permanent deepening in colour of the macula with some thickening of surrounding retina giving rise to the appearance of a 'hole at the macula.' This latter appearance is sometimes found in cases of arterio-sclerosis of retinal vessels with anæmia of the retina.

Circulatory Disturbances of the Retina.

eræmia of the Retina, when slight, is recognized by increased redness of the disc and by slight striation of its margins. Such a condition is often found in persons suffering from the effects of errors of refraction (asthenopia), and in those whose vocations expose the eye to excessive light or heat. Marked arterial hyperæmia is an accompaniment of inflammation of the retina and of surrounding ocular structures. Venous hyperæmia is seen as a result of pressure, in certain general diseases (especially heart disease), and in a most pronounced form in thrombosis of the central vein.

Anæmia of the Retina may be merely the ocular expression of a general condition, or it may be local. The latter form may be acute or chronic. Acute anæmia is known as ischæmia of the retina. It may result from occlusion (embolism of the central artery), compression, or spasm of the retinal arteries. There is extreme narrowing of the retinal arteries, pallor of the disc, and blindness. Such a condition is observed in cholera, and temporarily in migraine. Quinine poisoning furnishes an example of ischæmia in which some diminution in acuteness of vision and some contraction of the field of vision are permanent. The chronic form of anæmia is frequently seen after retinal disease causing atrophy. Here

the bloodvessels become narrower or even changed into slender, empty threads.

Hæmorrhages in the Retina (retinal hæmorrhages, Fig. 228, Plate XIX.) often occur without any signs of inflammation.

Objective Signs.—The hæmorrhages vary in size, shape, and position. They are found most frequently in the neighbourhood of the larger bloodvessels. When situated in the nerve-fibre layer, they have a striate or flame-shaped form; when deep, they are rounded or irregular in outline. Sometimes a large round extravasation is seen in the region of the macula, between the retina and vitreous. This is known as a subhyaloid hæmorrhage. Retinal hæmorrhages become absorbed slowly. The smaller ones may leave no traces. But more frequently white spots, sometimes pigmented, indicate their site.

Subjective Symptoms.—The interference with vision depends upon the size and particularly the situation of the hæmorrhage. If at the macula, vision is much diminished. A scotoma results if the retinal tissue has been injured. Subhyaloid hæmorrhage may cause no permanent change in vision after absorption, since the retina is not involved.

Etiology.—The *causes* of retinal hæmorrhages are: (1) Injuries. (2) Local disease of the vessels of the retina and choroid. (3) Diseased state of the bloodvessels, especially atheroma. This condition is commonly associated with heart and kidney disease, frequently found in old persons, and is often a warning of cerebral apoplexy. (4) Disturbances in the circulation, causing retinal embolism, thrombosis, hæmorrhages in the new-born, and after operations. (5) Valvular heart disease and cardiac hypertrophy. (6) Changes in the composition of the blood and in the walls of the bloodvessels, seen in anæmia, leukæmia, purpura, scurvy, pyæmia, and septicæmia, the malarial fevers, poisons, etc.

Treatment.—Management of the etiological factor is indicated. Locally, there is no treatment.

Changes in the Fundus in Arterio-Sclerosis are of importance in general prognosis, since the finding of such indicates similar

lesions in other parts of the body, especially the brain. Ophthalmological evidence may be the first to reveal the existence of this serious vascular lesion. The fundus may present any or all of the following changes (Fig. 229, Plate XIX.): Increased tortuosity and beaded appearance of the bloodvessels; greater opacity of the arteries and widening of the central light-streak; apparent interruption of continuity in the veins where they are crossed by arteries, and dilatation just beyond these points; white lines along the course of vessels (peri-vasculitis); retinal œdema near disc, along bloodvessels, or scattered in spots; hæmorrhages.

Obstruction of the Central Artery.—Plugging of the central artery of the retina by a non-infected embolus, or by a thrombosis due to arterio-sclerosis, causes *sudden blindness*, which is sometimes unrecognized by the patient, because it is usually unilateral and there is no pain. The left eye is the one generally affected.

Symptoms.—There are no external signs, but the ophthalmoscopic picture is very characteristic. Within a few hours the *fundus* becomes *pale* and *œdematous*, *grayish*, or even milky. This is most pronounced near the disc and macula, and fades out toward the periphery. In the situation of the fovea there is a bright *cherry-red spot*, which stands out in marked contrast to the neighbouring grayish-white retina. This is due to the red colour of the choroid seen through the very thin retina opposite this area. The *arteries are very thin*, and can be followed only a short distance from the disc. Beyond this point they may be lost entirely. The veins also contain less than the normal amount of blood. There may be small *hæmorrhages*. Pressure upon the eyeball does not produce arterial pulsation, but gives rise to the appearance of *broken columns* of blood with clear spaces between them.

There is sudden and complete blindness; even perception of light is lost. Occasionally good central vision is preserved. This is due to the existence of a small macular branch given off from the central artery below its bifurcation, where the embolus usually lodges, or to the presence of a cilio-retinal artery (anastomosis between retinal and ciliary systems);

PLATE XIX.




FIG. 228.—HÆMORRHAGES IN THE RETINA.



FIG. 229.—ARTERIO-SCLEROSIS.

but even in such exceptional cases most of the field is lost.

After embolism of the trunk of the central artery has lasted a few days, degeneration of the retina occurs, and after a few weeks atrophy sets in. The œdema subsides, the disc atrophies, and the bloodvessels become shrunken or are represented by white lines.

The foregoing description applies to cases in which the main trunk of the central artery is occluded. The embolus may, however, lodge in one of the branches of the central artery. In such cases the interference with sight and the changes in the background will be limited to the area supplied by the occluded branch. Occasionally the embolus can be seen, but usually its presence is shown by a swelling in the artery, beyond which the vessel is thin or obliterated.

Etiology.—Plugging of the central artery is most frequently due to valvular heart disease, less often to atheroma, aneurism, Bright's disease, and pregnancy. A thrombus of the central artery may give rise to the same signs and symptoms as embolism, and a differential diagnosis is difficult or impossible. In the rare instances in which the patient has recovered good vision, the blocking has probably resulted from temporary spasm of the walls of the artery.

Treatment is rarely effective. If the case is seen early, paracentesis of the cornea, massage of the eyeball, and inhalations of amyl nitrite have been employed for the purpose of driving the plug along into one of the smaller branches, where it will give rise to less serious results. In a few cases such treatment has been beneficial.

Thrombosis of the Central Vein (Fig. 23r, Plate XX.) may occur in old persons with atheroma and cardiac disease; it also follows cellulitis of the orbit; it may occur in younger people from thrombo-phlebitis of the vessel wall. It is one of the causes of hæmorrhagic retinitis. It may be complete or partial. There is diminution of vision, either corresponding to the entire field, or, if only a branch is affected, to the part of the retina supplied by it. The veins are greatly engorged and tortuous, the arteries very small; there are numerous

large hæmorrhages, and indistinctness of the margins of the disc.

The prognosis is bad, especially in cases with arterio-sclerosis, and vision is usually permanently affected, from atrophy of the optic nerve. But in younger patients the outlook, after some months have elapsed, may be much better than is usually supposed. In affections of a branch (Fig. 230) complete recovery is much more likely. A frequent complication is subsequent rise of tension, from the blocking of filtration angle by a gummy albuminous exudate.

Treatment.—In early stages complete rest; removal of septic foci; use of miotics. If tension rises, a trephine operation is indicated if any vision remains, otherwise enucleation for relief of pain.

Pigmentary Degeneration of the Retina, or Retinitis Pigmentosa.

A chronic form of retinitis, which has a constant tendency to become worse, and which consists of atrophy of the retina with migration of pigment from the pigment epithelium into the inner layers.

Symptoms.—*Night blindness* (hemeralopia), *concentric contraction of the field* of vision, *progressive diminution in sight*, terminating in advanced years in *complete blindness*.

In early life there is but slight reduction in the extent of the field with good illumination, and central vision is often perfect. But with feeble illumination the peripheral parts of the retina do not react, and on this account the patient cannot find his way about at night, because the field is small. With increasing years the field becomes contracted even with good illumination. Finally, in advanced life central vision becomes poor, and gradually complete blindness follows.

✓ The field of vision has been found to fail in a characteristic way. A ring scotoma, incomplete at first, develops between the 40° and 10° circles; the failure of field then progresses in the following order: (1) extreme lower part; (2) upper and outer

PLATE XX.




FIG. 230.—THROMBOSIS OF BRANCH OF CENTRAL
RETINAL VEIN.




FIG. 231.—COMPLETE THROMBOSIS OF CENTRAL
RETINAL VEIN.

portions; (3) lower and outer parts; (4) nasal half; and finally (5) the central area.

Ophthalmoscopic Examination (Fig. 232, Plate XXI.) shows *black spots in the periphery of the fundus*. These have the shape of *branching cells, like corpuscles with connecting bone processes*, and are found especially *along the bloodvessels*. In the course of years new spots form, and in this way the pigment circle gradually approaches the disc. Migration of colouring matter from the pigment layer of the retina allows the choroidal vessels to become plainly visible. The *disc and retina are atrophied*. The disc has a yellowish, waxy appearance. The retinal *arteries* are very *small*, and in the periphery are represented by mere threads.

There are cases of retinitis pigmentosa in which all the symptoms of this disease are present, and the ophthalmoscope shows all changes except the presence of pigment, and others in which the pigment is distributed in an atypical manner.

Syphilitic choroido-retinitis may give a picture similar to that of retinitis pigmentosa, but may be differentiated by the patches of choroidal atrophy.

Occurrence.—The disease affects both eyes. It is either congenital or develops in childhood. It is hereditary, and is often found in the offspring of consanguineous marriages. Not infrequently other congenital defects, such as deafness and defective intelligence, are present. It may be complicated with posterior polar cataract and other ocular anomalies.

Pathology.—This chronic, progressive degeneration of the retina is thought by some to start with degeneration of the chorio capillaris, depriving the outer layers of the retina of their nutrition, by others to be primarily a change in the pigment epithelium with migration of pigment cells into the retina and around the bloodvessels. The rods and cones degenerate and are replaced by neuroglia formed by extension, increase and hypertrophy of Mueller's fibres. With the progress of connective tissue and pigment invasion, the retina becomes atrophic and the ganglion cells are destroyed with degeneration of their axis cylinders and of the optic nerve; so that in the advanced stage there is disappearance of all nervous elements

and replacement with glial tissue embedded in which are masses of pigment; the retina is then adherent to the choroid. The bloodvessels of the choroid and of the retina suffer hyaline and endovascular obliterating changes. All these changes begin at or near the equator and then spread anteriorly and posteriorly, the macular region becoming affected only late in the disease.

Treatment is of no avail.

Retinitis Punctata Albescens is a rare disease, having all the subjective symptoms of retinitis pigmentosa. There is no pigmentation, but small white spots are scattered all over the fundus. It is slowly progressive, and incurable.

Detachment of the Retina.

Retinal detachment is a separation of the retina from the choroid. The name usually refers to a separation by serum, but detachment may also occur as a result of subretinal hemorrhage, exudation, or tumour.

Symptoms.—There is more or less complete loss of vision in that part of the field which is opposite to the detachment, and the appearance of a dark cloud before the eye. Early symptoms are metamorphopsia and flashes of light (photopsia). Central vision is preserved as long as the macula is not included.

Ophthalmoscopic Signs depend upon the degree and extent of detachment. When the detachment is flat, the retina appears but slightly changed. It is somewhat cloudy, and its vessels are dark and tortuous. The variation in level of the affected portion can be recognized by the difference in the refraction of a bloodvessel on the separated part. When the detachment is steep, as is generally the case, it is usually found near the periphery. It is at first limited in extent. It may commence at any part of the retina, but as a result of sinking of the subretinal fluid it is usually found below. It tends to enlarge and become total, then involving the entire retina. It presents a collection of grayish, bluish-gray, or greenish folds (Fig. 233, Plate XXI.) with white tops pro-

PLATE XXI.




FIG. 232.—RETINITIS PIGMENTOSA.




FIG. 233.—DETACHMENT OF RETINA.

To face page 290.

ecting a variable distance into the vitreous and shaking with movements of the eye. The bloodvessels pass over and follow these folds, and are therefore very tortuous, and hidden in places. They appear prominent, and of a dark red, almost black, colour. In most cases a hole, a rent (Plate XXII.), or a peripheral tear (disinsertion) can be found in the separated retina. The commonest site for the former is in the upper and outer quadrant; for disinsertion, the lower and outer quadrant. In the later stages opacities of the vitreous and cataract are often added. The rest of the fundus presents a normal picture. Externally the eye appears normal, but tension is usually lowered and the anterior chamber deepened.

✓ Etiology.—Serous detachment may be due to disease or injury. It may occur in the course of nephritis, and may complicate the toxæmia of pregnancy; in such cases it is preceded by a sub-retinal œdema, the source of which is the choriocapillaris.

When due to disease, it is generally found in myopia of medium degree—*i.e.*, 6 D. to 12 D. (less commonly in the more extreme degrees)—and after disease of the vitreous, iridocyclitis, and irido-choroiditis. In such cases the condition probably results from shrinking of the vitreous, which thus pulls the retina from its attachment to the choroid.

Traumatic detachment is usually the result of a blow. It may follow an accidental or operative wound, especially when there has been loss of vitreous.

Considerable stress is now laid on the occurrence of holes or tears in the retina, which are held to be the main factor in causing persistence of the detachment by allowing the fluid part of the vitreous to circulate freely beneath the retina. Such holes tend to occur in parts of the retina previously damaged by disease or injury.

Diagnosis is readily made, especially with the ophthalmoscope at a distance of a foot with an 8 D. or 10 D. convex lens in the sight-hole, but it is sometimes difficult to decide whether the detachment is serous or due to a tumour of the choroid (p. 208).

✓ Prognosis, which was formerly considered extremely un-

favourable, has become much more hopeful since the introduction of operations designed to close the holes and tears and to reattach the retina to the choroid by means of the production of an area of adhesive inflammation at the seat of its detachment. During the past few years from 30 to 40 per cent. of cases so treated have been reported as cured. Early operative treatment is essential, since the separated retina tends to atrophy and the detachment to become complete. The chances of reattachment after operation are best in recent cases and in those resulting from traumatism; they are less favourable in high myopia, and poor in cases of aphakia and old iridocyclitis; in old cases (over two years) there is no chance of success. Unless the retina can be reattached by operation, or if untreated, the detachment generally extends and becomes complete and blindness results, though rare cases of spontaneous reattachment have been reported. The outlook is much better when detachment is a complication of nephritis, and especially when it occurs with the toxæmia of pregnancy; in the latter case reattachment usually takes place spontaneously.

Treatment.—Operation is indicated in favourable cases and is the only means of reattaching the retina. Formerly, non-operative treatment was resorted to in recent cases, and consisted of absolute rest in bed in the supine position, instillation of atropine, a firm binocular bandage, iodides and sweating; this ordeal was continued for at least a month, but was rarely successful; equally unsuccessful were posterior sclerotomy, sub-conjunctival injection of salt solution, and trephining the sclera over the seat of detachment with aspiration of the sub-retinal fluid.

All operations now aim at creating a line of traumatic adhesive choroiditis around the hole or tear in the retina; when the sub-retinal fluid is allowed to escape, the retina falls back into position and reattaches itself to the choroid at these points.

In recent cases the patient should be put to bed at once, atropine instilled, and a careful search made for holes and tears. If these are found, their position should be localized.

PLATE XXII.




FIG. 234.—DETACHMENT OF RETINA, SHOWING
LARGE TEAR.



FIG. 235.—SAME FUNDUS AFTER DIATHERMIC
OPERATION.

The limit of ophthalmoscopic examination extends to a point just behind the ora serrata, and in normal eyes is taken as being 8 mm. behind the limbus; in highly myopic eyes it is calculated as 9 mm. The distance of the hole, or innermost part of the tear, from the limit of examination is estimated in disc diameters (*i.e.*, 1.5 mm.); to find its distance from the limbus 8 mm. (or 9 mm.) must be added. Thus a hole 3 disc diameters from the periphery is 4.5 mm. plus 8 mm., or 12.5 mm., from the limbus. The meridian in which it lies is given by a clock-like notation.

In the first of the recently introduced operations (Gonin's), the sclera was exposed and punctured opposite the hole with a small actual cautery, which was later replaced by the electric cautery, the purpose being to sear the edges of the hole and thus cause adhesion. This operation, though very successful in many cases, has the disadvantage that it is not always easy to locate holes and tears very accurately, and that there is a tendency for the scarred retina to develop secondary tears. There is also difficulty when the tear is large or the holes multiple.

Multiple trephining of the sclera around the hole or tear was then introduced by Guist, followed by the touching of each trephine hole with the tip of a tiny stick of caustic potash, immediately neutralized with $\frac{1}{2}$ per cent. acetic acid; the sub-retinal fluid was then evacuated by puncturing three or more holes with a punctum dilator. Very successful results can be obtained by this method, but it is very tedious.

The operative method now favoured is that of diathermic coagulation, using very weak currents. Two methods are employed, Larsson's and Safar's. Larsson uses a blunt electrode to pass the current (50 to 60 milliampères) through the exposed sclera at multiple points round the hole or tear. To prevent charring of sclera and diffusion of current, the surface is wiped dry of blood and then moistened with distilled water. With the newer, more rigid, electrodes this moistening has been found unnecessary. Each application is for 5 seconds. Several similar applications are then made in the area thus delimited, and the sub-retinal fluid then evacuated either by trephin-

ing over the hole or puncturing with an electric cautery. Safar uses fine needle-pointed electrodes which are passed through the dry sclera into the sub-retinal space, either single needles $1\frac{1}{2}$ mm. long or else a number of needles attached to a small plate. Each needle point requires 30 milliampères of current, which is applied for 1 to 2 seconds. When the points are withdrawn the sub-retinal fluid escapes through the minute openings. These two methods are sometimes combined as in Weve's operation. The conjunctiva is replaced over the exposed sclera and sutured; atropine is instilled; the patient is put to bed and kept quiet for two weeks with both eyes bandaged; he is then allowed special spectacles with a small central opening only, designed to compel him to keep the eyes in one position; after a week of this he is allowed out of bed, and wears the glasses for another week, making a month's post-operative treatment in all.

CHAPTER XX

DISEASES OF THE OPTIC NERVE

Anatomy.—The optic nerve may be divided into (1) an intra-ocular portion, the retina; (2) an orbital portion extending from the eyeball to the optic foramen; and (3) an intracranial portion situated between the optic foramen and the chiasm.

The intracranial portion of the optic nerve is short and flattened. The optic foramen forms an unyielding ring, which serves to compress the nerve in inflammation and injury.

The nerve pierces the sclera and choroid a little to the inner side of the posterior pole of the eyeball. At this point the outer layers of the sclera become continuous with the sheaths of the nerve, while the inner layers, together with the modified choroid, stretch across the foramen, presenting numerous openings for the passage of the separate bundles of the optic nerve. This sievelike arrangement is known as the lamina cribrosa. Here the nerve fibres lose their medullary layer and become transparent. Spreading apart before reaching the level of the retina, they leave a funnel-shaped depression at the middle of the disc (Fig. 43), the physiological excavation.

The lamina cribrosa represents the weakest portion of the layers of the eyeball, and in increased tension is the first to recede. It surrounds the bundles of the optic nerve with fibrous rings of connective tissue, which serve as constricting bands when swelling occurs.

The orbital portion of the optic nerve presents a sigmoid curve, permitting free movement of the eyeball. The nerve consists of bundles of nerve fibres separated by connective-tissue septa; between these there are lymph spaces. The optic nerve is surrounded by three sheaths originating from the three envelopes of the brain, and known as the pial, arachnoid, and dural sheaths. Between the pial and the dural sheaths is a space, the intervaginal space, divided into two parts by the arachnoid sheath. The two spaces thus formed are lymph spaces. They are lined by endothelium, and communicate with the corresponding cerebral spaces. Anteriorly, the intervaginal space ends in a blind extremity, and the sheaths unite with the sclera.

A short distance from the eyeball the central artery (a branch of the ophthalmic) enters, and the central vein emerges. The latter empties into the superior ophthalmic vein or directly into the cavernous sinus.

Affections of the Optic Nerve comprise (1) hyperæmia, (2) inflammation, (3) atrophy, and (4) tumours (very rare).

Hyperæmia, or Congestion of the Optic Disc.

The normal disc varies greatly in colour; hence it is often difficult to decide whether the papilla is congested or not. When congestion exists it shows itself in increased redness, due to capillary injection, slight blurring and striation of the margins of the disc, often limited to a portion of the circumference, and some fulness of the veins.

Such a picture is frequently presented in eye-strain from hypermetropia and astigmatism, excessive use of the eyes, or after work with insufficient or excessive light. It is also found with inflammations of the deeper portions of the eyeball. It may be the incipient stage of optic neuritis.

Inflammation of the Optic Nerve, or optic neuritis, is divided into:

1. *Papillitis*, or intra-ocular optic neuritis, in which the head of the optic nerve is the part affected, and in which there are marked visible changes in the disc.

2. *Retrobulbar Neuritis* affecting the nerve fibres behind the eyeball, and in which the changes of the disc are slight or absent, and the existence of inflammation is often inferred from subjective symptoms.

Papillitis, Intra-ocular Optic Neuritis, or Choked Disc.

Symptoms.—There is more or less disturbance of vision. It is usually considerable, but it is not always proportionate to the severity of the inflammation as revealed by the ophthalmoscope. There may be complete blindness. The field of vision is usually contracted peripherally, especially for colours. There may be hemiopia or scotomata. There is no pain, and there are no external signs.

Ophthalmoscopic Signs.—The papilla appears swollen and

projecting (Fig. 236), enlarged, of whitish or grayish colour, striated, and often presenting white spots and hæmorrhages. Its situation is recognized only by the convergence of the retinal bloodvessels, its margins having become indistinguishable, and extending gradually into the surrounding retina. The retinal vessels are altered, and are interrupted in places. The arteries are either thin or of normal calibre; the veins are greatly distended and exceedingly tortuous. The surrounding retina is usually œdematous, congested, and presents white patches and hæmorrhages. When a large portion of the surrounding retina is involved, the affection constitutes neuroretinitis (Plate XXIII.).

Clinical Forms.—Clinically we may distinguish between two types of papillitis: (1) Choked disc, in which there is marked swelling limited pretty sharply to the disc, with extreme dilatation and tortuosity of the veins. (Edema and engorgement are the predominating features. (2) Descending neuritis, in which there are less swelling and projection of the disc, less venous fulness and tortuosity, but more exudation and considerable extension of the latter into the surrounding retina. The picture in these cases points more to inflammation. No sharp line can, however, be drawn between these two forms, either from the standpoint of pathology or of etiology, and transition forms occur frequently.

Course.—Though occasionally acute, the course is usually a chronic one, extending over many months. It is possible for the changes to subside and the disc to regain its normal appearance, with the preservation of good sight (especially in syphilitic cases). But, as a rule, papillitis is followed by




FIG. 236.—PAPILLITIS
(CHOKED DISC).

The upper portion represents the ophthalmoscopic appearances; the lower half a longitudinal section.

post-neuritic atrophy. The disc becomes white or grayish-white, its margins are sharply defined but irregular, and surrounded by changes in the choroid, while the exudation becomes changed into connective tissue which covers the lamina cribrosa and fills up the physiological cup. The arteries are contracted and frequently bordered by white lines, but the veins remain dilated and tortuous (Fig. 241, Plate XXIV.).

Prognosis is always serious. The degree of atrophy depends upon the intensity of the preceding neuritis, and determines whether vision finally becomes useful, much impaired, or totally lost.

Etiology.—Papillitis is almost always bilateral, but one eye may be affected before the other. The causes are: (1) Diseases of the brain and its envelopes; (2) syphilis; (3) general diseases; (4) anæmia, either simple or the acute form, due to great loss of blood; (5) diseases of menstruation, pregnancy, and lactation; (6) lead-poisoning; (7) heredity; (8) idiopathic cases (when no cause can be found); and (9) orbital and periorbital affections.

Brain Tumour is the most frequent cause. Papillitis occurs in 90 per cent. of such cases, and then most commonly assumes the choked-disc type. Sometimes the neuritis is the first symptom of brain tumour. The occurrence or degree of papillitis does not depend upon the size or the situation of the tumour. It frequently exists with cerebellar tumours. The neuritis of brain tumour occasionally gives a picture resembling that of albuminuric retinitis with its star-shaped figure at the macula.

Next in frequency comes meningitis, especially basilar and tuberculous. In such cases the papillitis is apt to be of the descending neuritis type. Then come abscess and hydrocephalus.

Syphilis is a frequent cause, and acts either by direct implication or through the development of specific affection in the cranial or orbital cavities.

Acute Febrile Affections (measles, scarlatina, diphtheria, typhoid, grippé) are occasional causes.

PLATE XXIII.




FIG. 237.—NEURO-RETINITIS.



FIG. 238.—CHOKED DISC.

General Affections, such as rheumatism, nephritis, and arterial disease, are sometimes responsible; also exposure to cold.

Orbital and Periorbital Affections include inflammations of the orbit, tumours of the orbit and optic nerve, and diseases of the neighbouring cavities (sphenoid, ethmoid, frontal, and maxillary). These constitute the examples of unilateral cases.

Pathology.—The process consists of an inflammatory swelling, exudation of leucocytes, venous engorgement, hæmorrhages, and distension of the intervaginal space. The exact mechanism is still unsettled. Numerous hypotheses have been advanced. The most prominent are: (1) That it is due to increased intracranial pressure forcing cerebro-spinal fluid into the intervaginal space of the optic nerve, causing stasis in the region of the lamina cribrosa, compression of the vessels resulting in venous engorgement and œdema (choked disc); (2) transmission of inflammation from the brain along the optic nerve to the papilla; (3) inflammation excited by irritating substances which pass from the cranial cavity to the optic disc.

Treatment should be directed against the cause of the inflammation. In syphilis an injection of salvarsan followed by a vigorous course of mercury and iodide of potassium. Even in non-specific cases mercury and iodide of potassium are often prescribed. Locally, rest of the eyes, and shading from light. Orbital and periorbital affections may call for operative treatment.

Cerebral Decompression is frequently undertaken to relieve the intracranial pressure responsible for choked disc, thus enabling the optic nerves to resume their functions. In most cases this operation causes a subsidence of the papillary swelling and an improvement in vision, especially if it be resorted to before much degenerative change in the nerves has taken place; incidentally other symptoms of tumour of the brain are relieved and life is prolonged. The operation is performed in the right temporal region for pre-tentorial growths; but when the tumour is

sub-tentorial, the opening is made in the sub-occipital region.

Lumbar puncture also is occasionally resorted to to relieve intracranial pressure, but it is not free from risk.

Retrobulbar Neuritis.

This affection involves the orbital portion of the optic nerve. Hence, unless the case should progress to atrophy, there may be few or no changes in the disc. In most cases there is merely an implication of the papillo-macular fibres of the optic nerve, giving rise to a central scotoma, either absolute or relative. Retrobulbar neuritis may be acute or chronic.

Acute Retrobulbar Neuritis.

This rather uncommon affection is generally unilateral, occasionally bilateral.

Symptoms.—Neuralgia or headache on the same side, pain in and about the orbit, aggravated by movements of the eye, and tenderness on pressing the eye backward into the orbit. With these symptoms there is rapid impairment of sight, progressing in the course of a week to partial or complete blindness. Externally the eye appears normal.

Ophthalmoscopic Signs.—At first there are no changes. Later there may be slight haziness of the disc, with distension and sometimes diminished calibre of the retinal vessels.

Course.—The disease runs an acute course, and after a month or two the sight usually becomes normal; or recovery may be partial, and a central scotoma remain, in which case there will be a pallor of the temporal part of the optic disc corresponding to degeneration of the papillo-macular nerve fibres. Occasionally it terminates in permanent and total blindness.

Etiology.—The causes may be general or local. Multiple sclerosis is the most common cause, the disease occurring as an early symptom in about one-half of the cases; less frequent causes are general diseases (syphilis, so-called rheumatism,

diabetes), acute infectious diseases (influenza), septic foci (mouth, intestinal tract) and poisons (alcohol, especially methyl alcohol, lead, thallium used in depilatory creams).

Local causes comprise extension from the orbit (periostitis), extension from the nasal accessory sinuses (less importance is now attached to rhinological etiology than was the case formerly), hæmorrhage into the sheath of the optic nerve, fracture of the optic foramen and pressure upon the chiasm. Sometimes no cause can be found.

Prognosis depends upon the cause and the possibility of its removal. In general the prognosis is good; in multiple sclerosis the involvement of the optic nerve rarely leads to complete blindness.

Treatment.—Although there is a strong tendency to spontaneous recovery, the responsible cause should be attacked. In addition, complete rest of the eyes, protection from bright light, diaphoresis, potassium iodide and mercury, salicylates and later strychnine.

Leber's Disease (Hereditary Optic Neuritis).—Occasionally, bilateral acute retrobulbar neuritis occurs as a hereditary affection, generally attacking males, often several in the same family, commencing about the twentieth year; the descent is commonly through unaffected females. Vision usually fails rapidly at first and then remains stationary or improves, and there may even be complete recovery. There is commonly a central scotoma, either relative or absolute, which is permanent, with little contraction of the peripheral field. The fundus presents little if any change at first; later there is pallor of the temporal segment of the disc, rarely of the whole disc. The cause of the disease is unknown; it has been ascribed to pituitary disease. No form of treatment has been of any value.

Chronic Retrobulbar Neuritis, Toxic Amblyopia, Tobacco Amblyopia.

A chronic affection of the orbital portion of the optic nerve, of frequent occurrence, usually attacking both eyes, and due in the great majority of cases to excessive indulgence in

DISEASES OF THE OPTIC NERVE

tobacco, alcohol, or both combined. Some authorities, however, regard the toxic amblyopias as definite clinical entities due to poisoning of the ganglion cells of the retina, and not to be classed as retrobulbar neuritis.

Symptoms.—There is gradual diminution in acuteness of sight; foggy vision; the patient sees better in the evening, and the visual disturbance is more marked in bright light. The field of vision presents the normal peripheral boundary, but there is a central colour scotoma for red and green, corresponding to the distribution of the papillo-macular fibres of the optic nerve. This colour defect is usually small, and more marked for green than for red.




FIG 239. — THE FIELD OF VISION IN TOXIC AMBLYOPIA (CENTRAL COLOUR SCOTOMA).

The scotoma of tobacco-poisoning is at first limited to the area between Blind Spot and fixation point, and only in the later stages does it extend to the nasal side of fixation point.

That due to diabetes is usually very small and central.

Ophthalmoscopic Signs.—Sometimes there are no changes in the papilla, or merely a slight haze with increased redness. At a later period there is very often a pallor of the temporal side of the disc.

Course.—The course of the disease is slow. If poisoning continues, vision becomes more impaired, and may be reduced very much. If the cause be removed, there is usually gradual improvement, and sight is often restored to the normal. But in severe cases there may be some permanent reduction in the acuteness of vision, and the relative scotoma may continue indefinitely. Even though a man may have smoked heavily with impunity for many years, after one attack a small quantity of tobacco is liable to cause a relapse. A second attack is seldom completely recovered from.

Etiology.—The condition results most frequently, from over-indulgence in tobacco, whether in smoking or chewing, occasionally after snuff-taking. The stronger tobaccos used in cigars and pipes are the forms which are most frequently

responsible. Certain individuals are more susceptible than others. Impairment of the general health predisposes, as does also the practice of smoking when the stomach is empty. It occurs almost exclusively in middle-aged or elderly men. In most cases both alcohol and tobacco act together. It is sometimes an early symptom of disseminated sclerosis, though the acute form is the more common in the early stages of this disease. Other poisons which in toxic doses cause a condition resembling tobacco amblyopia are iodoform, lead, arsenic, the poison of diabetes, wood-alcohol, bisulphide of carbon, and nitro-benzol.

Pathology.—The process consists of an interstitial neuritis of the papillo-macular (axial) fibres of the optic nerve, with subsequent degeneration of these fibres and the ganglion cells in the macular region.

Treatment of tobacco amblyopia consists in total and permanent abstinence from tobacco. If some stimulant is required, it must be restricted to a small amount with the principal meals. The general health should be improved. Strychnine may be given by mouth or hypodermically. Iodide of potassium in small doses (gr. ii. to gr. v.) is believed to assist in eliminating the poison.

In other cases the exciting cause must be determined and dealt with.

Atrophy of the Optic Nerve.

This affection occurs either as a primary disease (simple, primary, non-inflammatory, or progressive atrophy) or secondary to some other affection of the nerve or retina (post-neuritic, secondary, or inflammatory atrophy).

Symptoms.—There are reduction in the acuteness of vision, concentric or irregular contraction of the field (Fig. 240), first for colours and then for form, diminution in the light sense, sometimes scotomata, and colour-blindness (first for green, then for red, then for blue). These symptoms tend to progress, and end in complete blindness.

In cases due to pressure on the chiasma and nerve due to pituitary tumour, there is typically a bitemporal contraction

of the field of vision, and frequently a bitemporal paracentral scotoma (see also p. 451).

Ophthalmoscopic Signs differ somewhat in the simple and post-neuritic forms:

In simple atrophy (Fig. 241, Plate XXIV.) the disc is white, grayish, or bluish-white, its edges are sharply defined and regular, its size is somewhat diminished, and it presents a saucer-shaped excavation (Fig. 178); the lamina cribrosa is often seen very plainly. The minute vessels of the disc have disappeared. The retinal vessels may appear normal, but the arteries are usually diminished in calibre.




FIG. 240.—MARKED CONCENTRIC CONTRACTIONS OF THE FIELD OF VISION IN OPTIC-NERVE ATROPHY.

In post-neuritic atrophy (Fig. 242, Plate XXIV.) the disc is covered by connective tissue resulting from the

previous neuritis. It has a dense white or grayish colour, with more or less irregularity and obscuration of the margins, its minute vessels have disappeared, and the lamina cribrosa is hidden by the organized exudation. The retinal arteries are narrow, enclosed in white lines, and the veins are usually diminished in size and tortuous.

In the secondary atrophy following retinitis pigmentosa the disc has a dirty, grayish-red or yellow, waxy look (Fig. 232, Plate XXI.), the vessels are exceedingly narrow, and many disappear entirely.

After a time the differences in the appearances of simple and post-neuritic atrophy become much less marked.

It should be borne in mind that the disc varies in colour in health, and may appear atrophied as the result of congenital or senile peculiarities, although vision is normal and the field perfect. Hence in many cases the diagnosis cannot be made from the ophthalmoscopic signs alone, especially when these signs are not pronounced.

Etiology.—Simple atrophy is frequently due to spinal diseases, especially locomotor ataxia. It develops in one-

PLATE XXIV.




FIG. 241.—PRIMARY, OR SIMPLE, OPTIC.
NERVE ATROPHY.




FIG. 242.—POST-NEURITIC ATROPHY.

third of the cases of this affection, and is an early symptom. It is common also in affections of the brain: disseminated sclerosis, general paralysis of the insane, and tumours, especially tumours of the pituitary body. It is also due to syphilis, malaria, diabetes, acromegaly, impaired nutrition, and certain poisons. Occasionally it is hereditary, and in some cases no cause can be found. It occurs chiefly in middle life.

Secondary atrophy follows papillitis, retrobulbar neuritis, retinitis, pigmentary degeneration of the retina, embolism of the central artery, and glaucoma. It may result from penetrating wounds of the optic nerve or injury to the nerve, due to fracture of the orbital canal, following a blow or other violence. In such cases the atrophy does not show itself for a number of weeks, though more or less reduction of vision and contraction of the field results immediately.

Pathology.—The process consists of increase in the interstitial connective tissue, with atrophy of the nerve fibres. In the secondary form the periphery of the nerve is changed as above described, but the centre of the nerve is less affected.

Prognosis is usually unfavourable. Simple atrophy generally progresses to absolute blindness. In secondary atrophy the prognosis is better, and depends upon how much sight has been destroyed by the antecedent inflammation.

Treatment consists in attempting to control the cause of the atrophy. For the atrophy itself very little can be done. Potassium iodide, strychnine, mercury, nitro-glycerine, and galvanism are the remedies most frequently employed. Thyroid extract may be tried in pituitary cases, or operation may be necessary; in some cases this is very successful.

Treatment of Syphilitic Primary Optic Nerve Atrophy.—Intravenous injections of arsphenamine, bismuth, mercury and the iodides in any combinations are unsuccessful. But subdural treatment, with arsphenaminized serum, neoarsphenamine, or bichloride of mercury dissolved in salt solution or spinal fluid, mercurialized serum, or air, injected intraspinally or intracisternally, has been found of some value; results are encouraging, but not conclusive.

Tryparsamide is absolutely contra-indicated in the treatment of such cases.

CHAPTER XXI

AMBLYOPIA AND FUNCTIONAL DISEASES OF THE RETINA

AMBLYOPIA is a reduction in the acuteness of vision which cannot be relieved by glasses, and which is not dependent upon any visible changes in the eye. The term is sometimes used in a less restricted sense to designate poor sight, even when some changes are found in the eye, as, for instance, toxic amblyopia, in which temporal pallor of the disc exists.

Amaurosis is the name applied to absolute blindness when unaccompanied by discoverable ocular changes; the use of this term has, however, been extended so as to include all cases of absolute blindness, including those which show ophthalmoscopic or external changes.

Congenital Amblyopia and Amblyopia ex Anopsia.

Congenitally defective vision; nearly always affects one eye, only rarely both. It is nearly always associated with a high degree of astigmatism. There is frequently hypermetropia or myopia also. Probably in most of the so-called congenital cases the amblyopia is really acquired. The errors of refraction have prevented perfect images from being focussed on the retina, and this lack of training has caused amblyopia. The most careful correction of the error of refraction fails to produce normal vision; in young patients, however, the sight can frequently be improved or brought up to the normal after suitable glasses have been worn for a time.

Any interference with vision dating from early life, which prevents perfect focussing upon the retina, causes amblyopia from non-use (amblyopia ex anopsia); hence the advisability

of operating upon congenital and infantile cataracts early. An obstacle to vision beginning after the age of seven or eight years does not usually interfere with the functional activity of the retina.

Unilateral amblyopia predisposes to squint by lessening the value of binocular vision. Very commonly amblyopia develops in an eye which has squinted from early life on account of its exclusion from the visual act, the retinal image in this eye being suppressed (p. 411).

Exercise of such an eye during youth, by forcing it to work while the sound eye is excluded, will frequently improve its visual power.

Bilateral congenital amblyopia of high degree is nearly always associated with nystagmus.

Congenital Word Blindness is a rather rare affection, said to be more common in males than in females. It is supposed to be due to a defect in the visual memory centre for words and groups of letters. There is a difficulty or inability to read words as a whole, though the words are recognized when spelt letter by letter. Special training often results in considerable improvement.

Colour Amblyopia (Colour-Blindness).

Congenital Colour-Blindness occurs in from 3 to 4 per cent. of males, and in only 0·3 per cent. of females. It affects both eyes, and is frequently hereditary; the functions of the eye are otherwise normal. The cause and pathology are unknown, and the defect is incurable, though the colour sense can be developed, largely through the appreciation of light intensities of various colours, if training is begun sufficiently early.

The absence of all appreciation of colours is very rare as a congenital condition, though not so uncommon in acquired cases due to atrophy of the optic nerve. As a rule there is inability to differentiate one, or two, of the three fundamental colours (red, green and blue).

Light consists of impulses of various wave lengths. The longest wave which can be perceived by any human eye gives

the sensation of red, and the smallest the sensation of violet; intermediate waves give the sensation of the other colours of the spectrum—orange, yellow, green, blue, and indigo.

The various theories which have sought to explain the observed phenomena of colour vision are conjectures. The principal ones are those of Young-Helmholtz, Hering, and Edridge Green.

1. The Young-Helmholtz theory assumes that there are three sets of colour-perceiving elements in the retina, each of which, if stimulated alone, would give rise to the sensation of one of the three fundamental colours, red, green and blue, and that all other colours arise from combinations of these. With a defect of one of these primary perceptions a colour will be seen as if composed of the remaining two only. According to the colour which is deficient a patient is said to be red blind, green blind, or violet blind. The more commonly recognized forms are red blindness, green blindness and red-green blindness.

2. The Hering theory is that the colour sense depends upon chemical changes in three different visual substances in the retina—white-black, red-green and blue-yellow—by the decomposition and restoration of which substances the sensations of colour are produced; for instance, red light produces destruction in the red-green substance, and thus a sensation of red; green light causes a restoration in the same substance and thus a sensation of green. According to this theory, colour-blindness is caused by the absence of one or two of these visual substances—if one is absent the patient is either red-green (frequent) or blue-yellow (rare) blind; if two are absent nothing but the white-black substance is left, and the patient has total colour-blindness.

3. The Edridge Green theory supposes that a photograph is formed in the retina by decomposition of the visual purple in the rods, during stimulation by light. This chemically stimulates the ends of the cones, causing a visual impulse to be transmitted through the nerve fibres to the brain. It assumes that this impulse differs in quality according to the wave-length (colour) of the rays of light producing it, and that there is a special centre in the brain to distinguish these differences.

Inability to distinguish these differences causes the various varieties of colour-blindness.

He classifies the colour-blind into those who see the whole spectrum as a more or less uniformly gray colour (monochromics); those who see the two ends of the spectrum (red and violet), but do not differentiate intermediate colours (dichromics); those who see red, green and violet only (trichromics); and those who can also differentiate yellow between the red and the green (tetrachromics). Others he finds who can see five colours only in the spectrum; then come the normal who can see six or seven.

He also differentiates a group who have shortening of one or both ends of the spectrum, so that either red or violet rays cannot be seen at all and appear black.

His classification varies from that in general use, and his 'dichromics' must not be confused with those commonly so classified by adherents of the Young-Helmholtz theory.

Good colour vision is essential in some occupations, and an inability to distinguish red from green and *vice versa*, or not to see those pure red rays which best penetrate a foggy atmosphere, is especially dangerous in those occupations which involve the use of coloured signals, such as the railway and steamship services and the navy; it may also be of disadvantage to many others now that coloured traffic signals are in general use.

The tests for colour vision which may be employed are:

1. The lantern. That devised by Edridge-Green is one of the best. The colours have been specially selected and checked, and are (1) pure red; (2) red of different density; (3) yellow; (4) green; (5) signal green (blue-green); (6) blue; (7) purple. Three rotating discs make various combinations possible. The size of the light can be varied by a diaphragm to represent a signal light at various distances. The candidate should be seated in a dimly lighted room at about 20 feet from the lantern and asked to name the colour of the light produced by the coloured glasses alone or in combination, or modified by another set of glasses which represent mist, rain, or varying degrees of fog. A candidate must be rejected if he calls red

green, or green red in any circumstances; if he calls the white light red or green or *vice versa*; or if he calls red green, or white lights black, or fails to see them.

2. A simple test is Edridge-Green's Bead Test, which consists of the sorting out of various coloured beads into four compartments with a cover to each and in each cover a hole. The compartments are labelled red, yellow, green, and blue.

3. A useful portable test is the card test in which various letters or numbers are picked out in regular or irregular spots of colour on a background of confusion colours. Those available are Edridge-Green's, Stilling's and Ishihara's. Green's is very good, though some of his cards are difficult even for normal educated people, and his classification is different from that in general use. Ishihara's is simpler and more convincing in that, on some of the cards, one set of letters or figures appears to the normal eye and a totally different set to the colour-blind.

4. Wool classification and matching tests are still official tests of some public bodies, that of Holmgren being the best known. They are less reliable than other tests in that a man may be a good matcher of colours, judging them by their light intensity, and yet be so colour blind that he cannot recognize a colour with certainty unless he has another with which to compare it. Holmgren's test consists of a large assortment of coloured worsteds. The collection consists of (1) certain colours called 'test colours' (a pale green, a light pink and a bright red); (2) lighter tints and darker shades of the same colours ('match colours'); (3) confusion colours (yellow, brown, gray, drab, fawn, mauve, pale blue, etc.) which the colour-blind may select as matching the test colours. The pale green is shown first and the candidate asked to match it; next the pale pink; finally the bright red. If he selects not only similar colours, but also confusion colours to match the pale green, his colour sense is defective; if he matches blue or violet with the pink skein he is *red-blind*; if he selects green or gray he is *green-blind*. Finally, if he matches the red skein with greens or browns *darker* than the red, he is red-blind; if he selects shades of these colours *lighter* than the red, he is green-blind.

5. A spectroscope is used in official tests at the Board of Trade and Admiralty, in which any part of the spectrum can be isolated by shutters.

6. An interesting and sometimes amusing test is to make the colour-blind person copy a coloured drawing with paints or crayons, thereby proving his disability to his friends.

Acquired Colour-Blindness is often found in diseases of the retina and optic nerve. It is generally present in optic-nerve atrophy when vision is markedly impaired.

Coloured Vision is occasionally complained of by patients with or without changes in the retina. The most frequent form is red vision (erythropsia) after cataract extraction.

Hysterical Amblyopia.

This condition usually occurs in young girls and women, sometimes in young persons of the male sex. It is usually bilateral, occasionally unilateral.

Symptoms.—The most constant symptom is a diminution in the acuteness of vision, which frequently amounts to complete blindness. The field of vision is contracted concentrically, both for white and colours; since the retina becomes exhausted rapidly, this limitation may become more marked with each succeeding test during the same examination. The colour-fields have not the same relative areas as with the normal eye; they may be larger than that for white; their order is often reversed—*i.e.*, green the largest, red next, and blue the smallest. There may be scotoma or hemiopia. A great variety of other ocular symptoms may be present, such as photophobia, flashes of light, blepharospasm, corneal anæsthesia, monocular diplopia, ptosis, and changes in the size and shape of images. The pupillary reflexes and ophthalmoscopic appearances are normal.

With these ocular manifestations there are usually other hysterical symptoms, especially hemianæsthesia of the affected side. It is sometimes difficult to distinguish between this affection and malingering. It sometimes follows injuries (traumatic hysteria) even when these do not involve the eye.

FUNCTIONAL DISEASES OF THE RETINA

Prognosis is good, although the affection may last many months or even years.

Treatment is directed to the hysterical condition.

Simulated Amblyopia (Malingering).

Patients sometimes pretend to be blind in one eye in order to recover damages for alleged injury; occasionally bilateral blindness is simulated.

The detection of pretended monocular blindness is usually easy. One of the following tests may be employed:

Tests.—1. Place a lighted candle 15 or 20 feet in front of the patient, and put a prism of 6 degrees, base upward or downward, before the sound eye; if the patient admits seeing double, it is an indication that the vision is present in both eyes.

2. With the lighted candle in the same position, cover up the supposed blind eye. Then produce monocular diplopia by moving a 6-degree prism, base upward or downward, until the apex corresponds to the centre of the pupil. Next uncover the blind eye, and at the same time move the prism until it covers the entire pupil. If now there is still double vision (binocular diplopia) it is evident that both eyes see. This test is difficult to apply.

3. Place a strong convex lens (12 D.) before the good eye and a weak concave lens (0.25 D.) in front of the supposed blind eye, and direct the patient to read the distant test types; if he succeeds, it is proof of malingering, since it is impossible for him to see with the sound eye when it is covered by the strong lens.

It is rare for a patient to simulate blindness in both eyes and more difficult to detect him in such cases. A diminution in acuteness of vision of both eyes is more frequently feigned, than binocular blindness. In such cases malingering is suspected when there is an absence of agreement in the results of the functional and objective examination of the eyes, contradictory statements regarding the different steps in the functional examination, or contraction of the pupils to light. In rare instances, the pupils react on exposure to

light in cases of absolute blindness, the lesion being situated in the visual centres or in the connexion between these centres and the corpora quadrigemina (3, Fig. 243A). In feigned binocular blindness a close watch must be kept on the patient when he thinks he is free from observation, and the following test may be employed: Place a lighted candle in front of the patient; hold a 6-degree prism, base outward, before one eye; if both eyes see, the one covered by the prism will move inward in order to avoid diplopia; on removing the prism, it will move outward, the other eye remaining fixed.

Blindness of one eye may be due to neglected squint, or rarely to congenital amblyopia. Blindness of both eyes without ophthalmoscopic signs may occur immediately after a head injury. In case of doubt repeat the examination a month later. Presence or absence of pallor of the optic discs will then decide the question.

Since the passing of the Workmen's Compensation Act, most of the cases which present themselves are those in which some injury has been sustained. The question to be decided is to what extent, if at all, the workman is exaggerating his symptoms. When preparing a report upon the case, it should be remembered that the report may be the subject of cross-examination in a court of law, and any careless wording may subject one to much annoyance. Facts should first be stated definitely. Opinions and inferences should then be stated as such, and one should be prepared to give reasons for them. In a law court, such a statement as 'The patient suffers from headaches' would certainly be met by the question, 'Do you know this is a fact? If so, how do you know?'

Amblyopia and Amaurosis from Various Causes.

Besides the forms of amblyopia already described, there are others, of less frequent occurrence, due to uræmia, reflex irritation, malaria, and quinine. A considerable number of drugs are occasionally responsible for more or less complete amblyopia.

Uræmic Amblyopia has been described on p. 280.

Reflex Amblyopia, due to reflex irritation, is rare and of rather doubtful occurrence, except in the case of the teeth, irritation from which has been found responsible for amblyopia in occasional instances.

Malarial Amblyopia has been observed in malarial diseases. It affects one or both eyes, lasts some hours or days, and usually disappears completely as a result of the use of anti-periodics.

Quinine Amblyopia, or amaurosis, occurs after large quantities of quinine have been taken, occasionally with moderate doses in susceptible individuals. Besides other symptoms of cinchonism, there are more or less complete blindness, often noticed suddenly, contracted fields, dilated pupils, and marked pallor of the disc, with extreme contraction of the retinal vessels. The condition is due to spasm of the retinal vessels causing anæmia of the fundus, degeneration of the ganglion cells and nerve fibres of the retina, and later atrophy of the optic nerve. After a time, central vision is restored completely or partially, and the field widens, but rarely regains its full extent. Treatment consists in discontinuing the drug, inhalations of nitrite of amyl, nitro-glycerine, the use of bromides, strychnine, and digitalis.

Night Blindness is a condition in which the sight is good by day or with good illumination, but more or less deficient at night, or with reduced illumination. It is a symptom of certain forms of retinitis, especially retinitis pigmentosa, but it also occurs without ophthalmoscopic changes. The latter form of diminished light-sense is caused by anæsthesia of the retina; it usually co-exists with xerosis of the conjunctiva, and is dependent upon the same cause—diminished ocular nutrition, due to a debilitated state of the system, such as exists in starvation, profound anæmia, scurvy, and the like. The affection usually disappears with improvement of the general health by good and sufficient food tonics (cod-liver oil, iron), and the use of dark glasses.

Day Blindness is the name given to a condition in which the sight is better at dusk or in feeble illumination than in bright

light. This symptom is found in tobacco amblyopia and with central scotoma in general. In cases in which there are central opacities of the lens or cornea the patient sees better in reduced illumination because the dilated pupil permits vision through the peripheral clear portion of the cornea and lens.

Hemianopsia.

Connexion between the Retinæ, the Fibres of the Optic Nerves and Tracts, and the Cerebral Cortex (Figs. 165 and 243A, also Plate XXV).—Familiarity with the course of the optic-nerve fibres from the eye to the cortex is of great practical value in the localization of various lesions causing defects in the field of vision.

The optic nerves terminate at the chiasm, which lies in the optic groove on the body of the sphenoid bone, where they semi-decussate; from the posterior border of the chiasm they are continued backward as the optic tracts. The optic tracts pass outward and backward, winding around the crura cerebri to the primary optic ganglia—the external geniculate bodies, the anterior corpora quadrigemina, and the pulvinar of the optic thalami (*POG*, Figs. 165 and 243A). The bulk, if not all, of the fibres are now believed to pass to the external geniculate bodies, 10 per cent. of these then passing to the anterior corpora quadrigemina, none connecting with cells of the optic thalami. The fibres from the anterior corpora quadrigemina pass to the nuclei of the oculomotorius and preside over the reflex action of the pupils and the movement of the ocular muscles. After forming connexions with the cells of the geniculate body, which is composed of laminæ of cells with which the different parts of the retina are connected in definite sequence, the relay fibres pass on through the posterior portion of the internal capsule, then form the optic radiations or fibres of Gratiolet, and end in the cortical ganglion cells of the mesial surface of the cuneus and the parts surrounding the calcarine fissure; this portion of the occipital lobe is known as the visual area of the cerebral cortex (*O*, Fig. 243A). In this visual area the different parts of the retina are represented in definite sequence,

the macula most posteriorly, the peripheral portions most anteriorly. The upper portion of the retina is represented in the roof of the fissure, the lower portion in the floor; finally the retina nearest the vertical meridian is represented in the




FIG. 243A.—SCHEMATIC REPRESENTATION OF VISUAL PATHS.

L, left eye; *R*, right eye; *TL*, temporal field of left eye; *NL*, nasal field of left eye; *NR*, nasal field of right eye; *TR*, temporal field of right eye; *ON*, optic nerve; *C*, chiasm; *POG*, primary optic ganglia; *OMN*, oculomotor nuclei; *O*, occipital lobe; *OR*, optic radiations. Division of fibres at 1 causes complete blindness of the left eye and loss of direct pupillary reaction; at 2, right homonymous hemianopsia with loss of reaction of the pupil when the left halves of the retinae are illuminated; at 3, right homonymous hemianopsia with preservation of the reaction of the pupil when the left (and right) halves of the retinae are illuminated; at 4, bitemporal hemianopsia; at 5, left nasal hemianopsia.

depth of the fissure and that nearest the horizontal meridian in the lip.

In the ganglion cells of the visual area an excitation in the optic-nerve fibres is changed into a sensory perception (sight) or into permanent changes (memories, optical memory pictures). After destruction of this area, excitation of the optic-nerve fibres either fails to arouse visual sensation of any kind (blindness), or fails to summon forth any recollection of objects or circumstances acquired through previous

PLATE XXV.




FIG. 243.—DIAGRAMMATIC REPRESENTATION OF THE
VISUAL AND PUPILLARY PATHS.

To face page 316.

education; in the latter case, objects are seen but not recognized (psychical or cortical mind-blindness).

Each retina is supplied by optic-nerve fibres passing to both sides of the brain. Each optic nerve is composed of an external set of fibres derived from the outer or temporal half of the retina, and an internal set derived from the inner or nasal half of the retina. In the axis of the optic nerve is found a special set of fibres which pass to the macula and the space between it and the disc. These macular fibres, when they reach the eyeball, are collected into a sector corresponding to the outer third of the disc, the apex directed toward the centre and the base toward the margin of the papilla. The external or temporal fibres are continued along the lateral part of the chiasm and tract, and pass to the primary optic centre of the same side. The inner fibres, derived from the nasal half of the retina, pass into the chiasm and decussate; they are continued in the tract of the opposite side, thus passing to the side of the brain opposite to the eye which they supply.

The chiasm presents laterally the direct or temporal fibres of both eyes, and in its centre the decussation of the inner or nasal fibres of both retinae. Consequently, the decussation in the chiasm is not complete, but partial—a semi-decussation.

Each optic tract contains fibres from both eyes. The right optic tract consists of non-decussating fibres from the right (temporal half) of the retina of the right eye, and decussation fibres from the right (nasal) half of the left eye. Hence the right halves of both retinae, and thus the left halves of both visual fields, are connected with the right tract (Plate XXV.). It follows, therefore, that the visual impulse, excited by objects placed to the left of the median line, passes to the cortex of the right hemisphere by means of the right optic tract; and that the perception of all objects placed to the right of the median line is conveyed by the left optic tract to the cortex of the left hemisphere.

Hemianopsia.—This arrangement of fibres in the chiasm explains the occurrence of a form of visual disturbance known as hemianopsia (hemianopia, hemiopia), by which we mean

the loss of vision for corresponding halves or sectors of the visual fields. If a lesion interrupts the continuity of the right optic tract, the right cortical visual area, or any portion of the visual path between these parts, there will be blindness of the right halves of both retinae; as a result the left halves of the fields of vision of both eyes will be lost, and only objects which are placed to the right of the median line will be perceived. This is known as homonymous or lateral hemianopsia, and in this particular case the condition is called left homonymous hemianopsia, because the left halves of the fields of vision are wanting. Homonymous hemianopsia (Fig. 244), therefore, always points to a lesion situated in the visual path or cortex on the central side of the chiasm, and upon the




FIG. 244.—THE FIELDS OF VISION IN RIGHT HOMONYMOUS HEMIANOPSIA.

same side as the blind halves of the retinae. It is the commonest form of hemianopsia.

If a lesion extends antero-posteriorly through the chiasm it will destroy all the decussating fibres which supply the inner or nasal halves of both retinae, and there will be a loss of vision in the outer or temporal halves of the field of both eyes, a condition called bitemporal hemianopsia (4, Fig. 243A). It is seen in acromegaly and pituitary tumours.

If a lesion attacks each side of the chiasm it will destroy the non-decussating fibres which come from the temporal halves of the retinae, and will, therefore, cause a loss of the nasal or inner half of the field of vision of each eye; this is known as binasal hemianopsia. Bitemporal and binasal

hemianopsia are known as crossed hemianopsia. These forms are rare, as will be inferred when the situation of the lesion necessary to produce them is considered; it is doubtful whether binasal hemianopsia ever occurs. Another rare form of hemianopsia is altitudinal (inferior or superior)—when the upper or the lower half of each field is wanting.

Hemianopsia is said to be complete when there is a symmetrical absence of the entire half of the field of vision. It is incomplete when there is an absence of a small portion or sector occupying a symmetrical position in the visual fields of the two eyes; the lesion then involves only a portion of the fibres of a visual tract or cortical visual area.

Even in cases of complete hemianopsia, the line between the absent and the preserved portion of the field seldom extends through the fixation point, the portion of the field corresponding to the macula being usually preserved. When both halves of the fields are lost successively (double homonymous hemianopsia), there will be blindness except at the situation of these macular fibres. This occurrence has been explained by supposing—(1) that the macula is represented in both hemispheres, and (2) that the cortical centre for the macula receives a special and abundant blood-supply from both the posterior calcarine and the middle cerebral arteries. The last view is the one now most generally accepted.

Hemianopsia is known as absolute when there is loss of light, form, and colour sense, and relative when only the colour sense or both the colour sense and form sense are destroyed over the symmetrically defective areas, while the light sense remains relatively intact. This is known as hemiachromatopsia; it was formerly believed to indicate the existence of separate cortical centres for colour, form, and light perception, but is now explained by assuming the existence of a lesion of less intensity than that which causes absolute hemianopsia.

Complete blindness in one eye only is always due to a lesion situated in front of the chiasm. The same applies to scotomata, which are defects in the visual field of one eye, or non-symmetrical defects in the fields of both eyes; when

central, they indicate an involvement of the papillo-macular sector of the optic nerve.

The Hemianopic Pupillary Reaction (Wernicke) may be of value in determining whether a lesion causing homonymous hemianopsia is situated behind or in front of the primary optic ganglia. If behind this point the pupillary light reflex will be preserved; if in front of these ganglia (in the optic tract) it may be diminished, when the blind half of the retina is illuminated (Fig. 165). This test is very difficult to apply in a conclusive manner.

Scintillating Scotoma (Transient Hemianopsia) is a form of temporary blindness of not infrequent occurrence, generally associated with migraine and probably due to a circulatory disturbance in the occipital lobe. The attack begins with a central dark spot before both eyes, which spreads by scintillating and coloured zigzag lines, until there is a considerable gap in the field, often hemianopic. Accompanying the attack there are headache, general malaise, vertigo, and sometimes nausea and vomiting. The attacks vary in frequency and last about fifteen minutes, after which the amblyopia disappears entirely. The affection occurs after excessive mental or physical exertion and following marked eye-strain. Unless associated with paralysis, aphasia, or other symptoms of cerebral trouble, it is not of serious import. Treatment consists in attention to the general health, correction of eye-strain, avoidance of fatigue of any kind, and the use of remedies suited to migraine.

CHAPTER XXII

GENERAL OPTICAL PRINCIPLES

FROM a luminous point, rays of light pass out in straight lines in every plane and in every direction; the lines of direction are called rays. These travel with a rapidity which diminishes with the density of the medium traversed.

The amount of divergence of the rays of light falling on a given area is inversely proportionate to the distance of the luminous source; the nearer this point, the more divergence. When proceeding from a point distant 20 feet or more, the divergence of rays is so slight that for practical purposes we may assume them to be parallel.

When a ray of light meets an opaque body, it is either absorbed or reflected. When it meets a transparent medium, some of it is absorbed and reflected, but the greater part traverses the medium (provided that the angle of incidence is not greater than the critical angle of the medium), being deflected in its course; this bending is called refraction.

Reflection occurs from any polished surface (mirror)—plane, concave, or convex. The ray striking the mirror is called the incident ray (I B, Fig. 245); that returning from the mirror, the reflected ray (B R, Fig. 245).

Laws of Reflection.—(1) The angle of reflection is equal to the angle of incidence; (2) the reflected and incident rays are both in a plane perpendicular to the reflecting surface. In Fig. 245, I B is the incident ray on the reflecting surface A C, B R the reflected ray, and P B the perpendicular. The angle of incidence, I B P, is equal to the angle of reflection, P B R. I B, P B, and B R lie in the same plane.

Reflection by a Plane Mirror.—The image is formed at a distance behind the mirror equal to the distance of the object in front of it; it is a virtual image, erect, and of the same size as the object. In Fig. 246, O is the object, I the image, and E the eye of the observer. The image of the candle, O, is formed behind the plane mirror, MM; the observer's eye, E, receives the rays from O as if they came from I.

Reflection from a Concave Mirror.—A concave surface may be considered as made up of a number of plane surfaces inclined toward one another. Parallel rays falling on a concave mirror are reflected as convergent rays which meet on the axis of the surface at a point called

the principal focus (Pf, Fig. 247); the latter is midway between the mirror and its optical centre, C. The distance of the principal focus from the mirror is called the focal length of the mirror.




FIG. 245.—REFLECTION BY A PLANE SURFACE.




FIG. 246.—FORMATION OF IMAGE BY A PLANE MIRROR.

The position of an image formed by a concave mirror varies with the distance of the object from the mirror. If the object be placed at the principal focus, Pf, the reflected rays are parallel to each other and to the axis of the mirror. If the object be placed at the centre of




FIG. 247.—REFLECTION BY A CONCAVE MIRROR.

concavity, C, the reflected rays return along the same lines. If the object is beyond the centre, at CF, the reflected rays focus between the centre and the principal focus at cf; and conversely, if the object

be moved between the principal focus and the centre, at cf , its focus will be beyond the centre, at CF . These two points, CF and cf , bear a reciprocal relation to each other, and are known as conjugate foci. The nearer the object approaches the principal focus, the greater the distance at which reflected rays meet. If the object be placed nearer the mirror than the principal focus at X , reflected rays will be divergent and never meet; if, however, these divergent rays are continued backward, they will unite at a point, Vf , behind the mirror. This point is called the virtual focus, and an observer placed in the path of the reflected rays will receive them as though they came from this point,

It follows, therefore, that concave mirrors produce an enlarged, erect, and virtual image if the object is placed nearer than the principal focus; no image of an object placed at the principal focus; an enlarged, inversed, real image if the object is placed between the principal focus and the centre; an inverted image of the same size when placed at the centre; and a small, inversed, real image if the object is placed beyond the centre.

Reflection by a Concave Mirror.—

Parallel rays falling on a convex surface are reflected divergent, and hence never meet; but if prolonged backward a negative image is formed at a point called the principal focus (Fig. 248, F).




FIG. 248.—REFLECTION BY A CONVEX MIRROR.



FIG. 249.




FIG. 250.

FIG. 249.—PASSAGE OF A PERPENDICULAR RAY THROUGH A TRANSPARENT MEDIUM.

FIG. 250.—REFRACTION BY A TRANSPARENT MEDIUM WITH PARALLEL SURFACES.

The image is always virtual, erect, and smaller than the object, independent of the position of the object before the mirror.

Refraction is the deviation in the course of rays of light in passing from one transparent (dioptric) medium into another of different density (refracting medium). The ray which falls perpendicular to the surface separating the two media is not refracted, but continues in a straight course (Fig. 249, $P'P$).

In passing from a rarer to a denser medium, a ray is refracted toward the perpendicular to the refracting surface; in passing from a denser to a rarer medium, the ray is refracted away from the perpendicular. In Fig. 250 the incident ray, IR , in passing from a rarer medium (air) into a denser medium (glass), is refracted toward the perpendicular

P P; in passing from a denser to a rarer medium, the emergent ray, E R, is refracted from the perpendicular, P P. The ray continues in a line parallel to its original course, but has suffered lateral deviation. The angle formed by the incident ray with the perpendicular, I R P, is known as the angle of incidence; the angle formed by the emergent or refracted ray with the perpendicular, P E R, is known as the angle of refraction.

Index of Refraction.—The relative density, or the comparative length of time occupied by light in travelling a definite distance in different transparent media, is known as the index of refraction. Air being taken as 1.00, the index of refraction of water is 1.33, of the cornea 1.33, of the lens 1.40, of crown glass 1.5, of flint glass 1.6, and of diamond 2.50.

Prisms.

A prism is a piece of glass or other refracting substance bounded by plane surfaces inclined toward each other (Fig. 251). The angle formed by the two surfaces is called the refracting angle of the prism (B A C), the thin edge where the intersecting surfaces meet is known as the apex (A), and the opposite thick portion as the base (B C).

Refraction by a Prism.—Rays of light passing through a prism are bent toward the base. In Fig. 251 the incident




FIG. 251.—REFRACTION BY A PRISM.




FIG. 252.—PASSAGE OF PARALLEL RAYS THROUGH A PRISM.

ray I R is refracted toward the perpendicular P R, at R, and assumes the direction R R in the prism; on emerging, it is refracted away from the perpendicular and continues as R E toward the base of the prism. To the eye placed at E the ray R E seems to come from X; hence an object seen through a prism appears displaced toward the apex. A prism has neither converging nor diverging power, and therefore has no focus and cannot form an image; rays that are parallel before entering the prism are parallel on emerging (Fig. 252).

The Numbering of Prisms.—The strength of a prism is expressed either in degrees or in prism diopters; a third method (the centrad) is not used much. In the first method, which is most commonly employed in practice despite certain faults, the value of the prism corresponds to the refracting angle (geometrical angle), and is expressed: Prism 1° , 2° , 10° , etc. The prism diopter is a deviation, the tangent of which is $\frac{1}{100}$ of the radius, and is expressed: 1 P. D., or 1^Δ ; 2 P. D., or 2^Δ , etc. A centrad corresponds to a deviation, the arc of which is $\frac{1}{100}$ of the radius and is expressed 1^Δ , 2^Δ , 10^Δ , etc. Within the limits of common use, the three scales can practically be considered alike. But the angle through which the prism bends a ray of light varies according to the kind of glass of which it is made. The degree of this angle is usually about half the number of the prism. The most rational method is to number prisms according to the number of degrees through which they deflect a ray of light.

The Position of a Prism when placed in front of an eye is indicated by the direction of its base; 'base out' means that the thick part of the prism is toward the temple; the base may be up, down, in, or out. In correcting muscle insufficiencies the base is placed towards the muscle it is to help.

The Use of Prisms.—(1) To counteract the effects of muscular paralysis or insufficiency; (2) for the exercise of weak muscles; (3) to test the extent to which the eyes can be diverted from parallelism; (4) as a test for muscular insufficiency; (5) for detecting simulated blindness.

Lenses.

A lens is a transparent refracting medium, usually made of glass, in which one or both surfaces are curved. There are two kinds: spherical and cylindrical lenses.

Spherical Lenses (abbreviated Sph. or S.) are so called because the curved surfaces are segments of spheres (Fig. 253); such lenses refract rays of light equally in all meridians or planes. There are two kinds of spherical lenses, convex and concave.

Convex Spherical Lenses may for the moment be considered as if they were formed of an infinite number of graduated

prisms with their bases at the centre of the lens and their apices radiating to the periphery (Fig. 254, A); the lenses are therefore thick at the centre and thin at the edge. They are known as




FIG. 253.—THE RELATION OF THE SURFACES OF LENSES TO SPHERES.

1, Plano-convex; 2, bi-convex; 3, convex meniscus; 4, plano-concave; 5, bi-concave; 6, concave meniscus.

converging, magnifying, positive, and plus lenses, and denoted by the sign $+$. They have the power of converging parallel rays and of bringing them to a focus (Fig. 257). There are three different forms: (1) Plano-convex, one surface plane, the other convex (1, Fig. 255); (2) biconvex or double convex, both surfaces convex (2, Fig. 255); (3) concavo-convex (convex periscopic, convex or converging meniscus), one surface convex, the other concave—the former having the shorter radius of curvature (3, Fig. 255).

The periscopic lens (whether $+$ or $-$) diminishes spherical aberration and enlarges the field of vision.

Concave Spherical Lenses may similarly be considered as formed of prisms with their apices together and toward the




FIG. 254.—THE FORMATION OF LENSES BY PRISMS.




FIG. 255.—CONVEX LENSES.

1, Plano-convex; 2, biconvex; 3, convex meniscus.



FIG. 256.—CONCAVE LENSES.

1, Plano-concave; 2, biconcave; 3, concave meniscus.

centre (Fig. 254, B); they are therefore thin at the centre and thick at the edge. They are known as diverging, reducing, negative, or minus lenses, and denoted by the sign $-$. Rays of light after passing through a concave lens are rendered

divergent; if prolonged backward they form an image on the same side as the object (Fig. 258). There are three different forms: (1) Plano-concave, one surface plane, the other concave (1, Fig. 256); (2) biconcave or double concave, both surfaces concave (2, Fig. 256); (3) convexo-concave (concave periscopic, concave or diverging meniscus), one surface convex and the other concave, the latter having the shorter radius of curvature (3, Fig. 256).

The Action of Spherical Lenses.—Since spherical lenses are formed of prisms with their bases (convex) or apices (concave) in apposition, and since rays in passing through




FIG. 257.—THE ACTION OF A CONVEX LENS ON PARALLEL RAYS.




FIG. 258.—THE ACTION OF A CONCAVE LENS ON PARALLEL RAYS.

a prism are refracted towards its base, it follows that convex lenses cause convergence (Fig. 257), and concave lenses produce divergence, of rays (Fig. 258).

A line passing through the centre of the lens (optical centre or nodal point, *o*, Fig. 259) at right angles to the surfaces of the lens is called the principal axis (*A B*, Fig. 259). A ray passing through this axis (axial ray) is not refracted; all other rays suffer more or less refraction. Rays passing through the optical centre of a lens, but not through the principal axis (secondary rays), are slightly deviated, but emerge in the same direction as they entered (*C D* and *E F*, Fig. 259); the deviation in thin lenses is so slight that practically they may be considered as straight lines, and are called secondary axes.




FIG. 259.—PRINCIPAL AND SECONDARY AXES OF A CONVEX LENS.

Foci of a Convex Lens.—The point to which rays con-

verge after refraction by a convex lens is called its focus. The principal focus is the focus for parallel rays (F, Fig. 260); the distance of this point from the optical centre is called the focal distance of the lens (XF, Fig. 260). Since the course of a ray passing from one point to another is the same, independent of the direction, it follows that rays from a luminous point placed at the principal focus will emerge as parallel after passing through the lens.

In Fig. 260 the rays A B C strike the surface of the lens at L M N; the axial ray B strikes the lens at M perpendicular to its surface, and consequently continues in a straight line to F. The ray A strikes the lens obliquely at L, and is bent




FIG. 260.—THE PRINCIPAL FOCUS OF A CONVEX LENS.

toward the perpendicular of the surface of the lens at that point, shown by the dotted line P R; on leaving the lens obliquely at S, it is deflected away from the perpendicular R T, being directed to F, where it meets the axial ray BF.

The ray C is refracted in a similar manner; it is bent upon entering the lens at N, and rendered additionally convergent when emerging from the lens, and finally it meets the other rays at F. If, in this same illustration, the rays proceed from F, the principal focus, they emerge parallel (L A, M B, N C) after passing through the lens.

Conjugate Foci of a Convex Lens.—Conjugate foci are interchangeable foci in which the image can be replaced by the object and the object by the image. When divergent rays (*i.e.*, rays coming from a point nearer than 20 feet) proceed from a point beyond the principal focus, they will meet at a point beyond the principal focus on the other side of the lens. The more distant the luminous point, the nearer the principal focus on the other side of the lens will the rays be focussed. If the luminous point is situated at a distance equal to twice the focal length of the lens, the rays will focus

at the same distance on the opposite side. These are *conjugate foci*.

In Fig. 261 the rays diverging from O and passing through the lens converge at I; if they diverge from I they would




FIG. 261.—CONJUGATE FOCI OF A CONVEX LENS.

return in the same path and meet at O; the points O and I are conjugate foci. In the preceding example the conjugate focus is positive or real.

Virtual or Negative Focus of a Convex Lens.—When rays diverge from some point between the lens and its principal focus (O, Fig. 262), they will continue divergent after refraction, but less so than before entering the lens; if prolonged backward they will meet at a point (I, Fig. 262) on the same side of the lens from which they diverged; this point is a negative or virtual focus.




FIG. 262.—VIRTUAL FOCUS OF A CONVEX LENS.

Foci of a Concave Lens.—After passing through a concave lens, rays of light, whether originally parallel or divergent, are always divergent, and the focus is, therefore, always negative or virtual; it is found by continuing these divergent rays backward until they meet at a point (Fig. 258).

Formation of Images.—The image of an object formed by a lens is a collection of foci, each corresponding to a point in the object. Such images are either *real* or *virtual*. A real image is formed by the meeting of rays; it can be projected on a screen. A virtual image is formed by the prolongation backward of diverging rays until they meet at a point; it can only be seen by looking through the lens.

To find the position and size of an image formed by a lens, it is necessary to obtain the conjugate focus of each extremity of the object. Two lines are drawn from each of these points, one parallel to the axis of the lens and then




FIG. 263.—REAL, INVERTED, AND REDUCED IMAGE FORMED BY A CONVEX LENS.

through the principal focus, and the other through the optical centre; the image will be formed at the point where these rays intersect (Fig. 263, 264, 265).

In Fig. 263 *AB* is the object, *o* is the optical centre of the lens, and *PF* its principal focus. From *A* two rays are drawn, one parallel to the axis of the lens and then through the principal focus *PF*, and a secondary ray through *o*; the image of the point *A* is formed at *a*, where these two lines intersect. The conjugate focus of *B* is found in the same manner.

The relation in size between image and object depends upon their respective distances from the optical centre of the lens.




FIG. 264.—REAL, INVERTED, AND ENLARGED IMAGE FORMED BY A CONVEX LENS.

In Fig. 263 the object is placed at a greater distance than twice the principal focus, hence the image is real, inverted, and smaller. If the object is situated at exactly twice the distance of the principal focus, the image will be real, of the

same size, and inverted. If the object is situated just beyond the principal focus, the image will be real, enlarged, and inverted (Fig. 264). If the object be placed at the principal focus, the rays will be parallel after refraction, and no image will be obtained. If the object be nearer than the principal




FIG. 265.—VIRTUAL IMAGE FORMED BY A CONVEX LENS.




FIG. 266.—VIRTUAL IMAGE FORMED BY A CONCAVE LENS.

focus, the rays will be divergent after passing through the lens (Fig. 265), and no real image will be formed; but by projecting these rays backward they would meet, and an eye placed at *PF*, Fig. 265, will receive the rays from *a b* as if they came from *A B*; the image will be enlarged, erect, and virtual; it is on the same side of the lens as the object, and is seen only by looking through the lens, which acts as a magnifying-glass.

Images formed by concave lenses are always virtual, erect, and smaller than the object; they are seen only by looking through the lens, which acts as a reducing-glass (Fig. 266).

Cylindrical Lenses.—A cylindrical lens or cylinder (abbreviated *Cyl.* or *C.*) is a segment of a cylinder parallel to its axis (Fig. 267). Cylinders are divided into convex and concave. Light passing through a cylinder in the plane of its axis is not refracted, and behaves exactly as though passing through a plate of glass with parallel sides; in this direction the surface of the lens is straight.




FIG. 267.—THE CONSTRUCTION OF A CONVEX AND A CONCAVE CYLINDRICAL LENS FROM A CYLINDER.

But when light passes through in a plane opposite or perpendicular to the axis of a cylinder, the rays are rendered

convergent or divergent, according as the cylinder is convex or concave; in this direction the surface of the lens is curved. Parallel rays of light after refraction by a cylinder are focussed in a straight line which corresponds to the axis of the cylinder (Figs. 268, 269). A spherical lens refracts equally in all planes; a cylindrical lens does not refract in the axial plane, but all other rays are refracted, those the most which pass at right angles to its axis. It is necessary to indicate the direction of the axis of a cylinder; in the lenses of the trial case, used for the estimation of the refraction of the eye, this is done by a short linear scratch on




FIG. 268.—THE ACTION OF A CONVEX CYLINDRICAL LENS UPON PARALLEL RAYS.




FIG. 269.—THE ACTION OF A CONCAVE CYLINDRICAL LENS UPON PARALLEL RAYS.

the lens at its margins, or by having a portion of the surface on each side ground parallel to its axis (Fig. 271).

The Numeration of Lenses.—The strength of a lens refers to its power of bringing parallel rays to a focus—*i.e.*, its refractive power; this is indicated by its principal focal distance, the interval between the optical centre of the lens and the principal focus. The shorter this distance the stronger the lens; the greater the principal focal distance the weaker the lens. The strength of a lens is the inverse of its focal distance.

✓ *The Metric or Dioptric System* of numbering lenses accepts as its unit a lens which has its principal focus at 1 metre

distance ($39\frac{1}{2}$ English inches, in round numbers 40 inches), this lens is known as 1.00 diopter (abbreviated D.). Every lens is numbered by its strength in whole numbers and in decimal fractions (0.25, 0.50, 0.75). A lens which has twice the strength of the unit is known as 2 D.; its focal distance is $\frac{1}{2}$ metre. If the lens has a strength four times that of the unit, it is called 4 D., and its focal distance is $\frac{1}{4}$ metre. If ten times as strong as the unit, it is known as 10 D., and its focal distance is $\frac{1}{10}$ metre. If one-quarter, one-half, or three-quarters as strong as the unit, it is known as 0.25 D., 0.50 D., or 0.75 D. respectively. In this system the number of the




FIG. 270.—SPHERICAL LENS FROM TRIAL CASE.




FIG. 271.—CYLINDRICAL LENS FROM TRIAL CASE.

lens does not express its focal distance; but the focal distance in centimetres is obtained by dividing 100 cm. by the number of the lens; for example, a 2 D. lens has a focal distance of $\frac{100}{2} = 50$ cm.; a 5 D. lens has a focal distance of $\frac{100}{5} = 20$ cm. The dioptric system is now universally adopted.

To convert the focal distance in inches into the focal distance in diopters, or *vice versa*, divide the number 40 by the number of inches or diopters expressed. For example; 3 D. = $\frac{40}{3} = 13\frac{1}{3}$ inches = $\frac{1}{13\frac{1}{3}}$; 0.50 D. = $\frac{40}{0.50} = 80$ inches = $\frac{1}{80}$; $\frac{1}{20}$ (20 inches) = $\frac{40}{20} = 2$ D.; $\frac{1}{10}$ (10 inches) = $\frac{40}{10} = 4$ D. The following table gives the commonly employed (approximate) equivalents in the inch and the dioptric systems:

COMMONLY EMPLOYED (APPROXIMATE) EQUIVALENTS OF LENSES
NUMBERED IN THE DIOPTRIC AND INCH SYSTEMS.

Dioptrs.	Inches	Dioptrs.	Inches.	Dioptrs.	Inches.	Dioptrs.	Inches.
0.25	1.60	2.25	18	5.50	7.0	13	3.0
0.50	.80	2.50	16	6.00	6.5	14	2.8
0.75	.50	2.75	14	7.00	5.25	15	2.6
1.00	.40	3.00	13	8.00	5.0	16	2.4
1.25	.32	3.50	11	9.00	4.5	—	—
1.50	.26	4.00	10	10.00	4.0	18	2.2
1.75	.22	4.50	9	11.00	3.5	—	—
2.00	.20	5.00	8	12.00	3.3	20	2.0

The Trial Case (Fig. 272) is a box containing + and - spherical and + and - cylindrical lenses, arranged in pairs. The spherical lenses (Fig. 270) usually correspond to those given in the preceding table (30 pairs), the weaker ones



FIG. 272.—THE TRIAL CASE OF LENSES.

separated by intervals of 0.25 D., those of moderate strength by 0.50 D., and the stronger ones by 1 D. The cylindrical lenses (Fig. 271) usually run from 0.25 D. to 6.00 D. The - lenses are mounted in nickelled rims, the + lenses in brass

rims. Besides these lenses, the trial case usually contains a set of prisms, various metal discs, one of which (obturator) is solid, and is used to exclude one eye in the examination and a trial spectacle frame (Fig. 280).

Recognition of the Kind of Lens and Estimation of its Strength.—By moving a spherical lens before the eye and looking at an object, the latter will appear to move, rapidly if the lens is a strong one, slowly if a weak one. If the object seems to move in the opposite direction and appears enlarged, the lens is convex. If the object appears to move in the same direction and seems smaller, the lens is concave.

When a cylinder is moved before the eye in the direction of its axis an object looked at does not appear to change its position; when moved in the opposite direction objects appear to move as with spherical lenses—in the opposite direction when the cylinder is convex, in the same direction when concave.

Having recognized the character of the lens, the strength can be determined by neutralizing. Lenses of opposite kind and known strength are taken from the trial case and placed in front of that to be tested, and the two lenses moved in front of the eye. The neutralizing lens is the one which stops all apparent movement of an object looked at when the combined lenses are moved in front of the eye. The Geneva Lens Measure (Fig. 273) furnishes a very quick and fairly accurate method of determining the character and strength of any lens.




FIG. 273.—GENEVA LENS MEASURE.

Finding the Centre of the Lens.—The optical centre of a lens in the trial case should coincide with the geometric centre. To find the optical centre we look at two lines at right angles to each other through the lens held a few inches away. The portion of the vertical and of the horizontal line seen through the lens is made continuous with the portion seen beyond the lens; then the two lines should cross at the geometrical centre of the lens.

Varieties of Lenses Used to Correct Errors of Refraction.—

(1) Simple spherical lens, convex or concave. (2) Simple cylindrical lens, convex or concave. (3) Sphero-cylinder, a combination of a spherical with a cylindrical lens. (4) Simple prism. (5) Prism combined with various lenses.

Abbreviations and Signs.—The following are in general use in ophthalmology:

Acc.	Accommodation.	O.D.	Optic disc.
Aq.	Aqueous humour.	O.P.	Optic papilla.
As.	Astigmatism.	P.	Pupil.
A.C.	Anterior chamber.	Pr.	Presbyopia.
Ax.	Axis.	P.L.	Perception of light.
C.	Cornea.	p.p.	Punctum proximum.
Ch.	Choroid.	p.r.	Punctum remotissimum.
cm.	Centimetre.	R.	Right eye (and L., left eye).
Cyl.	Cylindrical lens.	Ret.	Retina.
D.	Diopter or dioptric.	Scl.	Sclerotic.
E.	Emmetropia.	Sph.	Spherical lens.
F.	Field of vision.	T.	Tension of the eyeball.
H.	Hypermetropia.	T.n.	Tension normal.
H.l.	Latent hypermetropia.	T. + 1, T. + 2, T. 3, + } de- T. - 1, T. - 2, T. - 3, } de- degrees of increase and de- crease of tension.	
H.m.	Manifest hypermetropia.		
I.	Iris.	Vit.	Vitreous humour.
L.	Left eye (and R., right eye).	Y.S.	Yellow spot and (M.L., macula lutea).
m.	Metre.	V.	Visus, acuteness of sight, power of distinguishing form.
mm.	Millimetre.		
My.	Myopia.		
M.L.	Macula lutea (and Y.S., yellow spot).		
Oph.	Ophthalmoscope, ophthalmoscopic examination, ophthalmoscopic appearances.		

SYMBOLS.

+ Symbol for a convex lens.
- Symbol for a concave lens.

' Foot.
" Inch.
''' Line.

CHAPTER XXIII

OPTICAL CONSIDERATION OF THE EYE

THE eye may be considered as an optical instrument, often compared to the photographic camera, in which by means of a refracting (dioptric) system a small and inverted image of external objects is formed on the retina. The impression received by the rods and cones is conveyed through the optic nerve to the visual cortical area, where the visual act is completed and results in sight.

The eye is well adapted for its function of refraction. It is spherical in shape, about 24 mm. in diameter, and protected externally by the opaque sclera behind and the transparent cornea in front. The outermost portion of the retina consists of a layer of pigment cells which absorbs the excess of light and prevents dazzling.

Dioptric Apparatus of the Eye.—In passing through the eyeball rays of light traverse the cornea, aqueous humour, lens, and vitreous. The refracting surfaces of the eye are the cornea, the anterior surface, and the posterior surface of the lens; the refracting media are the aqueous humour, the substance of the lens, and the vitreous. These surfaces and media constitute the dioptric or refractive apparatus of the eye, a system which is represented by a convex lens of 23 mm. focus; hence in an emmetropic eye in a condition of rest parallel rays are brought to a focus on the retina. The greatest deflection of rays takes place at the anterior surface of the cornea; additional deflection occurs at the anterior and posterior surfaces of the lens. In each case the effect is one of convergence. By the term 'refraction of the eye' we mean the changes which the transparent ocular media exert upon rays of light when the eye is in a state of rest.

Cardinal Points of the Eye.—It is necessary to be acquainted with the cardinal points of the eye (Fig. 274) in order to understand the course of rays of light through this organ; they are the two principal points, the two nodal points, and the two principal foci, all situated on the optical axis.

The Principal Points (P, Fig. 274) are two points so related that when an incident ray passes through the first principal point the corresponding emergent ray passes through the second principal point. These two points are placed so close together in the anterior chamber that they may be con-




FIG. 274.—CARDINAL POINTS OF THE EYE.

sidered as one point, situated about 2 mm. behind the cornea.

The Nodal Points (N, Fig. 274) correspond practically to the optical centre of the dioptric system; they are so close together that they may be considered as one point situated near the posterior pole of the lens about 7 mm. behind the cornea. Rays passing through this point are not refracted, and form either the axial or secondary rays.

The First Principal Focus (A, Fig. 274) is that point on the axis at which parallel rays in the vitreous meet; it is situated about 14 mm. in front of the cornea.

The Second Principal Focus (F, Fig. 274) is that point on the axis at which parallel rays meet after being refracted by the dioptric system of the eye; it is situated to the inner side of the macula, between it and the optic disc, about 23 mm. behind the cornea.

The Centre of Rotation of the eyeball (R, Fig. 274) is situated in the vitreous, about 10 mm. in front of the retina.

The Optical Axis (A F, Fig. 274) is the line connecting the centre of the cornea, the nodal point, and the posterior principal focus on the retina.

The Visual Line (O M, Fig. 274) is the line passing from the object looked at through the nodal point to the macula.

The Line of Fixation is the line joining the object looked at with the centre of rotation; practically it corresponds to the visual line.

The Angle Gamma (γ , Fig. 274) is the angle formed by the optical axis with the line of fixation (practically with the visual line); it varies with the refraction of the eye, being about 3 degrees usually in emmetropia, larger in hypermetropia, and smaller in myopia.

The Angle Alpha is the angle formed by the visual line with the major axis of the corneal ellipse.

Refraction of the Eye.

Emmetropia.—When parallel rays are focussed exactly on the retina with the eye in a condition of rest, the refraction of the eye is normal or emmetropic (Fig. 275, A), and the condition is known as emmetropia.

Ametropia.—When with the eye in a condition of rest parallel rays are not focussed on the retina, but behind or in front of it, the eye is ametropic, and the condition is known as ametropia. The forms of ametropia (errors of refraction) are hypermetropia, myopia, and astigmatism.

Hypermetropia (or far-sightedness) is that form of ametropia in which the axis of the eyeball is too short or the refractive power of the eye too weak, so that parallel rays are brought to a focus behind the retina (Fig. 275, B).

Myopia (near-sightedness) is that form of ametropia in which the axis of the eyeball is too long or the refractive power too strong, so that parallel rays are focussed in front of the retina (Fig. 275, C).




FIG. 275. — A, EMMETROPIA. B, HYPERMETROPIA. C, MYOPIA.

Astigmatism is that form of ametropia in which the refraction in the several meridians of the eyeball is different (Figs. 297 to 301).

The Acuteness of Vision and the method of its determination for distance and near have been described with the functional examination of the eye in Chapter II.

Accommodation.

Accommodation is the act of altering the focus of the eye so that divergent rays (those coming from an object nearer than about 20 feet are appreciably divergent) are brought together on the retina. This is accomplished by means of an increase in the convexity of the lens and thus in its refractive power. The degree of accommodation must vary for every distance of object near.

In the emmetropic eye at rest parallel rays are brought to a focus on the retina (P F, Fig. 276), but rays coming from




FIG. 276.—THE EMMETROPIC EYE IN A STATE OF REST.




FIG. 277.—THE EMMETROPIC EYE DURING ACCOMMODATION.

a near object (divergent rays) are not brought to a focus at all, as they would tend to be focussed behind the retina (D X, Fig. 276); hence distant objects appear distinct and near objects blurred. If the refractive power of the eye is increased by accommodation, parallel rays will be brought to a focus in front of the retina (P F, Fig. 277), while divergent rays will be focussed on the retina (D X, Fig. 277); consequently near objects appear distinct and distant objects appear blurred during accommodation.

Mechanism of Accommodation.—The lens is an elastic structure, and when released from the flattening influence

of its suspensory ligament tends to assume a spherical shape. During accommodation the ciliary muscle (especially the circular fibres) contracts, drawing forward the choroid and relaxing the suspensory ligament; this diminishes the tension of the lens capsule, and allows the inherent elasticity of the lens to increase its convexity. The change in curvature affects chiefly the anterior surface of the lens (Fig. 278). This is Helmholtz's theory and the one usually accepted. Tscherning has advanced a different theory. He maintains that the ciliary muscle increases the tension of the suspensory ligament during contraction, and that this causes peripheral flattening of the lens with bulging anteriorly at its centre.




FIG. 278. — SECTION OF THE ANTERIOR PORTION OF THE EYEBALL.

The dotted lines illustrate the changes during accommodation.

The act of accommodation is accompanied by contraction of the pupil and, in an emmetrope, by convergence of the visual lines.

The Far Point.—When the eye is in a state of rest, with accommodation completely relaxed, it is adapted for its far point (*punctum remotum*). This is the farthest point of distinct vision, and in the emmetropic eye it is situated at infinity.

The Near Point (*punctum proximum*) is the nearest point at which the eye can see distinctly when employing its maximum amount of accommodation. It varies with the amount of accommodation possessed by the eye. The usual plan of determining the near point is to note the shortest distance at which the patient can read the smallest test type (Jaeger, No. 1, Fig. 18) with each eye separately.

The Range of Accommodation is the distance between the far point and the near point.

The Amplitude of Accommodation is the difference between the refractive power of the eye when at rest and when the accommodation is exerted to the utmost. It is expressed

in diopters representing that convex lens which it would be necessary to place before the eye to take the place of accommodation for the near point.

The amplitude of accommodation in diopters is found by dividing 40 by the distance of the near point in inches, or 100 by the near point in centimetres; for example, if the near point of an emmetropic eye is 8 inches or 20 cm., $\frac{40}{8}$ or $\frac{100}{20}=5$ D.=amplitude of accommodation; this rule applies to emmetropia.

In hypermetropia some of the accommodation is required for distant vision; hence we find the apparent amplitude of accommodation, and then add that lens which enables the patient to see distant objects without his accommodation; for example, if the near point of a hypermetropic eye is 8 inches or 20 cm., and the patient is compelled to use 2 D. of accommodation for distant objects, his amplitude of accommodation would be $\frac{40}{8}$ (or $\frac{100}{20}$) $=5+2=7$ D. With the same amplitude of accommodation the near point is farther away than in emmetropia, since some of the power of accommodation is expended in adapting the eye for distant objects; and if the near point were the same, the amplitude of accommodation would be greater in hypermetropia than in emmetropia.

In myopia, since a concave lens is necessary to enable the patient to see distant objects clearly, we must deduct the strength of this glass from that the focal length of which equals the distance of the near point from the eye; for example, if the myopia equals 2 D. and the near point is 4 inches or 10 cm., the amplitude of accommodation will be $\frac{40}{4}$ or $\frac{100}{10}=10$ D. -2 D. $=8$ D. With the same amplitude of accommodation, the near point is closer to the eye in myopia than in emmetropia; and if the near point were the same the amplitude of accommodation would be less in myopia than in emmetropia.

The power of accommodation gradually diminishes, and the near point recedes as age advances, owing chiefly to loss of elasticity of the lens. In the emmetrope at ten years the p.p. is at 7 cm.; at forty years it has receded to 22 cm.; at

sixty years to 100 cm.; and at seventy-five years to infinity, the accommodation being suspended and the p.p. coinciding with the p.r. The following table gives the amplitude of accommodation and the near point at various periods of life. The near point applies only to emmetropic eyes, but the amplitude of accommodation applies to all eyes, whether emmetropic or ametropic. There is a tendency towards increased amplitude of accommodation in hypermetropic, and diminished amplitude in uncorrected myopia.

Year.	Amplitude of Accommodation in Diopters.	Near Point in Centimetres.	Near Point in Inches.	Year.	Amplitude of Accommodation in Diopters.	Near Point in Centimetres.	Near Point in Inches.
10	14.0	7.0	2.8	45	3.5	28.0	11
15	12.0	8.5	3.3	50	2.5	40.0	16
20	10.0	10.0	4.0	55	1.75	55.0	22
25	8.5	12.0	4.7	60	1.0	100	40
30	7.0	14.0	5.6	65	0.75	133	53
35	5.5	18.0	7.0	70	0.25	400	160
40	4.5	22.0	9.0	75	0.0	∞	∞

Presbyopia.—When the near point of the emmetropic eye has receded to a distance at which the finer kinds of work become difficult, the condition is known as presbyopia. This state is the result of a physiological process which affects every eye, and must not be considered a disease. It is usually said to be present when the near point recedes to a distance of more than 22 cm. (9 inches) from the eye, an event which generally happens between the fortieth and the forty-fifth years.

The Association between Accommodation and Convergence.

—The preceding considerations of the subject of accommodation referred to monocular vision or sight with one eye. With binocular vision it is necessary to consider convergence as well as accommodation, for these two actions (together with the contraction of the pupil) are normally associated.

Convergence is the power of directing the visual lines of the two eyes to a near point, and results from the action of

the internal recti muscles. When we look at a distant object accommodation is at rest and the visual lines are parallel. When we look at a near object we are compelled both to accommodate and to converge for that distance; with a certain amount of accommodation a corresponding effort of convergence of the visual lines is associated.

The angle which the visual line makes in turning from a distant object to a near one is called the angle of convergence. The unit of convergence is the metre angle (M.A.), which is the angle formed by the visual line with the median line at a distance of 1 metre (Fig. 279). If the eyes look




FIG. 279.—DIAGRAM ILLUSTRATING THE UNIT OF CONVERGENCE, THE METRE ANGLE.

at an object $\frac{1}{2}$ metre distant the convergence is twice that of the unit, and convergence (C.) = 2 M.A.; if directed toward a point $\frac{1}{3}$ metre distant, C. = 3 M.A.; if toward an object 2 metres distant, C. = $\frac{1}{2}$ M.A.

The emmetropic eye requires for each distance of binocular vision as many metre angles of convergence as it needs diopters of accommodation. To see an object at 1 metre distance, 1 metre angle of convergence is required,

and also 1 diopter of accommodation; at 10 cm., 10 metre angles of convergence and 10 D. of accommodation would be required.

This harmonious relationship between accommodation and convergence is not, however, unchangeable. Within certain limits either of these actions may take place independently of the other.

The Range or Amplitude of Convergence.—The far point of convergence is the point to which the visual lines are directed when convergence is at rest; the near point of convergence is the point to which the visual lines are directed with the maximum amount of convergence. The distance between the far point and the near point of convergence is

the amplitude of convergence; it is expressed by the greatest number of metre angles of convergence of which the eyes are capable. In a state of rest the far point of convergence is at infinity and the visual lines are parallel. In cases of convergent squint the visual lines deviate inward even when convergence is relaxed as far as possible; convergence is then said to be positive. In a case of divergent squint convergence is a negative quantity. Normally the eyes diverge, as a rule, during sleep.

Methods of Investigating the Refraction of the Eye.

There are three principal methods of testing the refraction of the eye: (1) the *subjective method*, in which the refraction is estimated by the acuteness of vision with test types and trial lenses; (2) *retinoscopy*; and (3) the *ophthalmoscope*. The last two are objective methods.

Every examination should be undertaken in a systematic manner. We begin with the external examination of the eyes as described in Chapter I. Next the patient is taken into the dark-room, and the media and fundus are examined with the ophthalmoscope (Chapter III.). Then the refraction may be determined with the ophthalmoscope. The retinoscopic mirror is now employed to estimate the refraction with the ~~shadow test~~. Finally, the patient is examined by the subjective method with test lenses and test types. By employing this order we will save time, since the ophthalmoscopic examination may show changes in the media or fundus which convince us of the impossibility of improving the patient's vision with glasses, or lead us to be satisfied with a limited result. The objective methods of determining the refraction of the eye give very close and accurate results; the subjective method serves to verify these conclusions and sometimes perfects them.

The Determination of the Refraction by the Acuteness of Vision with Test Types and Lenses. The Subjective Method.

After having determined the acuteness of vision for distance as described on p. 11, we endeavour to ascertain which lenses are necessary to correct any error of refraction, and to bring the vision up to the normal $\frac{6}{6}$. The patient is placed in front of the test types, which must be well illuminated by daylight or artificial light, at a distance of 6 metres. The trial frame (Fig. 280) is worn by the patient, and the left eye excluded by means of a solid metal disc. After testing the right eye we proceed with the left.




FIG. 280.—TRIAL FRAME.

If the patient reads $\frac{6}{6}$ without a glass he is probably either emmetropic or has hypermetropia. With bright illumination and contracted pupils an intelligent patient may be able to read $\frac{6}{6}$, even though he have myopia or astigmatism of 0.5 D. A weak convex spherical lens (+0.50 D. Sph.) is held in front of the eye; if he is easily able to read the $\frac{6}{6}$ line he has hypermetropia, and the strongest convex spherical lens with which he can read $\frac{6}{6}$ is the measure of his manifest hypermetropia. Even though he accepts a convex spherical lens, this is probably not the measure of his total hypermetropia, which can be estimated in young persons only after the eye has been

placed under the effects of a cycloplegic. The difference between the manifest and the total hypermetropia is known as the latent hypermetropia: it is this portion which is discovered after accommodation has been paralyzed.

If the patient reads $\frac{6}{6}$, and a weak convex spherical lens blurs his vision, he is either emmetropic or has hypermetropia which is latent.

If the patient's vision is below normal, and instead of reading $\frac{6}{6}$ he reads $\frac{6}{12}$ or $\frac{6}{18}$, he either has considerable manifest hypermetropia, or else he may be myopic or astigmatic; or he may have a combination of these errors. If hypermetropic, spherical lenses will improve his vision. If such improvement does not result upon placing convex spherical lenses before the eye we may try a weak concave spherical lens; if this aids his vision probably he is myopic, and the weakest concave spherical lens that brings his vision to $\frac{6}{6}$ is the measure of his myopia. If concave spherical lenses do not improve the vision, we assume the existence of astigmatism, and cylinders, alone or in combination with spherical lenses, are placed in front of the eye for the purpose of estimating the kind, the axis, and the amount of astigmatism.

This is briefly the method pursued in determining the refraction by means of the acuteness of vision (subjectively); greater details will be supplied in discussing the errors of refraction. But, as already pointed out, it is better and saves times to precede this subjective test by the objective methods, using the former to confirm the findings of the others; this is especially advisable if the error of refraction be a difficult or complicated one.

The vision for near is also tested. To the patient is given a page of Jaeger's test types (Fig. 18), and we note the smallest type which he is able to read with each eye separately, the distance which he selects, and the nearest and farthest distances at which he is able to read. These data give us valuable information regarding the refraction. In myopia the patient may hold the print at a closer distance than normal. In presbyopia he will hold it at a greater distance than normal.

The Ophthalmoscope as a Means of Detecting and Estimating Refractive Error.

The Ophthalmoscope at a Distance (p. 27) gives us qualitative information regarding errors of refraction. When the patient is emmetropic, no details of the fundus will be seen when the light is thrown into the eye from an ophthalmoscope held at a distance of 15 inches. If some part of the disc or vessels is seen the patient is ametropic. If the examiner moves his head from side to side, and the vessels seem to move in the same direction, the case is one of hypermetropia (for in hypermetropia the rays emerge divergent and the image is a virtual erect one). If the vessels seem to move in the opposite direction the case is one of myopia (since in myopia the emerging rays are convergent, and form an inverted image). If the vessels of one meridian only are seen astigmatism is present; this is hypermetropic if the vessels move with the movements of the observer's head, myopic if they move in the opposite direction, and mixed if one set move with and the other against them.

The Indirect Method is not used for determining the amount of error of refraction, but we obtain information of the form of ametropia by noting the size and shape of the inverted image of the disc, and its behaviour upon withdrawing or approaching the lens before the patient's eye. If no change takes place in the shape and size of the image when we withdraw the lens, the eye is emmetropic. If the shape remains the same, but the image becomes smaller when the lens is withdrawn, it indicates hypermetropia. If the shape remains the same, but the image becomes larger on withdrawing the lens, the case is one of myopia. In astigmatism the disc usually appears oval, and the shape of its image changes on withdrawing the lens; one diameter decreases or increases while the other remains stationary in simple astigmatism; both increase or decrease unequally in compound astigmatism; and one increases and the other decreases in mixed astigmatism.

These two methods give no information as to the state of

the refraction which is not more readily obtained by retinoscopy, and are not generally used for the purpose. But, as they will be used in the general examination of the eye in any case, it is well that the student should train his power of observation by noting all there is to be seen.

The Direct Method is a very valuable means of determining the refraction, and, in case of error, the kind and the amount; reliable results are obtained, however, only after considerable practice. For accurate results it is necessary that the accommodation of both patient and observer be in abeyance. The beginner always has difficulty in relaxing his accommodation, and requires considerable training before he masters this necessary step (p. 34). The patient's accommodation is suspended by directing him to look at the wall or at a distant




FIG. 281.—THE ESTIMATION OF THE REFRACTION BY THE DIRECT METHOD OF OPHTHALMOSCOPY.

Both patient and observer are emmetropic.

object, or, better, by the use of a cycloplegic. The examiner, if ametropic, corrects his error by wearing suitable glasses, by having a special correcting lens applied to the sight-hole of the ophthalmoscope, or by subtracting the amount of his error from the result which he obtains in the examination. The examination is conducted in the manner described on p. 32; for accurate results it is essential that the shortest possible distance separate the eye of the patient from that of the observer.

Emmetropia.—The examiner selects a bloodvessel at the outer margin of the disc or between the disc and the macula. If the vessel appears distinct, and if upon rotating a $\times 0.50$ D. lens before the sight-hole it becomes blurred, the eye is emmetropic. Rays coming from an emmetropic eye at rest are parallel, and the observing eye will focus these rays on the retina (Fig. 281).

Hypermetropia.—If the image is blurred, we rotate the lens disc of the ophthalmoscope so as to place convex lenses in the sight-hole; if these render the image distinct the eye is hypermetropic. The strongest convex lens with which we get a distinct image is the measure of the hypermetropia. In Fig. 266 H is the hypermetropic eye under examination, and




FIG. 282.—THE ESTIMATION OF HYPERMETROPIA BY THE DIRECT METHOD OF OPHTHALMOSCOPY.

E the emmetropic eye of the observer. Rays from *a* emerge divergent as though coming from X. The convex lens +L makes them parallel, so that they focus at *b*, on the retina of E, the emmetropic eye of the observer.

Myopia.—If when the image appears blurred a convex lens makes it more indistinct, we rotate the disc of the ophthalmoscope so that concave lenses are brought opposite the sight-hole. If these render the image distinct the eye is myopic, provided that neither the patient nor the surgeon is accommodating.




FIG. 283.—THE ESTIMATION OF MYOPIA BY THE DIRECT METHOD OF OPHTHALMOSCOPY.

The weakest concave lens is the measure of the myopia. We stop at the weakest concave lens which accomplishes this, since stronger lenses of this sort would only encourage the observer to accommodate. In Fig. 283 M is the myopic eye under examination, and E the emmetropic eye of the observer. Rays from *a* leave the myopic eye convergent, and would meet

at X. The concave lens - L renders them parallel so that they are focussed at *b* on the retina of the observer.

Astigmatism.—We find the lens with which a small vertical vessel is seen distinctly, and then the lens which enables a small vessel at right angles to be seen clearly, always remembering that the lens which clears up the image of a vessel in one direction is the measure of the refractive error of the meridian at right angles to it.

Suppose the horizontal vessels appear distinct without any lens, then the vertical meridian is emmetropic; and that the vertical vessels require a convex or a concave lens to render them distinct, then the horizontal meridian is hypermetropic or myopic; the case is one of simple hypermetropic or myopic astigmatism (Figs. 297 and 298).

If both vertical and horizontal vessels are rendered distinct by convex lenses, but a stronger one can be used for the horizontal, the case is one of compound hypermetropic astigmatism (Fig. 299), with the vertical meridian the more hypermetropic; if both vertical and horizontal vessels are best seen with concave lenses, but of different strength, the case is one of compound myopic astigmatism (Fig. 300).

If the vertical vessels can be seen clearly with a convex lens and the horizontal vessels require a concave lens, the case is one of mixed astigmatism (Fig. 301), the horizontal meridian being hypermetropic, the vertical meridian myopic.

Retinoscopy.

Retinoscopy, the Shadow Test, or Skiascopy, is a very accurate objective method of determining the state of the refraction by illuminating the eye with a plane or concave mirror, and observing the direction of the movement of the retinal illumination and its bordering shadows when the mirror is rotated. The shadow test has many advantages. It can be used in children, illiterates, and in markedly defective sight; it is entirely objective, and hence requires no co-operation on the part of the patient; it is quick and accurate; and it requires no expensive apparatus.

The principle of retinoscopy is the finding of the point of reversal or the myopic far point. In myopia an inverted image is formed in the air in front of the eye at the far point—the distance from which rays would be focussed on the retina; this point is known as the point of reversal. If the eye is hypermetropic or emmetropic, a convex lens is placed before it so as to give it an artificial far point.

When light is thrown into the eye by means of a plane or concave mirror at a distance of 1 metre, the fundus is illuminated. By looking through the sight-hole of the mirror an observer will see the illuminated portion (red fundus reflex) and also the shadow bounding this bright area. On rotating the mirror the illuminated area and the shadow will move across the pupil.

The examination is conducted in the dark-room, the darker the better. The source of illumination is placed above, or to one side of, the head of the patient and somewhat behind, so that his face is in darkness (Fig. 285). An unfrosted electric focus-lamp may be used.

Either a plane or a concave mirror may be employed; the plane mirror has certain advantages and is more commonly used. The Lister mirror—a very slightly concave mirror which behaves as a plane mirror at 1 metre distance—is still more satisfactory. The retinoscopy mirror (Fig. 284) usually has a diameter of $3\frac{1}{2}$ cm. with a 3 mm. opening.




FIG. 284.—RETINOSCOPIC MIRROR.

Unless the observer has had great experience in retinoscopy, the patient's pupils should be dilated, and his accommodation paralyzed. He is directed to look at the light. Each eye is tested separately, and one eye is usually covered.

The observer is seated at 1 metre distance (Fig. 285); he should wear correcting lenses if ametropic; he need not relax his accommodation as in using the ophthalmoscope.

since this does not influence the result. If the surgeon is presbyopic he will find it a comfort to have a small lens, $+0.75$ D. or $+1$ D. sph., cemented behind the sight-hole of his mirror.

If now the mirror be rotated slowly from side to side on its vertical axis, so that the light moves across the pupil horizontally, the observer will see an illuminated area and a shadow coming from behind the pupil; if the mirror be




FIG. 285.—THE RETINOSCOPIC EXAMINATION.

rotated on its horizontal axis the light will move across the pupil vertically. The direction of movement of this light and shadow as compared with that of the mirror depends upon the state of the refraction of the eye. The light moves either in the same direction (with) or the opposite direction (against); if we turn the mirror toward the right and the light moves toward the right, we say it moves with the mirror; if upon turning the mirror toward the right the light moves toward the left, we say it moves against the mirror. The

illuminated area and the shadow appear to move with the mirror when the observer is within the point of reversal, and against the mirror when he is beyond this point. With the plane mirror the light moves with the mirror in hypermetropia, emmetropia, and in myopia of less than 1 D., and against the mirror in myopia of more than 1 D.

Besides the direction of the movement, we acquire information from the brightness, the form, and the rate of movement of the light and shadow. If the reflex is bright, its edge




FIG. 286.—RETINOSCOPIC ILLUMINATION AND SHADOW IN ASTIGMATISM.




FIG. 287.—RETINOSCOPIC ILLUMINATION AND SHADOW IN MYOPIA, HYPERMETROPIA, OR EMMETROPIA.

sharp, and the light and shadow move rapidly, the error of refraction is a low one; if the illumination is dull, its edge indistinct, and the movement of light and shadow slow, the error is a high one. If the shadow has a straight edge it is an indication of astigma-

tism (Fig. 286); in hypermetropia, myopia, or emmetropia, the shadow has a crescentic edge (Fig. 287).

We next find the correcting lens—*i.e.*, the lens which causes a reversal of the direction of movement of the light. This lens will be correct for the distance separating the observer from the patient, 1 metre. For infinity, we must add -1 D. to all results; this increases the myopia 1 D., and diminishes hypermetropia 1 D.

If with the plane retinoscope the light moves against the mirror, we place concave spherical lenses before the eye until we succeed in causing a reversal of the movement of the shadow—*i.e.*, cause it to move with the mirror; this lens, to which we add -1 D., is the measure of the patient's myopia. Suppose on placing -1 D. before the eye the light still moves against the mirror, the same with -2 D., but with -2.50 D. the movement of the light is reversed, then $-2.50 + 1 = -1.50$ D. is the correction.

If with the plane retinoscope the light moves with the mirror, the eye may be hypermetropic, emmetropic, or myopic less than 1 D. In such a case we begin by adding a convex lens of +0.50 D. If this causes a reversal of the light the eye is myopic 0.50 D., since $\frac{+0.50}{-0.50} = +0.50$ D. If the +0.50 D. lens does not alter the direction of the movement of the light, but the next lens (+1 D.) causes a reversal, the eye is emmetropic, since $\frac{+1.0}{-1.00} = 0 = E$.

If the +1.00 D. lens has no effect upon the direction of movement of the light, the eye is hypermetropic; we place stronger + spherical lenses before the eye until we find the one which causes a reversal of the movement of the light. Say this is +4 D., then the hypermetropia amounts to $\frac{+4}{+3} = 3$ D.

✓ In the previous examples the results were the same whether the mirror was rotated upon its vertical or its horizontal axis. In astigmatism upon correcting each of the two principal meridians separately, one meridian will require a different lens from the other to cause a reversal of the light. The most common positions of the two meridians in astigmatism are vertical and horizontal. But frequently the edges of the shadows lie more or less obliquely. In such cases the mirror must be rotated so that the light moves obliquely and parallel with the movement of the light. *shadow*

For example, suppose the light moves with the mirror in both meridians, but is more distinct and moves more quickly in one meridian than the other, we diagnose astigmatism. Then we correct the vertical meridian, and find it requires +2 D. for the reversal of the light. Next we find that in the horizontal meridian +4 D. is required for reversal. We add -1 D. to each of these results, and have +1 D. vertical and +3 D. horizontal. The case is one of compound hypermetropic astigmatism, and requires for its correction +1 D. spherical lens combined with +2 D. cylinder, axis vertical.

Now, in order to find the exact axis of the cylinder, the

correcting cylinder and the reversing sphere are put in the trial frame and the axis of the cylindrical lens adjusted until the shadow is even in all meridians.

In practice, it is not necessary to work at exactly 1 metre. The observer may take whatever distance he pleases, provided that it is always the same. Having decided upon the most convenient distance, he must find out by experiment what deduction is to be made from the reversing lens.

CHAPTER XXIV

ERRORS OF REFRACTION

IN emmetropia the eye in a state of rest, without accommodation, focusses the image of distant objects exactly upon the retina (Figs. 275, A, and 288, A); such an eye enjoys distinct vision for distant objects without effort or fatigue. Any variation from this standard constitutes ametropia, a condition in which the eye, in a state of rest, is unable to focus the image of distant objects (parallel rays) upon the retina. Ametropia includes hypermetropia, myopia, and astigmatism. The effects of ametropia are not only indistinctness of vision, but various pains and other symptoms comprised under the term *asthenopia* (weak sight, eye-strain).

Hypermetropia.

Hypermetropia is an error of refraction in which, with accommodation completely relaxed, parallel rays (rays from distant objects) tend to be brought to a focus behind the retina (Figs. 275, B, and 288, B); divergent rays (from near objects) tend to be focussed still farther back.

Etiology.—It is most commonly due to shortening of the antero-posterior diameter of the eyeball (axial H.), less frequently to diminished convexity of the refracting surfaces of the eye (H. of curvature), changes in the media, or absence of the lens (*aphakia*). It is by far the most frequent error of refraction, and is congenital. It is often hereditary. Children are usually hypermetropic at birth, and subsequently become less hypermetropic, emmetropic, or even myopic.

The Course of Rays.—The hypermetropic eye cannot, with-

out accommodation, see either distant or near objects distinctly (Fig. 288, A). In a condition of rest it is adapted for convergent rays, and these are not found in nature. To focus parallel rays on the retina it must either accommodate—*i.e.*, increase the convexity of its lens, as shown in Fig. 288, B—or a convex lens of such a strength that the rays are made sufficiently convergent to be brought to a focus on the retina (Fig. 288, C) must be placed in front of the eye.

To focus divergent rays—*i.e.*, rays from near objects—the




FIG. 288.—A, HYPERMETROPIC EYE IN A STATE OF REST. B, HYPERMETROPIC EYE DURING ACCOMMODATION. C, HYPERMETROPIC EYE CORRECTED BY A CONVEX LENS.




FIG. 289.—A, EMMETROPIC EYE ACCOMMODATING FOR NEAR VISION. B, HYPERMETROPIC EYE ACCOMMODATING FOR NEAR VISION.

hypermetrope must not only accommodate to the extent required of an emmetropic eye (Fig. 289, A), but to an additional extent to compensate for his error. In other words, he requires some accommodation constantly in order to see distant objects distinctly, and in addition the amount equal to that required by the emmetrope for near vision (Fig. 289, B). Such an eye (when the error is uncorrected) is never in a condition of rest as long as it enjoys distinct vision.

Changes in the Eye.—As a result of the constant strain and overaction of the ciliary muscle, the latter becomes hypertrophied, especially its circular fibres (Fig. 291); it remains in a greater or lesser condition of spasm. In high degrees of H. the eyeball may be diminished in size, the anterior chamber shallow, the sclera flat with a sharp curve at the equator, and there may be an apparent external squint owing to the high angle gamma (see pp. 339 and 412).

Hypermetropia is divided into (1) *manifest* and (2) *latent*, the sum of these being the (3) *total*.

The manifest hypermetropia is that which is detected without paralyzing the accommodation, and is represented by the strongest convex glass with which the patient sees most distinctly; it corresponds to the amount of accommodation which he relaxes when a convex lens is placed before the eye. Manifest hypermetropia may be either facultative, when it can be overcome by an effort of accommodation, or absolute, when it cannot be overcome in this manner.

The total hypermetropia is the entire amount of hypermetropia detected after the accommodation has been paralyzed or during complete relaxation of the ciliary muscle.

The latent hypermetropia is the difference between the Hm. and the Ht., and is the amount which is habitually concealed and is discovered only after the use of a cycloplegic.

The application of these terms can be illustrated by considering an example of H. of 2.5 D. in a young person. If in such a case $V = \frac{6}{1\frac{1}{2}}$, and, without the use of a mydriatic, a +1 D. spherical lens brings up the vision to $\frac{6}{8}$, and a stronger convex lens makes the vision again indistinct, we say Hm. = 1 D.; if now we paralyze the accommodation with a cycloplegic and find $V = \frac{6}{3\frac{1}{2}}$, and that a +2.50 D. spherical lens




FIG. 290. FIG. 291. FIG. 292.

FIG. 290.—SECTION OF THE CILIARY MUSCLE IN AN EMMETROPIC EYE.

FIG. 291.—SECTION OF THE CILIARY MUSCLE IN A HYPERMETROPIC EYE.

FIG. 292.—SECTION OF THE CILIARY MUSCLE IN A MYOPIC EYE.

increases this to $\frac{6}{8}$, the Ht.=2.50 D.; the difference between 2.50 D. and 1.00 D.=1.50 D.=Hl.

The ratio between the manifest and the latent hypermetropia is not constant; it depends more or less upon the age and vigour of the individual. In youth the amount of Hl. is apt to be considerable, and consequently a cycloplegic is essential in estimating the amount of hypermetropia. The older a person grows, the less accommodative effort he is able to make; hence the Hl. becomes less and the Hm. greater.

In old persons there is no Hl., the total hypermetropia becoming manifest.

Symptoms.—Unless the error be considerable or the patient be advanced in years, there is usually distinct vision for distance. A great many patients with hypermetropia present no symptoms whatever; this is likely to be the case when the hypermetrope is young, in good health, and indulges in plenty of outdoor exercise. Under these circumstances he is apt to accommodate for his optical defect without any evidence of overaction of the ciliary muscle. In other cases the accommodative effort will be unequal to the task imposed in near work, and as a result the hypermetropia will give rise to accommodative asthenopia (weak sight, eye-strain).

The Symptoms of Asthenopia show themselves particularly after reading, writing, sewing, and other forms of near application, especially in the evening and with artificial illumination. They comprise pain referred to the eyes or above the eyes; headaches, usually frontal, but also occurring in the occiput and other parts of the cranium; various neuralgias; congestion of the conjunctiva and margins of the lids; lacrymation, blinking, and slight photophobia; burning sensation in the lids; and blurring of near vision. These symptoms are more pronounced whenever the general health is below par.

With advancing years there will be greater difficulty in reading without correcting glasses.

In children H. shows a physiological tendency to diminish with the growth of the child. In the adult it remains stationary.

In early childhood hypermetropia often causes convergent squint in a patient whose fusion-sense is deficient (see p. 411).

The smaller and flatter cornea and shallower anterior chamber often associated with hypermetropia form a predisposing cause of glaucoma. Conversely, glaucoma is uncommon in myopic eyes.

Hypermetropic eyes are predisposed to conjunctivitis and blepharitis, phlyctenular affections, and internal squint.

Tests.—These have been described in the preceding chapter. They are the following:

The Subjective Test with Test Types and Test Lenses.—We first record the acuteness of vision and then place convex lenses before the eye, commencing with $+0.50$ D. The strongest lens with which the patient sees $\frac{6}{6}$ or better is the measure of the manifest hypermetropia. Then the accommodation is paralyzed and the test repeated; the strongest lens 'accepted' (i.e., with which the patient's vision is improved) is the measure of his total hypermetropia.

The Ophthalmoscope at a Distance.—The retinal vessels appear to move in the same direction as the observer's head.

The Ophthalmoscope, Indirect Method.—On withdrawing the lens in front of the patient's eye the apparent size of the disc diminishes.

The Ophthalmoscope, Direct Method.—The disc and vessels can be seen distinctly with a convex lens in the sight-hole, the strongest being the measure of the H.

Retinoscopy.—With the plane mirror held at 1 metre, the shadow moves with the mirror; the direction of movement is reversed by convex lenses placed in front of the patient's eye. The lens which causes a reversal, -1 D., is the measure of the H.

Treatment consists in prescribing such convex spherical lenses as will make vision distinct, and enable the patient to do near work without fatigue. The mere existence of hypermetropia is no indication for the use of correcting glasses unless these are worn in childhood for the cure of convergent squint. It is only when there is a diminution in the acuteness of vision or when symptoms arise indicating eye-strain that convex lenses should be used.

Though theoretically it would seem proper to prescribe the

full correction (for Ht.), practically there are many objections and exceptions to this. In every case of hypermetropia occurring in a young child the accommodation should be paralyzed with atropine before performing retinoscopy. An older child may require atropine. Each case should be considered on its merits. There may not be any object in finding out the Hl. if the patient sees perfectly and the correction of Hm. relieves all symptoms. A surgeon who is reasonably skilful in retinoscopy will very rarely need to use a mydriatic for an adult.

The symptoms of the individual give us reliable indications as to the proportion of the Ht. which ought to be corrected, and the constancy with which the glasses should be worn. In cases of squint, and when glasses are prescribed for the relief of conjunctivitis, blepharitis, and headaches, or the occurrence of which is independent of near use of the eyes, they must be worn constantly. In other cases glasses should be worn continuously or only for near work, according to whether the symptoms are always present, or follow only after using the eyes for reading and the like. When distant vision is perfect and comfortable, and the patient does not suffer from any symptoms except when engaged in near work, glasses need be prescribed only for such use; this is often the case in young adults who enjoy good health. Under such circumstances the correction of the Hm. may be sufficient; or we may add to this the correction for part of the Hl., or we may correct the Ht. In cases in which the correction is only partial the glasses may require changing from time to time. In the case of a child, if glasses are required at all a good general rule is to order 0.5 D. less than the Ht. In hypermetropes after forty-five convex lenses may be worn to improve distant vision, and a stronger pair for near; the weaker set is for the H., the stronger pair to correct both the hypermetropia and the presbyopia. Under such circumstances bifocal lenses (Figs. 310 and 310A) are very convenient, the upper segment corresponding to the weaker glass, the lower to the stronger.

Myopia.

Myopia (short-sightedness) is that refractive condition in which, with accommodation completely relaxed, parallel rays are brought to a focus in front of the retina. These rays cross in the vitreous; when they reach the retina they have become divergent, forming a circle of diffusion and consequently a blurred image (Fig. 293, P P F). Certain divergent rays, coming from the myopic far point, are focussed on the retina (Fig. 293, D X) without accommodation.

The greatest distance at which the patient can read fine print is the far point. This is always at a definite distance corresponding to the amount of M.; the higher the M., the closer to the eye is the far point; the distance of the latter is the measure of the M. For example, if the far point is at 20 inches (0.5 metre) the $M. = 2 \text{ D.}$ ($\frac{1}{20}$ or $\frac{1}{500} = 2$); if at 10 inches (0.25 metre) the $M. = 4 \text{ D.}$ In these two instances concave lenses of 2 and 4 D. respectively would render parallel rays as divergent as if they came from a distance of 20 and 10 inches (0.5 and 0.25 metre); and with these lenses the myope would be able to see distant objects distinctly (Fig. 294).

Etiology.—Myopia almost always depends upon a lengthening of the antero-posterior diameter of the eyeball (axial myopia); in M. of 3 D., for example, the eyeball measures 24 mm. in its antero-posterior diameter, and in M. of 10 D., 27 mm. from before backward, instead of 23 mm., the normal diameter. Much less frequently M. is due to increased curvature of the cornea (anterior staphyloma and keratoconus), increase in the refraction of the lens from swelling in incipient cataract (often referred to by the laity as 'second




FIG. 293.—THE FOCUSSED OF PARALLEL AND DIVERGENT RAYS IN MYOPIA.

FIG. 294.—THE CORRECTION OF MYOPIA BY MEANS OF A CONCAVE LENS.

sight,' because it sometimes enables an old person again to read print for a time without glasses), and spasm of accommodation. The determining causes are associated with the demands which civilization and education make upon near vision. It is rarely congenital, though there is often an hereditary tendency for its development. It is an acquired change which commences at an early age when, during the developing period, the eyes are used excessively or improperly for near work. Its occurrence is in direct proportion to the standard of education, and also bears a certain relation to the general health and strength of the individual. It is much more common in cities than in the country.

Excessive study with insufficient outdoor exercise, fine or indistinct print, poor illumination, opacities of the cornea and other lesions causing imperfect vision, faulty construction of desks, sedentary habits, and poor health, are among the frequent exciting causes of myopia, especially in those who are predisposed.

The cause of the lengthening of the eyeball is attributed (1) to pressure of the extra-ocular muscles during excessive convergence causing the posterior pole, which is the least resistant part of the eyeball, to bulge; (2) to congestion, inflammation, and softening of the layers of the eyeball, together with increased tension, produced by fulness of the veins of the head as a result of stooping postures and other predisposing causes; and (3) to the shape of the orbit in broad faces causing excessive convergence, as seen in the German race, which is especially subject to this error of refraction.

Clinical Forms.—In most instances myopia is of low degree, develops during youth, and then comes to a standstill or increases very little; this is known as stationary or simple myopia.

In other cases the error reaches a considerable height in youth, and increases steadily up to the twenty-fifth year or even later, resulting in a high degree of myopia; this is known as progressive myopia. These are the cases which are accompanied by destructive changes in the choroid and other parts of the eye, leading to a considerable impairment of vision, and even blindness. In these cases myopia may

properly be considered a disease. Extreme cases of progressive myopia are known as malignant myopia.

Symptoms depend on the degree of myopia.

In slight degrees and in many cases of moderate amount there are often no symptoms except indistinct vision for distance. Near work can be accomplished with comfort; in fact, since the myope requires less accommodation than the emmetrope, he may have an advantage in close application. It is on this account that the circular fibres of the ciliary muscle are less developed than in the emmetropic eye (Fig. 292).

In other cases of moderate myopia and in high degrees, distant vision is very indistinct; there is often pain in the eyes after near use; the patient will be unable to continue at work for any length of time on account of excessive convergence; the eyes tire easily, are sensitive to light, and irritable; there are black spots before the eyes (*muscae volitantes*), and sometimes bright flashes of light. In some cases there may be absolute scotomata.

In high myopia there are often prominence of the eyes, a deep anterior chamber, and dilated pupils; the patient is apt to screw the eyelids together; there is sometimes an appearance of convergence. The strain of excessive convergence is so great and painful that the effort is sometimes given up and divergent squint results.

Ophthalmoscopic Signs.—In low (less than 3 D.) or moderate (3 to 6 D.) degrees there are frequently no changes except a crescent-shaped patch of atrophy of the choroid of whitish or grayish colour, embracing the outer side of the disc; this is called a myopic crescent.

In high myopia (more than 6 D.), a well-marked crescent is usually found, often posterior staphyloma (bulging of the sclera, Fig. 172, Plate XV.), and there may be patches of choroidal atrophy with pigmented margins, exposing the sclera. In progressive cases there are frequently added to these lesions atrophic and pigmentary changes in the macular region, hæmorrhages, especially at the yellow spot, fluid vitreous floating bodies in the vitreous, and opacities of

the lens; sometimes there is detachment of the retina. Owing to these changes the vision is often very markedly reduced, and is sometimes lost in severe forms of progressive myopia.

Tests.—*The Subjective Test with Test Types and Test Lenses.*—Distant vision is below the normal, and the patient requires a concave spherical lens to bring the sight up to $\frac{6}{6}$. The weakest lens which accomplishes this is the measure of the myopia. In young persons it is important to paralyze the ciliary muscle, so that spasm of accommodation will not cause the patient to select too strong a lens. The results are recorded as follows: R.E. $V = \frac{6}{60}$, $c - 4$ D. Sph. $= \frac{6}{6}$. The reduction in distant vision generally is proportionate to the degree of M.

The myope is able to read the smallest print, but at a shorter distance than that which the emmetrope selects. The farthest distance at which he is able to read the finest print is his far point, and this is also a measure of his M. (p. 350).

The Ophthalmoscope at a Distance shows an inverted image of the fundus which appears to move in the opposite direction to the examiner's head.

The Ophthalmoscope, Indirect Method.—The disc appears small, and seems to increase in size upon withdrawing the objective lens.

The Ophthalmoscope, Direct Method.—The fundus cannot be distinctly seen until a concave lens is placed behind the mirror; the weakest concave lens with which the details are seen clearly indicates the amount of myopia.

Retinoscopy.—With the plane mirror and the observer at 1 metre distance, the shadow moves in the opposite direction (except when M. is less than 1 D.), and is reversed by the addition of concave lenses. The lens which causes reversal plus -1 D. is the measure of the M. In high M. the shadow is very faint, but becomes plainer when concave lenses are added.

Prognosis.—In low and moderate degrees of stationary myopia the prognosis is good. Progressive myopia is always a serious condition, especially when the choroïdal and vitreous

changes are marked; it may necessitate absolute cessation of all near work. In malignant myopia the prognosis is grave.

Treatment consists in prescribing glasses where they are required and in avoiding anything which may lead to an increase in the myopia.

In general terms, it is proper to give a full correction for low and moderate myopia in young persons as soon as discovered, and to direct these glasses to be worn for both distance and near work; this places the eyes under normal conditions of vision and accommodation. The glasses must be prescribed after the accommodation has been paralyzed, so that there will be no danger of over-correction on account of spasm of accommodation. Full correction corresponds to the weakest concave spherical lens which, with accommodation paralyzed, gives normal vision. In low degrees of M. an adult may be allowed to read without glasses if he finds this convenient.

In high myopia the full correction is prescribed for distance; 2 D. to 3 D. under-correction may be required for near work; the reading-glasses should be such as to enable the patient to read at a comfortable distance, say 13 inches (33 cm.). Suppose -10 D. gives the best vision for distance; then -10 D. + 3 D. Sph. = -7 D. will enable him to read at this distance without accommodation. Adults who have previously been much under-corrected often cannot wear their full distance correction. They are frequently more comfortable with 1 D. less than their full myopic correction in cases of about 10 D. of short sight or 2 D. less in cases of about 20 D.

After the age of forty-five the distance glasses cannot be worn for near work, since the convex lenses usually required for presbyopia must be added to the concave lenses, thus reducing the strength of the latter.

In prescribing glasses in M. every case must be considered on its merits. Many myopes wear strong lenses, representing the full correction, constantly and with absolute comfort; this is more usual when they have been kept fully corrected since childhood; others require two sets of lenses, one for distance and a weaker pair for reading.

In order to check any tendency to increase of M., rigid

hygienic rules, both local and general, should be carried out. These are of especial importance in the young.

The patient's habits should be regulated so that he will enjoy good health. He should have plenty of outdoor exercise and sufficient sleep. Constipation should be avoided.

In progressive myopia near work should be restricted, and the patient not be allowed to read too long at a time. The book should be held at not less than 13 inches (33 cm.). In most cases the full correcting lenses should be worn for near work. The illumination should be good, neither too bright nor too dim, and should come from behind; the myope should avoid reading at dusk or with feeble illumination; the amount of work done with artificial light should be limited. The print should be large and clear, with ample spacing. Desks should be constructed so that the sitting posture is comfortable, and so that the child is not encouraged to stoop over his books; the myope must be taught not to bend over his work, but to lift the latter to the required distance from the eyes. In London and some of the larger towns in Britain special 'myope schools,' in which all these recommendations are put into practice, are available for such children. Teaching is largely oral.

If, notwithstanding such precautions, myopia progresses rapidly, and, especially if there are choroidal changes it is necessary to forbid all near use of the eyes. It may be necessary to take the patient from school and send him to the country for a long period, during which he is instructed to be out of doors as much as possible, and to avoid all reading and near work. Young adults suffering from rapidly progressive myopia should give up sedentary occupations necessitating close application, and select those in which but little near use of the eyes is required. On the other hand, myopes usually prefer, and are best fitted for, sedentary occupations, so studies should not be entirely interrupted without real necessity.

Telescopic Spectacles are occasionally prescribed for improving the vision of myopes of very high degree, as well as for others with very poor vision which cannot be improved with ordinary lenses. They act on the principle of opera glasses, magnifying

about two diameters, and thus improving vision. But the number of individuals who can be aided by them is exceedingly small, since such spectacles limit the field of vision and often cause distortion and apparent movement of objects upon rotating the head; also they are heavy, and expensive.

Operative Treatment.—In children and young adults with high myopia, uncomplicated by excessive pathological changes in the fundus, the removal of the lens by discission and subsequent extraction used to be advocated and gave good results in some cases. The lens was needled, and after several days the swollen lens substance was removed by extraction. The operation was limited to M. of at least 15 or 20 D. After the removal of the lens the eye may be almost emmetropic, since the optical effect in such highly myopic eyes is quite different from that which would follow if the lens were removed in an emmetropic eye; a weak convex glass may be required for distance, and a stronger one for near work, since the accommodation has been sacrificed. Owing to the large number of cases of detached retina which have been seen to follow this operation, after some years have elapsed, it has been practically entirely abandoned in Britain. In any case it is never advisable to operate on both eyes.

Astigmatism.

Astigmatism is that refractive condition of the eye in which there is a difference in the degree of refraction in different meridians; each of the principal meridians has, therefore, a different focus (Figs. 297 to 301).

In E., H., and M., rays coming from a luminous point are brought to a single focus at a certain distance behind the cornea. In astigmatism, since the refracting surfaces are not spherical, rays from a luminous point are brought to a focus at different points; the shape of the image may be a line, an oval, or a circle, but never a point.

Varieties.—Astigmatism may be divided into (1) irregular, comparatively infrequent, and (2) regular, very common.

Irregular astigmatism consists in a difference of refraction

in different parts of the same meridian. It is generally due to changes in the cornea, such as opacities and cicatrices following ulceration, injuries, or surgical operations, and keratoconus. It may also result from partial dislocation of the lens, or from a congenital or acquired change in the refractive power of different sectors of the lens. The acuteness of vision is considerably diminished, and cannot be improved materially by glasses. Details of the fundus when seen with the ophthalmoscope appear distorted. An insignificant amount of irregular astigmatism is present normally, and accounts for our seeing the stars as stellate points instead of round dots.

Regular Astigmatism.

Regular astigmatism is that form in which, though the refraction in a meridian is the same throughout, there is a difference in the degree of refraction of the two principal meridians. In other words, the curvature of the cornea is different in the two meridians; these are called the principal meridians, and are always at right angles to each other; one exhibits the maximum and the other the minimum refraction. When the term 'astigmatism' is used without qualification it refers to regular astigmatism.

Etiology.—Astigmatism is usually due to a change in the curvature of the cornea, with or without some shortening or lengthening of the antero-posterior diameter of the eyeball. It is also caused, in part at least, by defects in the curvature of the lens; this lenticular astigmatism may partly neutralize that of the cornea. It is usually congenital, and there is often an hereditary tendency. It may, however, be acquired, and is then caused by corneal changes resulting from inflammation, injury, or operation. Pressure of the lids in ametropia is believed by some surgeons to be capable of producing permanent regular astigmatism.

Refraction of Rays in Regular Astigmatism.—Parallel rays refracted by a spherical surface form a circular cone, and come to a focus at a point. In astigmatism, those rays which pass through the meridian of greater curvature come to a

focus sooner than those which pass through the meridian of lesser curvature, and the resulting cone will not be circular, but more or less oval; hence the vision of astigmatic subjects is not simply indistinct, but the diffusion images are more or less elongated.

In looking at straight lines (which are made up of a succession of points), these may appear distinct or indistinct to astigmatic persons according to their direction. If an astigmatic eye, in which the vertical meridian is out of focus and the horizontal meridian normal, looks at a vertical line, this will be slightly elongated; but the sides will appear distinct, since each point of light will be seen as a small vertical line, and these overlap each other. But if such an eye looks at a horizontal line each point of light will again be seen as a small vertical line, and consequently the line will appear blurred (Fig. 295). There is, therefore, one direction in which straight lines appear most distinct, and another, at right angles to it, in which they appear most indistinct; this forms the basis for the construction of the astigmatic dial or fan (Fig. 302) commonly used as a test for this error. The lines parallel with the ametropic meridian are seen most clearly, and those parallel with the emmetropic meridian are seen most indistinctly (in simple As.).

Varieties of Regular Astigmatism.—According to the refraction of the principal meridians, astigmatism is divided into:

1. *Simple*, in which one meridian is emmetropic, and the other hypermetropic or myopic; it comprises simple hypermetropic astigmatism (Fig. 297), and simple myopic astigmatism (Fig. 298).

2. *Compound*, in which both meridians are either hyperopic




FIG. 295.

FIG. 296.

FIG. 295.—VERTICAL AND HORIZONTAL LINES AS SEEN BY AN ASTIGMATIC EYE IN WHICH THE HORIZONTAL MERIDIAN IS EMMETROPIC.

FIG. 296.—VERTICAL AND HORIZONTAL LINES AS SEEN BY AN ASTIGMATIC EYE IN WHICH THE VERTICAL MERIDIAN IS EMMETROPIC.

or myopic, but unequal in degree; it comprises compound hypermetropic astigmatism (Fig. 299), and compound myopic astigmatism (Fig. 300).

3. *Mixed*, in which one meridian is hypermetropic and the other myopic (Fig. 301).

In most cases of astigmatism the cornea presents its




FIG. 297. — SIMPLE HYPERMETROPIC ASTIGMATISM.




FIG. 298. — SIMPLE MYOPIC ASTIGMATISM.

maximum curvature in or near the vertical meridian, and the least curvature in or near the horizontal meridian, corresponding to the slight astigmatism of the normal eye; when this is the case it is said to be astigmatism with the rule; when the relative curvatures are reversed it is astigmatism against the rule. In astigmatism with the rule the axis of the cylinder is vertical or nearly so in hypermetropic astig-




FIG. 299. — COMPOUND HYPERMETROPIC ASTIGMATISM.




FIG. 300. — COMPOUND MYOPIC ASTIGMATISM.

matism, and horizontal or nearly so in myopic astigmatism. The chief meridians, though vertical and horizontal in the majority of cases, may occupy an oblique position; in such cases they are most frequently symmetrical—*i.e.*, inclined an equal number of degrees from the vertical or horizontal on each side.

Symptoms.—With very small degrees of astigmatism, there may be no reduction in the acuteness of vision, but with greater

degrees this is always found both for distance and near. There is commonly considerable asthenopia, especially upon use of the eyes for near work. These asthenopic symptoms are similar to those occurring in hypermetropia (p. 360), but are apt to be more pronounced and more continuous. They vary with the degree and variety of astigmatism, the amount of near work indulged in, and especially the state of the patient's health; a small amount (0.50 D. or even 0.25 D.) will, for instance, often give rise to severe asthenopic and nervous symptoms in a delicate, neurasthenic individual. The involuntary accommodative efforts of the ciliary muscle, made to diminish the effects of the error, cause continuous eye-strain and explain the frequency of asthenopia.

The Correction of Astigmatism.—Astigmatism is corrected by cylinders, sphero-cylinders, or crossed cylinders. The curve of




FIG. 301.—MIXED ASTIGMATISM.

the correcting cylinder corresponds to the ametropic meridian; consequently its axis is at right angles to this meridian.

Tests.—We usually suspect astigmatism when the vision cannot be brought up to $\frac{6}{6}$ with spherical lenses, notwithstanding the fact that the fundus is normal and the media are clear. In testing for astigmatism in children, it is necessary to use a cycloplegic. In young adults this is frequently advisable, in older patients less often required. The lower the degree of the astigmatism and the less the experience of the surgeon, the more often should a cycloplegic be used; results are otherwise apt to be unsatisfactory.

The Astigmatic Dial.—The diagnosis of astigmatism is made if the patient, when placed before the dial (Fig. 302), which is formed of radiating lines in different meridians, is unable to see all the lines with equal distinctness. To isolate the radial

lines of the plus axis with greater ease, the patient, if not already myopic, should be made so by putting up plus spheres before each eye separately sufficiently to make these lines clear,




FIG. 302.—ASTIGMATIC DIAL CHART.

while the others fade into grayness in contrast. To make it easier to determine the exact line which is the clearest, a movable **V** is attached to many dials. The patient's attention is directed to the two arms of the **V** and he is asked to say which appears to be the clearer of the two. The **V** is then rotated in this direction until

the two arms appear to be equally gray, the clearer arm fading after passing the position of the blackest axis on the dial, which is now at the point of the **V** and indicated at the side of the chart. The indicator at the bottom of the chart will now also automatically give the axis of the minus cylinder, whose strength, the measure of the astigmatism, brings the other principal radial axis in tone with the first. It now but remains to clear the vision for the test types by further defogging with minus spheres.

The metal disc with stenopæic slit (about 1 mm. in diameter) may be used to discover the two principal meridians. It is placed in front of one eye, the other being excluded, and is rotated slowly so that the slit occupies each meridian successively. The patient is placed at 6 metres before the distant test types, and the position of the slit in which the best vision is obtained is noted. Then convex or concave lenses are placed in front of the slit, and the strongest convex

or the weakest concave lens which gives the most improvement is the measure of the refraction in this meridian. The slit is then turned 90 degrees, and convex and concave lenses are again applied until one is found which improves vision most. In this way the refractive error of the two principal meridians is determined. If, for instance, when the slit is vertical the patient reads $\frac{6}{8}$, and convex lenses in front of the slit make the types indistinct, the vertical meridian is emmetropic; if, when the slit is horizontal, the patient reads $\frac{6}{18}$, but this increases to $\frac{6}{8}$ when +3 D. Sph. is placed in front, the horizontal meridian is hypermetropic 3 D. This case would be one of simple hypermetropic astigmatism corrected by a +3 D. cylinder, axis vertical.

The Subjective Test with Test Types and Test Lenses is best employed after the objective tests have furnished us with pretty definite conclusions regarding the correcting lenses. It then serves to confirm or improve upon the results obtained by objective methods. The lenses selected with the objective tests are placed in the trial frame and may require modification, either in the strength of the spheres or the strength and axis of the cylinder, so as to secure the most acute vision.

The Ophthalmoscope, Indirect Method.—The shape of the disc is oval instead of circular, and changes when the objective lens is withdrawn.

The Ophthalmoscope, Direct Method.—The disc appears oval, the elongation corresponding to the meridian of greatest refraction, and is at right angles to the long axis of the oval seen with the indirect method. To determine the kind and amount of error we estimate the refraction of a small vertical vessel, and then of a small horizontal vessel near the disc, by means of the strongest convex or the weakest concave lens with which these are distinctly seen. For instance, suppose a vertical vessel is seen clearly with +2 D. Sph., (indicating hypermetropia of horizontal meridian), and a horizontal vessel with +4 D. (indicating a greater amount of hypermetropia in the vertical meridian), the case is one of compound hypermetropic astigmatism. When the principal

meridians are oblique, we find a vessel the direction of which corresponds to one of the meridians, and then another at right angles to the first, and estimate the refraction of each.

Retinoscopy is the most rapid and reliable objective method of determining astigmatism. The principal meridians are clearly indicated by the edge of the shadow (Fig. 270). Each of the principal meridians is corrected separately by causing a reversal of the movement of the shadow by spherical lenses, and adding -1 D. (with plane mirror at 1 metre distance).




FIG. 303.—THE JAVAL-SCHIÖTZ OPHTHALMOMETER.

The Ophthalmometer (Fig. 303) is an instrument used for determining the principal meridians and the amount of *corneal* astigmatism. It is only of service when used in connexion with other tests. It consists of a telescope containing a combination of convex lenses and a bi-refracting prism, supporting a graduated arc upon which are two sliding objects called 'mires.' The latter are of white enamel, one

quadrilateral in shape, the other of similar size but cut out on one side into steps; both are divided in the middle by a horizontal black line. The patient's face is placed in a frame at the other end of the instrument, and steadied by chin and forehead rests. The mires are reflected upon the cornea, and the observer, looking through the tube and focussing, sees four images in a line. The two peripheral images are ignored; the two central ones are approximated until their inner edges touch, and the black lines subdividing the mires form one continuous straight line; it may be necessary to revolve the barrel of the telescope more or less of 45 degrees to the right or left to accomplish this. This position, indicated on a dial,




FIG. 304.




FIG. 305.



FIG. 306.



FIG. 307.



FIG. 308.

FIG. 304.—THE MIRES OF THE OPHTHALMOMETER INDICATING AN ABSENCE OF CORNEAL ASTIGMATISM.

FIG. 305.—THE OVERLAPPING OF THE MIRES OF THE OPHTHALMOMETER INDICATING 1 D. OF CORNEAL ASTIGMATISM.

FIG. 306.—CONVEX REFLECTION OF PLACIDO'S DISC IN EMMETROPIA.

FIG. 307.—THE SAME IN REGULAR ASTIGMATISM OF VERY HIGH DEGREE.

FIG. 308.—THE SAME IN IRREGULAR ASTIGMATISM.

gives the meridian of least refraction. Next the arc is turned at right angles to this meridian. If the images of the mires are still in apposition, the curvature of the cornea is uniform, and there is no corneal astigmatism (Fig. 304). If in the second meridian the relative position of the images of the mires has changed, each step which is overlapped by the quadrilateral figure indicates 1 D. of astigmatism (Fig. 305).

Placido's disc or keratoscope (Fig. 6, p. 5) may give a useful qualitative test. If no astigmatism is present the rings are circular. If regular astigmatism exists the rings will appear elliptical with the long axis corresponding to the meridian of least curvature. If the cornea is the seat of irregular astigmatism the rings will be distorted.

The Cross Cylinder is used by some as a contrast or checking lens to test the accuracy of the strength and axis of the cylinder chosen as a result of other objective and subjective methods. The cross-cylinder has the effect of a convex cylinder and of a concave cylinder of equal strengths with their axes at right angles to each other; the most useful combination is made by combining a $+0.25$ spherical with a -0.50 cylinder; this combination is mounted in a ring attached to a round handle, permitting easy rolling between thumb and index finger; the handle is attached at 45 degrees from the axis of either cylinder.

To check the strength of cylinder selected.—After correction by other tests, the cross cylinder is held an inch or two in front of the selected correction, with the axis of its cylinder corresponding to that of the cylinder in the trial frame; the patient reports whether this increases or diminishes the clearness of the lowest line of test-chart letters which he can see; then the cross cylinder is revolved so that the other cylinder faces towards the eye, and again the patient reports whether blurring or clearing results. In this way, information is obtained leading to an increase or decrease in the strength of the cylinder in the trial frame.

To check the selected axis of the cylinder in the trial frame, the cross-cylinder is placed before the eye with the axis of one of its cylinders 45 degrees from that of the selected cylinder from the trial case, then by rotating the handle the position of the axes is reversed. The patient now reports which axis of the cross-cylinder makes the letters more distinct and which makes them less distinct; the axis of the cylinder in the trial frame is then rotated in the direction of that cylinder which has made the letters most distinct, plus towards plus, minus towards minus, until the most perfect position of axis is obtained.

When the cross-cylinder renders the types equally indistinct in both positions, the strength and axis of the trial cylinder will have been proved correct.

The cross-cylinder may also be of use in arriving at the estimation of the amount and axis of astigmatism, where the usual means fail, such as cases of immature cataract, opacities

of the vitreous, etc., in which retinoscopy is unsuccessful, thus enabling us to improve poor vision to a certain, even though limited, degree; in such cases higher cross cylinders are used.

Treatment consists in prescribing glasses which correct the error. In many cases of moderate or high degree it is impossible to obtain V. $\frac{6}{8}$ with the full correction; we may have to be satisfied with $\frac{6}{9}$ or $\frac{6}{12}$. But the vision often improves after the lenses have been worn for a time. The glasses should be worn constantly if required to improve vision or relieve symptoms. When the correction has been estimated with the eye under the effects of a cycloplegic, a slight reduction may perhaps be necessary in cases of moderate or high degrees of astigmatism, but as a rule the full correction




FIG. 309—NOTATION OF THE AXIS OF CYLINDERS.

will be tolerated. The relief which cylinders give is usually very pronounced.

The direction of the axis of a cylinder should be indicated by the standard notation as adopted by the International Congress, and recognized all over the world.

The axis is denoted by the angle which it makes with the horizontal meridian, the angles being numbered from 0 degree on our left side (as we stand before the patient and counting downwards) to 180 degrees on our right side (Fig. 309)—*i.e.*, commencing at the temporal side of the right eye and the nasal side of the left eye of the patient.

Prescribing Glasses for Young Patients.

Having now discussed the correction of refractive error as it affects the eye, certain practical considerations demand attention.

The advantages one hopes to gain in prescribing glasses may be to prevent injury to the eye, to cure certain affections, such as squint or blepharitis, to relieve headaches or other subjective symptoms, or to improve sight. On the other hand, there are certain disadvantages; for example, the fact of a boy wearing glasses constantly at school handicaps him to some extent in sports. In the case of older girls the appearance may be a consideration. One should only prescribe glasses when the advantages to be gained outweigh any disadvantages.

Many boys and girls at school complain of headaches, and are very properly sent to an ophthalmic surgeon for an examination of the eyes. There may be a definite (often low) refractive error which needs correction; but frequently one finds an insignificant degree of hypermetropia which could not be the cause of the headaches. Many surgeons are in the habit of prescribing glasses for these children, whereas the cause of the headaches should be sought for in another direction.

In the lower degrees of myopia it is permissible to allow glasses to be discarded for many sports, and on special occasions—*e.g.*, a dance; otherwise they should be worn continuously to give the child all the advantages of normal sight.

If there is astigmatism the subjective symptoms should be the chief factor in deciding whether glasses are to be worn constantly or no.

Anisometropia.

This term is applied to cases of marked inequality in the refraction of the two eyes; slight differences are present in most cases of errors of refraction. Every possible combination may occur: (1) One eye may be emmetropic and the other ametropic; (2) both eyes may have the same variety of ametropia, but of unequal degree; (3) one eye may be myopic and the other hypermetropic, either simple or combined with astigmatism. Notwithstanding the unequal refraction, there is usually binocular vision; sometimes the eyes are used

alternately, and in other cases one eye is habitually excluded from vision.

In prescribing glasses ~~no arbitrary rules~~ can be followed; each case must be considered by itself. When one eye is emmetropic and the other ametropic no glass will probably be required, unless it be to prevent the ametropic eye from suffering from disuse, or for the relief of asthenopic symptoms. When the difference in the refraction is not great (1 to 2 D.) and there is good binocular vision we may give each eye its correction. When the difference is greater the full correction sometimes causes discomfort, then we must be satisfied with a partial correction. When there is no binocular vision we generally give a correcting glass for the better eye. In such cases, if the poor eye still possesses vision, the patient should be advised to exercise it daily with the aid of a suitable lens, the good eye being excluded, so that the amblyopic eye may retain its vision and the defect be prevented from becoming worse.

Asthenopia.

Asthenopia, weak sight, or eye-strain, is a convenient term which embraces the group of symptoms dependent upon fatigue of the ciliary muscle or of the extra-ocular muscles.

Symptoms.—The condition is of very frequent occurrence, and causes a great variety of symptoms. The most common manifestations of asthenopia are: (1) *Pain* in or around the eyes or headache, usually aggravated by use of the eyes for close work, and in some cases present only after near use. (2) *Fatigue and discomfort* upon use of the eyes for near work; this shows itself by inability to indulge in such work for more than a short period at a time (without the occurrence of dimness of vision and confusion of the lines of print), pain in and about the eyes, headache, drowsiness, lacrymation, photophobia, and congestion, and an irritable condition of the lids accompanied by itching and burning sensations. These symptoms are regularly worse at night, when the patient is tired, or when artificial illumination is employed. (3) *Vertigo* and a tendency to diplopia. (4) *Nervous affections*,

such as migraine, nausea, twitching of the facial muscles, chorea, etc.

The amount of asthenopia depends not only upon the degree of defect, but also upon the state of the patient's health, and is therefore pronounced in delicate, anæmic, and neurasthenic individuals.

Varieties.—(1) Accommodative; (2) muscular, (3) nervous or neurasthenic (reflex). Two of these varieties may be associated.

Accommodative Asthenopia is the most common variety. It is due to strain and fatigue of the ciliary muscle when used too constantly or excessively in ametropia. It is especially frequent in astigmatism and hypermetropia, but is common enough in myopia and in presbyopia. Treatment consists in the use of glasses correcting the error of refraction, as advised in preceding pages. In delicate and neurasthenic individuals attention to the general health is very important.

Muscular Asthenopia is due to a want of balance of the motor apparatus of the eyes (heterophoria). It may be associated with ametropia, and its existence be dependent upon the latter error, or it may occur in emmetropia. It is often due to myopia, in which, on account of the far point being close to the eye, the patient is obliged to converge excessively. Heterophoria is fully discussed in Chapter XXVIII.

Nervous, Neurasthenic, or Reflex Asthenopia is the variety which occurs in emmetropic patients or in ametropes in whom proper correcting lenses give no relief. It is a neurosis, and is dependent upon a general asthenic condition of the nervous system; consequently it is found most frequently in young women with hysterical tendency, who suffer from anæmia, neurasthenia, and often menstrual disorders; also in neurasthenic individuals in general, and in convalescents from debilitating diseases. The condition is often very troublesome and frequently obstinate. The more carefully one investigates the static and dynamic refraction and the motor balance of the eyes, the fewer cases one finds it necessary to class as neurasthenic. Treatment consists in removing the defect in

the general condition, rest of the eyes, and particularly attention to hygiene, such as the regulation of habits, outdoor exercise, etc.

Mydriatics and Cycloplegics.

The action of these agents and the method of obtaining the best results with them are described in Chapter XXXI.

A cycloplegic is indicated in the estimation of the refraction in all cases of children and frequently in young adults; much less often is it needed in an older patient. The frequency with which mydriatics become necessary in adults varies inversely with the experience of the surgeon. Before using these agents in elderly persons any suspicion of *glaucoma* must be excluded.

Homatropine (2 or 3 per cent. solution), or homatropine 2 per cent., combined with cocaine 1 per cent., is the agent most frequently employed; one drop is instilled every five or ten minutes for three or four doses, and the examination begun at the end of half an hour after the last instillation. Sometimes homatropine fails to produce complete paralysis of accommodation, as shown by more or less contradiction in the results of the objective and subjective tests. In such cases we may resort to atropine (1 per cent. solution), one drop being instilled three times daily for two or three days (smoked glasses may be worn during this period), and a final drop directly before the examination.

In the case of young children *atropine* drops or ointment (1 per cent.) should be used twice daily for three or four days before the examination.

The Fitting of Eyeglasses and Spectacles.

Much of the comfort and relief which lenses bring will depend upon the skill with which the glasses are fitted to the face. Whether the surgeon orders eyeglasses or spectacles, the lenses must be supported in their frames in such a manner that the distance between their geometric centres corresponds to the interval between the centre of the pupils (interpupillary distance).

If the glasses are to be worn constantly, the geometrical centre of the lenses should be slightly below the centre of the pupils, and the lenses tilted so that their surfaces form an angle of about 5 or $7\frac{1}{2}$ degrees with the plane of the face. If worn for distance only the level of the lenses should be the same, and the tilting about 5 degrees. If worn for near work only the lenses should be lower, and inclined about 10 or 15 degrees.

In every case the glasses should be worn as near the eyes as possible, just avoiding the lashes.

In cases of astigmatism it is necessary that the axis of the cylinder be constant. On this account spectacles are often preferable to eyeglasses, because with the latter the axis of the cylinder may vary according to the way in which the glasses are worn or in which they preserve their original adjustment. But eyeglasses can be worn in such cases if the optician exercises sufficient skill in fitting. Lenses for children, having a cylindrical element, should not be quite round, lest they become slack in the frame and allow the axis to rotate into a wrong position.

Lenses are usually made of crown glass. Periscopic lenses (p. 326) are preferred, since they give better definition of the peripheral part of the field when the eyes are moved from side to side. In simple cylinders one surface is generally plane and the other curved, and sphero-cylinders usually have the spherical power on one surface and the cylindrical power on the other; but lenses containing cylinders can be made in the concavo-convex form. They are worn with the convex surface away from the eye. Such lenses are known as torics. Modern toric lenses, such as isophane, give an enlarged field. At one time lenses were frequently cut from crystal and were known as pebbles. They were less easily scratched than ordinary lenses, and until recently were used for sports, as they were less likely to break. With the introduction of safety glass, lenses known as 'Sale' are now made which, when broken, do not fly into dangerous cutting splinters, as they are held together by the laminating material.

Bifocal lenses consist of an upper portion of one focus and a lower part of another. They are used principally in cases of

presbyopia associated with ametropia, the lower portion being used for reading and near work, and the upper for distance. The least expensive bifocal lenses are those in which the




FIG. 310.—CEMENTED BIFOCAL LENS, OVAL READING SEGMENT.




FIG. 310A.—CEMENTED BIFOCAL LENS, CIRCULAR READING SEGMENT.

addition consists of a thin oval or circular lens cemented to the lower portion of one surface of the distance glass (Figs. 310 and 310A). This form is comparatively cheap, but the edge of the cemented lens is very obvious both to the wearer and to others. Moreover, changes of temperature sometimes cause cloudiness in the cement joining the two lenses.

Modern bifocals, although more expensive than the old cemented form, are more comfortable to wear and are rightly known as invisible bifocals. They are made in two ways, one by fusing (Fig. 311) and the other from solid crown glass. Reference to the diagram of the fused bifocal will show that its surfaces are continuous. The power of the reading portion is obtained by using a flint glass of higher refractive index than the crown glass into which it is fused. The curvature of the fused surface controls the additional power. The solid bifocal has the power of the reading portion




FIG. 311. FIG. 311A.

FIG. 311.—SECTION OF FUSED BIFOCAL LENS.

FIG. 311A.—SECTION OF ONE-PIECE BIFOCAL LENS.

ground on to the surface of the lens, hence its humped appearance in the diagram (Fig. 311A).

Bifocal glasses need especial care in fitting. Some patients find great difficulty in wearing them constantly, as objects on the ground, seen through the reading segment, are blurred. Others readily accustom themselves to their use. To obviate this difficulty varieties of bifocal lenses are obtainable which leave a small portion of the distance correction below the reading segment.

Trifocal lenses are very occasionally ordered; in these the upper part is adapted for distant vision, the lowermost for close work, and a central portion for vision at an intermediate distance.

Protective Glasses.—These are of two kinds: (1) To prevent discomfort and ill-effects from excessive light; and (2) to guard against injury, especially from foreign bodies, but also against other mishaps in industrial occupations, or in sports.

Smoked glasses should not be prescribed without good reason, but are sometimes necessary. Some people, apart from refractive errors or disease of the eye, are hypersensitive to ordinary light; they may be more comfortable if they wear lenses which absorb a large part of the ultra-violet light without keeping out visible rays. Such special lenses relieve glare; the ones most frequently used are Crookes' A, having a faint gray tint, and Soft Lite No. 1, having a slightly pinkish hue. Deeper shades of these special lenses are prescribed to guard against very brilliant light such as is experienced in the tropics, amongst snow, etc.

In many diseases of the cornea, uvea, and retina the eye must be protected from light, especially when the pupil is kept dilated; this is accomplished by smoked glasses, worn alone, or over the distance lenses ordinarily used. The shade of these is designated by numbers, 1 being the lightest, 6 the darkest. No. 3 is the one usually prescribed. Yellow, greenish-yellow, and amber glass is also used for the same purpose, though less often; such coloured glass is known by trade names, 'Feuzal,' 'Euphos,' etc. With very intense light, such as is experienced with electric welding, deeply coloured glasses are worn, often

in the form of a plate of red glass covering a similar plate of green or blue.

For industrial uses great strides have been made of late in the development of reflecting glass as distinct from absorption glass. Reflecting glasses are made by thinly depositing metal (either platinum, gold, silver, or aluminium) on the surface of a glass which is protected by the addition of a cover glass. The layer of metal is so thin that it is possible to see through it.

2. To protect the eyes from flying particles of metal in industrial occupations, goggles should be worn; these should be light in weight, provide for replacement of scratched lenses, and have wire-mesh side-protection; the glass should be of the non-splinterable variety. Lenses made of this kind of glass are also useful for individuals exposed to risk of injury from broken spectacles in such sports as cricket, tennis, etc.; such lenses can now be made much more accurately to prescription than formerly, and their weight is little more than that of ordinary lenses.

CHAPTER XXV

ANOMALIES OF ACCOMMODATION

Presbyopia.

PRESBYOPIA, or old sight, is a physiological change which affects every eye, becoming evident usually at about the forty-fifth year, as a result of which the near point recedes beyond the distance at which we are accustomed to read ordinary print. The change is due chiefly to loss of elasticity of the lens, whereby it is prevented from responding to the action of the ciliary muscle; consequently the power of accommodation is lessened. As explained on p. 342, this diminution in the power of accommodation begins early, about the tenth year. When it becomes sufficient to interfere with the comfortable exercise of near vision, presbyopia is said to be present.

At the age of forty there are 4.5 D. of accommodation, and the near point is at 22 cm., or 9 inches. To read at 9 inches such an individual would require all of his accommodation, and the effort would soon become fatiguing, since only one-half or two-thirds of this power can be used for any length of time without causing asthenopic symptoms. Generally, however, the adult holds print at about 13 inches (33 cm.), requiring 3 D. of accommodation, and leaving a reserve of 1.50 D.; this is usually sufficient to ensure comfort. At forty-five his accommodation has diminished to 3.5 D.; all or nearly all of this would be required to read comfortably at 13 inches, leaving little or no reserve. If he keeps one-third of his accommodation in reserve he will have about 2.25 D. available for near work; with this his reading distance would be 45 cm., or 18 inches—too great for comfortable and continuous near

work. Hence we must supply the defect in the power of accommodation by a convex lens of sufficient strength to bring the near point back to a convenient distance.

Symptoms.—The presbyope is compelled to hold reading, writing, sewing, and other forms of near work farther away than the usual distance, making such efforts uncomfortable. With this recession of the near point beyond the usual situation the print becomes pale and indistinct, and fine type can be read only with great difficulty. The patient is apt to use strong illumination; this produces contraction of the pupil, and thus improves the definition by diminishing the circles of diffusion. If the condition be uncorrected he suffers from asthenopic symptoms, especially pain, fatigue, lacrymation, dimness of vision, and irritation of the lids, all of these symptoms being more marked in the evening with artificial illumination. Presbyopia has no effect upon distant vision in an emmetrope.

Treatment consists in prescribing convex spherical lenses for near work, so as to compensate for the lack of power of accommodation, and to bring the near point back to a comfortable working distance.

One can generally make a rough guess as to the correcting glasses according to age. The following represents the average for a healthy emmetropic Englishman:

Age 45	...	...	Glass required	+1 D.
" 50	...	...	" "	+1.5 D. to +1.75 D.
" 55	...	...	" "	+2.25 D.
" 60	...	...	" "	+2.75 D. to +3 D.
" 65 and over	...	...	" "	+2.75 D. to +3.5 D.

The numbers given are somewhat arbitrary. A tall, upright person will generally require a smaller presbyopia correction than a short person or one who stoops. The oft-quoted rule to allow +1 D. for every five years after forty has led many inexperienced surgeons astray. Apart from an error of static refraction, when a patient complains that he cannot read comfortably with his recently purchased glasses, one generally finds that they are too strong. The age at which patients are obliged to wear glasses varies within a few years, and is influenced to a certain extent by the vigour of the individual;

a delicate or neurasthenic person will require glasses for reading earlier than a robust individual.

The glasses must also be selected with reference to the occupation or the special use for which the patient wishes them. Thus in reading, writing, and sewing 13 inches (33 cm.) is a comfortable working distance for most persons; but a musician may prefer a distance of 20 to 25 inches, and consequently he will require a weaker glass.

The existence of ametropia will modify the strength of glasses required for presbyopia. Hence the patient's vision for distance, and his refraction, must be determined before estimating the glasses required for near work. In any case of ametropia the lenses required for distance must be added to those which would be selected for presbyopia in the emmetrope. This would have the effect of increasing the strength of the convex lens required for presbyopia in cases of hypermetropia, and of diminishing its power in myopia. For example, suppose a patient of fifty has hypermetropia of 1.75 D.; his glasses for reading would be H. 1.75 + Pr. 1.75 D. = +3.50 D. A myope of 1.5 to 2 D. will require no glass at fifty, since his myopia and presbyopia neutralize each other. If the myopia amounts to 4 D., the patient will never require glasses for reading, since his far point will never be more than 10 inches. In astigmatism the cylinders must be added to the convex lenses required for the correction of presbyopia.

Since presbyopia increases with age, the glasses will require changing for stronger ones every few years. *When the glasses have to be changed for stronger lenses very frequently, we should suspect glaucoma*, and examine the eye carefully for this disease.

Paralysis of Accommodation.

Paralysis of accommodation (cycloplegia) is a partial or complete loss of power in the ciliary muscle due to paralysis or paresis of the third nerve, or of that branch of the motor oculi which supplies the ciliary muscle and iris. Though occasionally confined to the ciliary muscle, the paralysis usually includes the sphincter pupillæ. When limited to the ciliary muscle and iris it is known as ophthalmoplegia interna.

Etiology.—The most frequent cause is the use of mydriatics, such as atropine and homatropine. It may be part of a complete paralysis of the third nerve. It occurs not infrequently after diphtheria. Other causes are contusions of the eyeball, debilitated states of the system, influenza, syphilis, diabetes, and cerebral disease.

Symptoms.—These are loss of power of accommodation and dilatation of the pupil. If emmetropic, the patient will have good vision for distance, but be unable to do near work without convex glasses. If hypermetropic, both near and distant vision will be impaired. If myopic, the patient will be able to see distinctly only at his far point; he may therefore be able to do without his accommodation if the myopia is considerable.

Prognosis is usually good, especially when the affection is due to syphilis, diphtheria, or the use of a mydriatic. In traumatic cases the condition may be permanent.

Treatment.—One should attempt to remove the cause of the paralysis. In syphilis, specific treatment is indicated. In post-diphtheritic paralysis, and in that due to debilitated conditions, tonics should be given, especially strychnine. Locally, the myotics (eserine or pilocarpine) may be employed. In a normal visual apparatus these cause contraction of the pupil and of the ciliary muscle. The local application of electricity is sometimes useful. In traumatic cases, complete rest is indicated, in addition to the remedies just mentioned. If the paralysis does not soon disappear, convex glasses may be given for near work. The glasses may be reduced in strength, or abandoned if the ciliary muscle should ultimately recover its function.

Spasm of Accommodation.

Tonic spasm of the ciliary muscle is frequently met with in children and in young adults; it occurs generally in hypermetropia, but it may accompany emmetropia or any error of refraction.

Etiology.—It is usually due to long-continued application

of the eyes for near work, especially when the young patient is in poor health, has uncorrected ametropia, and the work has been excessive and done with poor illumination.

Symptoms.—Both eyes are affected. There are asthenopic symptoms. In emmetropia, the spasm gives rise to the signs of myopia; in hypermetropia, it reduces the amount of manifest error and increases the proportion of latent hypermetropia, or it may even cause the patient to appear myopic; in myopia the error is increased. The diagnosis is made after instilling a cycloplegic; in most of these cases homatropine is insufficient, and atropine must be used.

Treatment consists in the abstinence from near work for a time, the correction of ametropia, attention to the general health, and the production of paralysis of accommodation for a few days by instillations of atropine.

CHAPTER XXVI

PARALYSES OF EXTERNAL OCULAR MUSCLES

Anatomy and Physiology.—The eyeball is moved by six muscles, *the extrinsic muscles*, consisting of the four straight and the two oblique. These arise from the wall of the orbit, and are inserted into the sclera.

The four recti (internal, external, superior, and inferior) arise from the circumference of the optic foramen at the apex of the orbit, run forward, surrounding the optic nerve and posterior portion of the eyeball, and are inserted into the sclera by means of flattened tendons about 10 mm. wide. The lines of insertion of these muscles are not equidistant from the cornea, but have somewhat the form of a spiral. That of the internal rectus is 5 mm., of the inferior rectus 6 mm., of the external rectus 7 mm., and of the superior rectus 8 mm., from the cornea.

The superior oblique arises from the border of the optic foramen, runs forward to the upper and inner angle of the orbit, at the anterior extremity of which it passes through a fibrous pulley; it then continues outward, passing beneath the superior rectus, and is inserted into the upper part of the sclera behind the equator. *The inferior oblique* arises from the superior maxillary bone at the inner portion of the lower border of the orbit, passes outward below the inferior rectus, and is inserted into the outer part of the sclera behind the equator.

The muscles are ensheathed by the fascia of the orbit which covers the sclera as *Tenon's capsule* and sends prolongations to the walls of the orbit. These prolongations are most prominent upon the internal and external recti muscles. They serve to restrain the excursions of the eyeball, and are known as 'check ligaments.'

Nerve-Supply.—The *third* nerve (oculo-motor) supplies all the muscles except the external rectus, which is innervated by the *sixth* (abducens), and the superior oblique, which is supplied by the *fourth* (trochlearis). The nuclei for these three nerves are found in the floor of the fourth ventricle.

Action of the Muscles.—The six extrinsic muscles serve to rotate the eyeball around a vertical, transverse, and antero-posterior axis, the centre of the rotation corresponding

approximately to the centre of the eyeball, and the movements being free in all directions, like a ball-and-socket joint. The movements which take place about the vertical axis are adversion (toward the nose) and abversion (toward the temple); about the transverse axis, elevation and depression; and about the antero-posterior axis, wheel rotation or torsion, by means of which the upper extremity of the vertical meridian is inclined inward or outward.

The external rectus moves the eyeball outward.

The internal rectus moves the eyeball inward.

The superior rectus moves the eyeball upward, inward, and turns the upper extremity of the vertical meridian inward.

The inferior rectus moves the eyeball downward, inward, and turns the upper end of the vertical meridian outward.

The superior oblique rotates the upper end of the vertical meridian inward, and moves the eyeball downward and outward.

The inferior oblique rotates the upper end of the vertical meridian outward, and moves the eyeball upward and outward.

Movements of the Eyeball.—In every movement of the eyeball several muscles act at the same time, as follows:

<i>Adversion</i>	{	Internal rectus. Superior rectus. Inferior rectus.
<i>Abversion</i>	{	External rectus. Superior oblique. Inferior oblique.
<i>Elevation</i>	{	Superior rectus. Inferior oblique.
<i>Depression</i>	{	Inferior rectus. Superior oblique.
<i>Rotation inward</i> of upper extremity of vertical meridian	{	Superior oblique. Superior rectus.
<i>Rotation outward</i> of upper extremity of vertical meridian	{	Inferior oblique. Inferior rectus.

The action of the superior and inferior recti and of the obliques varies according to the position of the eye. When the eye is averted, the superior rectus is almost a pure elevator,

and the inferior rectus a depressor; the more the eye is adverted, the more marked is the torsional action of these muscles. In the case of the superior and inferior obliques, the torsional action is greatest when the eye is abverted, and the vertical action greatest when the eye is strongly adverted. No muscle ever acts alone. Though one or two muscles may be the principal agent in a certain movement, all the other muscles are in action to steady the eye, and to eliminate undesired actions of these muscles.

Both eyes move always simultaneously (*associated movements of the eyes*). This association is regulated by centres of association which innervate certain muscles or groups of muscles of the two eyes simultaneously. The associated or conjugate movements occur either in the same direction, with the visual lines parallel, or with the lines inclined toward each other (convergence).

The Field of Fixation corresponds to the limits of movement of the eyeball in different directions, without moving the head. It is best estimated by the perimeter (Fig. 19). The patient's head is fixed so that the eye under examination is opposite the centre of the instrument. Small test-types are now moved along the arc of the perimeter, from the periphery to the centre, until the patient can name the letters. The movements must be made with the eye alone, without any change in the position of the head, and the other eye must be covered. The field of fixation in the normal eye is about 45 degrees upward, inward, and outward, and about 55 degrees downward.

Binocular Vision.—Under ordinary conditions, both eyes are concerned in the act of vision, and the relative directions of the visual axes are involuntarily adjusted, so that the image of an object is focussed on the macula of each eye. These two images are blended in the brain so that a single picture is perceived. The faculty of blending images is called the *fusion faculty*; the act of blending them is called *binocular vision*.

Diplopia.—When images fall on symmetrical points of the retinae of a normal pair of eyes, a single visual sensation is produced (*binocular single vision*). When the visual lines of

the two eyes are not directed toward the same object—*i.e.*, when one eye deviates—*diplopia* (double images) results, if the fusion faculty be perfect. But if the fusion faculty be very deficient, or absent, the image of the deviating eye is disregarded or suppressed. In a case of paralysis of an external ocular muscle, the angular displacement of the false image is equal to the angle of deviation of the eye. The image which corresponds to the eye which 'fixes' the object is distinct, because it lies at the macula, and is known as the *true image*; the image of the deviating eye is less




FIG. 312.—DEVIATION OF THE RIGHT EYE INWARD. HOMO-
NOMOUS DIPLOPIA.

TI, True image; *FI*, false image;
m, macula.




FIG. 312A.—DEVIATION OF THE RIGHT EYE. OUTWARD CROSSED
DIPLOPIA.

TI, True image; *FI*, false image;
m, macula.

distinct, because it is perceived by a peripheral part of the retina, and is known as the *false image*.

Objects which are situated to the right of the point of fixation throw their images to the left of the macula; and those placed to the left of the point of fixation form images to the right of the macula. In the same manner, objects above or below the point of fixation cast their images below or above the macula respectively. By reversing this process, we judge of the situation of an object, and place it at the extremity of an imaginary line drawn from the retinal image through the nodal point. This process is known as *projection*, and is learned by experience. It enables us to judge of the relative positions of objects. An object which forms its

image to the right of the macula is situated to our left; one which throws its image below the macula is situated above, etc.

Diplopia is said to be *homonymous* when the false image is on the same side as the deviating eye, and *crossed* when it is on the opposite side.

In Fig. 312 the right eye is turned in, and consequently binocular diplopia results. The patient sees a true image with the left eye, the image of the candle forming at the macula and being referred to its proper place, TI. In the right eye, on account of the deviation inward, the image is thrown upon the retina to the left of the macula, and consequently is projected to the right, at FI. The image of the right eye being to the right of the image of the left eye, the case is one of homonymous double images.

In Fig. 312A the right eye turns out and double images result. The image of the candle lies on the macula in the left eye, and this eye refers the image to its right position: a true image is seen at TI. In the right eye, because of its outward deviation, the image falls to the right of the macula, and is consequently projected to the left, at FI. The images having crossed in their relative positions, that of the right eye being seen to the left of the image of the left eye, the case is one of crossed diplopia.

Double images may also be produced without any deviation, by placing a prism in front of the eyes. The prism will deflect the rays, so that instead of falling upon the macula, they reach the retina to one side of it.

Varieties of Ocular Deviations.—A deviation may be (1) paralytic; (2) non-paralytic.

1. In paralysis the deviation is due to loss of function of one or more of the ocular muscles; the paralysis may be (a) complete or (b) partial (*paresis*).

2. Non-paralytic (concomitant) deviations are produced by anomalies of the power of convergence and divergence. In these cases the amount and character of the deviation does not vary in the different directions in which the eyes are turned, since we can converge or diverge our eyes with the same facility

on looking to the right as on looking to the left. Deviations may be (a) manifest or (b) latent.

(a) Strabismus (squint or heterotropia) is a *manifest* deviation in which binocular fixation is impossible. Fixation is maintained with one eye or the other, but not with both at the same time.

(b) Heterophoria is a condition in which the eyes have a constant tendency to deviate, but are forced into simultaneous fixation by muscular effort prompted by the desire for binocular single vision. Ordinarily the deviation is not apparent, hence it is said to be *latent*.

Paralysis of the Ocular Muscles.

Symptoms.—1. *Limitation of Movement* of the eye on the side and in the direction of action of the paralyzed muscle; this is pronounced in complete paralysis and less marked in paresis. It can generally be recognized when the patient keeps his head fixed and tries to follow the examiner's finger moved in different directions. If the paralysis be slight, the defective movement may require more elaborate tests for its detection.

2. *Deviation.*—When the eyes are turned in the direction of the normal action of the paralyzed muscle, the sound eye will be directed properly, but the affected eye will refuse to move, and will deviate. The deviation is generally obvious, and is more marked the farther the eyes are moved in the direction of the paralyzed muscle. When the eyes are turned in the opposite direction, in which the paralyzed muscle need not participate, there is no deviation.

The deflection of the deviating eye is known as the *primary deviation*; it is always in the direction opposite to the normal action of the paralyzed muscle.

If the affected eye be made to fix an object and the sound eye be covered, the latter will deviate in a corresponding direction, and much more than the affected eye; this deflection of the sound eye is known as the *secondary deviation*. The excess of secondary deviation over the primary is due

to the fact that the strong impulse of innervation required to enable the paralyzed eye to fix, being simultaneously transmitted to the associated muscle of the sound eye, produces an overaction of this muscle, and consequently a greater amount of rotation.

3. *Oblique Position of the Head.*—If the deviation be not of very high degree, the patient turns his head toward the side of the paralyzed muscle, and in the direction in which the paralyzed muscle would, if acting, move the eye. This is done so as to enable him to bring the visual axes into their normal *relative* directions. Hence there is a characteristic position of the head for every variety of paralysis.

4. *False Projection.*—The paralyzed eye does not see objects in their correct location. The false projection is due to markedly increased innervation, conveyed to the nerve supplying the paralyzed muscle in an effort to force it to act; this gives the patient an erroneous idea of the position of the eye. It can be demonstrated by closing the patient's sound eye and telling him to point quickly at an object in front of him. The finger will be directed to the side of the object corresponding to the paralyzed muscle.

5. *Diplopia* occurs when the patient looks at an object situated within the sphere of action of the paralyzed muscle, and becomes more marked the more the eyes are moved toward this side. The presence or absence of diplopia, the relative positions of the double images, and the increase or diminution of the distance between them in different parts of the field of fixation, form the most important means of determining the seat of the paralysis.

6. *Vertigo, nausea, and uncertain gait* are frequently symptoms dependent upon the diplopia and the false projection. They are relieved when the patient closes the paralyzed eye. On this account, patients frequently keep the affected eye closed or covered.

After paralysis has lasted a long time, there occurs contracture of the antagonist of the affected muscle, increasing the angle of the deviation, and so causing the false image to be formed on a more peripheral and less sensitive part of the

retina. On this account the diplopia and false projection become far less obvious.

When one muscle only is paralyzed, the diagnosis is easy; but when several muscles are involved, it is often difficult to determine the exact combination.

Method of Investigating a Case of Oculo-motor Paralysis.—

Apply the *mirror test* described on p. 412, throwing the light on to the eyes successively from the nine positions indicated by Fig. 313. It will immediately be seen which eye (if either) deviates, and in what direction. The direction in which the angle of the deviation tends to increase or diminish will also be noted.

The advantages of this test are its extreme rapidity and its reliability. It is a purely objective test, so that one is not liable to be deceived by a malingerer or by an hysterical or unintelligent patient.

The *diplopia test* is very commonly employed. It takes much time, and one is dependent upon the answers given by the patient. It is a very delicate test. In using this test care must be taken not to mistake a heterophoria for a paralysis.

For convenience of record and study a diagram is used, consisting of two horizontal and two vertical lines, forming nine spaces (Figs. 313 to 318). The patient must keep his head fixed and merely move his eyes. A red glass is placed before one eye, so as to distinguish its image. A candle or, preferably, an electric bar light, is moved about in different positions in the field of fixation, and the nature of the diplopia noted in each of the nine spaces. The data required are: (1) in which position of the field there is single vision, and in which diplopia; (2) whether the diplopia is homonymous or crossed; (3) the relative distances between the double images; (4) whether the two images are on the same or on different levels; and (5) whether the images are erect or inclined.

The false image is situated in the direction of the normal action of the paralyzed muscle, and the distance between the double images increases in this direction and diminishes in the opposite direction. In fact, most of the symptoms—

the limitation of movement, the false image, the turning of the face and oblique position of the head, the faulty projection, and the increase in the distance between the double images—are in the direction of the normal action of the paralyzed muscle. The deviation of the eye is the only symptom occurring in the opposite direction.

The anatomy of the extrinsic ocular muscles tells one all one need know about their actions. The student who has once thought out these actions for himself will have no difficulty in interpreting the data obtained by either of the tests just described. He has only to note in what direction movement is deficient, then consider what muscle or muscles should produce this rotation.

Varieties of Ocular Paralysis.—One muscle may be involved, or several muscles in various combinations may be affected. Paralysis of the external rectus is the most common; that of the superior oblique is frequent; isolated paralysis of the remaining four muscles is much less common. Combined paralysis of some or all of the four muscles supplied by the third nerve is exceedingly common.

Paralysis of the External Rectus (Sixth Nerve).—There is limitation of movement outward, the eye is adverted, and the face is turned toward the paralyzed side. Homonymous diplopia upon looking toward the paralyzed side; the images are on the same level and parallel (slightly tilted in the upper or lower portions of the field); the lateral separation increases with abversion of the paralyzed eye (Fig. 313).

Paralysis of the Internal Rectus.—There is limitation of movement inward, the eye is abverted, the face is turned toward the sound side. Crossed diplopia on looking toward the sound side; the images are on a level and parallel (slightly tilted in upper and lower portions of field); lateral separation increases with abversion of the paralyzed eye (Fig. 314).

Paralysis of the Superior Rectus.—There is limitation of movement upward and toward the sound side; deviation of the eye downward and a little outward, with the vertical meridian inclined toward the temple; the face is directed upward and toward the sound side, and the head inclined

toward the shoulder of the healthy side. Crossed and vertical diplopia upon looking up; the false image is higher and its upper end inclined toward the nose; the vertical




FIG. 313.—PARALYSIS OF THE EXTERNAL RECTUS.

The dotted outline refers to the false image.

distance between the images increases and the inclination of the false image diminishes, upon looking upward and toward the paralyzed side (Fig. 315).




FIG. 314.—PARALYSIS OF THE INTERNAL RECTUS.

The dotted outline refers to the false image.

Paralysis of the Inferior Rectus.—There is limitation of movement downward and toward the sound side; deviation of the eye upward and slightly outward, with the vertical meridian

inclined toward the nose; the face is directed downward and toward the sound side, and inclined toward the shoulder of the paralyzed side. Crossed and vertical diplopia on looking down;




FIG. 315.—PARALYSIS OF THE SUPERIOR RECTUS.
The dotted outline refers to the false image.

the false image is lower, and its upper end inclined toward the temple; the vertical distance between the images increases




FIG. 316.—PARALYSIS OF THE INFERIOR RECTUS.
The dotted outline refers to the false image.

and the inclination of the false image decreases, upon looking downward and toward the paralyzed side (Fig. 316).

Paralysis of the Superior Oblique (Fourth Nerve).—There is limitation of movement downward and toward the paralyzed side; the eye is deviated upward and slightly inward, with

the vertical meridian inclined toward the temple; the face is directed downward and toward the sound side, and the head is inclined over the shoulder of the sound side. The patient has great difficulty in moving about, especially in going downstairs. Homonymous and vertical diplopia on looking down; the false image is lower, and its upper end inclined toward the sound side; the vertical distance between the images increases, and the inclination of the false image decreases upon looking downward and toward the sound side (Fig. 317).

Paralysis of the Inferior Oblique.—There is limitation of movement upward and toward the paralyzed side; the eye is deviated downward and slightly inward, with the vertical




FIG. 317.—PARALYSIS OF THE SUPERIOR OBLIQUE.

The dotted outline refers to the false image.

meridian inclined toward the nose; the face is directed upward and toward the paralyzed side, and the head is inclined toward the affected side. Homonymous and vertical diplopia on looking up; the false image is higher and its upper end inclined toward the temple; the vertical distance between the images increases, and the inclination of the false image decreases upon looking upward and toward the sound side (Fig. 318).

Paralysis of the Third Nerve.—With complete paralysis of this nerve there is ptosis; the eyeball is almost immobile, the limitation of movement being upward, downward, and inward; the eye deviates outward and somewhat downward, with

the upper end of the vertical meridian inclined inward, especially upon looking downward; the face is directed upward and toward the sound side, and the head inclined to the shoulder of the paralyzed side. There is slight exophthalmos on account of the paralysis of the three recti which normally draw the eyeball backward; the pupil is dilated and immobile; accommodation is paralyzed; there is crossed diplopia—the false image is higher, and its upper end inclined toward the paralyzed side.

Paralysis of the third nerve is common; it is often incom-




FIG. 318.—PARALYSIS OF THE INFERIOR OBLIQUE.

The dotted outline refers to the false image.

plete, two or three of the muscles being affected. It may be associated with paralysis of other nerves.

When all the muscles of one eye are paralyzed, including the iris and ciliary body, the condition is known as *total ophthalmoplegia*.

When all the exterior muscles of the eyeball are paralyzed, but not the iris and ciliary body, the condition is known as *external ophthalmoplegia*. This form is more common than total ophthalmoplegia; the nuclei for the sphincter pupillæ and ciliary muscle being separate, they often escape involvement from destructive processes affecting the origin of the exterior ocular muscles. This form of paralysis is generally of central (nuclear) origin.

When only the sphincter pupillæ and the ciliary muscle

are paralyzed, the condition is known as *internal ophthalmoplegia* (p. 390).

Associated or Conjugate Paralysis involve associated muscles such as the external rectus of one eye and the internal rectus of the other. They are due to lesions in the association centres.

Etiology.—The lesions causing paralysis may be situated anywhere in the course of the nerve tract, from the cerebral cortex to the muscle. According to its site, the lesion is distinguished as central and peripheral. Central lesions may be situated in the cortical centres (cortical paralysis), the association centres, the nuclei of origin (nuclear paralysis), or in the fibres which connect these centres. Peripheral lesions may affect the nerves in some part of their course, either between the point where they issue from the brain and their entrance into the orbit (basilar paralysis), or in the nerve or its branches in the orbit (orbital paralysis).

The Differential Diagnosis between Central and Peripheral Paralysis is not always easy; it is based on the character of the paralysis and the accompanying symptoms. Complete paralysis, unaccompanied by any other symptoms, is generally peripheral. When central, the paralysis is generally less complete, more than one muscle is usually involved, there are apt to be cerebral symptoms, and there is commonly an absence of peripheral cause.

The Nature of the Lesion.—The lesion may be a neighbouring exudation, hæmorrhage, periostitis, tumour, injury, or vascular change, causing compression or inflammation of the nerves; less frequently it is a primary inflammation or degeneration.

The most common cause is *syphilis* (late symptom), which is responsible for one-half the cases. Rheumatism and exposure to severe cold sometimes cause oculo-motor palsies. Epidemic encephalitis is another cause. Muscle paralysis occur in various central nervous system diseases (tabes, general paralysis, disseminated sclerosis, etc.); after acute infectious diseases (diphtheria, influenza, etc.); in acute poisonings (alcohol, ptomaine, botulism, etc.); in diabetes; in rheumatism; in exophthalmic goitre.

Congenital paralyses are not uncommon, due to absence, abnormal insertion, or other structural defects of the muscles themselves. Congenital abnormalities most often affect the external rectus and the superior rectus.

Prognosis varies with the cause. Peripheral paralyses due to syphilis, rheumatism and cold usually recover under appropriate treatment, but there may be relapses. In the paralysis accompanying serious spinal and cerebral disease the prognosis is often bad. Long-neglected paralyses present an unfavourable prognosis, on account of the atrophy of the paralyzed muscle and the contracture of the antagonist. The course is always chronic, and even in favourable cases several weeks or months are required to effect a cure.

Treatment should, in the first place, be directed to the cause. In syphilis, mercury, iodides, and salvarsan are indicated. In rheumatism and gout, salicylate of sodium, iodide of potassium, and colchicum, singly or combined, are prescribed. In diphtheria, strychnine is indicated. In obscure cases, iodides with or without mercury should be resorted to. Hot baths and diaphoresis are sometimes employed.

Locally, we may use electricity, ocular-muscle exercises, prisms, and occlusion of one eye. In incurable cases, operative intervention may be resorted to.

Electricity may be tried; the constant current (3 milliamperes) is used, the negative pole being applied to the back of the neck, and the positive over the affected muscle.

The weakened muscle may be exercised by allowing the patient to look through a prism which almost corrects the diplopia, thus coaxing the paralyzed muscle into action. The same result can be achieved by directing the patient to move his head until the double images almost coalesce, and then to make a strong effort to fuse them without any further motion of the head. Such exercises are repeated ten times at each sitting, several times a day. When available, one of the instruments used for orthoptic training (p. 416), such as the myoscope or synoptophore, may be used in suitable cases.

~~In chronic cases with moderate paresis, prisms may neu-~~
tralize the diplopia, and thus add to the patient's comfort,

but they are seldom successful, because even in slight paralysis the diplopia changes in amount in whatever direction the eye is moved. The only satisfactory way to avoid double vision is to occlude the deviating eye by a patch, or by a ground glass in a spectacle frame.

If the condition persists for a long period in spite of all treatment, and the paralysis seems incurable, operative treatment is indicated. This consists in an advancement of the paralyzed muscle (p. 432), or musculo-capsular advancement, with or without tenotomy or recession of the antagonist.

Spasm of the Ocular Muscles is due to excessive innervation; it may be primary or secondary.

Primary Spasm is rare; it may be produced by meningeal or reflex irritation.

Secondary Spasm is common, and occurs with paralysis of one of the other ocular muscles, presenting excessive movement in field of action of the spastic muscle and spastic deviation of that eye; it appears frequently in the direct antagonist of a paralyzed muscle—*e.g.*, spasm of the internal rectus following paralysis of the external rectus of the same eye. Where the paralyzed eye is used for fixation, there is often a secondary spasmodic deviation of the other eye, due to spasm of the associate of the paralyzed muscle. The most common example of this type of deviation occurs in paralyses of the superior rectus followed by spasm of the inferior oblique of the other eye. The treatment of secondary spasms is operative—*i.e.*, tenotomy or recession to weaken the action of the over-active muscle.

Nystagmus.

Nystagmus is not a paralysis, but it is convenient to discuss it in this chapter.

Nystagmus is a short, rapid, involuntary oscillation of the eyeball, usually affecting both eyes and associated with imperfect vision. The movements are most frequently from side to side (lateral nystagmus), or around the antero-posterior axis (rotatory nystagmus), sometimes up and down (vertical nystagmus). There may be a combination of the

lateral or vertical with the rotatory movements (mixed nystagmus). The oscillations are similar in kind, duration, and frequency in the two eyes. They may be constant, or present or exaggerated only when the eyes are turned in certain directions. The patient is not, as a rule, inconvenienced by the existence of this condition; but when it commences in adult life there may be much annoyance from the apparent movements of objects.

Most cases exist from *infancy*. These may be caused by diminution in the acuteness of vision, as a result of opacities of the media, intra-ocular diseases, or congenital anomalies, such as albinism. But, as a rule, the motor inco-ordination is congenital, and causes amblyopia by rendering steady fixation impossible.

In *adults* it may develop with many cerebral affections, especially disseminated sclerosis, disease of the cerebellum, and Friedrich's disease. It is also found in coal-miners (miner's nystagmus). In this affection, which is an industrial disease notifiable to the Home Office, it is only one of the symptoms of a general functional affection of the nervous system, and is probably caused by poor illumination of the coal face which makes continuous fixation difficult. Many ~~other~~ causes have been suggested. Disease of the labyrinth is an occasional cause of nystagmus.

The infantile cases are not amenable to treatment, though the condition sometimes becomes less marked with advancing years; the correction of even high errors of refraction seldom improves the vision. Miner's nystagmus generally disappears with a change of occupation.

CHAPTER XXVII

COMITANT SQUINT

Concomitant Strabismus.

CONCOMITANT strabismus (concomitant squint, or heterotropia) is a manifest deviation of the visual line of one eye, the angle of deviation between the two visual axes remaining constant in whatever direction the eyes are moved. The range of movement of each eye is full. The eye which is directed toward the object gazed at is known as the *fixing eye*; the other as the *squinting eye*.

Concomitant strabismus differs from heterophoria in that, in the latter condition, there is no manifest deviation when both eyes are uncovered, the strong desire for fusion keeping the eyes straight. It differs from paralytic squint by presenting a normal range of movement of each eye and the same deviation in all parts of the visual field, while in paralysis the deviation is present only in the field of action of the paralyzed muscle and there is limitation of movement in the direction of action of that muscle; in concomitant squint the primary and secondary deviations are equal, while in paralytic squints the secondary deviation is greater than the primary (p. 398); double vision, a prominent symptom in paralytic squint, is not present in concomitant squint, as the image seen by the squinting eye is suppressed.

Concomitant squint may be (1) occasional, if the deviation is not always present; (2) constant, if present all the time and always manifested by the same eye; (3) alternating, if the patient fixes with either eye indifferently, while the other eye deviates.

According to the direction of deviation, concomitant squint is classified into (1) convergent strabismus (internal squint, eso-

tropia); (2) divergent strabismus (external squint, exotropia); (3) vertical strabismus (right or left hypermetropia according to the higher eye; (4) mixed strabismus, a combination of lateral and vertical squint.

Symptoms.—The most striking is the deviation of the visual axes. Absence of diplopia, from suppression of the image seen by the squinting eye, is almost invariable. Amblyopia is frequent, if the squint is constant, and has existed for some time in a young patient. This is usually acquired as the result of constant suppression of the image of the squinting eye; less commonly it is congenital, or due to injury of the optic nerve or retina from hæmorrhage at birth. A refractive error, commonly hypermetropia, with or without astigmatism, is frequently present.

Etiology.—There are several factors which may enter into the causation of concomitant squint, to any one, or all, of which any individual case may be attributable.

1. Probably the most important factor is some weakness of the inherent instinct in mankind to blend the images of the two eyes, which is known as the 'fusion sense.' This instinct develops very early in life and is fully established by the age of six years. So long as it is perfect a lot of disturbance from other factors is required to make one eye deviate, but when it is weak or absent, anything upsetting the equilibrium of the motor co-ordinations of the eyes may cause a permanent squint. Such influences may be:

2. An upset of the normal relationship between accommodation and convergence (see p. 343). In hypermetropia there is a *tendency* to over-convergence, and in myopia to under-convergence, so that in the former case the tendency is for any squint to become convergent, and in the latter case to be divergent. This tendency *may* be overbalanced by the factor of faulty muscle balance (heterophoria), so that a hypermetrope may develop a divergent squint, and, *vice versa*, a myope a convergent squint. On the other hand, the tendency may be increased by this factor. As hypermetropia is much more frequent than myopia in young children, convergent squints are much more common than divergent.

3. Anisometropia and congenital amblyopia, which make images dissimilar or blurred, and make fusion more difficult. Squints in which the normal relationship between accommodation and convergence has been disturbed by errors of refraction are known as accommodative squints. The non-accommodative squints comprise those associated with myopia and those in which no refractive error is found.

4. An upset of the normal muscle balance of the eyes. Any abnormal tendency to direct the eyes inwards will, in the absence of a strong desire for fusion, give rise to a convergent squint, and any tendency to deviate the eyes outwards will, under the same conditions, give rise to a divergent squint.

5. Debilitating illnesses, such as the specific fevers, or any violent mental disturbance, by making muscular control more difficult, may be the immediate exciting cause of a squint when other factors are present.

Methods of Investigating a Case of Squint.—The presence or absence of a deviation may usually be determined by simple inspection, but in slight cases the appearances may be misleading.

Test the corneal reflexes in a dark-room. Reflect the light from an ophthalmoscopic mirror into the patient's eyes from a distance of about 2 feet. An infant will instinctively look at the light; older patients are told to do so. A tiny image of the light is formed on the patient's cornea, usually slightly to the nasal side of the centre, owing to the angle gamma (p. 339). By flashing the light rapidly from one eye to the other any want of symmetry in the position of the reflexes can be seen and the degree of deviation roughly estimated. If the mirror is taken to a distance of 12 inches from the eyes each millimetre of deviation in the reflex represents roughly some 7 degrees of squint.

Test the movements of each eye separately by covering one eye at a time. These should be equal in both eyes.

Test the power of fixation in the squinting eye, after covering the fixing eye, by telling the patient to fix the light, or finger. If he fixes properly the eye has central fixation, and vision is probably not less than $\frac{6}{80}$; if not, central fixation is

lost and vision is less than $\frac{6}{80}$. Where the patient is old enough, carefully record vision in each eye, with any glasses that are ordered.

Measure the angle of deviation, which should be recorded at each visit. This may be done by examination of the corneal reflexes as described above, or, more accurately, by the 'screen test,' the perimeter, or the synoptophore, if this is available.

The screen test cannot be used when there is loss of fixation in one eye, otherwise it is an accurate method. With the patient fixing an object, a card is placed before one eye and then passed quickly to and fro from one eye to the other. The card is so passed that the patient has no chance to fix with both eyes at the same time, but must alternate his fixation. Each eye when covered deviates, and when uncovered turns back into the fixing position. A prism (base out for internal squint, base in for external squint) of sufficient strength to abolish this movement of correction represents the exact amount of deviation present. It is often more accurate to deduct 2 degrees from the weakest over-correcting prism. This test is done at 20 feet and 13 inches and the amount of deviation at both distances is noted.

The perimeter gives the angular measurement of squint. It cannot be used with small children, and the method is troublesome and not very accurate. The patient is seated with the squinting eye in the centre of the instrument and is directed to fix a distant object, placed in the median line. A light (small electric torch or a candle flame) is then moved along the inside of the perimeter arc till its reflection in the cornea is seen in the centre of the cornea of the squinting eye. The number of degrees on the arc at this point indicates the size of the angle of the squint.

The quickest and most accurate method is to read off the angle of deviation on a scale attached to one of the larger instruments used for squint training, such as the synoptophore. When the two corneal reflexes, given by the separate internal illumination in each tube of the instrument, are symmetrical, that reading of the scale gives the true angle of the squint.

In every case the state of the refraction must be ascertained by retinoscopy, after using a mydriatic, and measurements should be made both with and without glasses, if these are ordered.

Convergent Concomitant Strabismus.

In this form of squint there is deviation inward of the visual line of one eye (Fig. 319). It is generally associated with hypermetropia, with or without astigmatism; more rarely it



FIG. 319.—CONVERGENT SQUINT.

occurs in myopia and emmetropia. It usually commences in early life, between the first and fourth years, when the child begins to use his accommodation for near objects, such as toys and pictures; more rarely it is con-

genital. At first the squint may be noticed only at times, during near vision, or when there is any interference with the general health, but it is apt to become constant for both near and distant vision; sometimes it disappears at or about the age of puberty, but in the meantime, if left untreated, the vision may have deteriorated from disuse (*amblyopia ex anopsia*). Such amblyopia does not occur after the age of six or seven years. The acuteness of vision in the squinting eye is frequently greatly reduced, unless the squint is alternating. While sometimes the amblyopia is congenital, it is probably secondary to continuous suppression of the image seen by the squinting eye in the majority of cases.

Many cases of convergent squint start as a spasmodic esophoria for near vision, due to over-stimulation of convergence by the abnormal accommodative effort needed in hypermetropia for close work. This is much more likely to happen when the desire for fusion is not strongly developed. Once the child has learnt how much more comfortable it is to squint than to make any effort to keep the eyes straight, the deviation is likely to become manifest for both distance and near, and he develops a secondary weakening of his powers of divergence.

Sometimes an esotropia begins as a primary squint for distance, due to a divergence insufficiency with a later development of convergence excess.

Treatment.—This should be undertaken at the earliest opportunity. The objects to be kept constantly in view are: (a) to prevent deterioration of the vision of the deviating eye, and to restore, as far as possible, the sight of this eye in cases in which amblyopia has occurred; (b) to endeavour to remove one fundamental cause of the squint by training the sense of fusion at the earliest possible age; (c) to restore the visual axes to their normal relative positions.

There are five measures we may use in an endeavour to attain these objects:

1. Correction of refractive errors by glasses.
2. Exercising the squinting eye by occluding its fellow.
3. Instillation of atropine into the fixing eye.
4. Training the fusion sense (orthoptic training).
5. Operation.

1. The error of refraction should be estimated under atropine and nearly the total hypermetropia (and astigmatism if present) should be corrected by glasses prescribed for constant wear. In some cases, especially if squint is occasional, this alone effects a cure. Glasses may be worn by very young children, but are rarely practicable before the age of eighteen months. It is sometimes advisable to keep the eyes under the influence of atropine for a few days when the glasses are first worn, to ensure complete relaxation of the ciliary muscle. In the rare cases of myopia with convergent squint the myopia should be exactly corrected.

2. The fixing eye should be covered by a patch or bandage, or special occluder fitted to the glasses. This occlusion should be continuous at first, where practicable, if fixation is poor or visual acuity much below normal; it may be necessary to limit it to two or three hours a day, which is less effective. This compels the squinting eye to fix, exercises it, prevents amblyopia from disuse, and restores, as far as possible, the sight of the eye if amblyopia already exists. If no improve-

ment occurs after six weeks it is useless to continue. This treatment can be adopted at a very early age if the squint is constant in one eye.

3. Where fair vision exists in the squinting eye, the instillation of atropine into the fixing eye may compel the child to use the squinting eye for near vision, and so exercise the eye and prevent amblyopia. An unatropinized eye whose vision is only one-sixth, or even one-tenth, of the normal, is able to see near objects more clearly than a normal eye whose accommodation is paralyzed. This treatment must not be kept up for long if it turns the squinting eye into the fixing eye,




FIG. 320.—WORTH-BLACK AMBLYOSCOPE.

but must be intermitted and repeated again after a month or so if required.

4. Orthoptic training of binocular vision aims at encouraging the use of the two eyes at the same time, thereby overcoming the habit of suppression of the image of one eye. The desire for the simultaneous use of both eyes is then used to exercise the muscles which can keep the eyes straight. In favourable cases this treatment alone may correct the squint.

The most important thing in dealing with young children is to make the exercises sufficiently interesting to hold their attention, and to vary them frequently, to create interest and amusement.

The instruments nowadays used are the synoptophore

(Fig. 321) (or its modifications), which is an elaborate development of Worth's original amblyoscope; the cheiroscope, with which the child is unconsciously trained to use both eyes at the same time, and also to make use of his hands and fingers; the amblyoscope (Fig. 320), and various forms of stereoscopes. In addition, other special apparatus—*e.g.*, the myoscope—may be employed to strengthen any required muscular action by




FIG. 321.—THE SYNOPTOPHORE.

- | | |
|---|---|
| A. Lampholder for opaque cards. | F. Handle for adjusting angle of tubes. |
| B. Lampholder for transparencies. | G. Hyperphoria adjustment. |
| C. Fine adjustment for Fusion Training. | H. Cyclophoria scale. |
| D. Moving picture attachment. | I. Eyepiece. |
| E. Screw for altering distance between eyepieces. | |

the projection on to a screen of an object, or two objects requiring fusion (red and green seen through red and green glasses), which can be made to move backwards and forwards in any required direction.

One great advantage of the synoptophore is that the trainer has an unimpeded view of the patient's eyes, and that the illumination of the pictures is internal. Consequently the position of the corneal reflexes, given by the separate lights


in each tube of the instrument, can be watched, and the true position of the eyes noted. This is impossible with the amblyoscope and most stereoscopes. One can then determine whether the child is fixing the pictures with each macula (true binocular projection), or whether he is seeing the picture opposite the squinting eye with another part of the retina, adjacent to the macula. The importance of this may be understood when it is realized that more than 70 per cent. of cases have been found unable at first to superimpose the two images at the true angle of their squint, but to do so while still squinting—even to the extent of 40 degrees, but more commonly 10 degrees ('*binocular false projection*'). Hence instruments such as the amblyoscope or stereoscopes which are controlled by the child himself do more harm than good, by stimulating this false projection, if they are used before it has been (if possible) overcome.

The synoptophore (Fig. 321), a development of the amblyoscope, consists of two metal tubes; each tube is angled near the distal end at 135 degrees; a mirror is set inside at the angle, to reflect the image of a picture, placed in a holder at the distal end of the tube, through a convex lens, in the eyepiece at the proximal end of the tube, on to the patient's eye. Both tubes are fixed into a base and can be separately rotated inwards or outwards to suit any angle of squint. Vertical adjustments also can be made. Each picture is illuminated by a separate light inside each tube; the intensity of this can be varied by means of a rheostat.

The object-slides used with this instrument, as well as with the amblyoscope and stereoscopes, are of three classes: (1) Those which do not require any blending of images, but only simultaneous vision of dissimilar objects with the two eyes. Fig. 322 is an example. (2) Devices of the second class (*e.g.*, Fig. 324) require true fusion of images in order that the full picture may be seen. (3) Devices such as Fig. 325 can only be appreciated by persons who have a sense of perspective.

The first step in the treatment is to overcome the suppression of the image seen by the deviating eye. The instrument is adjusted to the angle of the squint, and exercises begun by

showing pictures of the first group, to stimulate simultaneous perception. The illumination of the picture before the deviating eye is increased and the relative brilliancy of the



FIGS. 322 AND 323.—DEVICES REQUIRING ONLY SIMULTANEOUS VISION.

lights adjusted, until the objects on both sides are seen simultaneously. The bird is made to go into the cage and out again, by slightly moving the two sides of the instrument. If projection has been found to be false, this first has to be corrected




FIG. 324.—DEVICE REQUIRING FUSION OF IMAGES.

by special exercises. Many similar devices are shown until the symptom of suppression is entirely overcome, under all conditions of illumination, while the child is looking into the instrument. The child is next taught to fuse images by using




FIG. 325.—TEST FOR THE SENSE OF PERSPECTIVE.

slides from the second group, and is then encouraged to keep them fused over as wide a range of movement (abduction and adduction) as possible. When the range, or amplitude, of fusion has been sufficiently increased, the sense of per-

spective is developed by slides from group three—*e.g.*, when seen stereoscopically the various geometrical figures in Fig. 325 should be seen in different planes.

If good amplitude of fusion has been obtained there will be a strong tendency to binocular single vision. This tendency will be sufficient to overcome a small deviation, or a deviation which has been approximately corrected by operation.

The most favourable age for training is between three years and six years. Over the age of seven years satisfactory results are much rarer. Good monocular fixation in the squinting eye is essential. If the child is old enough to be tested, the visual acuity should not be less than $\frac{1}{8}$ with correcting glasses. Cases which at first seem concomitant, but are found to be incomitant, are not usually suitable for training.

During the past few years orthoptic training has been largely developed, and in London and other large centres several hospital and private clinics have been established for this purpose. The treatment requires a good deal of time and patience and the selection of suitable cases. It is of great value in determining the selection of suitable cases for early operation, and in breaking down 'false projection,' which, unless overcome, will prevent a complete cure.

Many of the methods employed are also used to exercise muscles in cases of heterophoria, and also in the slighter cases of paresis of ocular muscles. In such cases binocular fusion is already well developed and is used to 'tie' the eyes while the range of movement is gradually increased.

5. When such non-operative treatment as is practicable has been given thorough trial for several months without correcting the squint, the question of operation arises. No child should be allowed to go through life with an obvious squint which can be remedied, as far as the appearance is concerned, by operation. But squints should not be operated upon for purely cosmetic reasons until the child is old enough to have this done under local anæsthesia, this age varying from nine to thirteen years. Exceptions to this rule occur when there is marked self-consciousness developing in the patient, or even in the parent, as the result of the squint,

and in such cases operation should be undertaken earlier, under general anæsthesia, though there is no expectation of obtaining binocular vision. But if there is a reasonable expectation of obtaining binocular vision after operation, as can be shown by the use of the synoptophore or amblyoscope, the sooner operation is undertaken the better, so that the habit of binocular vision may be formed early. After the age of seven it is much harder to develop the habit. When there is a good sense of fusion an approximate correction is usually sufficient to keep the eyes straight, the patient overcoming any slight tendency to deviation by his desire for binocular vision. When there is no desire for fusion adjustments have to be more accurate, and this accuracy can best be obtained under local anæsthesia.

✓ The operative procedures which are employed are tenotomy (or recession) of an internal rectus, and advancement of an external rectus, either singly or in combination.

The choice of operation depends partly on the degree of the squint and partly on the preference of the individual surgeon. ~~Simple unguarded tenotomies of the internus, which were once very popular, are seldom done, as they not infrequently gave rise to infrequent divergence of the eye, but recession of the internal rectus muscle, which fixes the completely tenotomized muscle farther back on the sclera, is coming more into general use.~~ It is specially indicated if the squint is present for near only.

As a rule advancement of one or both external recti, ~~with or without a guarded tenotomy of the tendon of the internal rectus, is the operation of choice.~~ If the squint is more than 30 degrees and present for both distance and near, a combination of the two operations is indicated; if it is present for distance only a simple advancement of the externus is called for. The operation is usually done first on the squinting eye, and subsequently, if necessary, on the other eye, after the final result of the first operation has been determined. Some surgeons, however, operate on both eyes at once, especially if they are doing simple advancement of the external recti. To obtain satisfactory results considerable experience and judgment are

necessary; the degree of advancement or recession required can be fixed by no arbitrary rules. The operations are described in Chapter XXIX.

Divergent Concomitant Strabismus.

This form of squint (exotropia) exists when one eye fixes an object and the other deviates outwards. It is much less common than convergent squint. Those cases which are associated with myopia are generally due to weakness of convergence, associated with the slight, or negative, accommodative effort which is required in this condition; there is an habitual deficiency of the stimulus for convergence; this power therefore gradually weakens, and the patient has a deviation at close range, but




FIG. 326.—DIVERGENT SQUINT.

none at a distance; there is then a gradual increase in the amount of deviation until it is present at all distances. Such squints usually begin about ten or twelve years of age, or in early adult life.

Other cases, which are primarily due to excess of divergence, are independent of refractive errors, and usually start early in life. The divergence is at first present only for distant vision; as time goes on the power of convergence weakens, and the deviation persists for both distance and near. Such patients, while usually bright and intelligent, are often of a 'nervous' temperament, and have a neuropathic family history.

Whereas in the first group there is usually marked desire for binocular vision, in the second this desire is feebly developed or entirely absent.

Whereas in the first group there is usually marked desire for binocular vision, in the second this desire is feebly developed or entirely absent.

Treatment.—In the first group the full correction of any existing myopia and astigmatism, at the earliest possible age, is indicated; this will cure those cases due to an uncorrected myopia, where this deviation is still periodic.

Fusion and prism exercises may help to complete the cure of

the squint in cases in which the use of glasses alone has proved insufficient.

In all other cases operation is required, an advancement of an internal rectus being indicated.

In the second group of cases there is little or no refractive error to correct. In constant squints amblyopia ex anopsia must be guarded against. Fusion exercises usually prove unsatisfactory. An accurate advancement of one or both internal rectus muscles will remove the deformity, but will rarely attain binocular fusion.

Non-comitant Divergent Squints other than Paralytic.

In *extreme myopia* the egg-shaped eyes tend to adapt their long axes to the divergent positions of the orbits. No treatment for the divergence should be attempted.

Blind eyes usually tend to diverge.

Tenotomy of an internal rectus in a certain proportion of cases produces an over-effect, so that the eye diverges. The retracted internal rectus muscle should be sought for and advanced.

CHAPTER XXVIII

HETEROPHORIA

IF a person with a perfectly normal pair of eyes looks steadily at any object both visual axes will continue to be accurately directed to that object even though one eye be shaded. In other words, his perfectly balanced motor co-ordinations are able to maintain the normal relative directions of the eyes even when the controlling influence of the fusion sense is temporarily withdrawn. This state of perfect oculo-motor equilibrium is called *orthophoria*.

Heterophoria is the name given to the condition of imperfect oculo-motor balance. There is here a *tendency* for the eyes to deviate from their normal relative directions. Ordinarily, however, this tendency is kept in check by the desire for binocular vision, so that there is no squint. But when the effort to maintain the eyes in the normal position becomes too great for any cause, the condition may readily become a *manifest* instead of a latent deviation—at first giving rise to a diplopia which passes off if the patient learns to suppress the image of the deviating eye. Distinctive names are employed to indicate the direction of the tendency:

Esophoria is a tendency to abnormal static convergence of the visual axes.

Exophoria is a tendency to divergence of the visual axes.

Hyperphoria is a tendency of the two eyes to rotate vertically in opposite directions, so that one visual axis tends to lie in a higher plane than the other. The eye which tends to turn up relatively to the other is called the hyperphoric eye. This may be associated with exophoria or esophoria.

Cyclophoria is a tendency of the vertical meridian of one eye to rotate from the vertical position.

The Methods of Testing the Motor Balance of the Eyes.—

In a case of heterophoria, under ordinary circumstances, the desire for binocular vision prevents the eyes from deviating from their normal relative directions. But if, by artificial means, the image formed in one eye be so altered in appearance or position as to make fusion with the other unaltered image impossible, the desire for fusion is suspended. The heterophoria then gives rise to a manifest deviation. The altered image in the deviating eye is not suppressed as in a case of squint. The diplopia, therefore, gives an easy means of ascertaining the direction and degree of the deviation. This is the principle on which all subjective tests for heterophoria are based.

The presence of heterophoria may be determined by noting that one eye deviates when it is covered, and that it straightens itself, so that its visual axis is parallel with that of the other eye, when the screen is removed. The exact amount of the deviation may be measured by means of prisms as described on p. 413, but other tests are more commonly employed for this purpose. These are (1) the Maddox rod; (2) the Maddox wing. Many other tests may also be used—such as the double prism, or Harman's diaphragm, to detect the muscle imbalance and to measure its degree.

The *Maddox rod* (Fig. 327) consists of one or more glass rods, usually coloured red, set in a hard rubber disc, so as to fit into the trial frame. A strong cylindrical lens will answer the same purpose. It converts the image of the light seen by one eye into a long streak of light, so that there remains no desire to fuse it with the image of the other eye. The line is always at right angles to the axis of the rod.

The test is used at 5 (or 6) metres and at $\frac{1}{3}$ metre, preferably in a semi-darkened room. A small electric light




FIG. 327.—MADDOX ROD.

(or a candle flame) is a satisfactory test object. The Maddox rod is placed horizontally before the right eye, converting the image of the light into a vertical streak. If orthophoria is present, this streak appears to pass directly through the image of the light seen by the left eye (Fig. 328). If the line




FIG. 328.

FIG. 329.

FIG. 330.

FIG. 328.—THE MADDOX ROD TEST IN ORTHOPHORIA.

FIG. 329.—THE MADDOX ROD TEST IN EXOPHORIA.

FIG. 330.—THE MADDOX ROD TEST IN ESOPHORIA.

of light appears to the *left* of the light there is crossed diplopia, indicating exophoria (Fig. 329); if to the right of the light there is homonymous diplopia, indicating esophoria (Fig. 330). The amount of heterophoria is measured by the prism, base in or base out, which serves to displace the streak until it runs




FIG. 331.

FIG. 332.

FIG. 333.

FIG. 331.—THE MADDOX ROD IN ORTHOPHORIA.

FIG. 332.—THE MADDOX ROD IN LEFT HYPERPHORIA.

FIG. 333.—THE MADDOX ROD IN RIGHT HYPERPHORIA.

directly through the flame. For distance a slight esophoria or exophoria, 1° to 2° , may be allowed as normal, and for near a slight exophoria 2° to 4° , may be similarly allowed.

The rod is then placed vertically before the right eye, converting the image of this eye into a horizontal line of light, which will pass through the image of the left eye if there is no

vertical imbalance. If the line of light is below the image of the light seen by the left eye, there is right hyperphoria (Fig. 333); if above, there is left hyperphoria (Fig. 332). The degree of hyperphoria is measured by the prism, base up or down, which causes the light streak to pass directly through the flame.

The Maddox wing is a rapid means of detecting the presence of heterophoria for near vision and of measuring its degree. The patient looks through the slits in the eyepieces; for horizontal deviations the upper oblique wing divides the visual field into two halves, so that the right eye sees the finger pointing upwards, while the left eye sees the scale towards which the finger




FIG. 334. — THE MADDOX WING TEST FOR HETEROPHORIA.

points. The character and degree of the deviations can then be at once determined by the apparent position of the finger on the scale. For vertical deviations the two vertical screens cut off the red arrow from the left eye and the red vertical scale from the right eye, so that any vertical deviation can be rapidly detected and measured by the apparent position of the arrow on the vertical scale.

Cyclophoria can be measured by adjusting the movable wire forming the shaft of the arrow till it appears to the patient to be parallel with the horizontal scale. The actual departure from parallelism is a measure of the cyclophoria.

To obtain accurate results when testing for muscular balance any error of refraction must be corrected.

Prism duction (the degree of prism which the eyes can overcome) should always be investigated in any case in which important heterophoria has been found. The patient, wearing a trial frame and having both eyes open, is seated 5 or 6 metres

from a light or a test card. Prisms increasing in strength are successively put in the cell before the right eye. The




FIG. 335.--ROTARY PRISM.

highest prism, apex up, which the patient can bear without seeing the object double gives the range of superduction of the right eye. The right subduction is tested with prisms apex down. The left superduction and subduction are similarly measured. Binocular abduction is measured with prisms apices out. Binocular adduction is so intimately associated with

accommodation that attempts to measure it with prisms (which cause the eyes to converge without accommodating) give misleading results. A rotary prism (Fig. 335) is very useful in making these measurements. The normal limits of prism duction, expressed in degrees of actual deflection, are as follows:

Superduction	-	-	-	$1\frac{1}{2}$	degrees to	$2\frac{1}{2}$	degrees.
Subduction	-	-	-	$1\frac{1}{2}$	„	$2\frac{1}{2}$	„
Abduction	-	-	-	4	„	5	„

No amount of practice appears to increase the duction power in these three directions. Convergence, on the other hand, can nearly always be much increased by practice. As the degree of prism duction does not vary from time to time, and is independent of voluntary effort on the part of the patient, the information obtained is reliable.

Symptoms.—In slight degrees of heterophoria there are very often no symptoms, but these may be marked in low degrees of hyperphoria. In more pronounced forms they are those of ‘eye-strain’ in general—frontal headache coming on towards the end of the day, pain in the eyes after watching anything intently, migraine, dizziness, conjunctival hyperæmia, etc.; momentary diplopia is not uncommon. These asthenopic symptoms are the result of the strain imposed upon the muscles in overcoming the deviation. There are

frequently periods of clear vision with strain, alternating with periods of diplopia with confused vision. Head tilting, or even torticollis (ocular torticollis) may be present as the result of the patient's endeavour to correct a diplopia, particularly if it is vertical. A characteristic feature of the symptoms due to heterophoria is their disappearance on the closure of one eye.

Etiology.—Heterophoria may be refractive or non-refractive in origin.

An error of refraction is a frequent cause for a disturbance of the normal relationship between accommodation and convergence. For example, a hypermetrope has to use an abnormal amount of accommodation to maintain clear vision; thus his power of convergence tends to be constantly overstimulated and an esophoria may result; conversely a myope uses too little accommodation, and is liable to develop an exophoria.

Heterophoria of non-refractive origin is common, since all the cases due to a primary disfunction (over-activity or under-activity) of the power of divergence are not influenced by the state of the refraction it is equally true that many cases of weakness of convergence result from non-refractive causes. Heterophoria is frequently seen in neurasthenia, hysteria, anæmia, focal affections, in connexion with nasal and accessory sinus disease, and in persons who are debilitated from any cause whatever; it is also found in perfectly healthy individuals, and in some cases is due to an anatomical defect of one of the external muscles.

Treatment consists in the correction of any refractive error, attention to the general health, prism exercises, the wearing of prisms, and, as a last resort, operation.

I. Correction of the refractive error is of the greatest importance, and frequently is curative, though some cases are uninfluenced by glasses. An esophoria due to convergence excess—*i.e.*, one which is greatest at close range, is usually corrected by the constant use of the full hypermetropic and astigmatic correction; if myopia is present it should be under-corrected. A convergence insufficiency causing an exophoria for near range calls for a full correction of the myopia and an

under-correction of the hypermetropia. An exophoria or an esophoria due to a divergence anomaly—*i.e.*, most pronounced for distance, is not materially affected by correction of the refractive error.

2. Attention to the general health is a necessary and valuable adjunct to local treatment, especially in neurasthenic and debilitated individuals who show a high degree of convergence weakness at close range and a remote near point of convergence, with no refractive error to account for the deviation (see below).

3. Prism exercises are used chiefly in cases of convergence weakness of non-accommodative origin. In esophoria and hyperphoria such exercises are less satisfactory. Starting with a weak prism, base out, before one eye, stronger and stronger prisms are inserted every few seconds till the patient can no longer fuse the images. He is gradually improved by exercises till he can overcome at least a 50-degree prism base out. The exercise is used at 20 feet or 13 inches, or at both distances. It is continued for several minutes two or three times daily, and must be persisted in for several weeks to give results.

In an uncomplicated case of weakness of convergence simple exercises should first be tried. The patient begins reading a book at the ordinary distance. Then, while still reading, he brings the book nearer his eyes till the print blurs. He then slowly removes the book to its first position, repeating the process ten or a dozen times, once or twice a day for a month. The amblyoscope is useful in such cases as well as in cases of over-convergence. The exercises given in squint training are reversed in order, starting with those in which a desire for fusion helps to increase the range of binocular fixation and finishing with the simplest exercises in which there is no such 'tie' for the eyes.

4. Prisms for wear may be used to correct deviations of low degree. The base of the prism is placed in the direction of the muscle it is desired to help—*i.e.*, in hyperphoria, base down; in esophoria, base out; in exophoria, base in. They are most satisfactory in hyperphoria in which low degrees of imbalance often give rise to severe symptoms requiring full

prism correction. In cases of greater degree a partial correction is often satisfactory. In esophoria, which is not corrected by glasses, prisms base out, which represent the deficiency of abduction (*not* the degree of esophoria), may be required. In exophoria they are seldom of use, but in persistent deficiency of convergence not corrected by other means they may be of great service when incorporated in the glasses for close work. Any prism over 1 degree should be divided between the two eyes.

If glasses are worn, the effect of a prism may be obtained by decentration of the lenses, if such lenses are of sufficient strength—*i.e.*, by displacement of the optical centre so that it no longer corresponds to the geometrical centre of the lens. Decentering a convex lens in, or a concave lens out, produces the effect of a prism with its base towards the nose; decentering a convex lens up or a concave lens down gives the effect of a prism with its base up. A lens of 1 D. must be decentered 8.7 mm. to produce the effect of a prism of 1 degree. To calculate the amount of decentration necessary to produce a certain prismatic effect, we multiply 8.7 by the value of the prism, and divide the result by the strength of the lens in diopters. For example a +4 D. lens with 2-degree prism base in equals $\frac{8.7 \times 2}{4} = 4.3$ mm.; such a lens should be decentered inward 4.3 mm. to have the added effect of a 2-degree prism base in.

5. Operation, if performed in carefully chosen cases, gives satisfactory results; it should not be resorted to until one is certain no other method will suffice. In esophoria a recession of the internal rectus, or an advancement of the external, may be done. In exophoria an advancement of the internal rectus, or in some cases a tenotomy of the externus, gives good results. As a rule hyperphorias should not be touched, but occasionally it may be necessary to advance the inferior rectus of the hyperphoric eye, to tenotomize (or recess) the inferior rectus of the opposite eye, or to recess the superior rectus of the hyperphoric eye.

CHAPTER XXIX

OPERATIONS ON THE EXTERNAL OCULAR MUSCLES

As mentioned in Chapter XXVII., the operations employed in the correction of a squint are: (1) Advancement, which brings the attachment of a muscle further forwards; alternatives are resection, which reattaches a shortened muscle to its original insertion, and tendon-tucking, which permanently shortens a muscle by folding a portion of it on itself without cutting through its insertion. (2) Tenotomy, with its modification of recession. These operations are employed either separately or combined.

Advancement.

Numerous methods have been designed for this operation, mainly varying with the manner in which the sutures are inserted. Many surgeons use three simple sutures, passing one through the edge of the overlying conjunctiva and the centre of the tendon, which has been cut through at its insertion and separated from its scleral attachments, and then through the sclera between the original insertion and the limbus; this central suture may be single, double, or a mattress suture, according to the preference of the individual surgeon. The other sutures are then passed through the conjunctiva and tendon, one on each side of the central suture, and then through the sclera and conjunctiva, a little obliquely, towards the upper and lower margins of the cornea.

Worth's operation was specially designed to get a firm, unyielding hold for the sutures at each end, so that any desired degree of rotation of the eyeball may be produced. No tenotomy of the opposing muscle is done. In squints of high degree it is necessary to operate on both eyes.

The anatomical relations of the advanced muscle are disturbed as little as possible. The immediate effect produced is the final result.




FIG. 336.—SQUINT SCISSORS.

FIG. 336A.—LANCE-HEADED CUTTING NEEDLE FOR SCLERA.



FIG. 337.—
HOOK.



FIG. 338.—
NEEDLE HOLDER.

The instruments required are speculum (Fig. 185), straight blunt-pointed scissors (Fig. 336), two fixation forceps (Fig. 186), advancement forceps (Fig. 339), needle-holder (Fig. 338), small curved needles, straight lance-headed cutting needles

for the sclera, with split eye (Fig. 336A), and silk thread. A general anæsthetic is required for young children or timid patients. In other cases local anæsthesia will suffice.

The eye is anæsthetized with cocaine. Adrenalin is instilled before and from time to time during the operation. The patient lies on a table with his feet towards the window. His lids are held open by the speculum. The surgeon, standing behind the patient's head, grasps the conjunctiva with the toothed forceps, while with the scissors he makes a curved vertical incision through it rather more than $\frac{1}{2}$ inch long. The convexity of the incision is close to the corneal margin. A similar incision is then made through the capsule of Tenon. The conjunctiva and capsule then retract, or, if necessary, they are pushed back so as to expose the insertion of the




FIG. 339.—ADVANCEMENT FORCEPS (IMPROVED PATTERN).

tendon. One blade of the advancement forceps is now passed under the tendon after the manner of a tenotomy hook, the other blade being superficial to the conjunctiva. The forceps is now closed, so that tendon, capsule of Tenon, and conjunctiva are all firmly clamped together, with their relations undisturbed except for the retraction of the membranes. The tendon and a few little fibrous bands beneath the tendon are now divided with scissors at their insertion into the sclerotic. The advancement forceps, holding the tendon, capsule, and conjunctiva, can now easily be lifted up so as to get a good view of the under-side of the muscle.

One of the needles is then passed inwards at A, through conjunctiva, capsule, and muscle, and brought out at the under-side of the muscle. It is then again passed through

muscle, capsule, and conjunctiva, and brought out at B. The bight of the thread thus encloses about the lower fourth of the width of the muscle, together with its tendinous expansions and capsule and conjunctiva. The other needle is similarly entered at A', passed through conjunctiva, capsule, and muscle, and brought out at the under-side of the muscle. It is then entered again at the under-side of the muscle, and




FIG. 340.—WORTH'S OPERATION OF ADVANCEMENT OF AN OCULAR MUSCLE.

brought out through the conjunctiva at B', the bight of this suture thus enclosing the upper fourth of the width of the muscle, etc. The object of inserting both sutures before proceeding further with either is that they may be symmetrically placed. The ends of the thread from A' and B' are then crossed over at C. The end bearing the needle is then entered at D, and passed through conjunctiva, capsule, and muscle, and brought out beneath the lower blade of the

advancement forceps. The first suture is then similarly dealt with. The anterior part of the muscle and capsule and conjunctiva are then removed by cutting them through with scissors behind where they are grasped by the advancement forceps.

The next stage in the operation, the insertion of the two sutures into the sclerotic at G and G, is one which requires great judgment and delicacy of touch. Take a lance-headed needle (Fig. 336A) in the holder. Remove the curved needle from a suture and insert the latter into the split eye. With the fixation forceps take a firm hold of the globe at the site of the old insertion. Lay the head of the needle flat on the sclerotic exactly in line with the position of the suture in the muscle, and about $\frac{1}{8}$ inch, or a trifle more, from the corneal margin. Press the point of the needle backwards so as to dimple the sclerotic a little. Then push it onwards so that it traverses at least half the thickness of the sclerotic, but taking the greatest care not to pierce the whole thickness. The longitudinal extent of the insertion is about $\frac{1}{8}$ inch. The other suture is then similarly inserted. No verbal direction can be given for gauging the depth. One judges by sight and touch. The gap is now closed, and the eyeball rotated into its correct position by tying each suture at HH.

In operating under cocaine, before the knots are tied at HH, an assistant holds the globe in the primary position with forceps, while the patient is told to try to look away from the operated muscle. This relaxes the muscle while it is being drawn forward by the sutures. The sutures are then temporarily secured at HH by the first hitch of the 'surgeon's knot.' The assistant then releases the globe. The fine adjustment is done by tightening or loosening the hitches at HH, the result being checked by the mirror test or by the reflection of a candle flame on the cornea. The surgeon's knots at HH are then completed.

The longitudinal position on the muscle of the knotted loops ABC, A'B'C' varies approximately according to the degree of rotation required.

Modifications are often employed to suit cases; for example,

several additional sutures are sometimes used. Care should be taken to avoid injuring the fascial covering of the under-surface of the muscle, lest the muscle form adhesions to its old insertion. If this should happen, the mobility of the eye in the opposite direction would be greatly impaired.

The patient should be kept in bed with both eyes bandaged for a week after the operation and sutures removed on the eighth or tenth day.

Tendon-Tucking.—Various operations have been employed for permanently folding tendon (and muscle) on itself. The muscle and tendon are exposed, freed from all attachments to the sclera, and then a portion of the muscle and tendon is folded on itself by means of a specially constructed double or triple hook; the folds of tendon are then sewn together with catgut and a permanent shortening of the muscle produced.

Tenotomy.

There are several slightly different methods of performing this little operation. The tendon may be exposed by free conjunctival incision or the operation may be done sub-conjunctivally through a small opening just sufficient to admit the scissors. The instruments required are speculum (Fig. 185), straight blunt-pointed scissors (Fig. 336), fixation forceps (Fig. 186), and tenotomy hook (Fig. 337). The patient is prepared as for advancement. Insert the speculum. Tell the patient to look in a direction opposite to that of the tendon to be divided, so as to bring its insertion well forward. With the forceps pick up the conjunctiva over the insertion of the tendon, and with the scissors make an incision about $\frac{1}{8}$ inch long, in a direction at right angles to that of the tendon. Now



FIG. 341.—TENOTOMY.

divide the capsule of Tenon in the same way. This brings the insertion of the tendon into view. While the forceps still hold up the cut edge of the capsule, make a few short snips with the scissors near one border of the tendon until the point of the scissors is felt to slip freely back without encountering any resistance. Now lay down the scissors, and take up the hook in the right hand. Pass the point of the hook into this incision, and hook it round the insertion of the tendon until it appears at the other border. During this ~~manœuvre~~ the point of the hook should be kept in contact with the sclerotic. Now lay down the forceps and transfer the hook to the left hand. Take care to avoid any dragging with the hook, as this causes pain. With the scissors cut between the point of the hook and the globe until the tendon is divided at its insertion and the hook comes away. It is usual to reintroduce the hook to seek for any fibres of insertion which may have escaped division. Where the effect produced by the tenotomy appears very small there is a temptation to divide the indirect attachments above and below. This should on no account be done. It is not necessary to suture the conjunctiva if the conjunctival incision is small. A restraining suture is sometimes passed through the muscle prior to tenotomy, so that, if an over-effect has been produced, the muscle can be drawn forward and sutured to the sclera so as to produce exactly the effect desired.

After the operation there is a very considerable defect of movement in the direction of action of the tenotomized muscle. This, to some extent, subsequently disappears. The *average* effect of a tenotomy of the internal rectus is 13 degrees, and of the other recti less than half this amount. But it varies within extremely wide limits. In some cases of unguarded tenotomy of an internal rectus the eye gradually deviates outwards in after years.

A pad and bandage should be worn for the first three or four days, after which it may be discarded. The eye should be bathed with boric lotion three or four times a day until the wound is healed.

The above description could be applied to tenotomy of any

of the recti muscles, but the technique of tenotomy of the inferior oblique, which operation is rarely performed, is quite different.

Tenotomy of the inferior oblique is indicated in paralysis of the superior rectus of the opposite eye with spasm of the inferior oblique of the opposite side, paresis of the superior rectus of the opposite eye, spasm of the inferior oblique, and in incurable paralysis of the superior oblique. A curved skin incision $\frac{3}{4}$ inch long is made at the intersection of the lower orbital margin with a perpendicular dropped from the supra-orbital notch. Dissection is made down to and through the septum orbitale close to the orbital margin. The tendon is engaged on a strabismus hook by keeping the hook in contact with the floor of the orbit and sweeping it inward. After being freed, the tendon is severed close to its periosteal attachment and a portion (10 mm.) is removed. No attempt is made to suture the ends. The skin wound is closed with interrupted sutures.

Recession is a modification of tenotomy which is nowadays frequently employed in place of tenotomy, since by this method the amount of retraction of the muscle and its loss of power are definitely limited and controlled. The severed tendon, which has been exposed by free vertical conjunctival incision, and separated, with its check ligaments, from the sclera, is sutured to the episcleral tissue at a point 2.5 mm. behind its original insertion; the distance depends on the degree of the squint. Sutures




FIG. 342.—RECESSION OF THE INTERNAL RECTUS.

of fine 000 ten-day chromic catgut, or fine paraffined silk, are passed, as shown in the illustration (Fig. 342), through the posterior lip of the conjunctival incision, the end of the cut tendon about 1.5 mm. behind its anterior extremity, the very superficial layers of the sclera at a point previously

determined behind its original insertion, and finally through the anterior lip of the conjunctival incision. A muscle must never be recessed beyond the equator (where the venæ verticosæ emerge), the position of which for the internal rectus is approximately 5 mm., and for the external rectus 2.5 mm., behind the insertion of the muscle.

CHAPTER XXX

THE OCULAR MANIFESTATIONS OF GENERAL DISEASES

THE systemic diseases which most frequently give rise to ocular symptoms and diseases are: Syphilis, tuberculosis, rheumatism, nephritis, diabetes, arterio-sclerosis, cardiac affections, diseases of metabolism, chronic intoxications, infective diseases, and affections of the nervous system.

This chapter should be read in conjunction with the earlier chapters, in which the ocular symptoms have been fully discussed.

Diseases of the Blood.

Anæmia and Chlorosis give rise to pale pink conjunctivæ and pearly white scleræ. There may be pallor of the disc and the rest of the fundus, the retinal vessels being pale, tortuous, and the retinal veins broader than normal. Occasionally retinal hæmorrhages are found.

Pernicious Anæmia often causes retinal hæmorrhages, occasionally retinitis. The fundus exhibits great pallor.

Hæmophilia predisposes to profuse hæmorrhage after injury to the eye, and under such circumstances may cause hyphæmia, or hæmorrhage into the retina or into the orbit.

Leukæmia.—Retinal hæmorrhages are very common, and a peculiar form of retinitis, leukæmic retinitis, is often present.

Purpura is often accompanied by hæmorrhage beneath the conjunctiva, in the retina, skin of the lids, and occasionally into the orbit.

Severe Hæmorrhage may be accompanied by amblyopia. The latter may be temporary and accompanied by little or no

ophthalmoscopic change, or it may be permanent and followed by atrophy of the optic nerve. Such sudden and severe anæmia may cause retinal hæmorrhages.

Diseases of the Circulatory System.

Heart.—Valvular heart disease and fatty heart are often accompanied by hæmorrhages into the retina, less frequently into the vitreous. Aortic insufficiency causes pulsation of the retinal arteries. Endocarditis may cause embolism of the central artery of the retina. The œdema dependent upon cardiac disease may involve the eyelids, being noticed especially upon rising in the morning.

Aorta.—Aneurism of the aorta may give rise to mydriasis, enlarged palpebral aperture, and exophthalmos, as a result of irritation of the cervical sympathetic; or to miosis, slight ptosis, and enophthalmos, through paralysis of the same. This condition may also cause thrombosis and embolism of the central artery of the retina, or of one of its branches.

Arterio-sclerosis gives rise to characteristic changes in the fundus, which are described on p. 285, and illustrated on Plate XVIII. It is a predisposing cause of glaucoma.

Diseases of the Digestive System.

Teeth.—The occurrence of ocular symptoms and diseases more or less dependent upon toothache and dental disease is not rare, and in such cases the ocular condition may improve when the offending tooth is filled or extracted. Periapical abscess is the dental condition most liable to cause such trouble. Such symptoms include—Conjunctival congestion, photophobia, epiphora, asthenopia, and weakness of accommodation. Iritis, keratitis, and cyclitis may be dependent upon dental disease; oral sepsis is a common cause of 'quiet cyclitis.'

Stomach and Intestines.—In dyspepsia and chronic constipation absorption of septic matter from the gastro-intestinal tract may give rise to iridocyclitis, choroiditis, and retinitis. The straining associated with constipation may cause sub-conjunctival, retinal, or vitreous hæmorrhage. Hæmorrhage from stomach or intestines may cause anæmia (*q.v.*).

Diseases of the Ductless Glands.

Acromegaly, due to over-activity of the anterior portion of the pituitary body, exhibits many ocular manifestations. There is hypertrophy of the margins of the orbit and thickening of the skin of the lids. Disease of the hypophysis may cause characteristic bitemporal hemianopsia, though other abnormalities of the field of vision are met with; and there is often reduction in the acuteness of vision. There may be optic neuritis and optic-nerve atrophy, and paralysis of one or more of the ocular muscles. Exophthalmos, hypertrophy of the lacrymal gland with epiphora, and sluggish reaction of the pupils, are also seen. Pain in the eyes and brow is sometimes complained of.

Myxœdema and Cretinism give rise to swelling of the eyelids, sometimes optic neuritis.

Exophthalmic Goitre (Graves' or Basedow's Disease).—Though this is a constitutional disease, and the ocular symptoms are not an essential part, the eye exhibits the most striking manifestations of the affection, some or all of the following ocular symptoms being present in almost every case: Exophthalmos is usually present. It varies in degree; it may be slight, or the proptosis may be so pronounced that the patient cannot cover the cornea with the lids. It is usually bilateral, but occasionally it affects only one eye. Von Graefe's sign consists in a failure of the upper lid to follow the eyeball normally when the patient looks downward; the upper lid lags behind. Stellwag's sign is the diminution in the normal involuntary power of nictitation, as a result of which winking is imperfect, less frequent, and more irregular than normal. Vision is not usually involved. The cornea may, however, suffer when the exophthalmos is extreme and causes much exposure; in such cases the lower part may become vascular, or dry, or ulcerated, and occasionally destruction of the eyeball is the outcome.

There is a brownish pigmentation of the skin of the lids in some patients. There may be arterial pulsation visible in the fundus. Dilatation and inequality of the pupils may be

present. The extrinsic ocular muscles, especially the abducens, may be the seat of paresis.

Diseases of the Ear.

Choked disc and congestion of the papilla are frequently observed in sinus thrombosis complicating mastoiditis. Nystagmus is common and of great diagnostic import in affections of the labyrinth.

Infective Diseases.

Cerebro-spinal Meningitis is often accompanied by ocular symptoms. Conjunctivitis occurs frequently. Œdema of the lids and conjunctiva may be seen. There may be paresis of the extrinsic ocular muscles, causing strabismus and ptosis. Nystagmus is encountered. There may be abnormalities of the pupils, keratitis, retinal hæmorrhages, optic neuritis, and optic nerve atrophy. Irido-choroiditis, and purulent choroiditis leading to pseudo-glioma, are not uncommon.

Diphtheria.—With the exception of diphtheritic conjunctivitis, which is now rather rare, the ocular manifestations of diphtheria occur after the acute stage of the disease has passed, and are therefore really post-diphtheritic symptoms. The latter include paralysis of one or more of the extrinsic muscles of the eye, usually the external rectus, and paralysis of accommodation. Occasionally optic neuritis occurs.

Erysipelas, when it spreads to the eye, causes great swelling and redness, so that the lids can be separated only with great difficulty. Following this there may be abscess of the eyelids, with sloughing of the skin. When the disease extends into the orbit, it causes orbital cellulitis, with exophthalmos and sometimes ulcer of the cornea. Thrombosis of the retinal veins, optic neuritis, and atrophy of the optic nerve, may follow under such circumstances. Glaucoma sometimes results, and occasionally inflammation of the lacrymal gland and sac.

Gonorrhœa is responsible for the local infection of the conjunctiva, resulting in purulent conjunctivitis in adults, and in

ophthalmia neonatorum in the new-born. It also gives rise to chronic iritis. The affection is analogous to gonorrhœal arthritis, and is due to the presence of toxins. For some time after an attack of gonorrhœa is thought to have been cured, the vesiculæ seminales may contain gonococci, which may continue the intoxication.

Influenza is almost always accompanied by congestion of the conjunctiva. There is frequently severe pain in and behind the eyeballs. A great many ocular manifestations credited to influenza are probably dependent upon the marked depression which follows the disease; in this category may be placed weakness of accommodation and severe asthenopia. Infrequent ocular complications include corneal ulcer, pareses of extrinsic ocular muscles, retrobulbar neuritis, optic neuritis, optic nerve atrophy, and orbital cellulitis.

Leprosy attacks the eyelids similarly to the skin of the face. The conjunctiva and cornea may also be attacked.

Malaria infrequently gives rise to the following ocular manifestations: Herpes corneæ febrilis, optic neuritis, retrobulbar neuritis, hæmorrhages into the retina and vitreous amblyopia, and paresis of accommodation.

Measles is regularly accompanied by a catarrhal conjunctivitis, with subjective symptoms of greater or less severity. In addition there are very frequently blepharitis, phlyctenulæ, hordeola, superficial corneal ulceration, and asthenopia.

Mumps is complicated by dacryo-adenitis in a small number of instances; this rarely leads to suppuration. Œdema of the lids and chemosis may be present.

Pneumonia may be complicated by herpes of the cornea, sometimes followed by corneal ulceration.

Scarlatina.—Catarrhal conjunctivitis is an ocular complication of scarlatina, but it is less frequent and milder than in measles. Corneal ulcer is sometimes seen. Both of these complications are more apt to be found in the convalescent stage than early in the disease. When this disease is complicated with nephritis, the characteristic fundus picture of albuminuric retinitis may be seen.

Septicæmia and Pyæmia give rise to retinal hæmorrhages, and sometimes to the lodgment of emboli in the choroid and retina. In the latter case the complications result either in purulent choroiditis, followed by pseudo-glioma, or in panophthalmitis.

Syphilis is frequently responsible for ocular disease. The primary sore may occur on the lids or conjunctivæ. Iritis is due to syphilis in perhaps 25 per cent. of cases. It is an early symptom of the secondary stage, at which time the anterior segment of the eyeball is the vulnerable part. The later stages of syphilis are more prone to attack the posterior segment, causing choroiditis, chorio-retinitis, optic neuritis, and diffuse opacity of the vitreous. In the tertiary stage, gummata may be deposited in the iris, ciliary body, and the periosteum of the orbital wall, and there may be optic neuritis and optic-nerve atrophy, rarely interstitial keratitis. During this tertiary period, paralysis and pareses of the ocular muscles, both extra- and intra-ocular, are quite common. Inherited syphilis is responsible for at least the great majority of instances of interstitial keratitis, and also for some congenital ocular defects.

Tuberculosis, though rather rarely involving the eye, may affect the iris, choroid, and sclera, presenting characteristic deposits; still more infrequently the conjunctiva and lids present tubercular disease. In acute general miliary tuberculosis and in tubercular meningitis it is not uncommon to find small tubercle deposits scattered over the fundus. In pulmonary tuberculosis the pupils are often unequal in size.

In the 'strumous diathesis' there is a predisposition to blepharitis, chronic conjunctivitis, phlyctenular conjunctivitis and keratitis, and perhaps interstitial keratitis.

Vaccinia.—There have been a number of examples of accidental inoculation of the eyelids and conjunctiva with vaccine virus. In such cases the pustules excite marked swelling and induration, involvement of the pre-auricular glands, and tendency to deformity of the lid from subsequent cicatrization.

Varicella may be complicated by conjunctivitis. The eruption may involve the conjunctiva and cornea, resulting in a superficial ulcer of little consequence.

Variola is responsible for destructive lesions of the eyelids and eyeball. The lids and conjunctivæ are often the site of pustules, and the subsequent cicatrices may cause deformity. Though pustules rarely appear upon the cornea, this part of the eye is not infrequently the seat of keratitis and of ulceration; the latter sometimes results in perforation, and may present as sequelæ opacities, adherent leucoma, or even destruction of the globe.

Whooping-Cough.—Sub-conjunctival hæmorrhage is often seen as a result of the severe paroxysms of coughing. Occasionally such an extravasation of blood takes place in the lid; rarely it involves the orbit, causing serious damage.

Yellow Fever, in its early stage, presents congestion of the conjunctiva. This redness is modified by the addition of yellowish discoloration at a later stage. Sub-conjunctival and retinal hæmorrhages are also found.

Diseases of the Kidneys.

Nephritis presents many ocular manifestations. (Edema is often present in the lids, and may also show itself in the conjunctivæ (chemosis). Albuminuric retinitis is common, occurring most frequently with the chronic parenchymatous variety, but also with other varieties, including the nephritis of scarlatina and pregnancy. During an attack of uræmia, amblyopia without ophthalmoscopic changes may be present; the pupils are dilated during this state.

Miscellaneous Diseases and Conditions.

Diabetes.—The common ocular complications of diabetes are cataract and hæmorrhages in the retina. Less frequently there occur retinitis, optic neuritis, retrobulbar neuritis, iritis, pareses of the external ocular muscles, and paralysis of accommodation. Diabetics occasionally present sudden and marked changes in the state of refraction of the eye, especially myopia,

accompanying an increase in the amount of sugar in the urine, or due to pre-cataractous swelling of the lens.

Gout is sometimes responsible for episcleritis and scleritis, and rarely for marginal ulcer of the cornea, glaucoma, and hæmorrhagic retinitis. Gouty individuals often complain of dry catarrh, a condition in which the conjunctiva is congested, and the patient experiences a hot feeling in the lids and a sensation as though a foreign body were present. Such patients are sometimes subject to attacks of transient periodic episcleritis.

Headache, when persistent or frequently recurring, should always lead to a careful examination of the eyes. Errors of refraction are common causes of headache and neuralgia. Not infrequently we find anomalies of the extrinsic ocular muscles (heterophoria); less often, presbyopia and accommodation weakness. The error of refraction which most commonly gives rise to headaches is astigmatism; less often hypermetropia is responsible. The amount of astigmatism may be very small; even 0.25 or 0.50 D. may cause trouble to a sensitive person who uses the eyes much in near work. The site of the pain produced by uncorrected errors of refraction varies, but it is often supra-orbital and frontal. Depreciation of the general health is in many cases a predisposing factor (and a result); thus, we often find that the glasses required to relieve headaches in individuals who were debilitated are no longer necessary when the system has regained its normal tone.

The constant effort to keep the eyes straight in a case of 'neuropathic divergence' usually causes a dull chronic headache. Similar symptoms may follow tenotomy of an internal rectus muscle.

Migraine.—This affection, thought to depend upon some disturbance in the circulation of the cerebral cortex, is characterized by periodic or irregular attacks, which begin with more or less blurring of vision. This visual defect occurs with or without scintillating scotoma, and is often more or less hemianopic in character. After a period varying from several minutes to half an hour, vision again becomes normal; then a very severe headache develops, accompanied often by nausea

and vomiting, and is followed by marked general depression. Though dependent, in part at least, upon depreciation in general health and excessive use of the eyes, the attacks are often aggravated by eye-strain. In such cases the seizures are prevented or made less severe by correction of errors of refraction or of heterophoria.

One rare form of migraine is associated with a temporary paralysis of the muscles of the eyeball, usually recovering quickly, sometimes lasting for two or three weeks. This is known as ophthalmoplegic migraine.

Rheumatism has been thought to be the etiological factor in some cases of scleritis, episcleritis, tenonitis, and palsies of the extrinsic ocular muscles. But it is more correct to say that rheumatism and these other diseases are caused by the same toxin. Most cases of iritis supposed to be due to rheumatism are probably gonorrhœal or toxic.

Rickets.—The subjects of rachitis often present congenital cataract (zonular), interstitial keratitis, and phlyctenular kerato-conjunctivitis.

Scurvy is often accompanied by hæmorrhages beneath the conjunctiva, in the retina, skin of the lids, and occasionally into the orbit. It not infrequently presents a form of night blindness, which disappears when the general health improves.

Vertigo, with or without nausea, is often dependent upon ocular refractive errors, or to insufficiencies of the extrinsic ocular muscles, or perhaps pareses of these muscles.

Diseases of the Nervous System.

The eye furnishes information of great importance in the diagnosis of diseases of the nervous system, the intimate relationship between this part of the human anatomy and the visual organs being evident. Particulars regarding the condition of the optic nerves, the pupils, the eye muscles, the acuteness of vision, and the fields of vision, are of great value.

Apoplexy gives rise to a number of ocular manifestations, varying according to the part of the brain involved. Retinal

hæmorrhages may precede the cerebral affection, and may serve as a warning of impending danger.

Encephalitis Lethargica frequently presents as an early symptom paralysis of the third cranial nerves, partial or complete, of one or both sides, giving rise to ptosis, squint, diplopia, and pupillary disturbances, especially eccentricity of the pupil. Sometimes the fourth or sixth nerve is involved. Nystagmus is common. Rarely there is optic neuritis.

Friedreich's Disease has no ocular disturbances, excepting a peculiar nystagmus, generally present, consisting of irregular twittings seen when the eyes are fixed upon a moving object in the horizontal direction. Ocular palsies, optic neuritis, and Argyll-Robertson pupils occur rarely.

Meningitis often presents optic neuritis, abnormalities of the pupils, and palsies or spasms of the ocular muscles, causing deviations. These ocular manifestations are seen most frequently in tubercular meningitis, in which variety tubercles of the choroid are not infrequently found.

Myelitis is in rare instances accompanied or preceded by retrobulbar optic neuritis, with severe pain in the brow and orbit.

Myasthenia Gravis is almost always accompanied by bilateral ptosis and weakness of the orbicularis; the ptosis increases with fatigue and is always least in the morning and most marked in the evening. In many cases there is also a partial or complete ophthalmoplegia externa; but the extrinsic muscles are not affected. Nystagmoid movements may be present, but are not common.

General Paralysis.—The subjects of this disease often present inequality and irregularity of the pupils, also miosis, and less frequently mydriasis. There is not uncommonly impairment or loss of the light reflex (Argyll-Robertson pupil); later there is added partial or complete loss of the reaction to accommodation. Sometimes atrophy of the optic nerve, with its attendant reduction in the acuteness of vision and restriction in the size of the field, is noted. Palsies of the third, fourth, and sixth nerves may occur, giving rise to diplopia, strabismus, and ptosis.

Disseminated Sclerosis presents many ocular manifestations; the latter are found in fully one-half of the cases. Nystagmus is a frequent symptom. The fields of vision often exhibit irregular peripheral contraction and central scotoma, either relative or absolute. An incomplete optic-nerve atrophy, generally unilateral, is of common occurrence, resulting from retrobulbar neuritis. There are also partial paralyses of the extra-ocular muscles, giving rise to diplopia.

Tabes is accompanied by many ocular signs. The Argyll-Robertson pupil, in which the reaction to light is lost, while that to convergence and accommodation is preserved, is present in the great majority of cases, and usually exists on both sides. A deviation from the circular shape, inequality and marked contraction of the pupil (miosis), are very common; much less frequently mydriasis is present, but it is then frequently associated with blindness. Atrophy of the optic nerve occurs frequently, is an early symptom, is progressive, and generally leads to blindness. With this change in the optic nerve there is reduction in the acuteness of vision and concentric contraction of the field. Ocular palsies are very common; they often occur early in the disease, involve the third and sixth nerves, rarely the fourth, appear suddenly in many instances, are generally transient, and are accompanied by diplopia; if the third nerve is involved, also by ptosis. Epiphora is sometimes observed, also inco-ordinated movements of the eyeballs.

Tumour of the Brain (including Abscess) gives rise to choked disc in the majority of cases; this is generally bilateral, and probably is in most instances more marked on the side of the growth. There may be palsies of the ocular muscles and alterations in the field of vision. The characteristics of these changes may be of great aid in the localization of the seat of the tumour.

Disease of the Pituitary Body—commonly the development of adenomatous growths or cysts—frequently causes impairment of vision associated with partial or complete atrophy of the optic nerve, alterations in the visual field are common, most characteristically consisting of bitemporal hemianopia, the

field generally contracting from the temporal side inwards and from above downwards; paracentral and central scotomata are also frequently found. Since there is usually a greater advance in the visual defects in one eye as compared with the other, there may be hemianopsia in the eye first involved and hemichromatopsia in the one which became affected later. Very rarely papillœdema is seen in this affection, but paresis of an ocular nerve, generally the third, is not uncommon. X-ray usually shows enlargement of the pituitary fossa, but some tumours arise in the stalk and are above the sella turcica; in such cases the fossa is not enlarged.

Functional Nervous Disorders.

Chorea.—Patients who suffer from choreic movements of the muscles of the lids and of the face and neck often suffer from errors of refraction, less frequently from lack of equilibrium of the eye muscles.

Coma.—Objective examination of the eyes may give important data in all forms of coma. If dependent upon organic brain disease, there may be choked disc, mydriasis, and deviation of the eyes. If due to cerebral hæmorrhage, there may be miosis, inequality of the pupils, and conjugate deviation. With increased intracranial pressure there may be dilated pupils. If accompanying uræmia, albuminuric retinitis may be found. When alcoholic, there may be dilatation of the pupils, and pareses of external ocular muscles. If due to poisoning by opium or similar drugs, there may be extreme miosis.

Epilepsy.—The seizure frequently begins with a visual aura: transient flashes of light, coloured sensations, and hemianopsia, or complete loss of vision. During the attack there may be narrowing of the retinal arteries; the pupils are generally dilated, and the light reflex is lost, and there is often spasm of the extrinsic ocular muscles, causing conjugate lateral deviation of the eyes. After the seizure there is distension of the retinal veins, often alterations in the size of the pupils, and not infrequently a temporary concentric con-

traction of the field of vision and a reduction in the acuteness of vision. Not very often, but certainly in some cases, epilepsy is made worse by eye-strain, and the number and severity of attacks are reduced by the wearing of proper glasses.

Hysteria is sometimes responsible for a great variety of ocular symptoms, the principal ones being diminution in the acuteness of vision (amblyopia, and even blindness), concentric contraction of the field of vision for form and colours becoming more marked with each repeated examination, and reversal in the relative size of the colour fields. Other ocular symptoms occurring in hysteria are scotoma, hemianopsia, photophobia, blepharospasm, and monocular diplopia. The pupillary reflexes and the ophthalmoscopic appearances are normal. The ocular manifestations are usually referred to one eye.

Diseases of the Nose, Naso-Pharynx, and Accessory Sinuses.

The communication between the nose and the conjunctival sac by means of the lacrymal duct explains the frequent occurrence of ocular symptoms and affections as a result of nasal disease. In coryza and 'hay-fever' there is very often conjunctival congestion or acute catarrhal conjunctivitis, with marked lacrymation. In chronic rhinitis, whether catarrhal or hypertrophic, chronic conjunctivitis, blepharitis, and phlyctenular affections are very common. In addition, the nasal swelling may obstruct the lower end of the lacrymal duct, and as a result there may be lacrymal stenosis, dacryocystitis, and lacrymal abscess. The lacrymal duct may be the means of conveying infective material from the nose to the conjunctival sac, and thus explain the occurrence of corneal ulcer.

Adenoids not infrequently give rise to catarrhal conjunctivitis, follicular conjunctivitis, epiphora, and asthenopia.

Diseases of the accessory sinuses (maxillary, ethmoid, sphenoid, and frontal) are not infrequently responsible for ocular symptoms and diseases, the principal of which are :

Exophthalmos, paresis or paralysis of the ocular muscles (both extrinsic and intrinsic), optic neuritis, and atrophy of the optic nerve. Abscess of the sphenoidal sinus may cause bitemporal hemianopia, simulating pituitary tumour.

Poisonings and Intoxications.

These conditions are not infrequently responsible for ocular symptoms and disease, especially retrobulbar neuritis (less frequently optic-nerve atrophy), which results from poisoning by tobacco, wood-alcohol, iodoform, lead, arsenic (atoxyl), bisulphide of carbon, and nitro-benzol.

Pregnancy and Parturition.

Pregnancy may be complicated by gravidic retinitis, which may be so marked as to justify premature delivery, in order to save sight.

Parturition is accompanied by danger to the eyes of the child. Conjunctival infection may give rise to ophthalmia neonatorum. The use of the forceps during delivery has resulted in bruising of the lids, injury to the sixth nerve, injury to the cornea, orbital hæmorrhage, causing exophthalmos, and even rupture of the eyeball. During this period the eyes of the mother may present, on rare occasions, retinal hæmorrhages; and if there has been great loss of blood, amblyopia without ophthalmoscopic changes, or reduction of vision with subsequent optic-nerve atrophy, may ensue. Puerperal infection may result in metastatic choroiditis or in panophthalmitis, with loss of the eye. Parturition may also be followed by optic neuritis, atrophy of the optic nerve, retrobulbar neuritis, retinal hæmorrhages, and embolism of the central artery of the retina, though all of these complications are rare.

CHAPTER XXXI

OCULAR THERAPEUTICS: GENERAL RULES FOR OPERATIONS UPON THE EYE

THE eye, being a very delicate and sensitive organ, may easily be injured by injudicious use of local applications. It is essential, therefore, to observe especial care as to the manner and strength in which such remedies are employed.

Constitutional Remedies.—When any disease of the eye presents itself, one should consider whether it may be due either wholly or in part to some general constitutional disease. Syphilis, tuberculosis, Bright's disease, diabetes, tabes, chronic intoxications, anæmia, and other disordered states often give rise to well-marked eye symptoms. In such cases it is obvious that one cannot hope to cure the ocular disease unless one treats the constitutional disease of which it is a symptom.

Local Remedies.—Drugs intended for local use to the eye are usually dissolved in water or oil, or used in the form of ointment or powder.

Cleansing and Antiseptic Solutions.

These are employed for washing out the conjunctival sac and removing secretion. They should be used freely, and should be lukewarm when employed. They may be allowed to run between the lids from an undine (Figs. 343 and 344), from a wad of sterilized cotton-wool, or from an eye-dropper (using two or three droppers full).

The eye bath or cup is a popular means of applying a lotion to the eye, for it can be used by the patient without help. The eye-bath should be quite filled with the lotion, and should fit the contour of the orbit so that it can be held vertically over

the eye with the head tilted backwards without spilling the lotion after it has been placed in position. The patient then opens and closes his eye several times and turns it in different directions so that the lotion reaches the whole of the conjunctival sac. All discharge that collects on the lid margins and at the inner canthus must be thoroughly wiped away and the lids dried. It is most useful in cases of blepharitis; it must not be used where there is ulceration of the cornea, as the movements of the lids necessary in using the cup are contra-indicated in this condition. It is not satisfactory in use when there is copious discharge, in which case irrigation should be employed.




FIG. 343. — UNDINE FOR IRRIGATING THE EYE.




FIG. 344. — METHOD OF IRRIGATING THE EYE WITH A SOLUTION POURED FROM AN UNDINE.

The cup must be thoroughly cleansed before treating the second eye.

The cleansing and antiseptic solutions which are used most frequently are:

1. Sterilized water.
2. Boric acid in saturated solution (about $\frac{1}{2}$ ounce to the pint).
3. Sodium chloride in physiological strength (0.6 per cent.; a teaspoonful to the pint).
4. Mercuric chloride, from 1 : 20,000 to 1 : 10,000.

Boric Acid is used more frequently than any other of these remedies. Though chemically an acid, it is neutral, bland, and soothing. It is often employed to irrigate the eye during operations. It is frequently prescribed with white vaseline (gr. x. ad ʒi.), in the form of an ointment, to prevent adhesion of the lids overnight, when there is considerable discharge.

It sometimes, especially in older people, gives rise to irritation of the skin, and should then be replaced by normal saline or an alkaline lotion—sod. bicarbonat., gr. x. ad ʒi.—or the following wash:

R	Sodii biborat.	}	-	-	-	-	āā	gr. xi.
	Sodii bicarbonat.							
	Sodii chlor.							
	Hazolini	-	-	-	-	-	ʒiv.	
	Aquæ destill.	-	-	-	-	-	ad ʒx.	

M. Ft. collyr. Sig.: Dilute with equal part of warm water for use in eye-bath.

This lotion will be found very soothing and useful for many mild conjunctival irritations, and after exposure of eyes to wind and dust.

Stimulating and Astringent Remedies.

The remedies of this class used most frequently in diseases of the eye are: zinc sulphate, tannic acid, alum, borax, potassium chlorate, camphor, silver nitrate, copper sulphate, yellow oxide of mercury, ammoniated mercury, and calomel. They are intended to cure abnormal conditions of the conjunctiva, and are used principally in various forms of conjunctivitis. For this purpose they are prescribed in small quantity. Two or three drops of a watery solution are allowed to fall upon the everted lower lid from an eye-dropper; the latter must not be allowed to touch the lids, since this would lead to contamination of the liquid. Copper sulphate and alum are frequently employed in the form of a solid pencil.

Zinc Sulphate is used very largely in astringent collyria.

R	Zinci sulph.	-	-	-	-	-	gr. ij.
	Aquæ destill.	-	-	-	-	-	ʒi.

M. Sig.: Two drops in each eye three times a day.

R	Zinci sulph.	-	-	-	-	-	gr. ss.
	Acidi borici	-	-	-	-	-	gr. v.
	Aquæ destill.	-	-	-	-	-	ʒi.

M. Ft. collyr.

Tannic Acid is frequently used in combination with other astringents. Solutions of 5 to 25 per cent. are painted on everted lids or instilled in drop doses by the patient in trachoma.

R	Acidi tannici	-	-	-	-	-	gr. ss.
	Zinci sulph.	-	-	-	-	-	gr. ss.
	Aquæ destill.	-	-	-	-	-	ʒi.

M.

Alum ($\frac{1}{4}$ to 1 grain to the ounce). Long-continued use is said to injure the cornea.

A crystal of alum, in the form of a flattened pencil, is applied to the everted lids in chronic conjunctivitis and in mild forms of trachoma.

Borax is used as a cleansing wash (ʒi. to Oi.), or in combination with other remedies.

R	Zinci sulph.	-	-	-	-	-	gr. ss.
	Sodii biborat.	-	-	-	-	-	gr. iij.
	Aquæ destill.	-	-	-	-	-	ʒi.

Potassium Chlorate is prescribed in solution, 10 grains to the ounce. A soothing lotion in conjunctival irritation.

Camphor.—Though feebly soluble in water, such solution (aqua camphoræ) is stimulating and astringent, and is often incorporated in collyria.

R	Acidi tannici	-	-	-	-	-	gr. $\frac{1}{4}$.
	Zinci sulph.	-	-	-	-	-	gr. ss.
	Aquæ camphoræ	-	-	-	-	-	ʒij.
	Aquæ destill.	-	-	-	-	-	ʒvi.

M.

Silver Nitrate, dissolved in distilled water, may be used in the strength of gr. $\frac{1}{16}$ to gr. $\frac{1}{8}$ to the ounce, dropped into the conjunctival sac. In stronger solution (1 to 5 grains to the ounce) it is brushed upon the everted lids in chronic conjunctivitis and in the papillary stage of purulent conjunctivitis. Solutions of nitrate of silver should be tightly

stoppered and kept in the dark. The brush or cotton applicator should not be dipped into the bottle, but some of the solution should be poured into a small vessel. The strong solutions should be applied only by the surgeon himself. Before the application the cornea should be protected by smearing with vaseline. Any excess of silver is neutralized by washing out the eye with normal saline solution. Silver solutions stain the conjunctiva (argyrosis); hence they should be used only for a limited period. Strong solutions act as caustics.

Copper Sulphate ('Bluestone') may be employed in solution (gr. $\frac{1}{2}$ to $\frac{3}{4}$), but its chief use is in the solid form. A flattened pencil (Figs. 124 and 127) is rubbed across the everted lids (not omitting the retrotarsal fold) in trachoma, and the excess washed off with water or solution of boric acid. The pencil should be flattened and have a blunt, rounded extremity.

It is prescribed, dissolved in glycerine ($\frac{1}{2}$ to 1 per cent.), for home use in trachoma, one drop instilled once or twice a day. It is an excellent remedy for the cure of obstinate cases of chronic catarrhal conjunctivitis; in such cases the pencil is applied very lightly and the conjunctiva flushed at once with boracic lotion.

Yellow Oxide of Mercury is insoluble in water; it is employed in the form of an ointment made with white vaseline, cold cream, or lanoline. The ointment must be thoroughly mixed, and should be preserved in a covered jar which is impervious to light, otherwise it spoils rapidly. It is frequently put up in collapsible metal tubes, or gelatine capsules. Its strength is usually 1 or 2 per cent.

This ointment is very useful in blepharitis, chronic conjunctivitis, phlyctenular keratitis and conjunctivitis, interstitial keratitis, and opacities of the cornea. In blepharitis the ointment is smeared along the margin of the lid; in the other affections a small amount is transferred from the end of a glass rod, probe, collapsible metal tube, or gelatine capsule to the everted lower lid, and thus into the conjunctival sac.

This ointment is often prescribed for many affections in which it is not only of no value but often harmful by producing

irritation; it should be remembered that this remedy is stimulating and somewhat irritating, and is therefore contraindicated in the more acute inflammatory conditions, in which a bland ointment such as 2 per cent. boracic vaseline is more soothing.

Ammoniated Mercury is a white insoluble powder prescribed in the same strength and under the same circumstances as the yellow oxide of mercury.

R	Hydrarg. ammoniat.	-	-	-	gr. i.
	Adepsis	-	-	-	3ij.
	M. Ft. ungt.				

Calomel, free from any trace of perchloride, in the form of an impalpable powder prepared by levigation, is dusted into the eye with a camel's-hair brush in cases of phlyctenular keratitis and corneal ulcers. This combination of mercury is thought to change slowly to corrosive sublimate as a result of contact with the tears. Calomel is apt to produce local irritation if the patient is taking an iodine salt internally, owing to the formation of iodides of mercury.

Ichthyol in 5 or 10 per cent. ointment, or mixed with zinc oxide, forms an excellent application in obstinate cases of ulcerative blepharitis.

R	Ichthyol	-	-	-	-	gr. vi.
	Vaseline	-	-	-	-	3ij.

M. Ft. ungt. Sig.: Apply to the edges of the lids after cleaning.

R	Ichthyol	-	-	-	-	gr. x.
	Ungt. zinci ox.	-	-	-	-	3ij.

M. Ft. ungt. Sig.: Apply to the edges of the lids after removal of crusts.

Lead Acetate *should not be applied to the eye.* It has the property of depositing an insoluble salt of lead upon any corneal abrasion; this stain cannot be removed. Lead and opium wash, so frequently used in other parts of the body, is not, therefore, a desirable application for the eye.

Disinfectants.

True disinfectants (capable of destroying germs) cannot be instilled into the conjunctival sac under ordinary circumstances, since they would injure the cornea. They are, however, applied to circumscribed areas, the excess being washed off by some bland solution. Corneal ulcers, especially when indolent or infected, and purulent conjunctivitis furnish common indications for such use. Some of the remedies classified under this head, though not, strictly speaking, true disinfectants in the strengths in which they are employed, have an inhibitory action upon the growth and development of micro-organisms, and thus act as practical disinfectants. The disinfectants used most commonly in connexion with the eye are mercuric chloride, alcohol, carbolic acid, formalin, tincture of iodine, silver nitrate, protargol, iodoform, and the cautery.

Mercuric Chloride (corrosive sublimate) is prescribed very frequently in purulent conjunctivitis. It may safely be used up to a strength of 1 : 10,000. A stronger solution might damage the cornea, and must consequently be limited in its application to the everted lids, and the excess carefully washed off. A strong solution, 1 : 500, may be rubbed into the conjunctiva after this membrane has been freed from trachoma follicles by the operation of expression. Solutions of corrosive sublimate attack the metal of instruments and dull their cutting edges.

Metaphen, an organic mercury derivative, is a new disinfectant, obtainable in solutions 1 : 500. It is diluted with 4 parts of distilled water and used for irrigation in purulent and other forms of conjunctivitis.

Absolute Alcohol is sometimes used in treatment of dendritic ulcers. It is also used for sterilizing the blades of cutting instruments.

Carbolic Acid (3 per cent. solution) is used for disinfecting instruments. Stronger solutions and pure carbolic acid are frequently applied to infected ulcers of the cornea.

Formalin.—Solutions of 1 : 1,000 and 1 : 2,000 are used in

purulent conjunctivitis; solutions of 1:500 are applied to infected ulcers.

Tincture of Iodine is an excellent remedy in the treatment of infected ulcers. It is applied upon a small piece of cotton rolled upon the end of a probe. Any excess is washed off with water.

Hydrogen Peroxide solution is sold in sealed bottles for dental use. It deteriorates after being opened. Diluted with 3 or 4 parts of water (or stronger), it is an admirable disinfectant and stimulant lotion for conjunctiva, lacrymal sac, and infected corneal ulcers. It is useful for cleansing lid margins in cases of squamous and ulcerative blepharitis, applied on cotton-wool wrapped round a glass rod.

Silver Nitrate is a very efficient and popular disinfectant. In 1 or 2 per cent. solution it is applied to the everted lids in purulent and in other forms of conjunctivitis, and the excess neutralized by a solution of sodium chloride. In Credes' method of prophylaxis against ophthalmia neonatorum a drop of 2 per cent. solution is instilled into the eyes of the new-born child, but a 1 per cent. solution is nowadays recommended. One to 4 per cent. solutions are sometimes applied to indolent corneal ulcers, after careful drying with blotting-paper, with good effect. In fissures at the outer canthus the 'mitigated silver stick' is very useful, but must be carefully applied.

For local anæsthesia preliminary to silver applications solutions of nitrate of cocaine are preferable to the hydrochloride which is generally used, as the latter salt precipitates chloride of silver.

Iodoform is a feeble disinfectant which is occasionally dusted upon corneal ulcers, or used as 2 to 4 per cent. ointment. It is not unfrequently dusted upon wounds after plastic operations. It should be very finely powdered, and may be used as a 2 to 4 per cent. ointment.

Protargol is an organic salt of silver, soluble in water, forming a brown solution. It is used in 5 to 25 per cent. solutions. It is not precipitated by fluids containing sodium chloride and albumin, and is devoid of the irritating qualities of silver nitrate. Its action is feeble. When used for a long time it may stain the conjunctiva.

Argyrol has similar properties, but its germicidal action is weaker than that of protargol.

Ethyl Hydrocuprein (Optochin), a derivative of quinine, is sometimes useful for pneumococcal ulcer in 1 per cent. solution or ointment.

The Cautery (p. 149) gives us the most certain means of limiting the spread of corneal ulcers by destroying the infecting micro-organisms. It is also used in conical cornea (see p. 160), and in Gonin's operation for detachment of the retina.

The Metri-Cautery is described on p. 150, and is useful in all forms of corneal ulceration, but especially in the infected varieties.

Mydriatics and Cycloplegics.

Mydriatics are remedies which produce dilatation of the pupil. Cycloplegics are agents which cause paralysis of the ciliary muscle (paralysis of accommodation). Mydriatics also produce more or less paralysis of the ciliary muscle.

The drugs which are commonly employed to induce mydriasis and cycloplegia are atropine and homatropine; very much less frequently duboisine, daturine, hyoscyamine, and scopolamine are used.

Cocaine and euphthalmine produce moderate dilatation of the pupil with only slight paresis of the ciliary muscle.

Indications.—Agents of this class are used (1) in iritis, for dilating the pupil, preventing adhesions, and exerting a sedative action; (2) in various diseases of the cornea and of the deeper structures of the eye; (3) in central ulcer of the cornea;




FIG. 345.—METHOD OF INSTILLING DROPS BY MEANS OF AN EYE-DROPPER.

(4) after certain operations; (5) to paralyze accommodation in order to investigate the refraction; (6) to dilate the pupil for ophthalmoscopic examination; and (7) to enlarge the pupil in lamellar and nuclear cataract.

Atropine, the alkaloid of belladonna, is the most commonly employed mydriatic; it is prescribed in the form of ointment of the alkaloid or solution of the sulphate. The strength of solution or ointment varies from $\frac{1}{2}$ to 2 per cent.; 1 per cent. is used most frequently. Atropine ointment made with soft paraffin is less irritating than the B.P. preparation, which must not be ordered in ophthalmic practice. In a case of iritis the addition of 1 per cent. cocaine helps the action of the atropine, but should not be continued for long owing to the action of cocaine on the corneal epithelium.

Atropine paralyzes the sphincter of the pupil and the ciliary muscle. Half an hour after the instillation of a drop of atropine mydriasis and nearly complete cycloplegia will be found; the effects last for a week or ten days. Atropine and other mydriatics have no effect on the tension of the normal eye, but may precipitate an attack of glaucoma in an eye which is predisposed to this disease. One should, therefore, carefully test the tension, and note the depth of the anterior chamber, in persons past middle life, before instilling atropine or any other mydriatic.

Atropine Poisoning.—In susceptible individuals atropine may cause general toxic symptoms: Dryness of the throat, flushing of the face, headache, vomiting, quick pulse, cutaneous eruption, excitability, and even delirium. These symptoms subside within a few hours of discontinuing the drug. In an extreme case it might be necessary to use the antidote morphine. In persons who show such an idiosyncrasy, or in others in whom we wish to push the remedy, it is well to instruct the patient to press the finger against the lacrymal sac during each instillation. When the susceptibility is very great one of the other mydriatics may be resorted to, or we may use a 10 per cent. solution of aqueous extract of belladonna in water; ophthalmic discs, which contain very small doses, may prove useful in these cases.

Atropine Irritation.—In some persons atropine may cause considerable local irritation, showing itself in œdema of the lids, eczematous condition about the lids, and conjunctival catarrh.

In using atropine or other solutions (myotics and anæsthetics) for the local effect upon the cornea or deeper portions of the eye, the drop is allowed to fall upon the cornea, the upper lid being raised, and the patient directed to throw the head back and to look down.

Duboisine sulphate (gr. $\frac{1}{2}$ to $\mathfrak{z}$ ij.), daturine sulphate (gr. $\frac{1}{4}$ to $\mathfrak{z}$ ij.), hyoscyamine hydrobromate (gr. $\frac{1}{2}$ to $\mathfrak{z}$ ij.), and scopolamine hydrobromate (gr. $\frac{1}{8}$ to $\mathfrak{z}$ ij.), are occasionally used as substitutes for atropine. They have similar, though less certain, actions, are contra-indicated in increased tension, and may also produce systemic poisoning.

Homatropine Hydrobromide resembles atropine in its action, but is milder. It is very largely used to paralyze accommodation during the examination for errors of refraction. Though this effect is not so perfect as with atropine, it is sufficient for most purposes, except in children, and lasts only from one to three days, thus exposing the patient to very much less inconvenience. For refraction cases it is usually used in 2 per cent. solution, one drop being instilled every five or ten minutes for three or four doses; half an hour after the final dose the eye is ready for examination. It is frequently combined with cocaine for this purpose:

R	Cocain. hydrochlor.	-	-	-	gr. i.
	Homatropin. hydrobrom.	-	-	-	gr. ij.
	Aquæ destill.	-	-	-	$\mathfrak{z}$ ij.

M.

Its effects may be neutralized within an hour by two drops of 0.5 per cent. eserine, which may cause some slight pain in the eyeball and annoying winking.

A solution of the alkaloids (not the salts) in liquid paraffin is even more effective, and the oil prevents drying of the cornea, which is a danger to be guarded against when using cocaine drops.

Euphthalmine is very useful for the purpose of dilating the

pupil for ophthalmoscopic examination. The hydrochlorate is used in 5 or 10 per cent. solution; one or two drops cause mydriasis in thirty minutes, and the effects pass off in a few hours. It has but a feeble action upon accommodation.

Cocaine Hydrochloride is frequently used to produce moderate dilatation of the pupil for ophthalmoscopic examination. One or two drops of a 4 per cent. solution cause sufficient dilatation in twenty minutes, produce insignificant interference with accommodation, and the effects disappear within an hour or two as a rule. Cocaine acts by constricting the bloodvessels of the iris. It diminishes intra-ocular tension (in rare cases the opposite effect has been observed). It is sometimes combined with other mydriatics, and then increases the action of the associated remedy.

Ephedrine, an alkaloid extracted from the Chinese plant ephedra, used in 5 per cent. aqueous solution of the muriate or sulphate, is a recent addition to mydriatics; in physiological action it resembles adrenalin. A drop of 5 per cent. solution dilates the pupil in thirty minutes without affecting the accommodation; it constricts the conjunctival bloodvessels; it does not increase intra-ocular tension. Mydriasis lasts half an hour. Its solutions can be boiled without injury and do not spoil with keeping.

Miotics.

Miotics diminish the size of the pupil. They produce tonic contraction of the sphincter and of the ciliary muscle, and diminish intra-ocular tension. These agents are employed chiefly in glaucoma, sometimes in ulcers of the cornea, especially when peripheral. Eserine salicylate ($\frac{1}{8}$ to $\frac{1}{2}$ per cent.) and pilocarpine nitrate, or muriate ($\frac{1}{2}$ to 1 per cent.), are described for these purposes. Eserine is stronger, and sometimes has a tendency to produce conjunctival irritation and iritis, and occasionally constitutional symptoms. Pilocarpine is milder and free from these drawbacks; it is sometimes given hypodermically to cause diaphoresis in certain diseases of the eye.

Local Anæsthetics.

Cocaine Hydrochloride is the most commonly employed remedy for producing local anæsthesia of the conjunctiva, cornea, and to a certain extent the iris during operations upon the eye. The strength of solution is usually 4 per cent. It is also of service as a *temporary* anodyne in corneal and iritic affections, and very useful as a mydriatic for ophthalmoscopic examinations. It assists the action both of mydriatics and myotics. It is frequently combined with atropine, homatropine, or with eserine. Cocaine has a tendency to cause desiccation of the cornea, and sometimes superficial ulceration; hence after the instillation of this remedy the patient should be directed to keep the lids closed; for the same reason it should not be used for any length of time, and it is generally unwise to order drops containing cocaine for home use. One drop of a 4 per cent. solution, and a second after a few minutes, is sufficient to anæsthetize the cornea for the removal of foreign bodies; for more penetrating effects the instillations are repeated three or four times at intervals of two or three minutes. Solutions of cocaine do not keep well, and should be freshly prepared previous to use in operations. Cocaine hydrochloride may be used in the form of powder.

Holocaine Hydrochloride is a local anæsthetic which is preferred to cocaine by some surgeons. It is usually employed in 1 per cent. solution. The advantages of this remedy as compared with cocaine are: it acts more quickly, is more penetrating, does not dilate the pupil, has no harmful effect on the corneal epithelium, and its solution keeps well; there is, however, more preliminary conjunctival irritation and it cannot be used hypodermically, since it causes toxic symptoms when employed in this way.

Novocain 2 per cent., in adrenalin chloride solution (1 in 10,000), is an excellent local anæsthetic in lacrymal sac and eyelid operations, from 20 to 60 minims being injected hypodermically. To produce anæsthesia in cases of congestion of the eyeball or sometimes in operations for removal of the globe, 2 c.c. of a 3 per cent. solution in 1 : 10,000 adrenalin

are often injected deep into the orbit through the skin of the lower lid just below the external canthus. A needle $1\frac{1}{8}$ inches long is required, and care must be taken that solutions are sterile and not injected too far back in the orbit, but into the cone of muscles just behind the eyeball to anæsthetize the ciliary ganglion.

A similar injection is often used to paralyze the orbicularis to prevent 'squeezing' in some cataract extractions. The solution is injected into the outer two-thirds of the lower lid and down the outer margin of the orbit from the brow to lower lid. The facial nerve may also be blocked by a deeper injection just below the lobe of the ear.

Other **Cocaine** substitutes. Especially since the passing of the Dangerous Drugs Act, a number of synthetically prepared drugs have been advocated as substitutes for cocaine. Cocaine, however, remains the choice for surface anæsthesia of the eyeball, and novocain is used almost exclusively for infiltration and deep injection. Apart from holocaine, the best-known substitutes for cocaine are butyn and pantocaine.

Butyn is used in 2 per cent. aqueous solution. It does not dilate the pupil nor influence tension or injure the corneal epithelium. It is more irritating to the eye and must not be used for injection, as some fatal results from such use have been reported.

Pantocaine has similar actions. It is used in 0.5 per cent. solution. Its toxicity when injected has not been definitely determined.

Other Therapeutic Measures.

Adrenalin is the active principle of the suprarenal gland. It is available in 1 : 1,000 water solution of the chloride as a colourless liquid, which can be diluted with salt solution of physiological strength. This remedy is a valuable astringent and hæmostatic. After instillation of solutions varying from 1 : 10,000 to 1 : 1,000, marked blanching of the conjunctiva occurs as a result of contraction of the bloodvessels, beginning in less than a minute, and lasting half an hour or longer. When the ocular structures are very much congested, cocaine

or holocaine produces unsatisfactory anæsthesia; if the instillation of these agents is preceded by that of adrenalin or suprarenalin solution, the anæsthetic effect is very much more pronounced. This remedy is used in some cases of conjunctivitis with marked congestion, in affections of the lacrymal passages to facilitate the removal of retained contents and the introduction of probes, glaucoma, and in operations to prevent bleeding and to improve the action of local anæsthetics. It must not be injected subcutaneously when general anæsthesia is employed, owing to its action on the circulatory system.

Glaucosan, a synthetic preparation resembling adrenalin, is used in two varieties: Lævorotatory glaucosan dilates the pupil and yet reduces tension. It has been found useful in cases of secondary glaucoma associated with iritis and posterior synechiæ. Dextrorotatory glaucosan, a powerful miotic, has been used for cases of acute glaucoma, but it has not been found as satisfactory as was at first expected.

Dionin is a derivative of morphine. It is a local vasodilator and lymphagogue. It is used to promote absorption of exudates in iritis and corneal inflammations. It increases the mydriatic effect of atropine. Dionin indirectly relieves deep-seated pain in the same way as hot fomentations, by dilating the vessels. When a 5 or 10 per cent. aqueous solution is instilled into the conjunctival sac, it causes very great chemosis in most cases, the swelling sometimes being so great that the eyelids cannot be opened. The swelling quickly subsides. After a few days the drug loses its effect, so that its characteristic reaction can no longer be produced. It is used in increasing strengths from 2 to 10 per cent. as tolerance is established. Dionin may be used also in the form of ointment, or the powder itself may be used.

Solid Carbon Dioxide.—A stick of solid carbon dioxide, nowadays readily obtainable by compressing the 'snow' which is formed by the escape of the liquid from its container on to a towel or piece of blotting-paper, may be used for destroying follicles in the upper tarsus in trachoma; for small capillary nævi and moles on the eyelids; or for small rodent ulcers when radium or X-rays are not available.

Several applications of some 30 seconds duration, repeated after a few days, will be required in the treatment of rodent ulcer.

Ionization.—When a weak electric current is passed through a solution of a metallic salt, the salt is slowly decomposed, the acid constituent seeking the positive electrode and the basic constituent the negative. By taking advantage of this fact, the ions of zinc or copper may be conveyed through the unbroken skin or conjunctiva. The electrodes are moistened with a solution of the salt, the positive being applied to the affected part and the negative elsewhere. A current of 1 or 2 milliampères for the cornea or conjunctiva and 3 milliampères for the skin is applied for one minute. The treatment has been strongly recommended in corneal ulcer, interstitial keratitis, pterygium, trachoma, rodent ulcer, etc. Results have not been very encouraging, and the treatment is painful.

Electricity is used in the form of the electro-cautery and diathermy, the chief use of the latter being in treatment of detached retina. It is also used for electric eye-warmers. Electrolysis may be employed for the removal of inturned lashes. The galvanic and Faradic currents are occasionally resorted to in paralysis of ocular muscles.

Radium is used in the treatment of rodent ulcer. It may be tried for intractable cases of spring catarrh. It has been used for selected cases of intra-ocular neoplasm, where removal of the eye was refused; in the form of radon seeds buried in the growth; in some cases shrinkage and temporary arrest of the growth have occurred. It has also been successfully employed for destroying hæmangiomas of the lids, palpebral conjunctiva and caruncle. It is used in the form of plaques, 'seeds,' and 'needles,' and the technique varies very much according to the condition for which it is employed. Even the various types and stages of rodent ulcer require different dosage and length of application.

It must be employed very cautiously by an expert, as must

X-rays, which are used in treatment of orbital neoplasms, in addition to their use in the diagnosis of intra-ocular foreign bodies and other conditions.

Ultra-violet Light Therapy ('artificial sunlight') has been used with considerable success in cases of phlyctenular ophthalmia and tuberculosis iridocyclitis. General therapy is employed for such cases and has to be continued over a prolonged period. Local therapy, in which the light rays are focussed on the cornea through a quartz optical system, has been employed in the treatment of various affections of the cornea and eyeball, but it requires a very special technique and is not generally available; it is still on trial.

Heat.—Hot compresses are prescribed in affections of the cornea, iris, and ciliary body. They are applied by means of flannel or absorbent cotton wrung out of water which is as hot as can be borne (115°); they are placed upon the closed lids, and renewed every minute or two. Heat may be usefully applied by the patient himself to his own eye by twisting a ball of cotton-wool round the end of a piece of wood (handle of wooden spoon, say), so that the wood makes a handle for a cotton-wool sponge. The wool is then dipped into very hot water, and applied carefully by the patient to his closed lids. Heat is also applied in the form of electric eye-warmers, which give more continuous heat than fomentations.

Cold.—Cold compresses are used in inflammatory affections of the conjunctiva. The best method of applying them is as follows: Strips of lint are folded so as to make pads of four thicknesses, about $1\frac{1}{2}$ inches square; a number of these are moistened and laid upon a block of ice; from the ice they are transferred to the closed eyelids, and changed as soon as they become warm. In the absence of ice the compresses may be wrung out of cold water. Ice should never be applied directly to the lids.

Local Blood-letting is of great benefit in affections of the deeper structures of the eye, especially in iritis, sometimes in glaucoma. Leeches are best for this purpose; from two to four are applied to the temple, midway between the outer canthus and the tragus.

Massage is employed for clearing recent opacities of the cornea following ulceration or interstitial keratitis, for reducing tension in glaucoma, and in the treatment of blepharitis. When

used to promote the clearing of corneal opacities, some medicated ointment, usually 2 per cent. yellow oxide of mercury, is placed in the conjunctival cul-de-sac and the upper lid moved gently over the cornea with the finger twice daily for a few minutes. In glaucoma massage is used both in unoperated cases and especially after operations in which filtration is not quite satisfactory; intermittent pressure is made by two fingers, as in testing the tension of the eyeball, some twenty times thrice daily. In blepharitis the edges of the lids are cleaned and then massaged with a small ball of cotton-wool on a glass rod after applying yellow oxide of mercury.

Sub-conjunctival Injections are employed in episcleritis, scleritis, iridocyclitis, choroiditis, corneal ulcer, and in detachment of the retina. After local anæsthesia by means of holocaine or cocaine, a fold of conjunctiva is seized with forceps about 8 mm. from the limbus, the needle of a hypodermic syringe introduced into this lifted tissue, and from 5 to 15 minims of fluid injected, or the needle may be introduced superficially under the conjunctiva without the aid of a forceps when the patient looks downward and the upper lid is raised. Various germicides (mercury bichloride 1:5,000 to 1:1,000, mercury cyanide 1:5,000 to 1:1,000, cinnamic acid 1:100) have been recommended, but a solution of sodium chloride of physiological strength is equally effective and much less painful.

Fluorescein, an orange-red powder, is used in a 2 per cent. aqueous solution (with 3 per cent. sodium bicarbonate) to detect abrasions, infiltrations, and ulcers of the cornea, and to define the limits of such lesions. A drop is instilled into the conjunctival sac, and after a minute the excess is washed off with a few drops of boric acid lotion; a green stain indicates loss, or disease, of corneal epithelium.

Salvarsan and its derivatives are used in the general treatment of syphilitic eye conditions. These arsenical preparations are of the greatest benefit in early cases, and must then be used with mercury or bismuth *at the same time*, as then only can late nervous sequelæ be avoided. In late cases care is necessary in using the various salvarsans, and it is generally

advisable to start with a course of mercury and iodides. It is not used in optic nerve atrophy, since no benefit has followed; there is no general agreement on the question whether salvarsan has a harmful effect when used in this affection, but some cases which have been reported indicate that this is possible. Subdural, intraspinal, or intra-cisternal injection of arsphenamized serum has been found of some value in the treatment of early optic nerve atrophy. Results in cases of syphilitic iritis are brilliant; in other affections (choroiditis, retinitis, paralysis of external muscles) the effects vary considerably. In interstitial keratitis it seems to be of greatest value when combined with repeated paracenteses of the anterior chamber. The arsenical preparations of the salvarsan type are often used successfully in the treatment of sympathetic ophthalmitis, especially if the case is treated early.

Bismuth may be given in the form of finely divided metal in suspension or as one of the bismuth salts.

Vaccines are valuable agents in suitable cases of ocular disease, and, when possible, an autogenous vaccine, made from the patient's own organisms, should be given. Where this is impracticable, stock polyvalent vaccines, made from virulent strains, may be used with advantage. Those most generally employed are gonococcal vaccines, staphylococcal vaccines, and tuberculin. Gonococcal vaccines give excellent results in gonorrhœal iritis, and are of some value in adult gonorrhœal ophthalmia, but have failed to show any beneficial effect in gonorrhœal ophthalmia neonatorum. Staphylococcal vaccines are of service in obstinate cases of relapsing hordeola and blepharitis, and in some cases of uveitis and corneal ulceration. In general, staphylococcal infections of the eye respond well to vaccine therapy.

Tuberculin is now, after a period of comparative disfavour, extensively used in tuberculous eye infections. It is used both as an aid to diagnosis and for treatment. Many preparations are employed, but for diagnosis Koch's Old Tuberculin T. is chiefly used. Of the methods in use for diagnostic purposes, the only one that is of any real value is the subcutaneous injection of Old Tuberculin. The patient should be in bed,

be free from all signs of pulmonary and surgical tuberculosis, and have a persistent normal temperature over several days. An injection of $\frac{1}{100}$ c.c. Old Tuberculin is given, and a positive reaction manifests itself by a rise in temperature, a local reaction at the site of inoculation, a focal reaction in the diseased area with exacerbation of symptoms, and a general feeling of malaise with nausea, headache, and loss of appetite, etc. If there is no reaction in two to three days a larger dose can be given, and even a third, up to 1 c.c. If this fails to produce a reaction, the patient is usually considered to have no active tuberculous condition present. It is a valuable test, but has fallen into disuse, as there is a certain small risk of damage to an already injured eye.

Von Pirquet's test is of little value except in children, but when a graduated test, using human and bovine Old Tuberculin, is performed on an adult, it is an excellent test of the sensitivity of the patient to tuberculin, and a valuable guide in the selection of the type of tuberculin to be used in treatment and the initial dose to be given. The modifications of von Pirquet's test, such as Moro's, Woodcock's and Mantaux's intradermal test, are of even less value. Calmette's ophthalmo-reaction has now been discarded as liable to lead to aggravation of the disease.

In treatment various preparations are employed, of which there are a large number. Old Tuberculin is favoured by a few, but the usual ones are TR. (Residual Tuberculin), BE. (Bacillary Emulsion, human), PBE. (Bacillary Emulsion, bovine), and the small dose method is most in favour. Combination of these tuberculins may sometimes be used with better results.

Treatment by tuberculin requires considerable skill and experience, and if given in too large doses or at too frequent intervals, harm may result. All tuberculins react very much in the same way—the important factor is to use a tuberculin with which one is absolutely familiar, and in small doses, so that very little, if any, general reaction is obtained.

Tuberculin is used in cases of tuberculous iridocyclitis, conjunctivitis, choroiditis, episcleritis and scleritis, and in

some specific cases of interstitial keratitis, Eales' disease, and other forms of intra-ocular hæmorrhage, and other rarer forms of ocular tuberculosis have been successfully treated.

Injection of Foreign Protein has been found of service in some conditions, especially in cases of severe iritis, also in some cases of severe scleritis, corneal ulceration, and sympathetic ophthalmia. Boiled milk in doses of 5 to 10 c.c. or gonococcal vaccine (1 c.c.) injected into the gluteal muscles are the most suitable. In some cases 2,000 units of diphtheria antitoxin injected subcutaneously, or typhoid vaccine injected intravenously, have been given.

General Rules for Ophthalmic Operations.

Preparation of the Patient.—When not operated upon in his own home, the patient should enter the hospital or nursing-home on the day previous to operation. He should take a bath and a mild cathartic on the night before, followed, perhaps, by an enema on the morning of the operation. He should be in good physical condition; old age, albuminuria, and diabetes are no contra-indications, but such patients require especial care.

It is imperative to examine the conjunctiva and the lacrymal sac before deciding to operate, especially if the eyeball is to be opened, as in iridectomy or cataract extraction. The presence of purulent or muco-purulent secretion from the conjunctiva or lacrymal sac renders an operation upon the eyeball extremely hazardous on account of the danger of infection. In such cases the conjunctival or lacrymal affection must be first cured by appropriate treatment. If there be the slightest doubt as to the condition of the conjunctival sac, a bacteriological examination should be made.

Preparation of the Hands of the Operator.—The hands should be scrubbed thoroughly with soap and warm water, and then immersed for a minute in 1 : 1,000 corrosive sublimate solution. Gloves are not usually worn for ophthalmic operations.

Preparation of Instruments.—Blunt instruments should be cleaned and polished, boiled in 1 per cent. solution of soda, rinsed with sterile water, and then dried and laid on sterilized gauze. Sharp instruments, especially cataract knives, should not be boiled for more than about one minute. If perfectly clean, they can be sterilized by immersing them in 90 per cent. alcohol for at least fifteen minutes. Previous to sterilizing, the cutting qualities are tested upon thin kid stretched in the testing-drum (Fig. 346), and the edge and point examined with a magnifying lens.

Position of the Patient.—The patient should lie on a narrow operating-table. The light, whether daylight or artificial illumination, must be good, and the field of operation must be



FIG. 346.—DRUM USED TO TEST THE CUTTING-EDGES OF EYE INSTRUMENTS.

well illuminated; for the latter purpose a large condensing lens, or better still an electric projection lamp, is used to throw the light upon the eye during operations upon the lens or iris.

Preparation of the Region of Operation.—The eyelids, including the lashes and the surrounding skin, should be washed with soap and warm water, and then with corrosive sublimate solution (1 : 5,000). The everted lids are cleansed gently. The conjunctival sac should be flushed with a large quantity of unirritating sterile fluid.

Anæsthesia.—In the great majority of adult cases local anæsthesia is sufficient. Two drops of a 2 or 4 per cent. solution of hydrochloride of cocaine are instilled every few minutes for three or four doses, the lids being kept closed in the intervals. Cocaine solutions must be freshly prepared, since they do not keep well.

In children and in nervous adults, also in enucleations, in glaucoma with very high tension, in blepharoplastic operations, and occasionally in other procedures, a general anæsthetic is often necessary. But even in acute congestive glaucoma and with enucleations and eviscerations of the eyeball, a painless operation can be performed after the injection of 2 c.c. of 3 per cent. solution of novocain with 1:10,000 adrenalin, deep into the orbit (p. 467), or into Tenon's capsule.




FIG. 347.—MONOCULAR BANDAGE. FIG. 348.—BINOCULAR BANDAGE.

In operations upon the eyelids, or in excision of the lacrymal sac, subcutaneous injection of novocain and adrenalin (see p. 84) gives good results.

Cleansing Solutions.—In the course of operations upon the eyeball it is necessary to cleanse the seat of operation, and to irrigate the cornea frequently to prevent desiccation. The solutions used for this purpose are boric acid 4 per cent., salt 0.6 per cent., and mercuric chloride 1:10,000. These cleansing solutions are applied by means of small wads of absorbent cotton-wool.

Dressings vary with the nature of the operation. Sterilized gauze, used dry or soaked in sterilized salt solution, are

generally applied next to the closed eyelids, covered by additional layers of absorbent gauze, and retained by a bandage covering one or both eyes, sometimes by strips of isinglass plaster.




FIG. 349.—THE MOORFIELDS HOSPITAL BANDAGE.

Eye bandages are $1\frac{1}{2}$ inches wide, 5 or 7 yards long, and made of gauze or muslin. If used for protection only, they are applied lightly; if for pressure, they are put on tightly; in the latter case, care must be taken that the depression between the supra-orbital margin and the nose is properly filled out.




FIG. 350.—THE MOORFIELDS BANDAGE APPLIED.

The Monocular Bandage (Fig. 347) is applied as follows: Begin over the temple of the same side as the affected eye (the right, for example); make one turn around the forehead, then pass across the occiput, below the right ear, and obliquely across the right eye; then another turn about the forehead,

below the right ear, across the right eye; and alternate in this way three or four times.

The Binocular Bandage (Fig. 348).—Begin over the temple of one side (the right, for example); make a full turn around the forehead, and continue to the left temple; then obliquely across the occiput, below the right ear, across the right eye, around the upper occipital region, above the right ear, downward over the left eye, below the left ear, across the occiput, below the right ear, across the right eye; and alternate in this manner for three or four turns.

The Moorfields Bandage (Figs. 349 and 350) is very useful for most eye operations. It is made of stout linen and tape. The figures show the exact dimensions and the manner of applying it. It enables one to change the dressing without lifting the patient's head from the pillow. After untying the knot, the bandage is lifted a little from the patient's face and passed upwards on to his forehead. The dressing having been changed, the bandage is replaced and the tapes tightened and tied.

CHAPTER XXXII

VISUAL REQUIREMENTS FOR BRITISH AND INDIAN PUBLIC SERVICES

Royal Navy.—Candidates for naval cadetship must possess full normal vision as determined by Snellen's tests, each eye being separately examined (*i.e.*, $\frac{5}{8}$ and Snellen's 0.6 or J 1). There must be no ocular disease and no squint. Colour vision must be normal.

(A high hypermetrope or a boy whose eyes are progressing towards myopia may pass the tests, but would be liable to be rejected as unfit in after years.)

Candidates for Other Branches of the Royal Navy.—Full normal vision is not required, but any defect of vision must be due to errors of refraction, which can be corrected to normal by glasses, and vision without glasses must in any case be not less than $\frac{5}{10}$ with each eye, and he must also be able to read Snellen's 0.6.

Imperfect colour-vision will disqualify a candidate, so also will any chronic disease of eyes or eyelids, lacrymal apparatus, squint, or any defect of the ocular muscles.

The following ratings are required to have full normal acuteness of vision without glasses: The seamen class, marines (except bandsmen), armourers, engine-room artificers, electricians, and boy artificers.

For candidates for other artisan ratings and for stokers the vision must be at least $\frac{5}{8}$. For all other ratings, including writers, ship-stewards' assistants, ship's cooks, sick-berth staff, officers' stewards and cooks, the vision must not be less than $\frac{5}{12}$.

Defects of vision must be due to errors of refraction only, and must be capable of correction by means of glasses. The candidate must be able to read Snellen's 0.6 without glasses. Marine bandsmen, sick-berth staff, writers, ship's stewards' ratings, ship's cooks' ratings, and officers' servants are allowed to wear glasses. Ship's cooks' ratings and officers' servants are not disqualified by colour-blindness; all the others are.

The British Army—Commissioned Officers.—The examination for determining the acuteness of vision includes two tests: one for distant, the other for near vision. The army test types will be used for the test for distant vision without glasses at a distance of 20 feet, and for

the test for near vision without glasses at any distance selected by the candidate. The standards of the minimum acuteness of vision with which a candidate will be considered fit are as follows:

STANDARD I.

Right Eye.

Distant vision, $\frac{6}{6}$.
Near vision, Snellen's 0.6.

Left Eye.

Distant vision, $\frac{6}{6}$.
Near vision, Snellen's 0.6.

STANDARD II.

Better Eye.

Distant vision, $\frac{6}{6}$.
Near vision, Snellen's 0.6.

Worse Eye.

Vision without glasses not below $\frac{6}{12}$, and after correction with glasses not below $\frac{6}{24}$.
Near vision, Snellen's 1.

STANDARD III.

Better Eye.

Distant vision without glasses not below $\frac{6}{24}$, and after correction with glasses $\frac{6}{6}$.
Near vision, Snellen's 0.8.

Worse Eye.

Vision without glasses not below $\frac{6}{24}$, and after correction with glasses not below $\frac{6}{12}$.
Near vision, Snellen's 1.

Each eye must have a full field of vision as tested by hand movements.

Squint or any morbid condition of the eyes or of the lids of either eye liable to the risk of aggravation or recurrence will cause the rejection of the candidate.

Each eye will be examined separately, and the lids must be kept wide open during the test.

Inability to distinguish the principal colours will not be regarded as a cause for rejection, but the fact will be recorded, and the candidate will be informed.

No relaxation of the standard of vision will be allowed under any circumstances.

Recruits—Vision.—In examining a recruit's vision he will be placed with his back to the light, and his visual acuteness will be tested by means of army test types placed under good illumination at a distance of 6 metres (20 English feet) from the recruit.

Each eye will be tested separately, and the lids must be kept wide open during the test.

A recruit for the regular army will be considered fit—

(a) If his vision *without* the aid of glasses is not less than $\frac{6}{18}$ with each eye.

(b) If his vision in the left eye *without* the aid of glasses is not less than $\frac{6}{36}$, provided that his vision in the right eye *without* the aid of glasses is not less than $\frac{6}{6}$.

(c) In the case of recruits enlisting into the Royal Artillery, Royal Engineers, Royal Corps of Signals (except operators, visual), Tank Corps, and R.A.S.C., with a view to appointment as tradesmen (except drivers), and in the case of recruits enlisting into the R.A.M.C., Army Dental Corps, R.A.O.C., R.A.V.C., and C.M.A.—

- (i.) If his vision in one eye (right or left) *without* the aid of glasses is not less than $\frac{6}{8}$, provided that his vision in the other eye *without* the aid of glasses is not less than $\frac{6}{12}$;
- (ii.) If his vision *without* the aid of glasses is not less than $\frac{6}{24}$ in each eye.

The vision of a recruit enlisting with a view to employment as a driver (Horse Transport or Mechanical Transport) in any corps *without* the aid of glasses must not be less than $\frac{1}{8}$ with each eye, provided that *with* the aid of glasses, if necessary, his vision is not less than $\frac{6}{8}$ in one eye (right or left) and $\frac{6}{12}$ in the other.

A candidate for the Army Educational Corps will be accepted if the medical examiner is satisfied that his vision, *with* or *without* glasses, is sufficiently good for the satisfactory performance of his duties.

Each eye must have a full field of vision as tested by hand movements. Squint or any morbid condition of the eyes or of the lids of either eye liable to the risk of aggravation or recurrence will cause the rejection of the recruit.

The visual acuity of each eye will be entered on the medical history sheet of approved recruits.

Board of Trade Tests for the British Mercantile Marine.—‘If he can read correctly nine out of the twelve letters in the sixth line ($\frac{5}{8}$) from the top, and eight of the fifteen letters in the seventh line ($\frac{5}{6}$) with one eye, and the whole of the eight letters in the fifth line ($\frac{5}{10}$) with the other eye, he may be considered to have passed the test. If he cannot do so, his case should be submitted to the Principal Examiner of Master and Mates.

‘A candidate who fails to pass the letter test may present himself for re-examination at intervals of three months.

‘Every candidate must undergo the lantern test on every occasion on which he presents himself for examination for his first certificate of competency; but if he then passes he will not be required by the Board of Trade to undergo the lantern test on any subsequent occasion.’

A candidate who fails to pass the lantern test may not be re-examined locally.

Home Civil Service.—There are no definite rules, but the candidate must have no defect of vision which is likely to interfere with his work.

Indian Civil Service (*i.e.*, Ecclesiastical, Education, Geological, Survey, Agriculture, Indian Finance, Customs, Civil Veterinary, and other departments not specially provided for).—1. The candidate will

be admitted if ametropic in one or both eyes, provided that with glasses he sees not less than $\frac{5}{8}$ with one eye and $\frac{5}{8}$ with the other, there being no morbid change in the fundus of either.

2. If myopic, the ametropia in either eye is not to exceed 2.5 D., with no active morbid changes in choroid or retina, though he may have a posterior staphyloma.

3. If the defect of vision is due to a corneal nebula, he will be disqualified if his vision is less than $\frac{1}{2}$ in either eye, and in such case the acuteness of vision in the better eye must equal $\frac{5}{8}$ with or without glasses.

4. Squint, or any morbid condition subject to risk of aggravation or recurrence in either eye, may cause the rejection of a candidate. The existence of imperfection of colour-sense will be noted.

The Departments of Forest, Survey, Telegraph, Factories, and for Various Artificers in the Indian Civil Service.—1. If myopia in one or both eyes exists, the candidate may be passed if it does not exceed 2.5 D., and if, with correcting glasses not exceeding 2.5 D., he sees $\frac{5}{8}$ with one eye and $\frac{5}{8}$ with the other, there being normal range of accommodation with the glasses.

2. If myopic astigmatism exists, the combined spherical and cylindrical glass must not exceed -2.5 D., and with that one eye must see not less than $\frac{5}{8}$ and the other $\frac{5}{8}$.

3. A candidate having total hypermetropia not exceeding 4 D. is not disqualified provided the sight in one eye (when under atropine) equals $\frac{5}{8}$ and the other $\frac{5}{8}$, with +4 D. or any lower glass.

4. Hypermetropic astigmatism will be allowed if the combined lenses required to correct the error do not exceed 4 D., and that with or without the glasses the sight is equal to $\frac{5}{8}$ in one eye and $\frac{5}{8}$ in the other.

5. If the defect is due to a corneal nebula, the sight of one eye must not be less than $\frac{1}{2}$. In such a case the better eye must be emmetropic. Defects of vision from pathological or other changes in the deeper structures of either eye which are not referred to in the above rules may exclude a candidate.

6. Squint, or any morbid condition subject to risk of aggravation or recurrence in either eye, may cause the rejection of a candidate. The existence of imperfection of colour-sense will be noted.

Artificers engaged in map and plan drawing may be considered separately, and this standard relaxed if it appears to be desirable.

Public Works Department and Superior Establishments, Railway Department of India.—1. If myopia exists, it must not exceed 3.5 D., but if with a 3.5 D. glass the candidate sees $\frac{5}{8}$ with one eye and not less than $\frac{5}{8}$ with the other, he will be passed. Range of accommodation must be normal.

2. If myopic astigmatism exists, the combined spherical and cylin-

drical glass must not exceed 3.5 D., and with this the vision must be equal to at least $\frac{5}{8}$ in one eye and $\frac{5}{8}$ in the other. Range of accommodation must be normal.

3. Hypermetropia must not exceed 4 D., and the sight of one eye (when under atropine) must equal $\frac{5}{8}$, and the other $\frac{5}{8}$ with a + 4 D. glass, or one of any less power.

4. Hypermetropic astigmatism is allowed if the combined correcting glass does not exceed 4 D., and that the sight, with or without lens, equals $\frac{5}{8}$ in one eye and $\frac{5}{8}$ in the other.

5. If a corneal nebula exists, the vision must not be less than $\frac{9}{12}$ in that eye, but the other eye must be emmetropic. Defects of vision arising from pathological or other changes in the deeper structures of the eye, which are not referred to in these rules, may exclude a candidate.

6. Squint, or any morbid condition subject to risk of aggravation or recurrence in either eye, may cause the rejection of a candidate. Any imperfection of the colour-sense is a disqualification for appointment to the engineering branch of the Railway Department, or as assistant-superintendent in the Traffic Department. In all other cases a note as to any imperfection of the colour-sense will be made.

The Indian Medical Service and Indian Police.—The same as the British Army.

Indian Pilot Service, and Candidates for Appointments as Guards, Engine-Drivers, Signalmen, and Pointsmen on Railways.—1. A candidate is disqualified unless both eyes are emmetropic, his acuteness of vision and range of accommodation being perfect.

2. A candidate is disqualified by any imperfection of his colour-sense.

3. Strabismus or any defective action of the exterior muscles of the eyeball disqualifies a candidate.

Indian Marine Service, including Engineers and Firemen.—1. A candidate is disqualified if he has an error of refraction in one or both eyes, which is not neutralized by a concave or convex 1 D. lens or some lower power.

2. He is disqualified by any imperfection of his colour-sense.

3. He is disqualified by strabismus or any defective action of the exterior muscles of the eyeball.

COMMISSIONS IN THE ROYAL AIR FORCE.

The regulations governing the examination of eyesight are as follows:

1. The examination for determining the acuity of vision will be conducted with well-illuminated standard test types at a distance of 20 feet. The candidate must be able to read the test types without hesitation. The minimal standard of acuteness of vision without correcting glasses

is: Distant vision in each eye tested separately - V.: $\frac{5}{8}$. Hypermetropia as tested by a + lens of 2 D. or more in either eye shall disqualify. Where a candidate is otherwise specially fit, the eye specialist may recommend acceptance—

- (a) For permanent commissions when the visual acuity is equal to $\frac{5}{8}$ in each eye;
- (b) For short service commissions when the visual acuity is equal to $\frac{5}{8}$ in each eye;

provided that in each case such vision is correctable by glasses to $\frac{5}{8}$ in each eye and the conditions in paragraphs 2, 3, and 4 are fulfilled.

2. Both eyes must have good fields of vision as tested by hand movements.

3. Colour vision must be normal.

4. There must be good (i.) binocular fusion, (ii.) convergence, (iii.) balance of eye muscles.

INDEX

ABBREVIATIONS, 336

Abscess, brain, 451

- cornea, 147
- eyelids, 55, 57
- lacrymal sac, 85
- orbital, 90

Abscission, 159

Absolute alcohol, 149, 461

Accessory nasal sinuses, 94

Accommodation, 340

- amplitude of, 341
- and convergence, 343
- anomalies of, 388
- mechanism of, 341
- paralysis of, 390
- range of, 341
- spasm of, 391

Accommodative squint, 412

Acromegaly, 443

Acuteness of vision, 11, 340

Adam's operation (ectropion), 68

Adrenalin, 84, 225, 434, 468

Advancement operation, 432

After-cataract, 261

Albuminuric retinitis, 278, 447

Alcohol, absolute, 149, 461

Alkaline wash, 457

Alternating convergent squint, 410

Alum, 458

Amaurosis, 306

quinine, 314

' Amaurotic cat's eye,' 209

family idiocy, 283

Amblyopia, 306

- after hæmorrhage, 441
- congenital, 306
- ex anopsia, 306
- hysterical, 311
- malarial, 314
- of squint, 411
- quinine, 314
- reflex, 314
- simulated, 312
- tobacco, 301
- toxic, 301
- uræmic, 280, 314

Amblyoscope, 417

Ametropia, 339

Ammoniated mercury, 460

Amplitude of accommodation, 341

- of convergence, 344

Anæmia, 441

- of the retina, 284
- pernicious, 441

Anæsthesia, 476

local, 467

Anatomy of choroid, 193

- ciliary body, 187
- conjunctiva, 101
- cornea, 129
- eyelids, 52
- glaucoma, 211
- innervation of pupil, 183
- iris, 172
- lacrymal apparatus, 77
- lens, 244
- normal fundus, 38, 273
- optic nerve, 295
- orbit, 87
- retina, 272
- sclera, 166
- visual paths, 315
- vitreous, 236

Aneurism, 92

of aorta, 442

Angioma, 93

Angioneurotic œdema, 55

Angle, alpha, 339

gamma, 339

iris, 220

metre, 344

minimum visual, 12

of convergence, 344

of the anterior chamber, 211

Angular conjunctivitis, 108

Aniridia, 173

Anisometropia, 380

Annular posterior synechia, 177

Anomalies of accommodation, 387

- of the eyeball, 96

Anophthalmos, 96

Anopsia, 306

Anterior chamber, 7

- Antiseptic solutions, 455
 Antrum disease, 95
 Aorta, aneurism of, 442
 Aortic insufficiency, 442
 Aphakia, 252
 Apoplexy, 449
 Aqueous chamber, 211
 humour, 212
 Arcus senilis, 140
 Argyll-Robertson pupil, 185, 451
 Argyrol, 463
 Argyrosis, 459
 Army, visual requirements for, 480
 Arsenic poisoning, 303
 Arteria centralis retinae, 39, 272
 'embolism of,' 286
 Arterial pulsation (disc), 216
 Arterio-sclerosis, 285, 442
 Artery, hyaloid, persistent, 236
 Artificial eyes, 99
 ripening of cataract, 251
 vitreous, 99
 Asteroid hyalitis, 237
 Asthenopia, 381
 accommodative, 382
 muscular, 382
 nervous, 382
 reflex, 382
 Astigmatic dial, 373
 Astigmatism, 340, 351, 369
 correction of, 373
 ophthalmoscope in, 375
 Placido's disc in, 377
 regular, 370
 refraction in, 370
 varieties of, 371
 retinoscopy in, 376
 tests, 373
 treatment, 379
 varieties, 369
 Astringent remedies, 457
 Atrophy, choroidal, 195
 of optic nerve, 281, 303
 Atropine, 465
 irritation, 465
 long use of, 250
 poisoning, 465
 Autogenous vaccines, 473
 Axis, optical, 338

 Bacteriological examination of
 eyes before operation, 475
 Bandage, binocular, 479
 monocular, 478
 Moorfields, 258, 479
 Barraquer's intracapsular extrac-
 tion, 263

 Basedow's disease, 443
 Bead test, Edridge-Green's, 310
 Beer's knife, 58
 Biconcave lens, 327
 Biconvex lens, 326
 Bifocal lens, invisible, 385
 Binasal hemianopsia, 318
 Binocular bandage, 479
 vision, 395
 Bisulphide of carbon poisoning, 303
 Bitemporal hemianopsia, 318, 451,
 454
 Bjerrum's screen, 20
 Blackboard test, 16
 Black cataract, 249
 'Black eye,' 75
 Blepharitis, 53
 Blepharoplasty, 68
 Dieffenbach's, 69
 Fricke's, 69
 Knapp's, 69
 Blepharospasm, 5, 71
 'Blight,' 105
 Blind spot, 19, 273
 Blindness, colour, 307
 day, 314
 embolic, 284
 night, 288, 314
 snow, 105, 283
 uræmic, 280
 word, 307
 Blood, diseases of the, 441
 letting, local, 471
 Bluestone, 459
 Board of Trade tests for Mercan-
 tile Marine, 482
 Borax, 458
 Boric acid, 457
 Bowman's membrane, 139
 Brain, abscess of, 451
 tumour, 298, 451
 Bright's disease, retinitis of, 278
 Broad needle, 228
 Buller's shield, 112
 Buphthalmos, 96, 235
 Burns of conjunctiva, 137
 cornea, 165
 eyelids, 76
 'Button-hole' in iris, 260
 Butyn, 468

 Calmette's ophthalmo-reaction,
 474
 Calomel, 466
 Camphor, 458
 Canal, hyaloid, 236
 of Petit, 244

- Canal of Schlemm, 188, 211
 Canaliculi, 77
 anomalies of, 78
 Canaliculus knife, 80
 slitting, 82
 Candle test, 16
 Canthoplasty, 65
 Canthotomy, temporary, 66
 Canthus, 52
 Canula, 258
 Capsule, extraction of cataract in,
 262
 of Tenon, 88
 of the lens, 244
 Capsulo-lenticular cataract, 245
 Capsulotomy, 262
 Carbolic acid, 149, 461
 Carbon dioxide snow, 75, 469
 solid, 469
 Carcinoma of eyelids, 75
 Cardiac disease, 442
 Cardinal points of the eye, 338
 Caruncle, 52
 Case-taking, 1
 Cataract, 245
 accidents, 258
 after-, 261
 after-treatment, 260
 anterior capsular, 144
 polar, 144, 246, 267
 artificial ripening of, 251
 black, 249
 canula 258
 capsular lenticular, 245
 central, 269
 chalky, 249
 complicated, 269
 concussion, 266
 congenital, 267, 268
 curette evacuation, 263
 delivery of the, 257
 dissection of, 267
 dressing after, 258, 260
 etiology, 246
 extraction of, 253
 complications of, 259
 fusiform, 269
 glass-blowers', 46
 intracapsular extraction, 262
 juvenile complete, 264
 lamellar, 268
 monocular, 252
 Morgagnian, 249
 needling of, 265
 nuclear, 246
 operation, modifications in,
 258
 Cataract, pathology, 249
 phakoerisis, 263
 physical signs, 247
 post-operative complications,
 260
 posterior polar, 246, 267
 prognosis, 252
 progressive, 246, 247
 punctate, 269
 pyramidal, 267
 ripe, 249
 secondary, 245, 269
 senile, 247
 shrunk, 249
 stages of, 248
 stationary, 246, 267
 symptoms, 246
 traumatic, 266
 treatment, 249
 varieties, 245
 uncommon, 269
 zonular, 268
 Catarrh, spring, 132
 Catarrhal conjunctivitis in new-
 born, 115
 ulcer (corneal), 145
 Caution, the, 463
 galvano-, 149
 metri, 150, 463
 thermo-, 160
 Cavernous sinus, thrombosis of,
 41
 Cellulitis, orbital, 91, 444
 Centrads (prisms), 325
 Cerebral decompression, 299
 Cerebro-spinal meningitis, 444
 Chalazion, 57
 scoop, 59
 Chalky cataract, 249
 Chemosis, 4, 103, 110
 Children, examination of, 6
 Chlorosis, 441
 Choked disc, 278, 296, 451
 in ear disease, 444
 Chorea, 452
 Chorio-capillaris, 193
 Choroid, anatomy of, 193
 atrophy of, 194
 coloboma of, 173, 199
 diseases of, 193
 rupture of, 200
 sarcoma of, 207
 tubercle of, 200
 Choroiditis, 193, 472
 central, 196
 diffuse, 196
 disseminated, 196

- Choroiditis, exudative, 193
 myopic, 197
 non-purulent, 193
 plastic, 193
 purulent, 198, 282
 senile central, 197
 syphilitic, 197
 Choroido-retinitis, 194
 syphilitic, 197
 Cicatrix, leaking, 229
 Cilia, 52
 forceps, 61
 misplaced, 60
 Ciliary body, anatomy of, 187
 diseases of, 188
 injuries of, 192
 injection, 102
 muscle, 187, 359
 processes, 187
 tenderness, 9, 188
 Circulatory system, diseases of, 442
 Cleansing solutions, 455, 477
 Cocaine as dilator, 29
 hydrochloride, 467
 nitrate, 462
 Cold compresses, 471
 Collyria, 108, 457
 Coloboma of choroid, 173, 199
 of iris, 173, 199, 227
 Colour blindness, 307
 dangerous cases, 309
 perception, theories of, 308
 sense, 11, 21
 vision, tests for, 309
 Colouring, 162
 Colour-perceiving elements in the retina, 307
 Colours, field for, 21
 in spectrum, 308
 Coma, 452
 Comitant squint, 397, 410
 Concave lens, foci of, 329
 meniscus lens, 327
 mirror, 321
 periscope lens, 327
 spherical lens, 326
 Concavo-convex lenses, 326
 Concussion cataract, 266
 Congenital anomalies of eyeball, 96
 cataract, 267, 268
 dislocation of lens, 269
 glaucoma, 96, 235
 word-blindness, 307
 Conical cornea, 159
 operation for, 160
 Conjugate foci, 328
 Conjunctiva, 52
 anatomy of, 101
 burns of, 137
 diseases of, 101
 foreign bodies on, 137
 hæmorrhage under, 137
 injection of, 102
 injuries of, 137
 lithiasis of the, 108
 wounds of, 137
 xerosis of, 122, 314
 Conjunctivitis, 103
 acute catarrhal, 103
 epidemic, 105
 muco-purulent, 103
 simple, 103
 adult purulent, 110
 angular, 108
 catarrhal, in new-born, 115
 chronic catarrhal, 106
 simple, 106
 croupous, 118
 differential diagnosis, 176
 diphtheritic, 116, 444
 exanthematous, 105
 follicular, 106, 108
 gonorrhœal, 110
 granular, 118
 infantile purulent, 113
 lacrymal, 79, 105
 membranous, 116
 Parinaud's, 128
 phlyctenular, 128
 purulent, 110
 pustular, 128
 trachomatous, 118
 traumatic, 104
 Constipation, 442
 Contact glasses, 160
 Contrecoup, injury by, 95
 Conus, 197
 Convergence, amplitude of, 344
 and accommodation, 343
 angle of, 345
 insufficiency of, 429
 range of, 344
 Convergent squint, 414
 etiology, 411
 operation in, 421
 Converging lens, 326
 meniscus lens, 326
 Convex lens, foci of, 327
 meniscus lens, 326
 mirror, 323
 periscopic lens, 326
 spherical lens, 325

- Convexo-concave lens, 327
 Copper sulphate, 124, 459
 Cornea, abscess of, 147
 anatomy of, 139
 burns of, 165
 conical, 159
 diseases of, 139
 examination of, 5
 in infants, 6
 fistula of, 144
 foreign bodies in, 163
 inflammation of, 140
 injuries of, 163
 opacities of, 5, 161
 paracentesis of, 150
 perforation of, 143
 protrusions of, 157
 sensitiveness of, 5
 staphyloma of, 157
 tattooing, 162
 ulcer of, 141
 wounds of, 165
 Corneal facet, 142
 magnifier, 24
 microscope, 43, 44
 precipitates, 50, 177, 190
 ulcer, atropine in, 148
 clinical forms, 145
 fluorescein in, 5, 142
 iodine in, 149
 serpent, 146
 vaccines in, 473
 Corneo-scleral trephining, 229
 Corrosive sublimate, 461
 Cortical visual area, 315
 Cowper's probes, 81
 Credé's prophylaxis, 115
 Cretinism, 443
 Croupous conjunctivitis, 118
 Crystalline lens, 244
 Cupping of disc, 39, 215
 Curette, 257, 263
 Cyclitis, 188
 and oral sepsis, 443
 mild, 189
 purulent, 192, 198
 'quiet,' 443
 severe, 190
 Cyclodialysis, 232
 Cyclophoria, 424, 427
 Cycloplegia, 391
 Cycloplegics, 383, 463
 Cylindrical lens, 331
 Cyst, Meibomian, 57
 tarsal, 57
 Cystoid cicatrix, 228
 Cystotome, sickle, 255
 Dacryoadenitis, 79
 Dacryocystitis, acute, 85
 chronic and corneal infection, 79
 Dangerous Drugs Act, 468
 'Dangerous zone,' 192
 Dark-room examination, 23
 Daturine, 465
 Day-blindness, 314
 Decentration (lenses), 431
 Decompression, cerebral, 299
 Degrees (prisms), 325
 Dendriform keratitis, 147
 ulcer (cornea), 147
 Dental disease, 443
 Dermoid cyst, 93
 Descemet's membrane, 140
 Descemetitis, 157, 189
 Desmarre's lid retractor, 7
 Detachment of retina, 290, 470
 De Wecker's iris scissors, 257
 Diabetes, 447
 Diabetic retinitis, 280
 Dial, astigmatic, 373
 Diathermy, 59, 61, 224, 293, 470
 Dichromics, 309
 Dieffenbach's blepharoplasty, 69
 Digestive system, diseases of, 428
 Dilatation of pupil, 29
 Dilator, Netteship's, 80
 Dionin, 151, 162, 167, 179, 469
 Diopters (prisms), 325
 Dioptic apparatus of the eye, 337
 system, lenses, 332
 Diphtheria, 444
 Diphtheritic conjunctivitis, 116
 Diplo-bacillus of Morax-Axenfeld, 105, 108
 Diplopia, 395, 399
 crossed, 397
 homonymous, 397
 monocular, 270
 test, 397, 400
 Direct method, ophthalmoscopic examination, 32, 349
 Disc, choked, 278, 296
 excavation of, 215
 optic, 31, 38, 272
 Placido's, 5, 377
 Discission for after-cataract, 262
 of lens, 265
 Disinsertion (retina), 291
 Disease:
 abscess of brain, 451
 apoplexy, 449
 Basedow's, 443
 blood, 441

Disease:

cardiac, 442
 cerebro-spinal meningitis, 444
 chorea, 452
 choroid, 193
 ciliary body, 188
 circulatory system, 442
 coma, 452
 conjunctiva, 52, 101
 cornea, 139
 dental, 442
 diabetes, 447
 digestive system, 442
 diphtheria, 444
 disseminated sclerosis, 451
 ductless glands, 443
 ear, 444
 encephalitis lethargica, 450
 epilepsy, 452
 erysipelas, 444
 Friedrich's, 450
 general, ocular manifestations
 of, 441
 paralysis, 450
 gonorrhœa, 444
 gout, 448
 Graves', 443
 headache, 448
 heart, 442
 hysteria, 453
 infective, 444
 influenza, 445
 intestines, 442
 intoxications, 454
 iris, 172
 kidneys, 447
 lacrymal apparatus, 77
 lens, 244
 leprosy, 445
 malaria, 445
 measles, 445
 meningitis, 450
 migraine, 448
 mumps, 445
 myelitis, 450
 naso-pharynx, 453
 nephritis, 447
 nervous system, 449
 nose, 453
 optic nerve, 295
 orbit, 87
 pneumonia, 445
 poisonings, 454
 pyæmia, 446
 retina, 272
 rheumatism, 449
 rickets, 449

Disease:

scarlatina, 445
 sclera, 166
 scurvy, 449
 septicæmia, 446
 stomach, 442
 tabes, 451
 teeth, 442
 tuberculosis, 446
 uveal tract, 187
 vaccinia, 446
 varicella, 447
 variola, 447
 vertigo, 449
 vitreous, 236
 whooping-cough, 447
 yellow fever, 447
 Disinfectants, 461
 Dislocation of lens, 269
 Dissector, blunt, 84
 Disseminated sclerosis, 451
 Distant vision, 11
 Distichiasis, operation for, 60
 Divergent squint, 422
 Diverging lens, 326
 meniscus lens, 327
 Dressings, 477
 Dry catarrh, 107
 Duboisine, 179, 465
 Ductless glands, disease of, 443
 Ear, diseases of, 444
 Ecchymosis, eyelids, 75
 Ectropion, Adam's operation for,
 68
 cicatricial, 66
 operations for, 68
 paralytic, 66
 senile, 66
 spasmodic, 66
 Edridge-Green's bead test, 310
 lantern test, 309
 theory, 308
 Egyptian ophthalmia, 123
 Electricity, 470
 Electric ophthalmoscope, 26
 welding, 283, 386
 Electrolysis, 60, 470
 Elliott's operation, glaucoma, 229
 'Embolism,' central retinal artery,
 286, 442
 Emmetropia, 339, 348
 Emphysema, eyelids, 76
 Encephalitis lethargica, 450
 Endocarditis, 442
 Enophthalmos, 89
 Entropion, 61

- Entropion, cicatricial, 61
 forceps, Knapp's, 62
 operations for, 62
 Jaesche-Arlt, 63
 Streatfeild-Snellen, 64
 senile, 61
 spasmodic, 61
 spastic, 65
 Enucleation of the eyeball, 96
 in glaucoma, 223
 Ephedrine, 466
 Epicanthus, 74
 Epilation, 60
 Epilepsy, 452
 Epiphora, 78
 Episcleral tissue, 101
 Episcleritis, 167, 168
 transient periodic, 168
 Epithelioma, eyelids, 75
 Erisophake, 263
 Errors of refraction, 357
 Erysipelas, 452
 Erythroptosis, 311
 Eserine salicylate, 466
 Esophoria, 424, 430
 Ethmoidal sinus, 87, 94
 Euphthalmine, 29, 465
 Eversion of eyelid, 2
 Evisceration of eyeball, 98
 Examination of the eye, 1
 colour sense, 21
 eyes of infants, 6
 field of vision, 15
 for heterophoria, 425
 for muscular paralysis, 400
 for squint, 412
 iris, 7
 lens, 25, 28, 248
 objective, 1, 23
 of anterior chamber, 7
 ophthalmoscopic, direct, 32
 distant, 27
 indirect, 29
 refraction, 345
 subjective, 11
 Exanthematous conjunctivitis, 105
 Excision, eyeball, 96
 lacrymal sac, 83
 Exenteration of the orbit, 100
 Exophoria, 424, 430
 Exophthalmic goitre, 443
 Exophthalmos, 89, 443
 pulsating, 92
 External ocular muscles, action
 of, 394
 operations on, 432
 paralysis of, 393
 Extraction by magnet, 241
 Eye, artificial, 99
 Snellen's, 99
 bath, 455
 cardinal points of, 338
 dioptric apparatus of, 337
 examination of, 1
 motor balance of, 424
 nodal points of, 338
 optical consideration of, 337
 principal points of, 338
 refraction of, 337, 339
 Eyeball, 87
 centre of rotation of, 338
 congenital anomalies of, 96
 enucleation of, 96
 evisceration of, 98
 movements of, 394
 tension of, 9
 Eyeglasses, fitting of, 383
 Eyelids, affections of the, 52
 anatomy of, 52
 burns of, 76
 carcinoma of, 75
 contused wounds of, 76
 ecchymosis of, 75
 emphysema of, 76
 incised wounds of, 76
 injuries of, 75
 insect bites, 76
 lacerated wounds of, 76
 physiology of the, 52
 rodent ulcer, 75
 sarcoma of, 75
 syphilis of, 56
 tumours of, 74
 vaccinia of, 56
 Eye-strain, 381
 Facet, corneal, 142
 Far point, 341
 Far-sightedness, 339
 Fascia, orbital, 87
 Fascicular keratitis, 129
 Fatty heart, 442
 Field for colours, 21
 of fixation, 395
 of vision, 11, 16
 perimeter in, 17
 tests for, 16
 Finger counting, 13
 Fistula of cornea, 144
 Fitting of eyeglasses, 383
 of spectacles, 383
 Fixation, field of, 395
 forceps, 225
 Flare, 50

- Fluorescein in corneal ulcer, 142,
 149
 in examination, 5
 Focal illumination, 23
 Foci of concave lens, 329
 of convex lens, 327
 Focus, conjugate, 328
 first principal, 338
 negative of convex lens, 329
 principal, 328
 second principal, 338
 virtual, of convex lens, 329
 Follicular catarrh, 108
 conjunctivitis, 108
 Fontana, spaces of, 211
 Forceps, advancement, 434
 cilia, 61
 curved iris, 225
 entropion, Knapp's, 62
 fixation, 225
 roller, Knapp's, 125
 Tyrrel's expression, 125
 Foreign bodies, conjunctiva, 137
 cornea, 163
 instruments for, 164
 magnet extraction of, 241
 protein, 179, 475
 vitreous, 239
 Form sense, 11
 Formalin, 461
 Fornix conjunctivæ, 101
 Fovea centralis, 272
 Fricke's blepharoplasty, 69
 Friedreich's disease, 450
 Frontal sinus, 94
 Functional examination of the
 eye, 11
 nervous disorders, 452
 Fundus oculi, examination of, 29,
 32
 normal, 38
 reflex, 28
 Fusion faculty, 395
 in squint, 411
 training of, 416
 Gait, uncertain, 399
 Galvano-cautery, 131, 149, 463
 General paralysis, 450
 Geneva lens measure, 335
 Germans, myopia in, 364
 Glands, lacrymal, 77
 Meibomian, 52
 of Moll, 52
 of Zeiss, 52
 Glassblowers' cataract, 46
 Glaucoma, absolute, 216
 acute congestive, 213
 stage, 214
 anatomy, 211
 chronic congestive, 217
 inflammatory, 217
 congenital, 96, 235
 cystoid cicatrix in, 228
 degeneration after, 216
 differential diagnosis, 222
 Elliott's operation, 229
 enucleation in, 223
 eserine in, 224, 234
 etiology, 219
 fulminans, 217
 hæmorrhagic, 235
 iridectomy, 224
 iridotaxis, 232
 iris angle, 211
 operations in, 224
 pathogenesis, 221
 pathology of, 226
 presbyopia, 390
 prodromal stage, 213
 prognosis, 222
 sclerotomy in, 232
 secondary, 234
 simple, 218
 tension in, 218
 treatment, 222
 varieties, 212
 Glaucomatous cup, 215
 halo, 216
 ring, 216
 state, 215
 Glaucosan, 469
 Glioma of the retina, 209
 pseudo-, 50, 199, 282, 444
 Glycosuria, retinitis in, 280
 Goitre, exophthalmic, 443
 Gold chloride, 163
 Gonin's operation, 293
 Gonococcal vaccine, 112, 181, 473
 Gonorrhœa, 444
 Gonorrhœal iritis, 181
 ophthalmia, 110
 vaginitis, 114
 Gout, 448
 Graefe knife, 254
 narrow, 254
 sign, 443
 Grafts, skin, 69, 135
 Granular conjunctivitis, 118
 lids, 118
 Gratiolet, fibres of, 315
 Grattage in trachoma, 126
 Graves' disease, 443
 Gravidic retinitis, 279, 447

Guist's operation, 293
 Gullstrand's corneal microscope,
 43, 44
 Gummata, 446
 Haab's magnet, 242
 Hæmophilia, 441
 Hæmorrhage after cataract opera-
 tion, 261
 amblyopia after, 441
 into the vitreous, 239
 retinal, 285
 subconjunctival, 137
 subhyaloid, 285
 Hæmorrhagic glaucoma, 235
 retinitis, 282
 Hand test, 16
 Hands of operator, preparation of,
 475
 Headache, 448
 Heart disease, 442
 Heat, 471
 Helmholtz theory, 341
 Hemeralopia, 288
 Hemiachromatopsia, 319
 Hemianopia, 317
 pupillary reaction in, 320
 Hemianopsia, 19, 315, 317
 altitudinal, 319
 binasal, 319
 bitemporal, 318, 451, 454
 crossed, 319
 homonymous, 318
 lateral, 318
 transient, 320
 Hemiopia, 317
 Herbert's operation, 65
 sclerotomy, 231
 Hering theory (colour), 308
 Herpes corneæ, 147
 zoster ophthalmicus, 55
 Herpetic ulcer, 144
 Hess's operation, 73
 Heterophoria, 382, 398, 424
 Hole at macula, 284
 (retinal), 291
 Holmgren's test, 310
 Holocain hydrochloride, 467
 Homatropine, 29, 383
 hydrobromide, 465
 Home Civil Service, visual require-
 ments for, 482
 Homonymous diplopia, 397
 hemianopsia, 318
 Hook, iris, blunt, 228
 tenotomy, 433
 Hordeolum, 57

Horn plate, 62
 Hot compresses, 471
 Hutchinsonian teeth, 155
 Hyaloid artery, persistent, 236
 canal, 236
 membrane, 236
 Hydrogen peroxide solution, 462
 Hydrophthalmos, 96, 235
 Hyoscyamine, 179, 465
 Hypermetropia, 339, 350, 357
 changes in the eye, 359
 symptoms, 360
 tests, 361
 treatment, 361
 Hyperphoria, 424
 Hyphæma, 7, 174
 Hypophysis, disease of, 443, 451
 Hypopyon, 7, 142, 174
 keratitis, 146
 Hysteria, 453
 traumatic, 311
 Hysterical amblyopia, 311
 Ichthyol, 460
 Illumination, focal, 23
 lateral, 23
 oblique, 23
 source of, 352
 Image, false, 396
 real, 329
 true, 396
 virtual, 329
 Images, 322
 formation of, 329
 Inch system, lenses, 334
 Index of refraction, 324
 Indian Civil Service, visual re-
 quirements for, 482
 Factory Department, 483
 Forest Department, 483
 Marine Service, visual re-
 quirements for, 484
 Medical Service, 484
 Pilot Service, 484
 Police Service, 482
 Public Works Department,
 483
 Railway Department, 484
 Survey Department, 483
 Telegraph Department, 483
 various artificers, 483
 Indigo, 308
 Indirect method, ophthalmoscope,
 29, 348
 Infant, examination of, 6
 Infantile purulent conjunctivitis,
 113

- Infective diseases, 444
- Influenza, 445
- Injection, ciliary, 102
 - conjunctival, 102
- Injections, subconjunctival, 472
- Injuries, ciliary body, 192
 - conjunctiva, 137
 - cornea, 163
 - eyelids, 75
 - iris, 182
 - orbit, 95
 - sclera, 176
- Insect bites, 75
- Inspection, 2
- Instruments, preparation of, 476
- Insufficiency, muscular. *See* Heterophoria
- Internal rectus, paralysis of, 401
- Interstitial keratitis, 152
- Intestines, disease of, 442
- Intoxications, 454
- Intracapsular extraction, 262
- Intra-ocular tumours, 207
 - optic neuritis, 296
- Invisible bifocal lens, 385
- Iodine in corneal ulcer, 149
 - tincture of, 462
- Iodoform, 462
 - poisoning, 303
- Ionization, 470
- Iridectomy, 226, 268
 - in cataract extraction, 256
 - indications for, 228
 - in glaucoma, 226
 - in staphyloma, 158
 - optical, 228
- Iridocyclitis, acute, 188
 - chronic, 188, 445
 - plastic, 190
 - purulent, 192
 - serous, 190
 - sympathetic, 201
- Iridocystectomy, 183
- Iridodialysis, 182
- Iridodonesis, 77, 270
- Iridotaxis, 232
- Iridotomy, 183
- Iris, anatomy of, 172
 - angle, 211, 220
 - bombé, 177
 - diseases of, 173
 - examination of, 7
 - forceps, curved, 225
 - hook, blunt, 228
 - injuries, 182
 - operations upon, 183, 224
 - repositor, 255
- Iris, scissors, 257
 - curved, 226
 - tumours of, 181
- Iritis, 173
 - clinical varieties, 180
 - complications, 177
 - diabetic, 181
 - differential diagnosis, 176
 - etiology, 177
 - gonorrhœal, 181
 - paracentesis in, 180
 - pathology, 177
 - rheumatic, 180
 - septic, 181
 - sequelæ, 177
 - spongy, 174
 - symptoms, objective, 174
 - subjective, 175
 - syphilitic, 180
 - treatment, 178
 - tuberculous, 181
- Irrigating apparatus, 258
- Jaeger's test-types, 14, 347
- Jaesche-Arlt operation, 63
- Javal-Schiötz ophthalmometer, 376
- Keratitis, 140
 - bullosa, 156
 - dendritiform, 147
 - disciformis, 157
 - due to exposure, 151
 - fascicular, 129
 - herpetic, 147
 - interstitial, 152
 - neuro-paralytic, 152
 - non-suppurative, 156
 - parenchymatous, 152
 - phlyctenular, 128
 - profunda, 156
 - punctata, 157, 159
 - (descemetitis), 189
 - ribbon-shaped, 157
 - sclerosing, 156, 168
 - superficial punctate, 156
 - suppurative, 141
 - vascular, 121
 - vasculo-nebulous, 121, 141, 156
 - vesicular, 141, 156
 - xerotic, 152
- Keratoconus, 159
- Keratoglobus, 161
- Keratomalacia, 144, 152
- Keratome, 228
- Keratoscope, Placido's, 6, 377

- Kidneys, 413
 Knapp's blepharoplasty, 69
 entropion forceps, 62
 knife-needle, 265
 roller forceps, 125
 Knife, Beer's, 58
 Graefe's, 254
 Took's, 230
 Weber's, 80
 Ziegler's, 262
 Knife-needle, Knapp's, 265
 Koch-Weeks bacillus, 105
 Krönlein's operation, 93, 100

 Lacarrère's intracapsular extrac-
 tion, 263
 Lacrymal apparatus, anatomy of,
 77
 diseases of, 79
 conjunctivitis, 79, 105
 gland, 77
 sac, 77
 abscess of, 89
 excision of, 83
 secretion, 78
 style, 83
 syringe, 81
 Lævo-glucosan, 224, 469
 Lagophthalmos, 71, 89
 Lagrange operation, 231
 Lamellar cataract, 268
 Lamina cribrosa, 166, 295
 vitrea, 193
 Lance-headed needle, 433
 Lang's eye-speculum, 225
 Lantern test, Edridge-Green's, 309
 Larsson's operation, 293
 Latent hypermetropia, 359
 Lateral illumination, 21
 Lawyer's questions, 313
 Lead acetate, 460
 poisoning, 298
 Leaking cicatrix, 229
 Leber's disease, 301
 Leeches, 179, 471
 Lens, anatomy of, 244
 architecture of, 46
 biconcave, 327
 biconvex, 326
 bifocal, 384
 capsule of, 244
 centre of, finding, 335
 complete dislocation of, 270
 concave, foci, of, 329
 meniscus, 327
 periscopic, 327
 concavo-convex, 326
 Lens, concavo-spherical, 326
 converging, 326
 convex, foci of, 327
 periscopic, 326
 spherical, 325
 convexo-concave, 327
 cylindrical, 331
 diseases of, 244
 dislocation of the, 269
 diverging, 326
 meniscus, 327
 Helmholtz theory, 341
 luxation of, 270
 magnifying, 326
 measure, Geneva, 335
 meniscus, 326
 minus, 326
 negative, 326
 neutralizing, 335
 nucleus, 47
 partial dislocation of, 270
 periscopic, 326
 physiology of, 244
 plano-concave, 327
 plano-convex, 326
 plus, 326
 positive, 326
 reducing, 326
 scoop, wire, 258
 spherical, 325
 structure of, 46
 subluxation of, 270
 sutures, 48
 Tscherning theory, 341
 Lenses, 325
 decentration of, 431
 numeration of, 332
 trial case, 334
 varieties of, 336
 Leprosy, eye affections of, 445
 Leucoma, 5, 161
 adherent, 143, 161
 tattooing in, 162
 Leukæmia, 407
 retinitis in, 280
 Lid retractor, Desmarre's, 7
 Lids. *See* Eyelids, 52
 œdema of, 55
 Ligament, suspensory, 244
 Ligaments, palpebral, 53
 tarsal, 53
 Ligamentum pectinatum, 211
 Light, excessive, 283
 sense, 11, 21
 Limbus corneæ, 140
 Lime burns, 137
 Line of fixation, 339

- Linear extraction, 263
 Lipæmia, retinal, 280
 Lister perimeter, 17
 Lithiasis of the conjunctiva, 108
 Local anæsthetics, 467
 blood-letting, 471
 Locomotor ataxy, 304
 Lost fixation, 413
 Lumbar puncture, 300
 Luxation of the lens, 270

 Macropsia, 194
 Macula, 5, 161
 lutea, 40, 272
 changes in infancy at, 283
 Maddox rod, 425
 wing, 427
 Magnet extraction, 241
 Haab's, 241
 Mellinger's ring, 243
 Magnifying lens, 326
 Malaria, 445
 amblyopia in, 314
 Malinger, 312
 Manifest hypermetropia, 359
 Marginal ulcer, cornea, 147
 Massage, 471
 May's electric ophthalmoscope, 26
 Measles, 445
 Mechanism of accommodation, 340
 Meibomian cyst, 57
 glands, 52, 53
 Melano-sarcoma, 207
 Membrana nictitans, 101
 Membranous conjunctivitis, 116
 Meningitis, 200, 298, 450
 cerebro-spinal, 444
 Meningocele, 93
 Meniscus lenses, 326
 Mercantile Marine, Board of Trade
 tests for, 482
 Mercuric chloride, 456
 Mercury, ammoniated, 460
 bichloride, 476
 yellow oxide, 459
 Metamorphosis, 194, 275
 Metaphen, 461
 Metastatic ophthalmia, 198
 retinitis, 282
 Methods of investigating field of
 vision, 16
 oculo-motor paralysis, 400
 refraction, 345
 test types, 346
 Metri cautery, 150, 463
 Metric system, lenses, 332

 Microphthalmos, 96
 Micropsia, 194
 Microscope, corneal, 43, 44
 Migraine, 448
 Milium, 75
 Mind-blindness, 317
 Miner's nystagmus, 408
 Minus lens, 326
 Miotics, 466
 'Mires,' 376
 Mirror, reflection by, 321
 Mitigated stick, 462
 Moll, glands of, 52
 Molluscum, 74
 Monochromic vision, 309
 Monochromics, 309
 Monocular cataract, 252
 bandage, 478
 Mooren's ulcer, 146
 Moorfields bandage, 258, 479
 Morax-Axenfeld bacillus, 105, 108
 Morgagnian cataract, 249
 spherules, 249
 Morton's ophthalmoscope, 26
 Motais' ptosis operation, 73
 Motor balance of eyes, 424
 imbalance, 412
 Movements of the eyeball, 394
 Mucocoele, 79
 Mueller's fibres, 273
 contact glasses, 160
 Mule's operation, 99
 Mumps, 445
 Muscæ volitantes, 20, 237, 365
 Muscle, ciliary, 187, 356
 levator palpebræ, 52
 orbicularis, 52
 Muscles, external ocular, 393
 operations on, 432
 paralysis of, 398
 'Mutton fat' deposits, 190
 Myasthenia gravis, 350, 363
 Mydriatics, 29, 383, 465
 Myelitis, 450
 Myopia, 339, 350, 363
 choroiditis in, 197
 clinical forms, 364
 etiology, 363
 in Germans, 364
 malignant, 365
 operative treatment, 369
 ophthalmoscopic signs, 365
 prognosis, 366
 progressive, 364
 simple, 364
 stationary, 364
 symptoms, 365

- Myopia tests, 366
 - treatment, 367
- Myopic crescent, 197
 - divergent squint, 422
 - far point, 352
- Myxœdema, 443
- Nasal duct, 78
 - probing, 83
 - sinuses, 94
- Naso-pharynx, diseases of, 453
- Nausea in diplopia, 399
- Naval cadets, vision of, 480
- Navy, visual requirements for, 480
- Near point, 341
 - sightedness, 339
 - vision, 14
- Nebula, 5, 161
- Needle holder, 338
 - tattooing, 162
- Needling of cataract, 265
- Negative focus, 329
 - lens, 326
- Nephritis, 447
 - retinitis of, 278
- Nerve paths, 184, 316
- Nerves of ocular muscles, paralysis of, 393
- Nervous disorders, functional, 451
 - system, diseases of, 449
- Nettleship's dilator, 80
- Neuritis, descending, 297
 - retrobulbar, 300
- Neuro-paralytic keratitis, 141, 152
- Neuropathic divergence, 448
 - divergent squint, 422
- Neuro-retinitis, 297
- New-born, conjunctivitis in, 115
- Night-blindness, 288, 314
- Nitrate of cocaine, 462
 - of silver, 458, 462
- Nitro-benzol poisoning, 303
- Nodal points of the eye, 338
- Non-comitant squint, 423
- Normal fundus, 38
- Nose, diseases of, 453
- Novocain, 467
- Nuclear cataract, 246
- Numeration of lenses, 332
- Nystagmus, 408
 - in ear disease, 444
 - lateral, 408
 - miner's, 409
 - rotatory, 408
 - vertical, 408
- Objective examination of eye, 23
- Oblique illumination, 23
- Oblique inferior, paralysis of, 404
 - superior, paralysis of, 403
- Obturator, 335
- Occlusion of pupil, 177
- Ocular deviations, varieties of, 397
 - manifestations of general diseases, 441
 - muscles, external, paralysis of, 393
 - paralysis of, 398
 - external rectus, 401
 - fourth nerve, 403
 - inferior oblique, 404
 - rectus, 402
 - internal rectus, 401
 - operation in, 408
 - prognosis, 407
 - sixth nerve, 401
 - superior oblique, 403
 - rectus, 401
 - third nerve, 404
 - treatment, 407
 - varieties of, 401
 - therapeutics, 455
- Oculo-motor paralysis, investigation of, 400
- Edema, angio-neurotic, 55
 - of the lids, 54
- Old sight, 388
- Opacities, cornea, 25, 161
 - media, 28
 - vitreous, 237
- Operations, Adam's (ectropion), 65
 - advancement of an ocular muscle, 432
 - blepharoplasty, 68
 - canaliculus slitting, 82
 - canthoplasty, 65
 - canthotomy, 66
 - cataract, 253
 - cicatrix, leaking, 229
 - conical cornea, 160
 - corneo-scleral trephining, 229
 - cyclodialysis, 232
 - Dieffenbach's, 69
 - discission, 265
 - distichiasis, 60
 - ectropion, 68
 - electrolysis, 60
 - Elliott's (glaucoma), 229
 - entropion, 62
 - enucleation of eyeball, 96
 - epicanthus, 74
 - epilation, 60
 - evisceration of eyeball, 98
 - excision of lacrymal sac, 83
 - exenteration of the orbit, 100

- Operations, expression, trachoma,
 125
 external ocular muscles, 432
 eyeball, 96
 Fricke's, 69
 general conditions, 475
 Gonin's, 283
 glaucoma, 224
 Guist's, 283
 Herbert's modification of
 Streatfeild-Snellen, 65
 sclerotomy, 231
 Hess's, 73
 intracapsular extraction, 262
 iridectomy for glaucoma, 224
 iridocystectomy, 183
 iridotaxis, 232
 iridotomy, 183
 iris, 183, 224
 Jaesche-Arlt, 63
 keratoconus, 160
 Knapp's, 69
 Krönlein's, 93, 100
 Lagrange, 231
 Larsson's, 293
 Motais', 73
 Mule's, 99
 needling, 239
 paracentesis, cornea, 150
 in iritis, 180
 preparations for, 475
 probing nasal duct, 83
 ptosis, 73
 recession, 439
 Saemisch's (ulcer), 151
 Safar's, 293
 sclerotomy, 232
 skin-grafting, 69
 spastic entropion, 65
 Streatfeild-Snellen, 64
 tarsorrhaphy, 70
 tenotomy, 421, 437
 Toti's, 85
 trichiasis, 62
 V Y (ectropion), 68
 West's, 85
 Wharton Jones's, 68
 Worth's, 160
 squint, 432
- Operator's hands, preparation of,
 475
- Ophthalmia, 103
 Egyptian, 123
 gonorrhoeal, 110
 neonatorum, 113
 prophylaxis, 115
 scrofulous, 128
- Ophthalmitis, sympathetic, 201
 symptoms, 202
 theories of, 203
 treatment, 204
- Ophthalmometer, 376
- Ophthalmoplegia, 405
 interna, 390, 406
- Ophthalmoscope, 25
 at a distance, 348
 examination by, 27
 direct, 32, 349
 indirect, 29, 348
- May's, 26
 Morton's, 26
 refractive error by, 348
 source of illumination, 352
 theory of, 35
- Ophthalmoscopy with red - free
 light, 37
- Optic disc, 31, 38, 272
 congestion of, 296
 cupping of, 39, 215
 nerve, anatomy of, 295
 atrophy of, 303
 diseases of, 295
 inflammation of, 296
 neuritis, 296
- Optical axis, 336
 considerations of the eye, 337
 iridectomy, 228
 optochin, 463
 principles, general, 321
- Ora serrata, 272
- Oral sepsis, 442
- Orbit, anatomy of, 87
 diseases of, 88
 injuries of, 95
 tumours of, 93
- Orbital cellulitis, 91, 444
 fascia, 87
 periostitis, 89
- Orthophoria, 424
- Osteoma, 93
- Palpation, 8
- Pannus, 121, 141, 156
- Panophthalmitis, 205
- Pantocaine, 468
- Papilla, 31, 38, 272. *See* Optic disc
- Papillitis, 296
- Paracentesis in iritis, 180
 of cornea, 150, 155
- Paralyses, accommodation, 390
 associated, 406
 basilar, 406
 central, 406
 congenital, 407

- Paralysis, conjugate, 406
 cortical, 406
 differential diagnosis, 406
 external ocular muscles, 398
 nuclear, 406
 ocular, 398
 oculo-motor, 400
 orbital, 406
 peripheral, 406
 Parenchymatous keratitis, 141,
 152
 Parinaud's conjunctivitis, 128
 Parturition, 454
 Patient, history of, 1
 position of, 476
 preparation of, 475
 Pediculi, 54
 Perception of light, 14
 Perforation of the cornea, 143
 Perimeter, 17, 395
 in squint, 413
 Periostitis, orbital, 89
 Peripheral vision, 16
 Periscopic lens, 326
 Peritomy, 127
 Pernicious anæmia, 441
 Persistent hyaloid artery, 236
 Phakoerisis, 263
 Phlyctenular conjunctivitis, 128
 keratitis, 120, 141
 Photophobia, 130
 Photopsia, 290
 Phthisis bulbi, 205
 Physiological cuppings of disc, 39
 Pilocarpin in glaucoma, 223
 muriate, 466
 nitrate, 466
 Pinguecula, 135
 'Pink eye,' 105
 Pituitary disease, 303, 451
 Placido's disc, 5, 377
 Plane mirror, 321
 Plano-concave lens, 327
 Plano-convex lens, 326
 Plastic cyclitis, 190
 Plica semilunaris, 101
 Plus lens, 326
 Pneumococcus, 106
 Pneumonia, 445
 Poisoning, atropine, 464
 Polyvalent vaccines, 473
 Positive lens, 326
 Potassium chlorate, 458
 Pregnancy, 454
 Preparation of hands, 475
 instruments, 476
 patient, 475
 Preparation, region, 476
 Presbyopia, 343, 388
 glaucoma in, 390
 symptoms, 389
 treatment, 389
 Prescribing glasses for young
 patients, 379
 Principal points of the eye, 338
 Prism duccion, 427
 position of, 325
 refraction by, 324
 rotary, 428
 Prisms, numbering of, 325
 use of, 325
 Probes, Cowper's, 81
 Teale's, 81
 Probing nasal duct, 81
 Projection, 253, 396
 false, 399
 Proptosis, 89
 Protargol, 462
 Protective glasses, 386
 Pseudo-glioma, 199, 282, 444
 Pterygium, 135
 Ptosis, 72
 operations for, 73
 Hess's, 73
 Motaïs', 73
 Pulsating exophthalmos, 92
 Puncta lacrymalia, 77
 anomalies, 78
 Punctum remotum, 341
 Pupil, the, 7, 183
 Argyll-Robertson, 185
 dilatation of, 29
 occlusion of, 177
 Pupillary membrane, 173
 reaction, hemianopic, 320
 reflex nerve paths, 183
 Purpura, 441
 Purulent conjunctivitis, 110
 infantile, 113
 cyclitis, 192
 retinitis, 282
 Pustular conjunctivitis, 128
 Pyæmia, 446
 Pyramidal cataract, 267
 Quinine amblyopia, 314
 ischæmia, 284
 Radium, 75, 93, 127, 208, 210
 rodent ulcer and, 75
 spring catarrh and, 133
 trachoma and, 127
 Radon seeds, 93, 210
 Range of accommodation, 341

- Range of convergence, 344
 Recession, 439
 Recti muscles, 393
 Rectus externus, paralysis of, 401
 inferior, paralysis of, 402
 internus, paralysis of, 401
 superior, paralysis of, 401
 Red-free light, ophthalmoscopic examination with, 37
 Reducing lens, 326
 Reflecting glass, 387
 Reflection, 321
 concave mirror, 321
 convex mirror, 323
 plane mirror, 321
 specular, 44
 Reflex paths, pupillary, 184
 senile, 245
 visual, 184
 Refraction, 321
 errors of, 357
 index of, 324
 methods of investigating, 345
 of the eye, 339, 341
 ophthalmoscope method, 348
 prism, 324
 subjective method, 346
 Remedies, astringent, 457
 constitutional, 455
 local, 455
 stimulating, 457
 Repositor (iris), 255
 Retina, 40
 anæmia of, 284
 anatomy of, 272
 central artery, obstruction of, 286
 vein, thrombosis of, 287
 circulatory disturbances of, 284
 colour perceiving elements in, 308
 contusion of, 283
 detachment of, 290
 diseases of, 274
 functional diseases of, 306
 glioma of, 215
 hæmorrhages, 282
 hyperæmia of, 284
 ischæmia of, 284
 œdema of, 276, 283
 physiology of, 274
 pigmentary degeneration of, 288
 section of, 273
 Retinal changes due to excessive light, 283
 uncommon forms, 282
 Retinal hæmorrhages, 282, 441
 hole, 291
 tear, 291
 lipæmia, 280
 Retinitis, 275
 albuminuric, 278
 Bright's disease, 278
 circinata, 283
 deep, 277
 diabetic, 280
 gravidic, 279, 447
 hæmorrhagic, 282
 leukæmic, 280, 407
 metastatic, 282
 parenchymatous, 277
 pigmentosa, 288
 proliferans, 239, 283
 punctate, 290
 purulent, 282
 renal, 278
 septic, 282
 serous, 277
 simple, 277
 striate, 283
 syphilitic, 281
 Retino-choroiditis, 194
 Retinoscopy, 351
 Retractor, 84
 Desmarre's lid, 7
 Retrobulbar neuritis, 300
 Retro-illumination, 44
 Retrotarsal fold, 3
 Rheumatism, 449
 iritis in, 177, 180
 Ribbon-shaped keratitis, 157
 Rickets, 449
 Ripe cataract, 249
 Rod, Maddox, 425
 Rodent ulcer of eyelids, 75
 Rods and cones, 273
 Roller forceps, Knapp's, 125
 Rotary prism, 428
 Rotation, centre of, 338
 Royal Air Force, visual requirements for commissions in, 484
 Royal Navy, visual requirements for commissions in, 480
 Rupture of choroid, 200
 Sac, lacrymal, 77
 Saemisch's method, 151
 Safety glass, 384, 387
 Salvarsan, 155, 472
 Sarcoma, choroid, 207
 eyelids, 75
 Scalpel, 66
 Scarlatina, 445

- Schiötz, tonometer of, 9
 Schlemm's canal, 188, 211
 Scintillating scotoma, 320
 Scissors, enucleation, 95
 iris, curved, 226
 de Wecker's, 257
 squint, 433
 Sclera, anatomy of, 166
 diseases of, 166
 injuries of, 170
 staphyloma of, 169
 Scleritis, 166, 168, 475
 Sclero-choroiditis, 197
 Sclerotomy, anterior, 232
 in glaucoma, 232
 posterior, 233
 Scoop, chalazion, 59
 wire lens, 258
 Scopolamine, 179, 465
 Scotoma, 19, 275
 colour, 302
 scintillans, 320
 Scrofulous ophthalmia, 128
 Screen test, 413, 425
 Scurvy, 449
 'Second sight,' 363
 Secondary glaucoma, 234
 Senile cataract, 247
 entropion, 61
 reflex, 245
 Septic retinitis, 282
 Septicæmia, 446
 Serous retinitis, 277
 Serpent ulcer, cornea, 146
 Service requirements, visual, 480
 Army, 480
 Home Civil Service, 482
 Indian Civil Service, 482
 Factory Department, 483
 Forest Department, 483
 Marine Service, 484
 Medical Service, 484
 Pilot Service, 484
 Police Service, 484
 Public Works Department, 484
 Railway Department, 454
 Survey Department, 483
 Telegraph Department, 483
 Various Artificers' Departments, 483
 Navy, 480
 Shadow test, 351
 Short-sightedness, 363
 Shrunk cataract, 249
 Siderosis bulbi, 240
 Sight-testing, 249, 346
 Sign, Stellwag's, 443
 von Graefe's, 443
 Signs used in ophthalmology, 336
 Silver nitrate, 458, 462
 Sinus, ethmoidal, 87, 94
 frontal, 87, 94
 nasal, 87, 94
 of the anterior chamber, 211
 sphenoidal, 87, 94
 Sixth nerve, paralysis of, 401
 Skiascopy, 351
 Skin-grafting, 69
 Thiersch's, 70
 Wolfe's, 70
 Slit-lamp, Gullstrand's, 43, 44
 Snellen's artificial eyes, 100
 test-types, 11
 Snow-blindness, 105, 283
 Sodium chloride, 456
 Solid carbon dioxide, 469
 Solutions, antiseptic, 455
 cleansing, 455, 477
 Spasm of accommodation, 391
 Spectacles, fitting of, 383
 telescopic, 368
 Spectrum, colours in, 308
 Specular reflection, 44
 Speculum, Lang's, 225
 Sphenoidal sinus, 87, 94
 Spherical lenses, 325
 action of, 327
 Sphincter pupillæ, 172, 183
 Spoon, 84, 258
 Volkmann's, 84
 Spring catarrh, 132
 and radium, 133
 Squint, accommodative, 412
 advancement for, 432
 alternating convergent, 410
 amblyopia of, 411
 amblyoscope in, 417
 angle of deviation, 413
 atropine in, 416
 comitant, 410
 etiology of, 411
 varieties of, 410
 convergent, operation in, 421
 treatment of, 415
 divergent, 422
 method of investigating, 412
 myopic, 422
 neuropathic, 422

- Squint, non-comitant divergent, 423
 occlusion of eye in, 415
 operations for, 432
 optical correction in, 415
 orthoptic training, 416
 perimeter in, 413
 scissors, 433
 tenotomy in, 421
 treatment of, 415, 422
- Staphylococcal vaccines, 473
- Staphylococcus, 106
- Staphyloma, anterior, 169
 ciliary, 169
 corneal, 157
 equatorial, 169
 intercalary, 169
 posterior, 198
 scleral, 169
- Stationary cataracts, 246, 267
- Stellwag's sign, 443
- Stenopæic slit, 160, 374
- Stimulating remedies, 457
- Stomach, disease of, 442
- Strabismus. *See* Squint
- Streitfeild-Snellen operation, 64
- Streptococcus, 105, 106
- Strumous conjunctivitis. *See* Phlyctenular keratitis
- Phlyctenular conjunctivitis
 keratitis. *See* Phlyctenular keratitis
- Stye, 57
- Style, lacrymal, 83
- Subconjunctival hæmorrhage, 137
 injections, 472
- Subhyaloid hæmorrhage, 285
- Subjective examination of the eye, 319
- Subluxation of lens, 270
- Superficial punctate keratitis, 156
- Suppurative keratitis, 156
- Suprachoroid, 193
- Suprarenalin, 469
- Suspensory ligament, lens, 244
- Symblepharon, 134
 in trachoma, 122
- Symbols used in ophthalmology, 336
- Sympathetic ophthalmia, 201
- Synchysis scintillans, 237
- Synechia, anterior, 7, 143
 posterior, 7, 174
- Synoptophore, 417
- Syphilis, 446
 and papillitis, 298
 of the eyelids, 56
- Syphilitic choroido-retinitis, 197, 289
- Syphilitic iritis, 180
 retinitis, 281
- Syringe, lacrymal, 81
- Tabes, 451
- Tannic acid, 458
- Tarsal cyst, 57
 tumour, 57
- Tarsorrhaphy, 70
- Tarsus, 53
- Tattooing cornea, 162
- Teale's probe, 81
- Teeth, diseases of, 442
 Hutchinsonian, 155
- Telescopic spectacles, 368
- Tenon's capsule, 88, 393
- Tenonitis, 92
- Tenotomy in squint, 421, 423, 437
- Tension of eyeball, 8, 158, 212
- Test, astigmatism, 373
 blackboard, 16
 Calmette's, 474
 candle, 16
 colour vision, 309
 diplopia, 400
 hand, 16
 Holmgren, 310
 malingering, 312
 mirror, 400
 in squint, 412
 motor balance of eyes, 424
 types for illiterates, 14
 Green's, 12
 Jaeger, 15, 347
 method, 346
 Snellen's, 11
 von Pirquet's, 472
- Testing drum, 476
- Tetrachromics, 309
- Theory, colour perception:
 Edridge-Green's, 308
 Helmholtz, lens, 341
 Hering's, 308
 ophthalmoscope, 35
 Tscherning, lens, 341
 Young-Helmholtz, 308
- Therapeutics, ocular, 455
- Thiersch's skin-grafting, 70
- Third nerve, paralysis of, 404
- Thrombosis of artery, central, of retina, 286
 of cavernous sinus, 91
 of central retinal vein, 287
- Tincture of iodine, 149, 462
- Tobacco amblyopia, 301
- Tonometer of Schiötz, 9
- Took's knife, 230

- Toric lenses, 384
 Toti's operation, 85
 Toxic amblyopia, 301
 Trachoma, 118
 clinical varieties, 120
 complications, 121
 etiology, 122
 expression in, 125
 granular form, 119
 grattage in, 126
 mixed form, 120
 papillary form, 119
 prophylaxis, 127
 radium and, 127
 sequelæ, 121
 treatment, 123
 Transient hemianopsia, 320
 Transillumination, 41
 Transparent ulcer, cornea, 147
 Traumatic cataract, 266
 conjunctivitis, 104
 hysteria, 311
 Trephine, Elliott's, 230
 Trephining, corneo-scleral, 229
 Trial case, 334
 frame, 346
 Trichiasis, 60
 operations for, 62
 Trichromics, 309
 Trifocal lenses, 386
 Tscherning theory, lens, 341
 Tubercles of the choroid, 200
 Tuberculin, 156, 168, 180, 473
 old, 473
 Tuberculosis, 446
 Tumours, intra-ocular, 207
 of brain, 269, 417
 of eyelids, milium, 75
 molluscum, 74
 xanthelasma, 74
 of iris, 181
 orbit, 93
 Tyrrel's forceps, 125

 Ulbrich's graduated drum, 45
 Ulcer of cornea, catarrhal, 145
 dendriiform, 147
 herpetic, 144
 marginal ring, 147
 Mooren's, 146
 of cornea, 141
 serpens, cornea, 146
 spontaneous perforation of, 143
 transparent, 147
 Ultra-violet light, 132, 180, 471
 Uncertain gait, 399

 Undine, 254, 258
 Uræmic amblyopia, 280
 Uvea, 172
 Uveitis, 188
 anterior, 168
 purulent, 192, 198, 205
 sympathetic, 192, 201

 Vaccines, autogenous, 473
 gonococcal, 473
 in ophthalmology, 473
 staphylococcal, 473
 treatment by, 55, 191
 Vaccinia, 446
 of the eyelids, 56
 Valvular heart disease, 442
 Varicella, 447
 Variola, 447
 Vasa vorticosa, 193
 Vascular keratitis, 121, 156
 Vasculo-nebulous keratitis, 121, 156
 Vectis, wire, 258
 Vein, central, of retina, 39, 272
 thrombosis of, 287
 Vernal catarrh, 132
 Verruca, 74
 Vertigo, 399, 449
 Vesicular keratitis, 141, 156
 Virtual focus, convex lens, 329
 image, 329
 Vision, acuteness of, 11, 340
 binocular, 395
 central, 11
 colour, 307
 coloured, 309
 direct, 11
 distant, 11
 field of, 11, 16
 monochromic, 309
 near, 14
 peripheral, 14
 suppressed, 411
 Visual angle, minimum, 12
 area, 315
 line, 338
 paths, 316
 requirements for public services, 480
 Vitreous, anatomy of, 236
 diseases of, 236
 foreign bodies in, 239
 hæmorrhages into, 239
 opacities, 237
 Von Graefe's sign, 443
 Von Pirquet's test, 474
 V Y operation, ectropion, 68

- Watery eye, 78
Weak sight, 381
Weber's knife, 80
Wernicke's hemianopic reaction, 320
West's operation, 85
Weve's operation, 294
Wharton-Jones's operation, 68
Whooping-cough, 258
Wire lens vectis, 258
Wolfe's skin-grafting, 70
Wood-alcohol poisoning, 301
Word-blindness, congenital, 307
Workmen's Compensation Act, 313
Worth's operation, 160, 432
Wounds of cornea, 163
 of eyelids, 75
 of sclera, 170
Xanthelasma, 74
Xerosis of conjunctiva, 122, 314
Xerotic keratitis, 141, 152
X-rays, 124, 165, 210, 240, 242, 470
Yellow fever, 447
 oxide mercury, 459
 spot, 272
Young-Helmholtz theory, 308
Y suture of lens, 48
Zeis, glands of, 52
Zeiss contact glasses, 160
Ziegler's knife, 262
Zinc sulphate, 457
Zonular cataract, 268
 lamella, 46

A Selected List
of
Standard Medical Books
Published by
BAILLIÈRE, TINDALL & COX

Partners: A. A. TINDALL, R. F. WEST

*Owing to the fluctuations in the exchange the
prices of American books and periodicals in this
catalogue are subject to alteration without notice*

Telegraph:
BAILLIÈRE-PHONE
LONDON



Telephone:
TEMPLE BAR 8568
(2 lines)

Established 1826

LONDON
7 & 8, HENRIETTA STREET, COVENT GARDEN

September, 1934

ANAESTHESIA

- HADFIELD—Practical Anaesthesia.** By C. F. Hadfield, M.B.E., M.A., M.D.
Second Edition. Pp. 352. Figs. 41. Price 7s. 6d.

ALLERGY, DIET AND FOOD

- COCA—Asthma and Hay Fever in Theory and Practice.** Pt. 1 by A. F. Coca, M.D. Pt. 2 by M. Walzer, M.D. Pt. 3 by A. A. Thommen, M.D.
Pp. 875. Figs. 97. 2 Plates. Price 38s.
- ROWE—Food Allergy.** By A. H. Rowe, M.S., M.D. Pp. 442. Price 25s.
- SURE—The Vitamins in Health and Disease.** By B. Sure, Ph.D. Pp. 220.
Price 9s.
- TIBBLES—Foods: Their Origin, Composition and Manufacture.** By W. H. Tibbles, M.D., M.R.C.S. Pp. 958. Price 20s.
- WALKER—Food Inspector's Encyclopaedia.** By A. H. Walker, M.R.San.I.
Pp. 310. Price 8s. 6d.

ANATOMY, EMBRYOLOGY AND HISTOLOGY

- Aids to Anatomy.** By C. H. Fagge, F.R.C.S. 9th Ed. Pp. 333. Price 5s.
- Aids to Embryology.** By R. H. Hunter, M.D. Third Edition. Pp. 184.
Figs. 30. Price 3s. 6d.
- Aids to Histology.** By A. Goodall, M.D., F.R.C.P. Third Edition. Pp. 156.
Figs. 20. Price 3s. 6d.
- Aids to Osteology.** By PHILIP TURNER, B.Sc., M.B., M.S.(Lond.), F.R.C.S.,
and N. L. ECKHOFF, M.S.(Lond.), F.R.C.S. Third Edition. Pp. 230.
Price 4s. 6d.
- BAILLIÈRE'S Synthetic Anatomy.** An atlas of the whole human body, printed
in colours on transparent paper. By J. E. Cheesman. In 13 parts in binding
case. Price 42s.
Single parts, containing 12 Plates and Text. Price 3s.
- BUCHANAN—Manual of Anatomy.** Fifth Edition. Edited by E. Barclay
Smith, M.D., J. E. Frazer, F.R.C.S., F. G. Parsons, F.R.C.S., and W.
Wright, M.Sc., F.R.C.S. In three volumes. Pp. 2022. Figs. 820. Price 42s.
Without Vol. III., Dissection Guide. Price 35s.
- CAJAL—Histology.** By S. Ramon Cajal, F.R.S., LL.D. Pp. 752. Figs. 335.
Price 36s.
- CLARK—Early Forerunners of Man.** By W. E. Le Gros Clark, D.Sc.
(Lond.), F.R.C.S.(Eng.). Pp. 312. Figs. 89. Price 15s.
- DRINKER—Lymphatics, Lymph and Tissue Fluid.** By Cecil K. Drinker,
B.S., M.D. Pp. 270. Figs. 14. Price 13s. 6d.
- FRAZER—Manual of Embryology.** By J. E. Frazer, F.R.C.S. Pp. 494.
Figs 282. Price 30s.

BACTERIOLOGY AND LABORATORY

- Aids to Bacteriology.** By W. Partridge, F.I.C. Fifth Edition. Pp. 320. Price 5s.
- BERGEY—Manual of Determinative Bacteriology.** By D. H. Bergey, M.D.
Fourth Edition. Pp. 680. Price 27s.
- BIGGER—Handbook of Bacteriology.** By J. W. Bigger, M.D. Third Edition.
Pp. 476. Figs. 84. 5 Coloured Plates. Price 12s. 6d.
- BUCHANAN and FULMER—Physiology and Biochemistry of Bacteria.**
By R. E. Buchanan, Ph.D., and E. I. Fulmer, Ph.D. In three volumes.
Pp. 1848. Figs. 137. Price 102s.

- EYRE—Bacteriological Technique.** By J. W. H. Eyre, M.D. Third Edition. Pp. 632. Figs. 238. 2 Coloured Plates. Price 21s.
- HARTMAN—The Anatomy of the Rhesus Monkey.** By C. G. Hartman. Pp. 392. Figs. 122. Price 27s.
- KENDALL—Bacteriology: General, Pathological and Intestinal.** By A. I. Kendall, B.S., Ph.D. Third Edition. Pp. 733. Figs. 103. 8 Coloured Plates. Price 31s. 6d.
- LEVINE and SCHOENLEIN—Compilation of Culture Media for the Cultivation of Micro-organisms.** Pp. 985. Price 67s. 6d.
- PARK and WILLIAMS—Pathogenic Micro-organisms.** By W. H. Park, M.D., and Anna W. Williams, M.D. Tenth Edition. Pp. 877. Figs. 216. 11 Plates. Price 35s.
- PARTRIDGE—Dictionary of Bacteriological Equivalents.** By W. Partridge, F.I.C. Pp. 154. Price 10s. 6d.
- SMITH—Oral Spirochetes and Related Organisms in Fusio-Spirochetal Disease.** By D. T. Smith, B.A. Pp. 256. Figs. 53. Price 18s.
- THOMSON—The Pathogenic Streptococci.** Their Role in Disease. Vols. III.-X. of the Annals of the Pickett-Thomson Laboratories. Price per vol. 42s.
- WADSWORTH—Standard Methods of the Division of Laboratories and Research of the New York Department of Health.** By A. B. Wadsworth, M.D. Pp. 724. Figs. 56. 11 Plates. Price 34s.
- WILLIAMS—Streptococci in Relation to Man in Health and Disease.** By Anna W. Williams, M.D. Pp. 272. 8 Plates. Price 22s. 6d.

BIOCHEMISTRY

- Aids to Biochemistry.** By A. Ashley Cooper, D.Sc., F.I.C., and S. D. Nicholas, F.I.C., B.A. Pp. 196. Figs. 11. Price 4s. 6d.
- BROWNING—The Vitamins.** By E. Browning, M.D. Pp. 807. Price 42s.
- KOCH—Practical Methods in Biochemistry.** By Frederick C. Koch. Pp. 289. Figs. 17. Price 10s.
- MATHEWS—Physiological Chemistry.** For Students. By A. P. Mathews, Ph.D. Fifth Edition. Pp. 1252. Figs. 109. Price 31s. 6d.
- MUNCH—Bioassays: A Handbook of Quantitative Pharmacology.** By J. C. Munch. Pp. 968. 239 Tables. Figs. 15. 3 Coloured Plates. Price 45s.
- PETERS and VAN SLYKE—Quantitative Clinical Chemistry.** By J. P. Peters, M.A., M.D., and D. D. Van Slyke, Ph.D., Sc.D. Vol. I. Interpretations. Pp. 1280. Figs. 124. Price 54s. Vol. II. Methods. Pp. 978. Figs. 95. Price 45s.
- Both volumes purchased together 90s.

BIOLOGY

- Aids to Biology.** By R. G. Neill, M.A. Pp. 258. Figs. 27. Price 3s. 6d.
- ARTSCHWAGER—Dictionary of Biological Equivalents.** German-English. By E. Artschwager. Pp. 240. 6 Plates. Price 20s.

BLOOD

- FISHBERG—Hypertension and Nephritis.** By A. M. Fishberg, M.D. Third Edition. Pp. 684. Figs. 39. Price 32s. 6d.
- SNYDER—Blood Groupings in Relation to Clinical and Legal Medicine.** By L. H. Snyder, D.Sc. Pp. 166. Figs. 28. 1 Coloured Plate. Price 22s. 6d.

BRAIN SURGERY

- BAILEY—Intracranial Tumours.** By Percival Bailey, Professor of Surgery, University of Chicago. Pp. xiv + 476. Figs. 155. Price 27s.
- CUSHING—Papers Relating to the Pituitary Body, Hypothalamus and Parasympathetic Nervous System.** By Harvey Cushing. Pp. 234. Figs. 99. Price 22s. 6d.
- CUSHING—Intracranial Tumours.** By Harvey Cushing. Pp. xii + 150. Figs. 111. Price 22s. 6d.
- DANDY—Benign Tumours in the Third Ventricle of the Brain.** By Walter E. Dandy, M.D. Pp. 171. Figs. 120. Price 22s. 6d.

CHEMISTRY

- Aids to Chemistry.** By W. Partridge, F.I.C. Fifth Edition. Pp. 288. Price 5s.
- Aids to Organic Chemistry.** By S. F. Smith. Pp. 122. Price 3s.
- Aids to Qualitative Inorganic Analysis.** By R. G. Austin, B.Sc., A.I.C., F.R.M.S. Pp. 214. Figs. 7. Price 3s. 6d.
- FISCHER The Lyophilic Colloids.** By Martin H. Fischer. Pp. 254. Figs. 84. Price 20s.
- MUTER—Short Manual of Analytical Chemistry.** By J. Muter, Ph.D. Eleventh Edition, Edited by J. Thomas, B.Sc. Pp. 252. Figs. 56. Price 8s. 6d.
- STARCK—Volumetric Analysis.** By H. P. Starck, M.A. (Cantab.). Pp. 236. Figs. 11, and Appendix. Qualitative Analysis Tables, pp. 32. Price 7s. 6d.

CHILDREN

- Aids to the Diagnosis and Treatment of Diseases of Children.** By J. McCaw, M.D. Sixth Edition. Pp. 338. Price 4s. 6d.
- Aids to Orthopaedic Surgery.** By E. A. Crook, M.Ch., F.R.C.S. Pp. 240. Figs. 17. Price 3s. 6d.
- ALLEN—Handbook of Diseases of Infants and Children: For Students and Practitioners.** By F. M. B. Allen, M.D. Pp. 604. Price 15s.
- CHAPIN and ROYSTER—Paediatrics.** By H. D. Chapin, M.A., M.D., and L. T. Royster, M.D. Seventh Edition, Pp. 776. Figs. 184. 10 Coloured Plates. Price 35s.
- SPAAR—Care of Children in the Tropics in Health and Disease.** By E. C. Spaar, M.D. Pp. 280. Price 7s. 6d.

DIABETES

- McKITTRICK and ROOT—Diabetic Surgery.** By L. S. McKittrick, M.D., and H. F. Root, M.D. Pp. 270. Figs. 79. 2 Coloured Plates. Price 21s.
- WARREN—Pathology of Diabetes Mellitus.** By S. Warren, M.D. Pp. 223. Figs. 83. 2 Coloured Plates. Price 21s.

DERMATOLOGY

- Aids to Dermatology and Venereal Disease.** By R. M. B. Mackenna, M.A., M.B. Pp. 244. Figs. 7. Price 3s. 6d.
- ARTZ and FUHS—Rontgen Rays in Dermatology.** A Handbook for Practitioners and Students. Pp. 216. Figs. 57. 5 Coloured Plates. Price 18s.
- JACOBSON—Fungous Diseases.** By H. B. Jacobson, M.D. Pp. 336. Figs. 153. Price 25s.
- GILCHRIST—Outlines of Common Skin Disease.** By T. C. Gilchrist, M.D. Pp. 53. Figs. 60. 5 Plates. Price 7s.
- MACKENNA—Diseases of Skin.** By R. M. B. Mackenna, M.A., M.B. Third Edition. Pp. 464. Figs. 143. 36 Coloured Plates. Price 20s.
- PUSEY—The History of Dermatology.** By W. Allen Pusey, M.A. Pp. 242. Price 13s. 6d.

DIAGNOSIS

- Aids to Case-Taking.** By H. L. McKisack, M.D. Pp. 176. Price 4s. 6d.
- Aids to Medical Diagnosis.** By A. J. Whiting, M.D. Fourth Edition. Pp. 188. Figs. 16. Price 3s. 6d.
- Aids to Surgical Diagnosis.** By C. P. G. Wakeley, F.R.C.S. Pp. 176. Figs. 6. Price 3s. 6d.
- BASS and JOHN—Practical Clinical Laboratory Diagnosis.** By C. C. Bass, M.D., and F. M. Johns, M.D. Third Edition. Pp. 206. Figs. 134. Plates 20. Price 34s.
- CABOT—Physical Diagnosis.** By R. C. Cabot, M.D. Eleventh Edition. Pp. 552. Figs. 279. 6 Plates. Price 22s. 6d.
- FRENCH and NUTTALL—Medical Laboratory Methods and Tests.** By H. French, M.A., M.D., and E. E. T. Nuthall, M.D. Fourth Edition. Pp. 256. Figs. 62. 3 Coloured Plates. Price 7s. 6d.
- ROWE—Differential Diagnosis of Endocrine Disorders.** By Allan Winter Rowe. Pp. 230. Price 18s.

DICTIONARIES

- ARTSCHWAGER—Dictionary of Biological Equivalents.** German—English. By E. Artschwager. Pp. 240. 6 Plates. Price 20s.
- Baillière's Nurses' Complete Medical Dictionary.** Fourth Edition. Revised by C. F. Marshall, M.D. Pp. 200. Price 2s. 6d.
- PARTRIDGE—Dictionary of Bacteriological Equivalents.** By W. Partridge, F.I.C. Pp. 154. Price 10s. 6d.
- STEDMAN—Practical Medical Dictionary of Words used in Medicine.** By T. L. Stedman, M.A., M.D. Twelfth Edition. Pp. 1268. Figs. 400. 22 Plates. Price 35s.

EAR, NOSE AND THROAT

- LAKE and PETERS—Handbook of Diseases of the Ear.** By R. Lake, F.R.C.S., and E. A. Peters, M.D., F.R.C.S. Fifth Edition. Pp. 320. Figs. 128. 4 Coloured Plates. Price 12s. 6d.
- LAMB—Practical Guide to the Diseases of the Throat, Nose and Ear: For Senior Students and Junior Practitioners.** By W. Lamb, M.D. Fifth Edition. Pp. 466. Figs. 73. X-Ray Supplement of 31 Plates. Price 12s. 6d.
- POLITZER—Textbook of Diseases of the Ear.** By A. Politzer, M.D. Sixth Edition. Pp. 790. Figs. 202. Price 31s. 6d.
- STUART-LOW—Care of the Nose, Throat and Ear.** By W. Stuart-Low, F.R.C.S. Third Edition. Pp. 108. Figs. 18. Price 5s.

GENITO-URINARY

- HEITZMANN—Urinary Analysis and Diagnosis by Microscopical and Chemical Examination.** By L. Heitzmann, M.D. Sixth Edition. Pp. 407. Figs. 131. Price 22s. 6d.
- IRWIN—Urinary Surgery: A Handbook for the General Practitioner.** By W. K. Irwin, M.D. Second Edition. Pp. 280. Price 10s. 6d.
- LE COMTE—Manual of Urology.** By R. M. Le Comte, M.D., F.A.C.S. Pp. 323. Figs. 46. Price 18s.

LEWIS—Urology in Women. By E. Catherine Lewis, M.S. Lond. Pp. 84.
10 Figs. in the text. 4 Coloured, 11 Plain Plates. Price 6s.

RANDALL—Surgical Pathology of Prostatic Obstructions. By A. Randall,
M.A., M.D. Pp. 282. 78 Plates. Price 31s. 6d.

GYNAECOLOGY AND OBSTETRICS

Aids to Gynaecology. By R. E. Tottenham, M.D. 7th Ed. Pp. 140. Price 3s. 6d.

Aids to Obstetrics. By Leslie Williams, M.D., M.S. (Lond.), F.R.C.S. (Eng.),
M.C.O.G. 10th. Ed. Pp. 227. Price 3s. 6d.

BLAIR-BELL—Principles of Gynaecology. By W. Blair-Bell, B.S., M.D.,
F.R.C.S., F.C.O.G. Fourth Edition. Pp. 861. Figs. 506. 11 Coloured Plates.
Price 35s.

BERKELEY and DUPUY—Pictorial Midwifery. By C. Berkeley, M.A.,
M.C., M.D., illustrated by G. M. Dupuy, M.D. Second Edition. Pp. 168.
Figs. 256. 2 Plates. Price 7s. 6d.

DE GARIS—Theory of Obstetrics. By M. De Garis, M.D. Price 12s. 6d.

FRANK—The Female Sex Hormone. By R. T. Frank, M.D. Pp. 335.
Figs. 86. Price 25s.

JELLETT and MADILL—Manual of Midwifery. By H. Jellett, B.A., M.D.,
and D. G. Madill, B.A., M.B. Fourth Edition. Pp. 1294. Figs. 570.
20 Plates. Price 55s.

KERR—Operative Midwifery. By J. Munro Kerr, M.D. Third Edition.
Pp. 742. Figs. 308. Price 30s.

SOLOMONS—Handbook of Gynaecology. By B. Solomons, B.A., M.D.,
F.C.O.G. Third Edition. Pp. 380. Figs. 250. 2 Coloured Plates. Price 15s.

HEART

(See also under Blood)

BOAS—The Heart Rate. By E. P. Boas, M.D., and E. F. Goldschmidt.
Pp. 178. Figs. 65. Price 16s.

GÉRAUDEL—Mechanism of the Heart and its Anomalies. By E. Géraudel.
Pp. 286. Figs. 200. Price 25s.

GROLLMAN—The Cardiac Output of Man in Health and Disease. By
A. Grollman. Pp. 340. Figs. 14. Price 18s.

HYGIENE AND PUBLIC HEALTH

Aids to Mathematics of Hygiene. By R. B. Ferguson, M.A., M.D. Fifth
Edition (reprinted). Pp. 198. Price 3s. 6d.

Aids to Public Health. By W. G. A. Robertson, M.D., D.Sc. Third Edition.
Pp. 215. Price 3s. 6d.

Aids to Sanitary Science and Law. By C. F. White, M.B., D.P.H., and
H. Willoughby, M.R.C.S., F.R.C.P. Second Edition. Pp. 330. Price 4s. 6d.

HOPE—Textbook of Industrial Hygiene and Medicine. By E. W. Hope,
O.B.E., M.D., in collaboration with W. Hanna, M.A., M.D., D.P.H., and
C. O. Stallybrass, M.D., D.P.H. Pp. 774. Figs. 123. Price 25s.

**KIRK—Hints on Equipment and Health for Intending Residents in the
Tropics.** By J. Balfour Kirk, M.B. Second Edition. Pp. 133. Price 3s. 6d.

NEWSHOLME—Evolution of Preventive Medicine. By Sir A. Newsholme,
K.C.B., M.D., F.R.C.P. Pp. 242. 6 Plates. Price 13s. 6d.

NEWSHOLME—Story of Modern Preventive Medicine. By Sir A.
Newsholme, K.C.B., M.D., F.R.C.P. Pp. 308. Figs. 10. Price 18s.

OSTERTAG—Textbook of Meat Inspection. By Robert V. Ostertag.
Translation by C. F. Marshall, M.Sc., M.D., F.R.C.S. Pp. 755. Figs. 247.
10 Coloured Plates. Price 45s.

MEDICINE*(See also under Treatment)*

- Aids to Medicine.** By J. L. Livingstone, M.D., M.R.C.P. Fourth Edition. Pp. 424. Price 5s.
- COBB—The Organs of Internal Secretion, their Diseases and Therapeutic Application.** By Ivo Geikie Cobb, M.D. Fourth Edition. Pp. 328. Price 10s. 6d.
- COBURN—Factor of Infection in the Rheumatic State.** By A. F. Coburn, M.D. Pp. 298. Figs. 45. 4 Coloured Plates. Price 27s.
- ENGELBACH—Endocrine Medicine.** By William Engelbach, M.D. In Four Volumes. Pp. 1,800. Figs. 933. Price 67s. 6d.
- FRENCH and NUTHALL—Medical Laboratory Methods and Tests.** By H. French, M.A., M.D., and E. E. T. Nuthall, M.D., B.S. Fourth Edition. Pp. 256. Figs. 62. 3 Coloured Plates. Price 7s. 6d.
- GILLESPIE—Sleep and the Treatment of its Disorders.** By R. D. Gillespie, M.D. Pp. 276. 5 Figs. Price 7s. 6d.
- GORDON—Systemic Infections, Their Diagnosis and Treatment.** By A. Knyvett Gordon, M.B., B.C. Pp. 186. Figs. 7. Price 10s. 6d.
- HULL—Diseases Transmitted from Animals to Man.** By T. G. Hull. Pp. 364. Figs. 21. Price 25s.
- LIPSCOMB—Diseases of Old Age.** By F. M. Lipscomb, M.R.C.P. Pp. 480. Price 12s. 6d.
- STRÜMPPELL and SEYFARTH—Practice of Medicine.** By C. Seyfarth, M.D., Ph.D. Translated from 30th German Edition. In 3 vols. Pp. 2379. Figs. 392. 17 Plates. Price 105s.
- WILLIAMS—Minor Maladies and Their Treatment.** By Leonard Williams, M.D., M.R.C.P. Sixth Edition. Pp. 432. Price 10s. 6d.
- WYATT—Chronic Arthritis and Fibrositis, Diagnosis and Treatment.** By Bernard Langdon Wyatt, M.D. Pp. 214. Figs. 17. Price 16s.

MEDICINE (Miscellaneous)

- BLAIR-BELL—Some Aspects of the Cancer Problem.** By W. Blair-Bell, B.S., M.D., F.R.C.S. Pp. 558. Figs. 37. 94 Plates. Price 63s.
- GILL—Genesis of Epidemics.** By C. A. Gill, M.R.C.S., D.P.H. Pp. 576. Price 21s.
- MACLAREN—Medical Insurance Examinations.** By J. P. Maclaren, M.A., M.B. Second Edition. Pp. 662. Price 30s.
- SPENCER—Lightning, Lightning-Stroke and its Treatment.** By H. A. Spencer, M.R.C.S. Pp. 102. Price 5s.

MENTAL DISEASES AND INSANITY

- Aids to Psychiatry.** By W. S. Dawson, M.A., M.D. Third Edition. Pp. 328. Price 4s. 6d.
- CLARK—Nature and Treatment of Amentia.** By L. Pierce Clark, M.D. Pp. 322. Price 12s. 6d.
- EWEN—A Handbook of Psychiatry.** By John H. Ewen, M.R.C.P., D.P.M. Pp. 275. Price 12s. 6d.
- Handbook for Mental Nurses.** Seventh Edition. Pp. 629. Figs. 90. Price 6s.
- Manual for Mental Deficiency Nurses.** Pp. 532. Figs. 61. Price 6s.
- HENRY—Essentials of Psychiatry.** By G. W. Henry. Pp. 318. Price 18s.
- HINSIE—Treatment of Schizophrenia.** By L. E. Hinsie, M.D. Pp. 224. Price 13s. 6d.
- NICOLE—Psychopathology: A Survey of Modern Approaches.** By J. E. Nicole, L.M.S.S.A., D.P.M. Pp. 284. Second Edition. Price 12s. 6d.

- SHERLOCK—Minds in Arrear.** By E. B. Sherlock, M.D. Pp. 190. Price 6s.
PEARN—Mental Nursing (Simplified). By D. P. Napier Pearn, M.R.C.S.,
 L.R.C.P. Pp. 312. Figs. 22. Price 5s.
TREDGOLD—Mental Deficiency (Amentia). By A. F. Tredgold, M.D.,
 F.R.C.P. Fifth Edition. Pp. 552. 36 Plates. Price 25s.
YOUNGER—Insanity in Everyday Practice. By E. G. Younger, M.D. Fifth
 Edition. Pp. 146. Price 5s.

NERVES AND NERVOUS DISEASES

- Aids to Neurology.** By E. A. Blake Pritchard, M.A., M.D., B.Ch. (Camb.).
 Pp. 383. Figs. 42. Price 5s.
CADWALADER—Diseases of the Spinal Cord. By W. B. Cadwalader,
 M.D. Pp. 222. Figs. 72. Price 22s. 6d.
**FORD, CROWTHERS and PUTNAM—Birth Injuries of the Central
 Nervous System.** By F. R. Ford, M.D., B. Crowthers, and M. Putnam.
 Pp. 178. Figs. 57. Price 18s.
GASK and ROSS—The Surgery of the Sympathetic Nervous System. By
 George E. Gask, C.M.G., D.S.O., F.R.C.S. (Eng.), and J. Paterson Ross,
 M.S. (Lond.), F.R.C.S. (Eng.). Pp. 175. Figs. 43. Price 16s.
GLOBUS—Neuroanatomy. By J. H. Globus, B.S., M.D. Pp. 255. Figs. 36.
 Plates 52. Price 16s.
GRINKER—Neurology. By Roy R. Grinker, Pp. 988. Figs. 401. Price 38s.
HASSIN—Histopathology of the Peripheral and Central Nervous Systems.
 By George B. Hassin, M.D. Pp. 504. Figs. 229. Price 27s.
MUSKENS—Epilepsy. By L. J. J. Muskens, M.D. Pp. 450. Figs. 55.
 Price 27s. 6d.

OPHTHALMOLOGY

- Aids to Ophthalmology.** By N. Bishop Harman, M.A., F.R.C.S. Seventh
 Edition. Pp. 244. Figs. 115. Price 3s. 6d.
ALEXANDER—Ocular Dioptrics and Lenses. By G. F. Alexander, M.B.,
 C.M. (Edin.). Pp. 224. Figs. 62. Price 12s. 6d.
CLARKE—Refraction of the Eye and its Treatment. By E. Clarke, C.V.O.,
 M.D., F.R.C.S. Fifth Edition. Pp. 256. Figs. 95. Price 8s. 6d.
GRIMSDALE and BREWERTON—Ophthalmic Operations. By H. B.
 Grimsdale, B.A., M.B., F.R.C.S., and E. W. Brewerton, F.R.C.S. Pp. 446.
 Figs. 129. Price 21s.
MAY and WORTH—Manual of Diseases of the Eye: For Students and
 Practitioners. Seventh Edition. Revised by Montague Hine, M.D., F.R.C.S.
 Pp. 514. Figs. 351. 22 Coloured Plates. Price 15s.
RAMSAY—The Eye in General Medicine. By A. M. Ramsay, M.D. Second
 Edition. Pp. 265. Price 12s. 6d.
WORTH—Squint: Its Causes, Pathology and Treatment. By C. A. Worth,
 F.R.C.S. Sixth Edition. Pp. 256. Figs. 42. Price 10s. 6d.

PATHOLOGY

- Aids to Pathological Technique.** By D. H. Haler, M.B., B.S. (Hons., Lond.).
 Pp. 200. Figs. 18. Price 3s. 6d.
Aids to Pathology. By H. Campbell, M.D., F.R.C.P. Sixth Edition. Pp. 258.
 Figs. 17. Price 4s. 6d.
Aids to Practical Pathology. By F. W. W. Griffin, M.A., M.D. Pp. 260.
 Price 4s.
GREEN—Manual of Pathology. Fifteenth Edition. Revised by H. W. C. Vines,
 M.A., M.D. Pp. xii+928. Figs. 425. 8 Plates. Price 25s.
LONG—History of Pathology. By E. R. Long, Ph.D., M.D. Pp. 315.
 48 Plates. Price 22s. 6d.

LONG—*Selected Readings in Pathology from Hippocrates to Virchow.* Pp. 315. 25 Plates. Price 18s.

WARREN—*Pathology of Diabetes Mellitus.* By S. Warren, M.D. Pp. 223. Figs. 83. 2 Coloured Plates. Price 21s.

PHARMACY

Aids to Dispensing. By A. O. Bentley, Ph.C. Pp. 206. Figs. 7. Price 3s. 6d.

Aids to Materia Medica. By G. H. Newns, M.D. Pp. 124. Price 3s. 6d.

Aids to Mathematics of Pharmacy. By A. W. Lupton, M.C., Ph.C. Pp. 104. Price 3s.

Aids to Pharmaceutical Latin. By G. E. Trease, Ph.C. Pp. 176. Price 3s. 6d.

BENTLEY—*Textbook of Pharmaceutics.* By A. O. Bentley, Ph.C. Third Edition. Pp. 924. Figs. 234. Price 15s.

CARTER—*Business Building for Pharmacists.* By W. G. Carter, M.P.S. Pp. 208. Figs. 13. Price 6s.

GADD—*Synopsis of the British Pharmacopoeia, 1914,* and of the Poison Laws. By H. W. Gadd, F.C.S. Twelfth Edition. Pp. 205. Price 2s. 6d.

MUNCH—*Bioassays: A Handbook of Quantitative Pharmacology.* By J. C. Munch. Pp. 968. 239 Tables. Figs. 15. 3 Coloured Plates. Price 45s.

PENNETHORNE—*Guide to Assistant's Examination in Materia Medica and Pharmacy, Apothecaries' Hall.* By G. Pennethorne. Pp. 276. Price 6s.

WHITLA—*Pharmacy, Materia Medica and Therapeutics.* By Sir Wm. Whitla. Revised by J. A. Gunn, M.D., F.R.C.P., assisted by H. Berry, B.Sc., and J. Clifford Hoyle, M.D. Twelfth Edition. Pp. 655. Figs. 16. Price 12s. 6d.

PHYSIOLOGY

Aids to Physiology. By H. Dryerre, Ph.D., M.R.C.S. Pp. 264. Figs. 57. Price 3s. 6d.

BARCLAY—*Elementary Anatomy and Physiology for Nurses.* By H. C. Barclay, M.D. Third Edition. Pp. 422. 86 Plates. Price 12s.

BROWN—*Physiological Principles in Treatment.* By W. Langdon Brown, M.A., M.D., F.R.C.P. Sixth Edition. Pp. 474. Price 10s. 6d.

DRINKER and FIELD—*Lymphatics, Lymph and Tissue Fluid.* By Cecil K. Drinker, B.S., and Madeleine E. Field, B.A. Pp. 269. Figs. 14. Price 13s. 6d.

FEARING—*Reflex Action.* By F. Fearing, Ph.D. Pp. 364. Figs. 7. 8 Plates. Price 30s.

FULTON—*Muscular Contraction and the Reflex Control of Movement.* By J. F. Fulton, M.A., Ph.D., B.Sc. Pp. 660. Figs. 216. Price 11s. 6d.

FULTON—*Selected Readings in the History of Physiology.* By J. Fulton, M.A., Ph.D., B.Sc. Pp. 338. Figs. 60. Price 22s. 6d.

PHYSIOTHERAPY AND MASSAGE

BJÖRKEGREN—*Handbook of Anatomy of Students of Massage.* By M. E. Björkegren, M.R.C.S. Second Edition. Pp. 243. Figs. 73. Price 6s.

CLAYTON—*Physio-Therapy in General Practice,* and for the Use of Masseuses. By E. B. Clayton, M.B. Second Edition. Pp. 241. Figs. 53. Price 12s. 6d.

PALMER—*Lessons on Massage.* By M. D. Palmer. Sixth Edition. Pp. 328. Figs. 71. Price 5s.

PHELPS—*Diagnosis and Treatment of Postural Defects.* By W. M. Phelps. Pp. 189. Figs. 108. Price 18s.

PROTOZOOLOGY, PARASITOLOGY, ETC.

- BAYLIS—Helminthology: Medical and Veterinary.** By H. A. Baylis, M.A., D.Sc. Pp. 316. Figs. 200. Price 30s.
- CALKINS—Biology of Protozoa.** By G. N. Calkins, Ph.D. Second Edition. Pp. 634. Figs. 238. Price 37s. 6d.
- CHOPRA and CHANDLER—Anthelmintics and their Uses in Medical and Veterinary Practice.** By R. N. Chopra, M.A., M.D., and A. C. Chandler, M.Sc., Ph.D. Pp. 304. 5 Plates. Price 22s. 6d.
- EWING—Manual of External Parasites.** By H. E. Ewing. Pp. 231. Figs. 96. Price 20s.
- KUDO—Handbook of Protozoology.** By R. R. Kudo, D.Sc. Pp. 462. Figs. 175. Price 25s.
- MacGREGOR—Mosquito Surveys.** By M. MacGregor. Pp. 282. With 59 Maps. Price 15s.
- MATHESON—Handbook of Mosquitoes of North America.** By R. Matheson. Pp. 286. Figs. 23. 25 Plates. Price 25s.
- MATHESON—Medical Entomology.** By R. Matheson, Ph.D. Pp. 503. Figs. 211. Price 22s. 6d.
- MÖNNIG—Veterinary Helminthology and Entomology.** By H. O. Mönnig, B.Sc., Dr.Phil., B.V.A. Pp. 420. Figs. 264. Price 30s.
- THOMSON and ROBERTSON—Protozoology: A Manual for Medical Men.** By J. G. Thomson, M.A., M.B., and A. Robertson, M.B. Pp. 392. Figs. 214. 4 Coloured Plates. Price 30s.
- WENYON—Protozoology.** A Manual for Medical Men, Veterinarians and Zoologists. By C. M. Wenyon, C.M.G., C.B.E., M.B. 2 vols. Pp. 1590. Figs. 565. 20 Coloured Plates. Price 84s.

PSYCHO-ANALYSIS

- GLOVER—Technique of Psycho-Analysis.** By E. Glover, M.D. Pp. 150. Price 6s.
- JONES—Papers on Psycho-Analysis.** By E. Jones, M.D., M.R.C.P. Third Edition. Pp. 742. Price 25s.
- JUNG—Collected Papers on Analytical Psychology.** By C. G. Jung, M.D., LL.D. Second Edition. Pp. 520. Figs. 15. Price 18s.
- JUNG—Two Essays on Analytical Psychology.** By C. G. Jung, M.D., LL.D. Pp. 298. Price 10s. 6d.
- LONG—Collected Essays on the Psychology of Phantasy.** By C. E. Long, M.D. Pp. 228. Price 10s. 6d.
- NICOLE—Psychopathology.** A Survey of Modern Approaches. By J. E. Nicole, D.P.M. Second Edition. Pp. 300. Price 12s. 6d.
- RICKMAN—Development of the Psycho-Analytical Theory of the Psychoses, 1893-1926.** By J. Rickman, M.A., M.D. Pp. 114. Price, paper, 6s.

RÖNTGENOLOGY

- ARTZ and FUHS—Röntgen Rays in Dermatology.** By L. Artz, M.D., and H. Fuhs, M.D. Pp. 216. Figs. 57. 5 Coloured Plates. Price 18s.
- GLASSER—The Science of Radiology.** By Otto Glasser, Ph.D. Pp. 463. Figs. 108. Price 25s.
- KÖHLER—Röntgenology.** By A. Köhler. Translated from Fifth German Edition. Pp. 574. Figs. 324. Price 42s.

SERUM THERAPY

- BESREDKA—Immunity in Infectious Diseases.** By A. Besredka, M.D.
 Pp. 372. Price 18s.
- BURROWS—Some Factors in the Localization of Disease in the Body.** By
 Harold Burrows, C.B.E., F.R.C.S. Pp. 312. Figs. 6. 8 Plates. Price 15s.
- HARROW and SHERWIN—Chemistry of the Hormones.** By Benjamin
 Harrow, Ph.D. and Carl P. Sherwin, D.Sc., M.D. Pp. 234. Price 11s. 6d.
- KOLMER—Serum Diagnosis by Complement Fixation, with Special
 Reference to Syphilis.** By J. A. Kolmer, M.S., M.D. Pp. 603. Figs. 65.
 Price 31s. 6d.

SEXOLOGY AND BIRTH CONTROL

- ALLEN—Sex and Internal Secretions.** By E. Allen, M.D. Pp. 973. Figs. 16.
 Price 45s.
- ANOMALY—The Invert and his Social Adjustment.** Pp. 192. Price 5s.
- COKKINIS—Reproduction of Life.** By A. J. Cokkinis, B.A., M.B. Pp. 304.
 Figs. 108. 3 Plates. Price 5s.
- DICKINSON and BEAM—The Single Woman.** A Medical Study in Sex
 Education. By Robert L. Dickinson, M.D., and L. Beam, M.D. Pp. 488.
 Price 21s.
- DICKINSON and BEAM—A Thousand Marriages.** By Robert L. Dickinson
 and Louis S. Beam, M.D. Pp. 508. Figs. 14. Price 21s.
- DICKINSON—Control of Conception.** Pp. 302. Figs. 72. Price 20s.
- DICKINSON—Human Sex Anatomy.** By Robert L. Dickinson, M.D.,
 F.A.C.S. Pp. 158. Figs. 175. Price 45s.
- KONIKOW—Physicians' Manual of Birth Control.** By A. F. Konikow,
 M.D. Pp. 260. Figs. 21. Price 12s. 6d.
- MEAGHER—Study of Masturbation and the Psycho-Sexual Life.** By J. F. W.
 Meagher, M.D. Second Edition. Pp. 130. Price 9s.
- MEAKER—Human Sterility, Causation, Diagnosis and Treatment.** By
 Samuel Raynor Meaker, M.D. Pp. 276. Price 18s.
- SANGER and STONE—The Practice of Contraception.** By Margaret
 Sanger and Hannah M. Stone, M.D. Pp. 334. Figs. 22. Price 18s.
- VON GRUBER—Hygiene of Sex.** By M. von Gruber, M.D. Second Edition.
 Pp. 186. Figs. 6. Price 7s.

STOMACH, INTESTINES AND RECTUM

- ARAFI—Modern Aspects of Gastro-Enterology.** By M. A. Arafi, M.R.C.P.
 Pp. 392. Figs. 79, including 32 Plates, of which 8 are coloured. Price 27s. 6d.
- BROWN and SAMPSON—Intestinal Tuberculosis; Its Importance, Diagnosis
 and Treatment.** By L. Brown, M.D., and H. L. Sampson. Second Edition.
 Pp. 390. Figs. 122. 2 Coloured Plates. Price 22s. 6d.
- CAWADIAS—Diseases of the Intestines.** By A. P. Cawadiaz, O.B.E., M.D.,
 M.R.C.P. Pp. 314. Figs. 62. Price 16s.
- LOCKHART-MUMMERY—Diseases of the Rectum and Colon and their
 Surgical Treatment.** By J. P. Lockhart-Mummery, F.R.C.S. Second
 Edition. *In the Press.*

SURGERY

- Aids to Operative Surgery.** By Cecil P. G. Wakeley, D.Sc., F.R.C.S., F.R.S.,
 Second Edition. Pp. 233. Figs. 3. Price 3s. 6d.
- Aids to Surgery.** By J. Cunningham, M.B., F.R.C.S., and C. A. Joll, M.S., F.R.C.S.
 Sixth Edition. Pp. 442. Price 4s. 6d.
- BAILEY—Intracranial Tumours.** By Percival Bailey, M.D. Pp. 490.
 Figs. 155. Price 27s.

- BURROWS—Pitfalls of Surgery.** By H. Burrows, C.B.E., M.B., F.R.C.S. Second Edition. Pp. 536. Price 12s. 6d.
- CUSHING—Intracranial Tumours.** By Harvey Cushing, M.D. Pp. 162. Figs. 111. Price 22s. 6d.
- CUSHING—The Pituitary Body, Hypothalamus and Para-sympathetic Nervous System.** Pp. 234. Figs. 99. Price 22s. 6d.
- CUSHING and BAILEY—Tumours Arising from the Blood-Vessels of the Brain.** By H. Cushing and P. Bailey. Pp. 229. Figs. 159. Price 34s.
- GASK and ROSS—The Surgery of the Sympathetic Nervous System.** By George E. Gask, C.M.G., D.S.O., F.R.C.S. (Eng.), and J. Paterson Ross, M.S. (Lond.), F.R.C.S. (Eng.). Pp. 175. Figs. 43. Price 16s.
- KANAVEL—Infections of the Hand.** By A. B. Kanavel, M.D. Sixth Edition. Pp. 568. Figs. 196. Price 30s.
- LOCKHART-MUMMERY—After-Treatment of Operations.** By J. P. Lockhart-Mummery, F.R.C.S. Fifth Edition. Pp. 292. Figs. 37. Price 7s. 6d.
- McKITTRICK and ROOT—Diabetic Surgery.** By L. S. McKittrick, M.D., and H. F. Root, M.D. Pp. 270. Figs. 79. 2 Coloured Plates. Price 21s.
- ROSE and CARLESS—Manual of Surgery.** Edited by C. P. G. Wakeley, F.R.C.S., and J. B. Hunter, F.R.C.S. Fourteenth Edition. Pp. 1547. Figs. 721. 16 Coloured Plates. In 1 or 2 volumes. Price 30s.
- STRAUB—Surgery of the Chest.** By George F. Straub, M.D. Pp. 494. Figs. 341. Price 47s. 6d.
- WHARTON—Minor Surgery, including Bandaging.** By H. R. Wharton, M.D. Ninth Edition. Pp. 648. Figs. 450. Price 18s.
- WHEELER—Handbook of Operative Surgery.** By Sir Wm. I. de C. Wheeler, B.A., M.D., F.R.C.S. Fourth Edition. Pp. 458. Figs. 298. Price 15s.
- WILSON—Fractures and Their Complications.** By G. E. Wilson, M.B., F.R.C.S. Pp. 424. Figs. 306. Price 36s.

TREATMENT

(See also under Medicine)

- Aids to Medical Treatment.** By J. T. Lewis, M.D., and T. H. Crozier, M.D. Pp. 252. Price 3s. 6d.
- BROWN—Physiological Principles in Treatment.** By W. Langdon Brown, M.A., M.D. Sixth Edition. Pp. 474. Price 10s. 6d.
- CAWADIAS—Modern Therapeutics of Internal Diseases.** By A. P. Cawadias, O.B.E., M.D. Pp. 168. Figs. 3. Price 10s. 6d.
- COKE—Colds and Hay Fever.** By Frank Coke, F.R.C.S. Pp. 158. Price 5s.
- HUNT—Chronic Indigestion, Differential Diagnosis and Treatment.** By Thomas C. Hunt, B.A. Pp. 349. Figs. 16. Price 12s. 6d.
- Modern Treatment in General Practice.** Edited by Cecil P. G. Wakeley, D.Sc., F.R.C.S., F.R.S.E. Pp. 434. 16 Plates. Price 10s. 6d.
- THORNHILL—Varicose Veins, Haemorrhoids, Varicocele, Hydrocele, and their Treatment by Injection.** By Ronald Thornhill, M.B. Pp. 126. Second Edition. Price 5s.
- WHITLA—Dictionary of Treatment:** Including Medical and Surgical Therapeutics. By Sir Wm. Whitla, M.A., M.D. Seventh Edition. Pp. 1108. Price 25s.

TROPICAL MEDICINE

- Aids to Tropical Hygiene.** By R. J. Blackham, C.M.G., C.I.E., D.S.O., M.D. Second Edition. Pp. 248. Price 4s. 6d.

- Aids to Tropical Medicine.** By G. E. Brooke, M.A. Third Edition. Pp. 238. Figs. 30. Price 4s. 6d.
- CARTER—Yellow Fever.** By H. R. Carter, M.D. Pp. 320. Price 22s. 6d.
- CASTELLANI and CHALMERS—Manual of Tropical Medicine.** By Sir A. Castellani, K.C.M.G., M.D., and A. J. Chalmers, M.D., F.R.C.S. Fourth Edition. *In the Press.*
- JEWELL and KAUNTZE—Handbook of Tropical Fevers.** Pp. 486. Figs. 90. Price 16s.
- KIRK—Hints on Equipment and Health for Intending Residents in the Tropics.** By J. B. Kirk, M.B. Second Edition, Pp. 140. Price 3s. 6d.
- MINETT and SEVERN—Practical Tropical Sanitation: A Pocket Book for Sanitary Inspectors in the Tropics.** By E. P. Minett, M.D., and A. G. M. Severn, M.A., M.D. Second Edition. Pp. 188. Figs. 55. Price 5s.
- PLATT—Home and Health in India and the Tropical Colonies.** By K. Platt, M.D. Pp. 226. Price 5s.
- SPAAR—Care of Children in the Tropics in Health and Disease.** By E. C. Spaar, M.D., B.S. Pp. 280. Price 7s. 6d.

TUBERCULOSIS

- BALDWIN—Tuberculosis Bacteriology, Pathology and Laboratory Diagnosis.** By E. R. Baldwin, M.D., S. A. Petroff, Ph.D., and Leroy S. Gardner, M.D. Pp. 358. Figs. 82. 4 Coloured Plates. Price 21s.
- BROWN and SAMPSON—Intestinal Tuberculosis: Its Importance, Diagnosis and Treatment.** By L. Brown, M.D., and H. L. Sampson. Second Edition. Pp. 390. Figs. 122. 2 Coloured Plates. Price 22s. 6d.
- MYERS—Tuberculosis Among Children.** By J. A. Myers, M.D., M.S., Ph.D. Pp. 228. Figs. 43. Price 16s.
- THOMSON and FORD—Tuberculosis of the Lungs: A Practical Guide for General Practitioners.** By H. H. Thomson, M.D., and A. P. Ford, M.R.C.S. Pp. 189. Figs. 30. Price 10s. 6d.
- TOMSON—Some Methods for the Prevention of Tuberculosis.** By W. B. Tomson, M.D. Pp. 162. 3 Plates. Price 6s.
- WELLS—The Chemistry of Tuberculosis.** By H. G. Wells, M.D. Second Edition. Pp. 496. Figs. 8. Price 31s. 6d.
- WILLIS—Laboratory Diagnosis and Experimental Methods in Tuberculosis.** By Henry Stuart Willis. Pp. 356. Figs. 25. Price 16s.

VENEREAL

- KAHN—Serum Diagnosis of Syphilis by Precipitation.** By R. L. Kahn, M.S., D.Sc. Pp. 249. Tables 56. Price 13s. 6d.
- KLINE—Microscopic Slide Precipitation Tests for the Diagnosis and Exclusion of Syphilis.** By B. S. Kline, B.A. Pp. 118. Figs. 28. Price 11s. 6d.
- KOLMER—Serum Diagnosis by Complement-Fixation, with Special Reference to Syphilis.** By J. A. Kolmer, M.S., M.D. Pp. 603. Figs. 65. Price 31s. 6d.
- LUYS—Textbook on Gonorrhoea and its Complications.** By G. Luys, M.D. Third Edition. Pp. 416. Figs. 212. 5 Coloured Plates. Price 21s.
- MARSHALL and FFRENCH—Syphilis and Venereal Disease for Students and Practitioners.** By C. F. Marshall, M.D., F.R.C.S., and E. G. Ffrench M.D., F.R.C.S. Fourth Edition. Pp. 445. Figs. 90. 7 Coloured Plates. Price 15s.
- MOORE—The Modern Treatment of Syphilis.** By Joseph Earle Moore, M.D. Pp. 535. Illustrated. Price 22s. 6d.
- STANNUS—A Sixth Venereal Disease.** By Hugh S. Stannus, M.D. Pp. xii + 270. Price 12s. 6d.

PERIODICALS

TITLE	ISSUES PER ANNUM	COST OF SUBSCRIPTION		
		£	s.	d.
Acta Rheumatologica - - -	Irregular	18	3	
American Journal of Roentgenology and Radium Therapy - - -	12	2	14	0
American Journal of Tropical Medicine -	6	1	5	0
American Journal of Clinical Pathology -	6	1	5	0
Annals of the Pickett-Thomson Research Laboratory - - -	1	3	3	0
Ars Medici - - -	12		16	0
British Journal of Inebriety - - -	4		10	6
British Journal of Tuberculosis - - -	4		10	6
Bulletin of the Johns Hopkins Hospital -	12	1	9	6
Child Development - - -	4	1	5	0
International Journal of Psycho-Analysis	4	1	10	0
Journal of Aviation Medicine - - -	4	1	5	0
Journal of Bacteriology - - -	12	2	10	0
Journal of Biological Chemistry - - -	12	5	0	0
Journal of Comparative Psychology - - -	6	1	17	6
Journal of Immunology - - -	12	2	5	6
Journal of Nutrition - - -	12	2	10	0
Journal of Medical Association of South Africa - - -	26	1	11	6
Journal of Pharmacology and Experi- mental Therapeutics - - -	12	3	15	0
Journal of Urology - - -	12	2	10	0
The Laryngoscope - - -	12	1	15	0
Medical Press and Circular - - -	52	1	1	0
Medicine - - -	4	1	5	0
Occupational Therapy and Rehabilitation	6	1	5	0
Personnel Journal - - -	6	1	5	0
Quarterly Review of Biology - - -	4	1	5	0
Social Forces - - -	4	1	0	6
South African Nursing Record - - -	12		10	6
Surgery, Gynaecology and Obstetrics	12	3	0	0

