UNIVERSAL
LIBRARY
OU_174155
UNIVERSAL
LIBRARY

A REVIEW OF WORK

ON

INDIAN MEDICINAL PLANTS

Indian Council of Medical Research

SPECIAL REPORT SERIES

No. 30



A REVIEW OF WORK
ON
INDIAN MEDICINAL PLANTS
(Including Indigenous Drugs & Poisonous Plants)

By

Bt.-Colonel R. N. CHOPRA

I. C. CHOPRA

NEW DELHI
1955

PRICE Rs 3/-

Printed at the Cambridge Printing Works, Kashmere Gate, Delhi

FOREWORD

The use of indigenous drugs for the alleviation of human suffering is of considerable economic importance to our country With this end in view the Indian Research Fund Association, now the Indian Council of Medical Research, has been sponsoring extensive researches in indigenous drugs for over two decades The pioneering work in this field was done by Colonel Ram Nath Chopra at the School of Tropical Medicine, Calcutta and it may be said that we owe to him and to his students such knowledge as we today possess about the indigenous drugs of India A review of the work done in this field was published by the Indian Research Fund Association in 1939. This review was greatly in demand. The Council therefore decided that it was essential to bring the subject up to date and requested Colonel Ram Nath Chopra to undertake the task. That he has done it very ably will be apparent from a perusal of the book. It gives in simple and non-technical language a brief account of the indigenous drugs so far investigated in India The review should therefore be useful not only to practitioners of the indigenous systems of medicine in the country but also to scientists and to the lay public as well It will indicate to the scientists which drugs require further investigation. The bibliography given at the end is a comprehensive one which, apart from its scientific value, will show the extent of interest which scientists both in India and abroad have taken in this subject

Much yet remains to be done and many more drugs will have to be investigated scientifically by the application of modern methods of botanical, chemical, pharmacological and clinical research. I would, however, like to congratulate Colonel Ram Nath Chopra and his collaborators for their painstaking labour and for putting this material in such a concise and interesting form.

AMRIT KAUR,

September 16, 1954. *Minister for Health, India.*

PREFACE

For many years past, the Indian Council of Medical Research, formerly Indian Research Fund Association, has given large grants to research workers all over India for the study of Indian Indigenous Drugs In fact it was this body which was chiefly responsible for initiating work on this subject of great economic importance to India. The senior author was one of the first recipients of these grants when he was Professor of Pharmacology at the Calcutta School of Tropical Medicine and Medical College, Calcutta As Professor at the former Institution, one of the duties laid down for him, was the study of Indian Indigenous Drugs Financial resources from the School itself for this work were very limited then and it was the generous grants given by the Indian Council of Medical Research, extending over a period of more than three decades, both at the School and later at the Drugs Research Laboratory, Kashmir which enabled him and his co-workers to carry out screening studies of a large number of indigenous drugs Considerable financial help was also received later from Indian Council of Agricultural Research and the Council of Scientific and Industrial Research

In 1938 the senior author was asked to write a review of the work done on Indigenous Drugs under the Council upto that time, in simple non-technical language and a brochure was published in 1939 Large demand came from the public for this pamphlet and the edition was soon exhausted In 1953 he was again requested to write a review of the work on Indian Indigenous Drugs, but this time it was not to be limited to work done under the Council but was also to include work done by other independent investigators A thorough study of the whole literature on the subject was, therefore, undertaken by the present authors It transpired that the work on Indigenous Drugs was being chiefly directed in the channel of investigation of Indian Medicinal Plants Hence the change in the title of the present edition

The general arrangement of the subject matter in the brochure of 1939 has been again adopted Brief summaries of the work done on various drugs have been given in "telegraphic language". A comprehensive bibliography has been given, arranged subject wise and with full titles of papers Lists of plants used in the treatment of various diseases in the indigenous systems of medicines have also been given after a careful analysis of the literature.

It is hoped that this review will be read by those interested in indigenous drugs and their practical use for alleviation of human ailments As the descriptions are given in simple non-technical language and common vernacular names have been included, the present volume will be of interest not only to research workers but also to the public in general. The bibliography of nearly two thousand publications on the subject, collection of which has involved immense labour, will we trust, be useful to research workers and others.

In writing the present volume we wish to gratefully acknowledge the assistance we have received at every stage from Mr K L Handa who has helped in collecting references from literature and has checked up the chemical data. Messrs. L D. Kapoor and B K Abrol have checked up botanical terms and synonomy and Dr Kartar Singh Pharmacological Assistant in Indigenous Drugs Enquiry has rendered valuable assistance

To the Governing Body of the Indian Council of Medical Research, its worthy President Rajkumari Amrit Kaur and its able Secretary Dr C G Pandit, the authors are grateful for the generous grants for work on Indigenous Drugs These grants are being continued even now and are enabling the authors to carry on the work at the Drug Research Laboratory

Finances were also kindly provided by the Council for the preparation of the manuscript and the printing of this volume.

Drug Research Laboratory,
Jammu April, 1954

R. N. CHOPRA
I C. CHOPRA

CONTENTS

PAGES

CHAPTER I

INTRODUCTION . Climate and topography Features of Indian Vegetation Relationship to other floras ... 1

CHAPTER II

MEDICINAL PLANTS OF INDIA Remedies used in Indigenous Medicine Collection of Vegetable drugs 3

CHAPTER III

INDIAN INDIGENOUS DRUGS Main lines of basic research. Indigenous Drugs of India Historical review ... 6

CHAPTER IV

BASIC REQUIREMENTS FOR STUDY OF INDIGENOUS DRUGS A Botanical Unit A Chemical Unit A Pharmacological Unit A Clinical Unit. 10

CHAPTER V

SCOPE OF WORK ON INDIGENOUS DRUGS To make India self-supporting To discover remedies from the claims of Indian Medicine How to effect economy Indian Pharmacopoeia. 13

CHAPTER VI

A RETROSPECT OF RESULTS ACHIEVED Value of negative results Summaries of work done on commonly used 183 drugs. 19

CHAPTER VII

I TOXICOLOGICAL ASPECTS OF INDIAN PLANTS Cryptogams (Flowerless Plants) Bacteria Algae Fungi—Parasitic on man and animals Those attacking foodstuffs Smuts Rusts Ergot Moulds. Mushrooms Poisonous and edible. Lichens Bryophyta. Pteridophyta. 52

CHAPTER VIII

II Toxicological Aspects of Phanerogams · (Flowering Plants) Plants poisonous to man and livestock Medicinal plants Food Poisons. List of 386 plants with a reference to their habitat, toxicological, insecticidal & piscicidal properties 59

CHAPTER IX

Medicinal Plants used in Indigenous Medicine Against Tuberculosis, Bowel Diseases (Dysentery & Cholera), Fevers Emenagogue & Abortificient Irritants producing dermatitis Insecticidal and Piscicidal plants ... 96

CHAPTER X

Bibliography 122

Supplement to the Bibliography 254

Appendix of Scientific Names 1

Appendix of Vernacular Names 39

A REVIEW OF WORK ON INDIAN MEDICINAL PLANTS

(including Indigenous Drugs and Poisonous Plants)

INTRODUCTION

The Indian Materia Medica, including the common household remedies at present includes about 2,000 different drugs Many of these are in actual use in different parts of India Of this number approximately two hundred are of mineral origin, about two hundred are of animal origin and the remainder are of vegetable origin The mineral drugs are mostly compounds whose chemical composition is generally known and whose actions are in most cases not difficult to understand The drugs of animal origin are a very varied assortment, few of which have been shown to have any rational usage The study of indigenous drugs, therefore, chiefly means a study of Indian medicinal and allied plants in which there is a vast and very rich field for investigation

Climate and topography The extreme variability that India presents in its meteorological and climatic conditions as also in its topographical features is perhaps unrivalled in the world The tremendous range of variation of temperature from 126°F in summer in desert areas to minus 46°F in winter in Dras in Kashmir, and the prevalence of arctic conditions throughout the year in some of the Himalayan ranges are well known The annual rainfall varies from 430 inches at Cherrapunji in the hills of Assam to less than 5 inches per annum in the desert areas of Rajputana The air is saturated with moisture in the coastal districts and in the hills during the south-west monsoon and there are periods of practically zero relative humidity in the dry weather These are some of the interesting contrasts presented by this vast country in which we live.

No less striking are its physical features It has the most massive and the loftiest range of mountains in the world, the Himalayas There are also lower hills and plateaus, extensive rich alluvial plains, sandy wastes and deserts, hills, streams, mighty rivers with their extensive deltaic and estuarine systems, numerous lakes, canals, ponds, tanks and extensive marshy tracts, sandy or rocky coasts spread over an area of more than a million and half square miles The country is in fact an epitome of almost all climates, seasons, and soils of the world Under these conditions it is not surprising that India possesses one of the richest and certainly the most varied flora of any area of similar size on the surface of the globe

Features of Indian Vegetation Roughly speaking, one-fifth of the Indian sub-continent is occupied by forests of different kinds There are ever-green forests occupying the west coast of the peninsula up to the ridges of the Western mountain chain, and the lower slopes of the

Note .— This work was initiated by the Indian Council of Medical Research formerly Indian Research Fund Association.

Eastern Himalayas, with their magnificent and majestic trees, the 'sky scrapers' many centuries old A large part of the peninsula is covered by the deciduous type of the forests which are very valuable to the country There are the dry forests of Rajputana and the Punjab which are also spread over a large area of Indian States There is very little vegetation of any kind in the Great Indian Desert itself

Many tidal creeks and backwaters along the coasts of India, chiefly the deltas of rivers, such as the Sunderbans are covered with another type of evergreen forest, the tidal or littoral, where many plants send up from their underground roots a number of aerial roots for respiration The massive chain of the Himalayas with its tropical, temperate and alpine flora presents unique features The eastern Himalayan flora differs remarkably from the western in several respects in addition to the disappearance of the tropical type as the extreme north-west is reached It is in this chain that the highest limit of vegetable life on earth has been recorded The tree limit in the Western Himalayas corresponds roughly with the snow-line, i e 12 000 ft. above sea level, above which a variety of beautifully coloured flowers flourish Aquatic or semi-aquatic types of vegetation are found in the tanks, jhils, ponds, water courses and swamps all over India, particularly in Bengal Lastly, there are areas in the plains, as well as in the hills, especially in the former, which have, through the influence of man in search of more and more agriculturally exploitable land, lost much of their original characters Another feature of the Indian vegetation is that whereas in one season of the year, vast areas in the plains of India are the scene of plenty for man and animals, in another they become a dreary brown sunburnt waste where herbivores starve by the thousands and where it passes human understanding how any creature dependent on plant life can survive the dry season

Relationship to other floras The Indian flora is closely related to that of some other countries, and it is interesting to remark that none of families of flowering plants is peculiar to this country There is a very great preponderance of those genera and species which are also found in the adjacent countries Malayan, Burmese and Chinese genera are specially well represented in Eastern India Next in order come the European and Middle East flora which are particularly strongly represented in Western India Of the typically European elements there are twice as many in the Western as in the Eastern Himalayas Nearly 600 European genera are represented in India, many of them by a single species, and the Middle Eastern element is certainly, as is only to be expected, no less prominent African, Australian and American elements follow these in decreasing proportions, the African flora being more noticeable in Western India Recently there has been an influx of American species with their remarkable tendency to spread Tibetan and Siberian floras only reach India in the alpine regions of the Himalayas, while Chinese and Japanese elements are strongly represented in its temperate belts All these factors tend to make the Indian flora very rich and cosmopolitan and full of possibilities of economic importance

CHAPTER II

MEDICINAL PLANTS OF INDIA

India abounds in all kinds of food plants, spices, perfumes, timber, fibres, gums, etc which have been known all over the world from ancient times There are more than about 700 important fodder plants including about 260 species of valuable fodder grasses It is not surprising, therefore, that plants containing active and medicinal principles also grow abundantly within its bounds More than 2,000 plants, out of a total of about 11,000 species found in India, are alleged to have medicinal properties of some description or other and have been enumerated in the literature of indigenous medicine Nearly three-fourths of the drugs mentioned in the British and other Pharmacopoeias grow here in a state of nature and others can be easily grown Indeed, this country is a veritable emporium of drugs The families to which the larger numbers of medicinal plants belong are Leguminosæ, Compositæ Labiatæ, Euphorbiaceæ, Rubiaceæ, Rosaceæ, Gramineæ, Liliaceæ, Rutaceæ, Ranunculacæ, Umbelliferæ, Cucurbitaceæ, Solanaceæ, Apocynaceæ and Malvaceæ

It stands to reason that all these 2,000 odd plants cannot possess the wonderful virtues attributed to them, but it is believed that there are some of these which might rightly deserve the reputation they have earned as cures In order to determine what these were and what medicinal properties they possessed the study of these plants was begun in the early part of the last century

The early studies were mainly confined to collection of available data from the literature of indigenous medicine as well as getting information about their popular use as household remedies by the people A rough survey with regard to the occurrence of these plants in different parts of the country was undertaken by various workers and in some cases chemical analysis of a preliminary nature was also carried out Modern medicine however, is intimately related to chemistry and detailed examinations of active principles of plants and other products form an essential part of it Experimental work on the pharmacological side to determine the action of these active principles needs laboratories well equipped with modern apparatus and appliances which were not available in those days Work on medicinal plants on modern scientific lines was thus started in the early part of this century with the main objective of making India self-sufficient and self-supporting by enabling her to utilize drugs produced in the country and by manufacturing them in a form suitable for administration There are a number of drugs of established therapeutic value which are in use in the pharmacopoeias of different countries Many of these grow wild and some in great profusion in different parts of India and a certain number are even cultivated If these resources could be developed and utilized and the finished products manufactured, treatment of many diseases could be brought within the means of the Indian masses whose economic condition is unfortunately of a very low order

A number of important drug plants extensively used by the medical profession, however, are neither found growing in a state of

nature nor have so far been cultivated in India. Cultivation of such plants is very important from an economic point of view, and scientific research in this direction, as is being carried out in other countries, would be very fruitful It is a matter of regret that India is still importing quantities of crude drugs in spite of the fact that practically every conceivable pharmacopoeial drug can be grown within her bounds. The history of the cultivation of cinchona, eucalyptus, digitalis, etc. in India clearly shows that the cultivation of medicinal plants is pregnant with rich possibilities if taken up on proper lines It is gratifying to note that in recent years a growing and increasing interest is now being taken in this direction

Remedies used in Indigenous Medicine. The second objective has been to discover remedies from the claims of Ayurvedic, Tibbi and other indigenous resources suitable for employment by exponents of Wetsern medicine This is only possible if these are shown to have effective active principles in them Since the period of decay and resuscitation of old systems of Indian medicine, knowledge of many of the effective remedies was lost while a number of uncertain ones crept in Belief in their efficacy originates in some cases from the teachings of the ancient commentators and is based on clinical data In others, however, there is no basis whatever The object of research initiated by the Indian Council of Medical Research (formerly Indian Research Fund Association) was to determine which of these plants had the therapeutic efficacy claimed and which had not Since this work was started about thirty years ago, several hundred medicinal plants have been partially screened and investigated by workers in this country Some of these have been shown to be of great practical utility and have been actually brought into use Many of the others were found to have effective medicinal properties and could be brought as cheaper substitutes for pharmacopoeial and Extra pharmacopoeial drugs A large number of others were found to be entirely useless. Much, however, remains to be done and hundreds of plants with alleged medicinal properties, remain to be investigated

Collection of drugs A few words may be said here with regard to the time of collection of drugs, for purposes of examination This should be done at the proper time when the plant matures and the active principles are at their maximum If this is not done even effective drugs may be found to have no activity During the last twenty five years several drugs of Indian origin have assumed considerable importance from the point of view of foreign trade Many firms of drug manufacturers in this country also use the locally produced raw materials for the manufacture of the finished products It is a matter of very great concern, therefore, that the crude drugs collected locally are often not up to the required standard and this has resulted in considerable economic loss Fortunately, this state of affairs is being gradually rectified It must be stated however, that collection of drugs is not so simple a matter as most people seem to imagine Collectors of medicinal drugs growing in a state of nature, and the present and prospective cultivators should bear in mind that there are certain factors which have to be considered in order to obtain standard medicinal products There is a good deal of variation in the active principles in the different parts of a plant and in different seasons in the same part of the plant Even the same part and at the same time of the year shows remarkable variations

in the contents of its active principles For example, the young and the old leaves of a plant and unopened and opened flowers differ materially despite the fact that they are collected from the same plant and during the same season.

The soil also has an important influence on the active principles of plants It has often been observed that plants collected at the proper time, when the active principles have reached maturity and are at their maximum, give very effective results while the same plants when collected under other conditions are utterly useless Conditions for the collection of drugs in the case of plants under cultivation are more favourable and strict control over various factors can be exercised with greater ease than is the case with plants growing in a state of nature. It has, however, to be admitted that ideal conditions for the collection of even many of the common and important medicinal plants are not known with precision and research is urgently needed to determine the time when the active principles are at their maximum under the environments existing in this country, as was done by the Dutch in Java in the case of cinchona If this could be systematically carried out, India could supply itself and other countries with medicinal 'herbs'

CHAPTER III

INDIAN INDIGENOUS DRUGS

Main lines of basic research Having given a general account of the potentialities of this country with regard to medicinal plants, we will now briefly describe some of the important lines of basic research taken up by workers in this connection The first of these is study of Indigenous Drugs of India

Indigenous Drugs of India It is desirable to point out at the outset that the term 'Indigenous Drugs' has been used, for the purposes of research in its widest sense so as to include within its scope not merely those drugs which were originally the natives of India, but also those which have been introduced from outside and have become completely naturalised Drugs which are cultivated in India, whether used in the indigenous system of medicine or in the pharmacopoeias of various western countries, have also been brought within the purview of this investigation The term also includes locally manufactured preparations of mineral and animal origin which are used in medicine as practised in India

Historical The study of Indian indigenous drugs was first begun in the early part of the last century and it was then confined chiefly to the collection of available information with regard to various medicinal plants growing in different parts of the country The earliest contributions were from the writings of Sir William Jones whose memoir entitled *"Botanical Observations on Select Plants"*, is well known This was followed in 1810 by John Fleming's *"Catalogue of Medicinal Plants"*, Ainslie's *"Materia Medica of Hindustan"* in 1813, and Roxburgh's *"Flora Indica"* in 1820 In 1844 O Shaughnessys published his *"Bengal Pharmacopoeia"* which was the first book of its kind which dealt exclusively with the properties and uses of the medicinal plants used in Bengal In 1868 a *"Pharmacopoeia of India"* was published under the able editorship of Waring It signalized a new epoch in establishing and recording the value of indigenous medicinal products on modern lines The more important drugs were officially recognised with a view to their eventual adoption in the British Pharmacopoeia As a large number of the drugs, especially those in local use by practitioners and as household remedies, were not included in this work, Mohideen Sheriff published his *"Supplement to the Pharmacopoeia"* in the year 1869 He is also the author of "Materia Medica of Madras" which was edited and published after his death by Hooper U C Dutt's translation of Sanskrit Materia Medica brought into prominence the drugs used in the old Hindu Medicine even now largely practised in India Then Fluckiger's and Hanbury's *"Pharmacographia and Materia Medica of Western India"* 1883 was published These were followed by the publication of that very comprehensive book on the Indian Medicinal plants the *"Pharmacographia Indica"* in 1885 under the joint editorship of Warden and Hooper This treatise contains a mass of information regarding the uses of the indigenous materia medica in the Eastern and Western medicine The most elaborate work of all is *"A Dictionary of the Economic Products of India"* published in 1895 by Sir George Watt, the Reporter on the Economic Products to the Government of India. This monumental work, compiled with the help of a large team of workers, refers

to all the previous work on medicinal plants and other plants of economic importance Its pages are full of information of every description regarding the use of different barks, roots, flowers, leaves and woods for different medicinal and other purposes Works published still later such as Kanai Lal Dey's "*Indigenous Drugs of India*" and Kirtikar and Basu's "*Indian Medicinal Plants*" are largely summaries and compilations from the above mentioned literature in a more systematised and elaborated form In the last named book, plates illustrating various important medicinal herbs are given which greatly help the reader in differentiating them from plants with which they are apt to be confused

The literature mentioned above is very valuable, as it contains not only information from Ayurvedic and Tibbi sources, but also gives the results of personal observations and experiences of some of the writers There is no doubt that a considerable amount of botanical investigation into the scientific names of many medicinal plants was accomplished More, however, remained to be done in the case of a large number of plants to clear up many points with respect to their exact botanical sources Many drugs that had escaped the previous investigators require to be explored in all their details

Admirable as all these attempts were, the pharmacology of most of the indigenous remedies remained an unexplored field till recent years The reason for this is not far to seek Investigations of this nature require a considerable outlay of money in the form of well-equipped chemical and pharmacological laboratories, while a liberal staff of competent chemists and pharmacologists is another essential prerequisite Medicine we have observed is now intimately related to chemistry, and the ultimate solution of most problems, whether physiological or biological, rests on some physical or chemical basis This is forcibly presented to us in the study of the action of drugs The importance of the cooperation of chemists at every stage of research work can only be realised by the workers themselves If satisfactory results have to be achieved and if the work is to be carried out on the same standard as in other civilised countries, the cooperation of competent chemists with experience is the first essential

It must be pointed out that the time and labour required to work out the chemical composition of a single drug are enormous This may be judged from the fact that it would take an experienced chemists several months, perhaps a year or more, to isolate in a pure state and roughly describe the nature of the different chemical constituents of a single crude drug The determination of the chemical constitution of the active principles concerned would take a considerably longer time. The isolation of a sufficient quantity of the active principles and the testing of them pharmacologically would occupy many months The magnitude of the task of working out all the drugs used in the indigenous systems of medicine in detail transcends all imagination There is such an enormous scope for research in this field, and so little has been done, that it is impossible for any one individual or any one institution to cope with it adequately. The cooperation and intimate association of a large number of sincere and devoted workers of ability is needed to find the truth This is now being gradually done. Chairs in Pharmacology have been founded by the various Universities and Medical Colleges and facilities for research work on modern scientific lines are now being made increasingly more available.

As the action of these drugs or their active principles can only be established by a careful chemical, pharmacological and clinical study, the investigation in all the three aspects should be carried on side by side. The experimental work on the pharmacological side can be done only in laboratories well equipped with all modern appliances The first laboratory of its kind established in this country was at the School of Tropical Medicine, Calcutta in 1921 One of the main duties of the Professor of Pharmacology laid down was investigation of the indigenous drugs on scientific lines The chemical department of this institution had a small team of chemists who worked out the chemical composition of drugs, isolated the active principles and handed them to the pharmacologist for determination of their action on the animal organism The clinical testing of the drug was made possible by the Carmichael Hospital for Tropical Diseases a research hospital attached to this institution In this way it was found possible to go through a number of drugs in all the varied phases of their investigation, i.e. from the isolation of their active principles to the testing of their action on animals and finally to the making of suitable preparations for trial on patients, and for recording the results of therapeutic trials.

The staff of the Departments of Pharmacology and Chemistry of this institution was only a nucleus to start this work It was wholly inadequate to deal with such an immense task in addition to its ordinary duties of post-graduate teaching Although a modest beginning was made in 1921 when the School started functioning, it was not really till 1926, when the Indian Research Fund Association, now known as the Indian Council of Medical Research, gave a grant to develop this work Investigations were then taken up in right earnest and on proper systematic lines. During the years that followed, the importance of this work was appreciated with greater emphasis and the grants were considerably increased It was realised by this body that this was the only effective reply to attempts at the wholesale revival of the indigenous system of medicine which were then in evidence In 1935 the Imperial Council of Agricultural Research (now Indian Council of Agricultural Research), appreciating the importance of basic research of this type which was being done gave a grant for the investigation of the closely allied group of Poisonous Plants and Food Poisons of India which are such a menace to man and live-stock in this country. By this grant the research facilities for work on indigenous drugs were also further augmented and certain deficiencies which existed in the technical staff were made good An efficient Botanical Unit was thus provided, the want of which had been very keenly felt in connection with the research work on medicinal plants

With all these facilities made available it was possible to undertake the study of a large number of drugs in all the varied phases of their investigation i e., from the isolation of their active principles to the testing of their action on animals and finally making suitable preparations for trials on patients and for recording results of therapeutic trials.

During the three decades that have followed, the research work on indigenous drugs has received considerable encouragement and has made satisfactory progress. The example of the Indian Council of Medical Research and Indian Council of Agricultural Research was followed by the Council of Scientific & Industrial Research. This Council

gave very generous grants to various Medical Institutions and other research bodies for this work. It also established in 1950, the Central Drug Research Institute at Lucknow as one of the eleven major National Laboratories of India One whole division of this great Institution is devoted entirely to the study of Indian Indigenous Drugs With the dawn of Independence, therefore, this research has been put on a sound and firm basis Much has been accomplished, in the study of Indian Medicinal Plants on systematic and scientific lines, though much more remains to be done. A brief review of this work will be found in the following pages.

It is obvious from what has been stated above that the Indian Council of Medical Research with its predecessor the Indian Research Fund Association were the pioneers in encouraging the study of Indian Indigenous Drugs on scientific lines It was the example of this body which stimulated interest on this important subject of vital importance to the country.

CHAPTER IV

BASIC REQUIREMENTS FOR STUDY OF INDIGENOUS DRUGS

We have said above that detailed study of Indigenous Drugs is not a one man job It needs close collaboration and association of scientific workers in different allied subjects The prerequisites for the work to be done properly on scientific lines are —

1. A Botanical Unit This Unit should consist of experienced botanists and technicians They help not only in the collection and identification of genuine drugs, but also in carrying out a survey of the medicinal and poisonous plants growing in different parts of the country The importance of this section must not be under-rated, as on the correct identification of the drugs depends the utility and authenticity of the whole work In the early stages, when expert botanical help was not available, many mistakes were made in connection with the identity of medicinal plants, thereby causing much confusion and waste of effort An extensive herbarium of these plants is being gradually built up by our botanists This is a unique collection of its kind in India, and is rendering the correct identification of medicinal plants for any trained worker an easy matter

2 A Chemical Unit This should consist of a band of expert research chemists with special experience in plant chemistry They should be trained for work in different aspects of plant chemistry involving extraction and identification of active principles, such as alkaloids, glucosides, essential oils, neutral principles, antibiotics etc The medicinal or poisonous plants sent by different forest officers or collected by the botanists, are first dealt with in this department and an attempt is made to determine the active constituents that may be contained in them The preliminary chemical examination, the isolation of the different constituents which may have pharmacological action, their purification and elementary analysis all require considerable time, patience and skill It may be stated here, that the isolation of the active principles of plant constituents is not an easy task of a routine nature, as may perhaps be imagined Each plant has its own peculiar problems and difficulties which have to be solved This can only be done by chemists with considerable practical experience and technical skill Besides being engaged in the isolation and identification of active principles of plants, the chemical section is also required to assist in chemical and biochemical assays of different drugs, in the preparation of galenicals, extracts, etc. for purposes of clinical trial in the hospitals The work involved, therefore, is of a varied and comprehensive character Efficient teams of workers, have now been brought into existence after many years of training, and the analytical work is being considerably facilitated as more experience is being gained

3 A Pharmacological Unit. This is composed, preferably of medical men especially trained in the technique of experimental pharmacology, who work under the guidance of a pharmacologist with large practical experience of this type of work They test the different preparations or the purified active constituents isolated by the chemists by properly planned animal experimentation They also carry out biological assays, test the toxicity of drugs and suggest suitable doses for therapeutic trials. Here also experience assists a great deal in assessing the

value of the reactions produced in animal tissues by drugs under investigation and their practical application to therapeutics in man

4 A Clinical Unit. This section carries out testing of drugs on patients in the out-patients department and in the hospital This work is carried out under the direct guidance of experienced pharmacologists and clinicians It has often happened that drugs have been tried in a haphazard fashion by medical practitioners and wrong conclusions have been drawn The proper clinical evaluation of these drugs needs a well equipped hospital with a specially trained house staff of medical men and nurses, in which patients can be kept and watched for weeks and months if necessary The clinician in charge of such investigation should also have experience of this type of work which needs considerable patience and sound judgment


The existence of such teams all working in close cooperation, makes it possible to carry out this type of work on proper systematic lines. Research work of this nature started in institutions or university laboratories is often cut short at an early stage as the workers have to depend on institutions such as Hospitals for trials Again some of the universities and colleges do not possess properly equipped laboratories for pharmacological studies The procurement of all these facilities is expensive and make this work difficult If such organisations as Indian Council of Medical Research, Indian Council of Agricultural Research and Council of Scientific & Industrial Research had not come forward with generous grants, it would not have been possible to make any headway

It will thus be seen that from the stage of empirical knowledge available of a crude drug to bring it to its rational usage in modern medicine, it must pass through the following stages —


(1) Botanical identification (2) Chemical examination and isolation of active principles (3) Determination of their pharmacological action and toxic effects (4) Chemotherapeutic studies in animals. (5) Clinical trials in man

In case of well-known drugs of established value in indigenous medicine, it may some time be convenient to start with clinical trials first and proceed to analysis and laboratory tests afterwards A schematic idea of the work is given below —

PROCEDURE I



PROCEDURE 2



In whatever way the investigation is carried out on an indigenous drug, the work is time-consuming and can only be done by team-work of several groups of scientists each expert in his own field. The work should be co-ordinated by a medical scientist of experience who can draw logical conclusions from the data obtained. Haphazard approaches to this work have led to wrong conclusions and much confusion.

CHAPTER V

SCOPE OF WORK ON INDIGENOUS DRUGS

Work on indigenous drugs on modern scientific lines was initiated by the senior author when he was appointed Professor of Pharmacology at the School of Tropical Medicine, Calcutta with the following main objects in view :

1. To make India self-supporting by enabling her to utilise the drugs produced in the country, by manufacturing them in a form suitable for administration
2. To discover remedies from the claims of Ayurvedic, Tibbi and other indigenous sources suitable to be employed by the exponents of Western medicine
3. To discover the means of effecting economy, so that these remedies might fall within the means of the great masses in India whose economic condition is very low.
4. To prepare an Indian Pharmacopoeia including indigenous remedies of value

(1) To make India self-supporting so far as drugs and their preparations are concerned

Pharmacopoeial and Allied Drugs The first proposition is likely to lead to great results, because a large number of drugs which grow in this country are known both to Eastern and Western medicine and the properties and actions in many cases are also known The research here has been diverted into two main channels Firstly, there are many drugs of established therapeutic value which are in use in the pharmacopoeias of different countries The majority of these grow wild and in abundance in many parts of India and a certain number are even cultivated Some of these are collected and exported, though only a small fraction of the quantity produced, to foreign countries and come back to us in the form of standardised pharmaceutical preparations and active principles in pure condition, probably at a price many times that of the original crude product. A host of others grow, mature and eventually die without being put to any practical use whatsoever. There are numerous examples which have been dealt with in detail elsewhere. A brief description of those investigated is given hereafter in this review with short notes with regard to the work done

Besides these, there are a number of pharmacopoeial drugs which are widely used by the medical profession, but which do not naturally grow in this country They thrive, however, when they are cultivated under proper conditions in suitable parts of the country. Examples of such drugs are numerous but a few of the important ones such as digitalis, ipecacuanha, eucalyptus, cinchona, jalap, etc., may be cited. They were introduced into India many years ago and are doing well. On account of the great demand for these drugs in medical practice, their production

in this country is of great economic importance. India, we have stated, possesses most wonderful variability so far as the temperature and general climatic conditions are concerned, and every conceivable drug ranging from those growing in the hotest tropical and damp climates to those growing in dry, temperate and very cold climates can be grown and acclimatised in some part or other. From the geological point of view also every grade of soil from alluvial deposits to hard rocky formation and sandy deserts are met with Professor Greenish of the London School of Pharmacy rightly said, "India owing to the remarkable variations she possesses of climate, altitude and soil, is in a position to produce successfully every variety of medicinal herb required by Europe".

It should be remembered, however, that the soil, the season and the gathering time are some of the important variable factors with plants, and it can hardly be expected that the amount of active constituents would be constant under all conditions In some cases the quality is good and constant, but in the majority of instances the percentage composition of active principles has to be determined by careful methods of chemical and biological assay, to show that these remedies, growing in a state of nature, are as good in quality as those required by the standards laid down in the Pharmacopoeias If they do not come up to the required standard, the best method of bringing them into general use by improving the quality of the active principles by suitable cultivation, in parts of the country where this can be done economically, has to be determined

The work done so far in connection with these two groups of plants, has been responsible for bringing into use a large number of crude drugs, which were formerly imported from foreign countries by the drug manufacturing industry, by showing that the active principles of many of the indigenous plants came up to the standards laid down by the pharmacopœia.

Secondly, a large number of plants grow in India which, though not exactly the same, have properties and actions similar to the imported and often expensive remedies and would form excellent substitutes Not infrequently there are some closely allied species which are pharmacologically just as active The properties of these plants are being worked out on scientific lines and they are being brought into use Some examples may be cited here *Colchicum luteum* growing on the slopes of the western temperate Himalayas has been shown to be an excellent substitute for the official *C autumnale*. *Scilla indica* grows extensively on the sea-coast and on the drier hills of the lower Himalayas and the Salt Range and is being used as a substitute for *S. maritima* *Ferula narthex*, growing in Kashmir, yields a gum resin resembling asafœtida The properties of *Picrasma quassioides* and *Gentiana kurroo* resemble those of *Picrasma excelsa* and *Gentiana lutea* which are official in the British Pharmacopoeia. Many other examples could be cited

In both these groups there is a large field for research and development and considerable work has been done in connection with recognition of their active ingredients, determination of their percentage composition and establishing their pharmacological action. Pharmaceutical preparations are now manufactured from these indigenous products to the immense advantage of the country

(2) To discover remedies from the claims of Ayurvedic, Tibbi, and other indigenous sources.

The Drugs used in the Indigenous Medicine. The second proposition of popularising and introducing new drugs to Western medicine is a more difficult one. It is believed that out of the very large number of drugs used in the indigenous systems of medicine for centuries past, and still in use, there must be some at least which might deserve the reputation they have earned as cures. Since the period of decay and recompilation of old Indian systems of medicine, many of the effective remedies have been lost and a number of uncertain ones have come in The result is that in the indigenous systems, at the present time, almost every plant and shrub growing in the country has ascribed to it some medicinal virtue These beliefs in some cases originate from the teachings of the ancient commentators and are based on clinical data, but in others have no foundation whatever Their introduction was empirical and often a drug was used simply because a single case appeared to have derived some benefit from it In this way remedies have multiplied without proof but by belief and as they hail from all parts of India, no one seems to have a correct notion about their uses and properties. The employment of a large number of them would thus appear to have been based on empirical evidence handed down from generation to generation A thorough and complete research into all these drugs would constitute the life long work of innumerable chemists, pharmacologists and physicians For practical purposes the method adopted by workers has been to make use of the experience of Vaids, Hakims and others, and to take up for investigation those drugs which have a great local reputation before touching the less known remedies Besides, many of these drugs have been clinically tried by medical men practising Western medicine, who have expressed their opinion regarding their efficacy, this has also been helpful in the selection of drugs to be investigated. In this way a large number of commonly used drugs have been scrutinised.

(3). How to effect economy and bring the treatment within the means of the masses.

The third proposition relates to the devising of expedients for effecting economy, so that these remedies may reach the masses This is only possible if the price of the drugs can be considerably reduced, for in a poor country like India, there are millions of people who cannot afford any kind of treatment, whether cheap or expensive, and have consequently to depend upon charitable medical relief institutions The cost of drugs is so heavy that most of these institutions, which have only a limited annual budget for drugs, are not able to cope with the demand for such common and essential drugs as quinine, castor oil, magnesia, etc., to say nothing of the expensive medicines which are sometimes necessary and even indispensable.

The only way in which drugs can be cheapened and brought within the means of the masses is to utilise the local resources and substitute the indigenous products for the more expensive imported preparations of Western medicine. This can be done by encouraging the production, collection and manufacture of the local materia medica by preparing pharmaceutical preparations in a systematic manner. By local production and substitution of equally potent drugs of Indian origin for the imported drugs, the cost of treatment can be considerably reduced.

We have already made a reference to some of these remedies and the possibilities of their development. Their active principles are now being isolated, and standardised preparations such as tinctures, extracts, powders, etc , are being prepared by a number of reliable manufacturers If this aspect is further developed it will be possible to affect large scale economy

It is a matter of satisfaction to note that a great deal has been done towards the development of a drug manufacturing industry in India, and towards encouraging the use of raw material produced in the country for the preparation of galenical and other preparations The price of preparations has already come down considerably during recent years and by further development of this industry it is hoped that there will be further reduction in prices The factors which were militating against the bringing down of the prices of medicinal preparations to the level of the low economic conditions of the masses were fully investigated by the Drugs Enquiry Committee (1930-31). These were firstly, the excise duty on alcohol used for bonafide medicinal purposes by the manufacturers, and secondly the high railway freight for purposes of transportation of drugs from one part of the country to another Attention is being given to all these and drug industry in India is being put on a sound basis

(4). INDIAN PHARMACOPOEIA.

The fourth proposition of preparing a Pharmacopoeia is by no means an easy one. A national Pharmacopoeia is primarily meant to meet the claims and satisfy the needs of a particular group of physicians at a particular time. There must exist, and there does exist, a great difference not only between the Pharmacopoeias of various countries, but also between various editions of the same Pharmacopoeia.

The object of a Pharmacopoeia is, in the words of the founders of the United States Pharmocopoeia, 1820, "to select from among substances which have medicinal power those the utility of which is most fully established and best understood, and to form from them preparations and compositions in which their powers may be exerted to the greatest advantage."

The modern Pharmacopoeia is above all a book of standards. Its fundamental object and scope is, "to provide standards for the drugs and medicines of therapeutic usefulness or pharmaceutic necessity sufficiently used in medical practice; to lay down tests for the identity quality, and purity, to insure, as far as possible, uniformity in physical properties and active constituents " In other words, usage, rational usage and scientific usage are the basis of judgement.

The evidence placed before the Drug Enquiry Committee 1930-31 showed that India should have an official publication which would record what she recognizes as a trustworthy and approved materia medica upon which can be established modern food and drug acts, poison laws, systems of taxation, and the modernization of legal medicine Her best scientific men should be enlisted to shoulder the responsibility of working out India's own standards and to develop the necessary analytical laboratories, biological institutes and pharmaceutical schools It involves not only the development of machinery for the administration of laws based upon official standards, it means also finding an authoritative standard

which every doctor and pharmacist in India will hail as a real guide in every-day work

A consideration of what is said above will show that although the preparation of an Indian Pharmacopoeia is very desireable, a very large amount of work must necessarily be done before such a work can be prepared. The Drug Enquiry Committee 1930-31 carefully considered the question of the compilation of an Indian Pharmacopoeia and thought that the utility of such a work for the country cannot be overrated.

The pharmacopoeia which is in view ought to include the therapeutically active substances and, to find admission to it, a drug must be of known composition, of definite pharmacological action, and of well-established therapeutic use, and fully investigated for its toxicity and necessary standard for determining a safe maximum dose, with a chemical or biological standard The large mass which do not satisfy this condition should be left out. Necessary tests have to be developed for the protection of doctor, pharmacist, and patient India ought to set a standard of strength and purity for the material which is to appear on her markets.

This work should be on the lines of the British and the United States Pharmacopoeias including only drugs of known composition, of definite pharmacological action, of well-established therapeutic properties, with known toxicity and the necessary standards of chemical and biological assay for determining the safe maximum doses

The work which is being done on the Indian Indigenous Drugs is exactly the type of work which will eventually lead to the preparation of an Indian Pharmacopoeia But it is obvious that much more will have to be done before such an authoritative publication can be prepared. The Drug Control Laboratory at Calcutta and later the Central Drug Research Institute, Lucknow have helped greatly towards evolving standards suitable for this country A large number of drugs commonly used have been fully worked out, and the conditions above stated are being satisfied with regard to them

It is a matter of great satisfaction to note that work on these drugs has sufficiently advanced to lay the foundation of an *Indian Pharmacopoeia* As a first step towards it an *Indian Pharmacopoeia List* was prepared by the Committee set up by the Government of India under the auspices of Drugs Technical Advisory Board in 1947 This was published in 1949 and is an authoritative document laying down standards for Indian climatic and other conditions for drugs sufficiently worked up to be included in the List The Health Ministry then set up a Permanent Pharmacopoeia Committee This Committee has been at work since and with the help of number of sub-committees composed of experts from all parts of India has now practically completed the labour of compiling the first edition of Indian Pharmacopoeia which will be published in the near future The pioneering work done in connection with the preparation of Indian Pharmacopoeia by the Indian Council of Medical Research (formerly Indian Research Fund Association) cannot be overrated. As early as 1926 this body appreciated the importance of the study of Indigenous Drugs on scientific lines as a preliminary to compilation of an Indian Pharmacopoeia and started financing a number of investigations in this

connection with the object of working out of the indigenous drugs on scientific lines .

Recently an *Indian Pharmaceutical Codex* has been published under the auspices of Council of Scientific & Industrial Research by Dr B Mukerji, Director, Central Drug Research Institute, Lucknow, which gives about 200 monographs and a similar number of formulary giving pharmacognostic, chemical, pharmcological and therapeutic data on indigenous drugs with their preparation and dosage for the medical and pharmaceutical professions.

CHAPTER VI

A RETROSPECT OF RESULTS ACHIEVED

During the period of three or four decades since this work was started, much has been accomplished though only the fringe of this vast problem has yet been touched A number of important medicinal plants commonly used by the Kabirajes, Hakims and as popular household remedies by the people have been investigated on modern scientific lines The chemical composition of many plants has been determined, the pharmacological action of the active principles of a large number has been worked out by animal experimentation, and finally suitable preparations made from a number of drugs have been tested on patients in properly equipped hospitals It is only by such thorough investigations that the real merits of these drugs can be proved and a demand created for them not only in India but in other parts of the world This laborious work has brought into prominence the merits and qualities of certain drugs and it has been shown that they may prove to be valuable additions to the armamentarium of the medical man to relieve the sufferings of humanity, if brought into general use Such drugs unfortunately are not many A few examples may be cited here *Holarrhena antidysenterica* (Kurchi) has proved useful in the treatment of amoebic dysentery especially the chronic form, *Rauwolfia serpentina* has proved to be an effective sedative which brings down the blood pressure particularly the diastolic This plant has attracted much attention of the Pharmacologists and clinicians in Europe and America There are a number of others *Plantago ovata*—Isabgul is an excellent sedative in irritative conditions of the guts, *Adhatoda vasica* is a good expectorant; *Psoralea corylifolia* is useful in leucoderma, *Cissampelos pariera* is likely to be a useful smooth muscle relaxant, *Deamia extensa* as smooth muscle stimulant

A large number of those examined showed activity more or less the same as drugs already possessed by the pharmacopoeias These can be brought into use as cheaper indigenous substitutes Many others were effective enough to be used in the treatment of common minor ailments which form about 80% of total morbidity in this country A fourth group of drugs which though used in indigenous medicine were found to have little or no activity as judged by our standards. Many drugs of questionable value and doubtful utility crept into the indigenous systems during the period of decay

Full details of work done in this connection upto the end of 1953 will be found in the 2nd Edition of "*The Indigenous drugs of India*" which is a compendium of information with the regard to the whole subject of indigenous drugs and constitutes a leading work of reference. A Glossary of Indian Medicinal Plants is in the course of publication by the Council of Scientific & Industrial Research. In this book which has involved immense labour, brief reference has been made to scientific name of plants, their common vernacular names, the active principles contained,

the diseased conditions in which they are popularly used and the important references in literature on the work done This will form a handy volume for those interested in medicinal plants growing in India

The *Wealth of India* 1949-53 is a very comprehensive treatise which is being published under the auspices of the Council of Scientific & Industrial Research. This book is really a new edition of a Dictionary of Economic Products of India, published in 1895, brought up-to-date. It will eventually come out in many volumes The first five are already out and do credit to the Editorial Board of Council of Scientific & Industrial Research. In these books the results of investigations of many drugs on the lines indicated above are given.

A list of drugs which have been investigated with a brief summary of work done will be found hereafter. Those which have been shown to be of proved utility are marked with an asterisk

Value of Negative results. Apart from establishing the value of many useful remedies there is another aspect of this work which should not be neglected in this review At the present time most of the drugs used in the indigenous medicine are supposed to be specific for some particular diseases and lay people will wax eloquent in their descriptions of the wonderful cures said to have been produced by some of these remedies. Glowing statements of this nature, supported by insufficient evidence, have also some times appeared in medical journals, as in the case of effectiveness of *Vitex peduncularis* in malarial fevers and black water fever This has done a great deal of harm and distinguished pharmacologists and clinicians of the West are beginning to doubt if there is really anything of much value in the vast array of the materia medica of the indigenous systems of medicine Many are inclined to take the view that an investigation into the properties of these drugs is not likely to lead to much material results In this way the reputation of these remedies has grievously suffered in Western medicine, the good ones being indiscriminately classed with the bad Only systematic research of this kind can establish the value of the useful ones.

It should also be remembered, however, that the indigenous systems of medicine, good, bad or indifferent as they might be, minister to the needs of about 80 per cent of the population of this vast country even at the present time It is, therefore, the duty of research organisations in this country to investigate the materia medica which are in use in order to prove or disprove their effectiveness and their practical utility in the treatment of disease

In the following pages a brief summary of the work done on commonly used drugs is given first and then its common vernacular names, habitat, the parts of the plant used in indigenous medicine and the ailments in which they are prescribed After this the active principles found in the plant, their pharmacological action and results of clinical trials are recorded. Indications are also briefly given of the direction in which further work may be carried out.

Abbreviations used in description of Plants.

Afgh	..	Afghanistan
B	...	Bengali
Baluchi	...	Baluchistan
Baz	..	Bazar
Bhutia.	...	Bhutan and other Himalayan countries
Bo	...	Bombay
B P	...	British Pharmacopoeia
B P C	...	British Pharmaceutical Codex
Burm	·	Burmese
Dec	..	Deccan
H	..	Hindi
I P	...	Indian Pharmacopoeia
I P C.	...	Indian Pharmaceutical Codex
I P L.		Indian Pharmacopoeial List
K	..	Kanarese
Kash	...	Kashmiri
Kumaon	..	Kumaon Hills
M	...	Malayalum
Nep	...	Nepalese
P	..	Punjabi
Pers	..	Persian
S		Sanskrit
Singh	.	Singhalese
Santh.		Santhal
Vern		Vernacular

Abroma augusta Linn Devils cotton (Vern —**H** & B Ulatkambal Bo —Olaktambol) I P C

Grows wild in Uttar Pradesh, Sikkim, Khasia Hills and Assam Both root and bark are used as emmenagogue in menstrual disorders and as uterine tonic Fresh viscid sap is used in dysmenorrhea in doses of 30 grains a day An alkaloid 0 01%, a fixed oil, resins and water soluble basis have been found Preliminary investigations of the alkaloid and water soluble bases revealed no remarkable pharmacological activity No proper clinical trials have been carried out

Abrus precatorius Linn. (Indian or wild Liquorice root) (Vern —S & Bo —Gunja, H —Gaungchi, B - Kunch, M —Gundumani)

Grows wild in plains of India and Ceylon and along the Himalayas up to altitudes of 3,000 ft Seeds are used as tonic and aphrodisiac Roots and leaves contain glycyrhizin, an alkaloid abrine, a glycoside abralin and a fatty oil Preliminary investigations of the alkaloid revealed its marked poisonous properties No systematic study has been carried out with regard to its therapeutic efficacy

Aconitum (Vern —Bachnag)

A paste prepared from the root is used as a remedy for neuralgia and other painful affections The drug is used in the treatment of fever, rheumatism, cough, asthma and snake-bite

Aconitum balfourii Stapf (Vern -Nep -Gobari, Darmiya gobriya)

It is found in sub-alpine Himalayas from Gharwal to Nepal at altitudes of 12,000 ft to 14,000 ft. Root contains total alkaloids 1 2% of which pseudo-aconitine is 0.4%.

Aconitum chasmanthum Stapf. Indian Nepellus (Vern -Mohri, Kash -Banbalnag). I P. L & I P C

Grows in Western Himalayas of Kashmir at altitudes of 7,000 ft to 12,000 ft Root contains 4 3 per cent of total alkaloids, aconitine being the principal

Aconitum deinorrhizum Stapf (Vern -Safed bikh, Mohra, Maurabikh)

It is found throughout the Central Himalayas from Kumaon to Nepal and Upper Bushahr in Himachal Pradesh Roots contain total alkaloids 0 9 per cent of which pseudo-aconitine is 0 4 per cent.

Aconitum ferox Wall (Vern -S -Visha, H & B -Bish, Bo -Vachnag, M -Vashanavi)

It is found in Northern Himalayas, Nepal and Kashmir. Root is used as a remedy for neuralgia and rheumatism and as a sedative, antipyretic and diaphoretic

Aconitum heterophyllum Wall (Vern -Atis, S -Ativisha, H -Atis, B -Ataicha, M -Ati vadayam)

It grows in sub-alpine and alpine zones of the Himalayas It is used as a bitter tonic and febrifuge in the treatment of fevers, diarrhoea, dyspepsia, cough and as an aphrodisiac Roots contain alkaloids atisine 0 4 per cent , heteratisine and hetisin

Aconitum laciniatum Stapf (Vern - Kalo bikhmo),

Grows in the sub alpine and alpine Himalayas of Sikkim and Tibbet at altitudes of 10,000 to 14,000 ft.

Aconitum spicatum Stapf (Vern -Bikh, Kalo bikhoma donghi)

Grows in Nepal, Sikkim, Chumbi at altitudes of 10,000 to 12,000 ft It is used as a poison Roots contain 0 4 per cent of alkaloid named bikhaconitine

Aconitum violaceum Jacq (Vern -Tilia Kachnag)

Found in alpine zones of the Himalayas from Gilgit to Kumaon at altitudes of 10,000 to 15,000 ft The alkaloid aconitine is a cardiac irritant Locally it produces tingling followed by numbness Internally it slows the heart.

Acorus calamus Linn (The sweet flag) (Vern -H & B -Bach, Bo -Vaj, M.-Vashambu). I P C

Native of Europe and North America, but cultivated in India and Burma at altitudes of 3,000 to 6,000 ft Common in Manipur and Naga Hills. The rhizome is used as emetic, antispasmodic, carminative and expectorant. It is an old remedy against chronic, diarrhoeas Dried rhizome contains 1.5% of an essential oil, starch and tannin. No systematic pharmacological investigations have been carried out.

Actinodaphne hookeri Meissn (Vern.-Pichli, Bo -Pisa).

Grows in Sikkim and on the eastern and western Ghats. Infusion from leaves is used as a remedy for urinary disorders and oil from seeds is used as an external application in sprains of joints Bark contains a crystalline alkaloid named as actinodaphnine. Leaves contain amorphous salts and seeds an essential oil (1 0%), fat and a fixed oil No pharmacological or clinical works have been carried out

Adhatoda vasica Nees (Malabar nut tree. Vern -Arusha, S.-Vasaka, H. & B.-Adulsa, B.-Bakas, M Adhatodai) B. P C., I. P. C. & I. P. L.

Grows all over the plains of India and in the lower Himalayan ranges ascending to a height of 4,000 ft Leaves, roots and flowers are used as a remedy against chronic bronchitis, asthma and pthisis Juice of leaves is used in diarrhoea, dysentery and malarial fevers. It is also used locally as counter-irritant It contains an alkaloid vasicine and an essential oil The alkaloid vasicine has no very marked pharmacological action and produces only slight bronchodilation The essential oil acts as an expectorant. Fluid extract from leaves relieves coughs It is said to be effective in asthma and pulmonary tuberculosis An antibiotic principle having a powerful action on tubercle bacillus has been discovered.

Aegle marmelos Correa (Bael fruit Vern -S.-Sriphal, H, B. & Bo -Bael). B P C, I P C & I P L

Grows wild all over sub-Himalayan forests, in Bengal, South India and in Burma Root, bark, leaves and fruit are used to cure gastro-intestinal disorders Roots, bark and leaves contain reducing sugars and tannin and fruit contains marmelosin, the most potent active principle. Marmelosin acts as laxative and diuretic in doses of 0 05 gm Larger doses depress the heart Extracts made from the fresh or dried unripe fruit are used effectively in treatment of chronic diarrhoeas and dysenteries

Alangium salviifolium (Linn f) Wang Syn -*A. lamarckii* Thwaites (Vern -S -Ankota, H -Akola, B -Akar Kanta, Bo -Ankola, M.-Alangi).

Grows in forests throughout India and Burma Root bark is used as an anthelmintic and purgative also as a remedy for leprosy and skin diseases Bark contains an alkaloid alangin (0 82%) which produces a fall in blood pressure and depression of the heart Respiration becomes irregular Parasympathetic mechanism is stimulated No clinical trials have been carried out

Allium sativum Linn (Garlic Vern -S -Lasuna, H & Bo -Lasan, B -Rasun, M.-Vallaipundu). B P C & I P C

Is grown all over India and is used as a spice in cooking. It is considered to be hot and stimulant and useful in fevers, coughs and other debilitating conditions Externally it acts as rubefacient in skin diseases. It contains a volatile oil (yield from 0 06 to 0 1%), allacin and a crystalline substance isolated from the alcohol-insoluble fraction of the residue Garlic has an antiseptic action and slightly tonic effect on frogs' heart and a paralysing effect on isolated rabbit intestine Garlic juice is used as an antiseptic for wounds It is considered useful in digestive

disorders and respiratory diseases such as pneumonia, pulmonary tuberculosis and bronchitis Contains a powerful antibiotic principle effective against tubercle bacillus

Aloe vera ex Linn. (Vern.-Musabbar, S & B -Ghrita Kumari, H.-Ghi kanvar, M -Kattalai) Syn -*A. barbadensis* Mill B P, B P C., I P. C & I P L

Native of North Africa but grows in East and West Indies, India and China In India it is found in the dry valleys upto Cape Comorin It is used as stomachic, purgative and emmenagogue, and in the treatment of piles, rectal fissures and as a poultice for application to inflamed parts It contains a mixture of glucosides called aloin It is valuable in the treatment of constipation

Alpinia galanga Willd (The greater Galangal Vern -H -Bare kulinjan, B -Sugandha-vacha, M -Pera-rattai) I P C

Is a native of Sumatra and Java but commonly found in East Bengal and South India Rhizome is used in bronchial catarrh and rheumatism, tubers and seeds are carminative and useful for impotence and nervous debility Root contains compheride, galangin and alpinin From the green rhizomes a pale yellow oil is obtained The drug is depressant to cardio-vascular system and stimulates respiration in smaller doses but larger doses depress it. Bronchioles are dialated It is used as an expectorant in respiratory troubles of children and in asthma, intestinal and biliary colics

Alstonia scholaris R Br (Dita-bark Vern -S -Sapta parna, H -Chatium, B -Chhatim, M -Edakula-pala) I P C

Is cultivated throughout India and found in sub-Himalayan tract upto 3,000 ft Also found in Bengal and in Southern India Bark is used as tonic and alterative It is considered useful in fevers and skin diseases It contains a bitter principle ditain, alkaloids ditamine, echitamine, and acid and fatty resinous substances The alkaloids are not protoplasmic poisons. They relax isolated strips of rat intestine In the intact cat, the alkaloids produce a sharp fall of blood pressure Larger doses produce irregularities of heart beat Drug is used in the treatment of malarial fever, chronic diarrhoea and advanced stages of dysentery Recently shown to have no specific curative action in malaria

Amomum subulatum Roxb The greater Cardamom (Vern -S -Brihat-upa-kunchika, H & B -Bara-elachi, M -Periya-yelakkay) I P C

Grows in Nepal and Darjeeling Hills Used for flavouring purposes Seeds contain an oil rich in cineole

Andrographis paniculata Nees (The Creat Vern -S -Bhunimba, H -Kiryat, B -Kalmegh, M -Nila vembu) B P C, I P C & I P L

Is found throughout the plains of India from Lucknow to Assam and is also cultivated in gardens in some parts of India Leaves and juice are used to relieve griping, irregular stools and loss of appetite in children. Roots and leaves are used as febrifuge, tonic, alterative and anthelmintic Leaves contain two bitter substances and traces of an essential oil No systematic pharmacological investigations have been carried out. It has an intensely bitter taste.

Antiaris toxicaria Letsh. (The Upas tree Vern -Chanduka, Bo - Chandla, M -Nettavil, Sing -Riti, Burm -Hmyaseik)

Grows in Java, Malaya and Burma In India it is found in Concan, Canara, Travancore and Coorg Juice from leaves and bark is used as an arrow poison in Java, Malaya and Burma Bitter seeds are used as a remedy for dysentery and as a febrifuge in doses of half a seed three times a day. Milky juice of the plant contains antiarol, large amounts of potassium nitrate, antiaresin, a crystalline protein and three glucosides α antiarin, β-antiarin, γ-antiarin Preliminary toxicity tests on guinea pigs revealed that the drug is highly poisonous and the M L, D is 6 mgm per kilo body weight Death occurs due to heart failure It also produces marked tonic contraction of the isolated and intact intestines No therapeutic trials have been carried out

Arachis hypogaea Linn (Vern -S -Buchanaka, H -Mungphali, B -Chiner badam, Bo -Bhui-chane, M -Nila kadalai) Ground Nut B P., B P C , I P C & I P L

Originally a native of Brazil but now cultivated in all tropical or sub-tropical countries India is the largest producer The seeds are used as food, and oil as nutritive, laxative and emollient Seeds yield 40 to 50 per cent of a clear fixed oil which is used as a substitute for olive oil in India, as a basis for liniments and ointments Also used in the manufacture of soap and cosmetics, leather dressing and as substitute for tallow and diesel oil

Areca catechu Linn (The areca or betelnut palm) (Vern -S.- Gubak, H & B -Supari, Bo -Sopari, M -Kumugu) B P C , I P C & I P L

Grows in Sunda Islands but extensively cultivated in Southern India, Assam and Eastern Archipelago It is widely used in India and China as anthelmintic in man and animals, and as a masticatory to prevent the decay of teeth The seed contains alkaloids arecaine 0 1%, arecoline 0 07 to 0.1%, arecaidine, guvacoline, guvacine and choline, tannin 15% and fat 14% Arecoline is a highly toxic substance and its action resembles muscarine The seeds are used as anthelmintic in doses of 6 drams and for astringent action in relaxed condition of bowels.

Argemone mexicana Linn (Mexican Poppy) (Vern -Bharbhand, Kandiari, S -Srigala-kantaka, H & B -Sialkanta, M -Birama-dandu)

Grows wild all over India Juice of plant is used to cure dropsy, jaundice and cutaneous affections The seeds and seed-oil is used as a remedy for dysentery and intestinal affections The plant contains alkaloids, berberine and protopine The seeds yield about 22% of argemone oil which is used as a purgative but has no special advantage over other purgatives It is used as an adulterant of mustard oil and is believed to produce epidemic dropsy.

Aristolochia indica Linn. (Vern.-S.-Rudrajata, H. & B.-Isharmul, M.-Ichchura-mula). I P.C. & I.P.L.

Grows all over tropical portions of India, Bengal, Konkan, Travancore and Coromandel Coast. Root is used in the treatment of bowel complaints in children and in fevers. It possesses emmenagogue and anti arthritic properties. Constituents of the root are an essential oil,

a bitter substance and an alkaloid aristolochin The alkaloid produces cardiac and respiratory paralysis in frogs and mice Skeletal muscles are stimulated in small doses More detailed pharmacological investigations and clinical trials are indicated

Artemisia Sp (Vern —Kirmala)

Grows abundantly in the Himalayas from Kumaon to Kashmir at altitudes of 4,000 to 12,000 ft , also in Baluchistan, Chitral and Afghanistan. Flowering tops are used as anthelmintic in 2 to 4 dram doses The drug is also used as a remedy for dropsy and as cardiac and respiratory stimulant. It contains a volatile oil, santonin and an allied body—artemisin.

Artemisia absinthium Linn Wormwood (Vern -H. & Dec - Vilayati afsantin).

Grows in Kashmir at altitudes of 5,000 to 7,000 ft The oil has a tonic and stimulating effect on the digestive system Contains 0 3 per cent. of an essential oil, a bitter glucoside absinthin and a crystalline compound

Artemisia dracunculus Linn

Found in western Tibet at altitudes of 14,000 to 16,000 ft and in Lahul Leaves are used for flavouring vinegar and as a spice. They contain about 0 3 per cent of an essential oil

Artemisia maritima Linn (Vern -S -Gadadhar, H -Kirmala, Bo. Kiramani owa) I P C & I P L

Found in Western Himalayas from Kashmir to Kumaon at altitudes from 7 000 to 9,000 ft in Kashmir, Kurrum, Kagan, Bushher, Waziristan and Chamba.

Artemisia pallens Wall ex-DC (Vern -Davana)

Grows in certain parts of South India in Mysore State and is also cultivated near Poona. Leaves are used as floral decoratives and the oil from the herb is a popular article in perfumery.

Artemisia sacrorum Ledele (Vern P.-Tatwen)

Grows in Western Tibet, Kanawar and in Tibetan regions of Kumaon at altitudes of 10,000 to 12,000 ft Contains 1 per cent. of an essential oil

Artemisia vulgaris Linn (Vern -Tithwan, S -Nagadamani, H -Nagadouna, B -Nagdona).

Grows throughout the mountainous regions of India ascending to an altitude of 12,000 ft., in western Himalayas upto 5,000 to 8,000 ft , in Sikkim and Khasia Hills, also in Mount Abu and Western Ghats, Used as antiseptic expectorant and anthelmintic Contains 0.2 per cent of a volatile oil which is a good larvicide.

Asteracantha longifolia Nees Syn.-*Hygrophila spinosa* T Anders, (Vern -S.-Kakılakshya, H -Talmakhana, B. Kuliakhara, Bo.-Talım khana, M -Vallarai) I P C

Grows commonly in moist places throughout India from Himalayas to Ceylon Roots are used as cooling, diuretic, stimulating and to remove dropsical fluids and kidney stones They contain phytosterol, traces of an essential oil, a yellowish green wax, gum and maltose Seeds contain an oil 23 per cent and an alkaloid. No pharmacological and clinical studies have been carried out

Atropa acuminata Royle (Vern -Sag-angur) B P , B P C & I P C.

Grows in abundance in the Himalayan ranges extending from Simla to Kashmir, at altitudes of 6,000 to 12,000 ft In Jammu & Kashmir State, it is found in the forest mountains round Kashmir valley. It is also found in Himachal Pradesh, Kulu, Parbati valleys and Narkanda forests of Simla Hills Belladonna and its alkaloid atropine are largely used as a sedative, antispasmodic and mydriatic in diseases of the eye, and as antidote in opium and muscarine poisoning Alkaloids (as hyoscyamine) are present from 0 15 to 0 7 per cent in the leave sand 0 3 to 0 66 per cent in the root Traces of an inactive volatile alkaloid is also present The plant is now cultivated and large amount of belladonna preparations in India come from this source

Azadirachta indica A Juss (Vern -Nim) The Neem. I. P C. & I P L

Grows throughout the greater part of India and Burma Bark and leaves are used as a bitter tonic and astringent It is considered useful in skin diseases Fruit is used as purgative, emollient and for expelling intestinal worms Considered beneficial in urinary diseases and in the treatment of piles Bark contains an alkaloid margosine and seeds yield a fixed bitter oil Margosates are toxic to protozoa and bacteria Clinical trials revealed that the drug has a beneficial effect in skin diseases

Bacopa monnieri (Linn) Pennell, Syn *Herpestis monniera* (Linn) H B & K (Vern -S -Brahmi, H -Braehmbhi, B -Brihmisak, M -Nirbrami) I P. C

Grows throughout India on the borders of streams and tanks Roots, leaf and stalks are used as nerve tonic and as a remedy for insanity and epilepsy It contains an alkaloid, herpestine, which is highly toxic It produces a fall of blood pressure and stimulation of respiration. The smooth muscle of intestines and uterus is stimulated. In therapeutic doses it resembles strychnine Powdered dried leaves gave satisfactory results in cases of asthenia and nervous break down Further clinical trials are needed

Bambusa bambos Druce, Syn.—*B. arundinacea* Willd. (Vern.-Kantabans, S.-Vansa, B. & H -Bans, Bo.-Mandgay).

Grows wild throughout the greater part of India particularly in the hilly forests of Western and Southern India upto an altitude of 3,000 ft. Leaves are considered useful in diseases of blood, leucoderma and

inflammatory conditions Tabashir found as concretions in it is used against blood diseases, tuberculosis, asthma and leprosy Seeds contain water 11 0 percent , starch 73 7 percent , albuminoids 11 8 per cent , oil 0 6 per cent, fibre 1 7 per cent and ash 1 2 per cent Young shoots contain a cyanogenetic glucoside No systemmatic work has been done to determine its proper therapeutic efficacy

Berberis aristata DC (The barberry) (Vern -Chitra, S -Daru haridra H -Dar-hald) I P L. & I P C

Grows in the temperate Himalayas at an altitude of 6,000 to 8,000 ft A tincture made from root bark was used as a bitter tonic, stomachic, cholagogue, antiperiodic and alterative Berberine is the chief alkaloid, also umbellatine is present It is used as diaphoretic and antipyretic in malarial fever but has little specification

Berberis asiatica Roxb. (Vern.-Sumlu, H -Kilmora, Nepal -Chitro). I P.L.

Grows in dry valleys of the Himalayas at an altitude of 3,000 to 7,500 ft in Bhutan, Garhwal and Behar Its uses are similar to those of *B aristata* and its active principle is berberine

Berberis coriacea—Brandis (Vern -Simla-Kashmal) I P L.

Grows in N W Himalayas at an altitude of 8,000 ft

Berberis floribunda Hort

Grows in Nepal The roots contain eight alkaloids namely Oxycanthine, bebamine, berberine, epiberberine, palmatine, dihydrocarydeline, jatrorrhizine and columbamine No detailed pharmacological studies have been carried out but the action of these alkaloids would appear to resemble berberine.

Berberis himalaica

Found in Bhutan Bark contains berberine (0 03%), jatrorrhiszine (0 01%) and an alkaloid limanthine It has not been used in medicine although it belongs to berberis group of plants.

Berberis insignis Hook

Grows in humid forests of the Eastern Himalayas from Nepal and Sikkim to Bhutan at altitudes of 8,000-10,000 ft Stem bark contains 1 52 per cent and root bark 2 5 per cent of total alkaloids consisting chiefly of umbellatine

Berberis lycium Royle (Vern H -Kashmal, Bo -Darhald)

Grows in the Western Himalayas at altitudes of 3000-9000 ft from Garhwal to Hazara It is used as febrifuge carminative and aperient and is considered beneficial in piles, enlargement of liver and spleen and as external application to eyelids in conjunctivitis. It contains alkaloid umbellatine.

Berberis umbellata Wall.

Occurs on Himalayan ranges at altitudes from 9,000 to 12,000 ft. from Kashmir eastward to Bhutan. Bark contains Umbellatine which like berberine has a specific inhibitory action on *Leishmania tropica*. The cardiovascular system is depressed and para-sympathetic nervous system is stimulated. Respiration is depressed.

Berberis vulgaris Linn (The true barberry) Vern. Zirishk, P - Kashmal) I P L

Grows in the Himalayas from Nepal westwards at an altitude of over 8,000 ft It is used as diuretic, astringent and refrigerant In small doses it acts as a tonic and in large doses as a purgative It contains berberine which has an inhibitary action on *leishmania*.

Blepharis edulis Pers (Vern —H -Utanjan, Bo -Utangan).

Grows in the Punjab and Baluchistan. Leaves used as astringent in bowel complaints It has aphrodisiac and alterative properties It is useful in fevers, urinary discharges and leucoderma. Seeds contain a bitter glucoside belapharine (1 2 per cent) and dl-allantoin (2 1 per cent.) No detailed pharmacological studies have been carried out.

Boerhaavia repens Linn (Vern Thikri, S - Shothaghni, H -Sant, B -Punarnaba, Bo -Ghetuli, M -Mukuk-rattai) B P , I P C. & I P L.

Grows all over India It is used as laxative and diaphoretic in oedema, anaemia, heart disease, cough and intestinal colic. It contains an alkaloid named punarnavine and an oily amorphous matter Sulphates, chlorides, nitrates and chlorates occur in ash The alkaloid is not toxic. It has a diuretic effect and raises blood pressure The movements of intestines are depressed The liquid extract in doses from one to four drams produces definite diuresis in patients with oedema and dropsy

Bregantia wallichii R Br (Vern —Chakrani).

Grows in southern half of Bombay State near the Coast, Madras and Ceylon. Used to cure chronic sores and ulcers Considered useful in cholera and diarrhoea. Roots contain a fatty oil and a yellow substance identical with isoaristolochic acid No pharmacological studies or clinical trials have been carried out.

Butea monosperma Lam -Kutze (Vern. Palas) I P C

Grows throughout India and Burma extending in the N W. Himalayas as far as Jhelum Gum (Kino) is used as astringent, diuretic and aphrodisiac. Seeds are used as an anthelmintic and contain 18 per cent. of a fixed oil, small quantities of resin and large amounts of water-soluble albuminoids The oil does not possess any marked anthelmintic action No systematic investigation has been carried out.

Butea superba Roxb. (Vern.-Palarvela, S Lata palas, H. & B.- Paas lata, M.-Kodi-murukkan).

Grows in the forests of Oudh and Bundhelkhand, Chotanagpur, Central and Southern India. It is used as a remedy against eruptions in children. Decoction of stem and leaves is emollient and is used as a

local application for piles. Root contains an esterogenic substance and a poisonous substance Flowers contain butrin, butein and butin An aqueous extract equivalent to 0.002 per cent of the dried powder causes estrus in spayed female mice Further pharmacological studies are indicated

Caesalpinia crista Linn Syn - *C. bonducella* Fleming. Fever nut (Vern -S -Kuberakshi, H -Kat-karanj, B -Nata-karanja, Bo -Sagaragoth, M -Gajega) I P.C.

Grows near the sea-coast, all over Bengal, Bombay and S India Seeds, leaves and roots are used as anthelmintic, emmenagogue and febrifuge Seeds contain starchy matter, a bitter principle called bonducin and an oil The plant is not of much therapeutic value

Calotropis gigantea Linn.- (Gigantic swallow root) (Vern Ak-Madar, S -Arka, H -Ak , B & Bo -Akanda, M -Erukku) I P C.

Grows in waste lands all over India upto altitudes of 3,000 ft Milky juice is used for tanning and dyeing purposes An intoxicating liquor is prepared from the juice Latex contains water soluble matter 86 to 95 5 per cent Stem bark contains calotropeolis, B-amyrin and giganteol No marked therapeutic value.

Calycopteris floribunda Lam (Vern H -Kokoranj, Bo.-Ukshi, M -Marsada boli)

Grows in Central and Southern India in plains and upto an altitude of 2,500 ft Leaves used as laxative, anthelmintic and the juice given in puerperal fevers Also used in the treatment of dysentery and malarial fevers The juice contains, chlorides, sulphates, nitrates, lime, ferric oxide, tannin, albuminoids and gummy matter Leaves contain a yellow crystalline flavone, calycopterin. The drug is toxic to fish and earthworms , more toxic to earthworms than santonine or chenopodium oil No detailed pharmacological and clinical investigations have been carried out

Camellia sinensis Linn (O Kunfize) Vern -Cha Tea plant B P & B.P.C.

In India, tea and coffee grow luxuriantly. Coffee is grown principally in Madras, Coorg, Mysore, Travancore and Cochin. Tea is also found in Ceylon and is grown in Assam, Bengal, Madras and Travancore Tea is used as a beverage in India and other parts of the world Tea leaves contain from 2.5 to 3% of Caffeine Coffee contains 1 5 per cent of caffeine Caffeine is used in medicine for stimulation of central nervous system and circulation and as a diuretic.

Cannabis sativa Linn. (*Cannabis indica*) (Vern S.-Ganjika, H & Bo -Ganja, M -Bhangi, Bhang) B P C , I.P C. & I.P L

Grows wild throughout the Himalayas from Kashmir to East of Assam and also in Southern India. Plant and its products are used for narcotic purposes, by smoking or taking internally It contains a terpene 1 5 per cent , a sesquiterpine 1 75 per cent., small amount of paraffin carbon and a resin named as Cannabinol (33 per cent.). Hemp drugs are consumed all over the world for euphoric purposes Their use alleviates the feeling of fatigue, encourages sleep and soothes restlessness.

Sensation of time and space is lost. Larger doses produce mental derangement and even insanity.

Carica papaya Linn. (The Papaya) (Vern Papita, H.-Papaya, B.-Papey, Bo -Papai, M -Pappayi). B.P , B.P.C , I P.C & I.P L.

Grows all over India and is used as anthelmintic, emmenagogue and digestive. The fruit pulp contains sucrose, invert sugar, resinous substance, papain, malic acid, salts of tartaric and citric acids and vitamins. Seeds contain, proteins, carbohydrates, fatty oil, volatile oil, an alkaloid carpaine, a glycoside carposide and an enzyme myrosin. Carpaine is a heart poison, it lowers blood pressure and depresses heart No proper clinical trials have been carried out

Carum carvi Linn The Caraway seed (Vern Zira, H.-Shiajira, B -Jira, M Shimai-shombu) B.P., B P C & I P C

Widely distributed in the temperate regions of both the hemispheres Grows in North and Central Europe, extending to the Caucasus, Persia, Tibet and Siberia In India it is cultivated on the hills of Baluchistan, Kashmir, Kumaon, Gharwal, Chamba at altitudes of 9,000 to 12,000 ft. Used as a cookery condiment Oil is used for flavouring purposes and as a carminative Seeds contain an essential oil with a marked carminative action

Cassia absus Linn (Vern H —Chaksu, N —Karun Kanam).

Grows in lower parts of the Western Himalayas and Ceylon Leaves are applied locally to tumours and internally in cough, asthma and bronchitis Seeds are used as bitter, astringent and diuretic and contain two water soluble bases, chaksine and Isochaksine, as carbonates (1 5 per cent) Premliminary studies show, that chaksine sulphate is depressant to heart, respiration and nerve centres More detailed studies are indicated Considered useful in eye diseases. No clinical trials have been carried out

Cassia angustifolia Vahl (Vern H -Hindisana, B.-Sonamukhi, M -Nila vakai) B.P , B P C & I P.C.

Cultivated in S India in Madras State (Madura) Leaves and pods are used as purgative. Leaves and pods contain cathartic acid, emodin and chrysophanic acid Senna is useful in habitual constipation. It increases the peristaltic movements of the colon. The tendency to gripe is overcome by mixing it with aromatics

Cedrela toona Roxb (Vern Mahalimbu, S. & B.-Nandibriksha, H -Tun, Bo.-Tuni, M -Tunumaram, P -Khushing).

Grows in Sub-Himalayan tracts from river Indus eastwards, Chittagong, Assam, Burma, Chotanagpur, Western Ghats of Bombay to the Nilgiris and other hills of the Deccan Peninsula The plant is used as astringent in the treatment of ulcers, gleet and scabies Flowers are used in menstrual disorders and contain a red colouring matter, nyctanthin, flavone and a glucoside. Bark contains, tannic acid, a bitter resin, citric acid and starch. No pharmacological or clinical investigations have been carried out.

Celastrus paniculatus Willd (Vern. S -Kanguni, H -Malkanguni, M -Valuluwai)

Grows in Sub-Himalayan tracts extending from Jhelum to Assam ascending to an altitude of 6,000 ft and in hilly parts of Bombay State, South Gujrat, Central India and Madras Seeds and oil are used in the treatment of rheumatism, gout, paralysis and epilepsy Oil is considered as nerve stimulant and brain tonic Leaves contain an alkaloid, a glucoside and colouring matter Seeds contain 52 2 per cent of a thick brownish oil No pharmacological studies have been carried out.

Centella asiatica (Linn) Urban, Syn -*Hydrocotyle asiatica* Linn (Vern -Brahmi, S -Vanduka paini, H -Brahmo manduki, B -Tholkuri Bo.-Karivana, M -Vallarai) I P C

Grows throughout India from Himalayas to Ceylon at an altitude of 2,000 ft Plant is used as tonic and is considered beneficial in diseases of the skin, nervous system and blood Also used in the treatment of leprosy and syphilitic affections. It contains an alkaloid, hydrocotyline and fresh leaves contain a glucoside asiaticoside Hydrocotyle is shown to be a circulatory stimulant Further work is indicated.

Centratbrum anthelminticum Kuntze, Syn -*Vernonia anthelmintica* Willd (Vern -S , H & B -Somaraj, Bo -Kalijiri, M -Kattu-Shiragam). I P C & I. P. L.

Is found in waste lands near villages throughout India The seeds are used as a remedy for leucoderma and other skin diseases, and as anthelmintic They contain a bitter principle 1 per cent, a fixed oil 18 per cent and a small amount of an essential oil No systematic pharmacological investigations have been carried out Clinical trials revealed that it is effective in thread-worm infections Further work is needed

Cephaelis ipecacuanha A Rich Syn -*Psychotria ipecacuanha* stokes B P , B P C & I P C.

Grows wild in Brazil In India it has been cultivated in South India and Bengal—in Darjeeling hills Contains emetine and other alkaloids used in the treatment of amoebic dysentery

Cerbera odollam Gaertn Syn -*C. manghas* Linn. (Vern B -Dhakur, M -Kataralı).

Grows all over India particularly in salt swamps in the Malabar Coast, Ceylon and Burma Plant is intensely poisonous The leaves and sap have emetic and purgative properties Seeds contain three glycosides named as cerberin, odollin and cerbeside Cerbrin is non-irritant to the skin and mucous membranes and produces vomiting, diarrhoea and even syncope when given subcutaneously Stimulates cardovascular system and smooth muscles Further clinical trials and experimental work is indicated.

Chenopodium ambrosioides Linn B P , B.P C , I.P.C. & I.P.L.

Grows in Central America, East Indies. In India found in Bengal, Sylhet, the Deccan (Coimbatore) Infusions from leaves and seeds used as a remedy against intestinal worms Seeds contain a volatile oil which

contains ascaridole from 45 to 70% Small portions of glycol-anhydride, mixture of hydrocarbons and traces of fatty acids also occur Clinical trials have shown its efficacy against hookworms and round worms

Chenopodium botrys Linn

Grows in the Himalayas from Kashmir to Sikkim Used as a substitute for *C ambrosioides* Contains a volatile oil 0 03 to 0.04 per cent

Cichorium intybus Linn (Vern -Kasni)

Grows in North Western India and is cultivated in Nadiad, Broach and Amalsad in Bombay State Used as fodder Cultivated sweet variety is considered beneficial in acne, opthalmia and throat inflammations Roots used as diuretic and to enrich and purify blood Wild variety used as emmenagogue and to cure asthma Plant contains a glucoside cichorin Seeds contain, a semi-drying oil, water, gum, glucose, bitter substances, inulin and fibre No systematic studies of its therapeutic action carried out

Cinchona Cinchona bark B P., B P C , I P.C & I P L.

Cinchona grows on the Eastern slopes of Central Western chain of the Andes mountains in South America, at altitudes of 2,500 ft to 9,000 ft. It is cultivated in India in the Nilgiris Hills, Mungpoo in Ranchi Valley and the Karen Hills in Burma The bark contains quinine, cinchonidine, quinidine and cinchonine It is used in the treatment of malaria

Cinnamomum Blume.

Grows in Asia and Australia About 20 species occur in India Used as an aromatic and spice

Cinnamomum camphora Nees B.P , & B.P.C.

Cultivated in India at Dehradun, Saharanpur, Calcutta, Nilgiris and Mysore

Cinnamomum zeylanicum Breyn (Vern —Dalchini, S -Gudatreaka) B P , B P C & I P C

Grows wild on Western Ghats from Konkan southwards and in forests of Tenasserim in Burma Found abundantly in Ceylon Used as carminative, astringent, stomachic, flavouring agent and as a spice. Bark contains an essential oil o 5 to 1% Leaves yield a dark coloured oil 1%

Cissampelos pareira Linn (Vern —Akanadi, S -Patha, H & B -Nirbisi) I P C

Grows throughout tropical and sub-tropical parts of Asia, East Africa and America Root is used to relieve pain and is considered useful in febrile conditions, dysentery, heart troubles and urinary disorders. Contains an amorphous alkaloid named pelosine, saponin and ammonium salts, also hayatin and hayatinin Hayatin methiodide, methochloride have curariform activity equal to tubocurarine chloride They produce fall of blood pressure which is prevented by antihistaminic drugs Further studies are in progress. A plant likely to be of great therapeutic value.

Citrullus colocynthis Schrad. (Vern.—Indrayan, S.-Indra-varuni, H. & Bo-Indrayan, B.-Makhal, M.-Peyt-tumatti). B.P., B.P.C. & I.P.C

Grows in arid tracts of N West, Central and South India and is met with in the Punjab, Sind and on the Coromandel coast. Fruit is used as cathartic to relieve biliousness and constipation, used in treatment of fever and intestinal parasites Plant contains a bitter principle colocynthin and traces of an alkaloid Used in medicine as a drastic purgative

Citrus aurantii-folia (Christm) Swingle Syn -*C medica*, var *acida*. The Lime Tree.

Citrus limon (Linn) Burm f *Syn,-Citrus medica*, var. *limonis*. The Lemon Tree B P., B P.C., I P C & I P L

Lime and lemon are found growing wild in the warm valleys of the North West Himalayas Also largely cultivated in the plains and up to altitudes of 4,000 ft Used as flavouring agent in medicine and also as a carminative and stomachic Peel contains an essential oil and juice citric acid upto 5 9 per cent

Cleome icosandra Linn. (Vern —Aurhur).

Grows throughout India but commonly found in Bengal and S. India Seeds are used as anthelmintic, rubificient and vesicant They contain a fixed oil (36 59 per cent) and a new flavone called 'Viscosin' No work on its pharmacological action and therapeutic properties has been carried out

Clerodendron infortunatum Linn (Vern.—H & B.-Bhant, Bhat)

Grows in waste lands throughout India and in damp forests of Ceylon, upto an altitude of 5,000 ft. Used as a vermifuge, anthelmintic and a substitute for Chireta Leaves contain ash 8.04 per cent, protein 21 1 per cent., crude fibre 18 84 per cent, total sugars 17 05 per cent and a bitter substance clerodin The last named has no haemolytic and bactericidal action It is toxic to earthworms, small fish and mosquito larvae. No clinical trials carried out

Coccinia cordifolia Cogn Syn - *Cephalandra indica* Naud, (Vern,- S -Bimba, H -Kanduri-ki-Bel, B -Telakucha Bo -Bhimb)

Grows wild in Bengal and other parts of India It is said to have antidiabetic properties It contains an enzyme, a hormone and traces of an alkaloid Did not show any action on blood sugar in animals nor by clinical trials in diabetic patients.

Colchicum luteum Baker (Vern —Surinjan) I P C. & I.P L.

Grows extensively in the Western temperate Himalayas, extending from Murree hills to Kashmir and Chamba. Used as an alterative and aperient in gout, rheumatism and diseases of liver and spleen. Bitter variety contains alkaloid colchicine, starch and oily resinous matter Galenical preparations of the crude drug and the alkaloid colchicine are used in the treatment of gout. Recently colchicine has been widely used in plant breeding to induce polyploidy.

Commiphora mukul Engl. (Gum gugul) (Vern -S -Guggulu, H ,B. & Bo.-Guggul, M.-Gukhulu).

Grows in Rajputana, Khandesh, Sind, East Bengal and Assam The gum resin from bark is used as demulcent, aperient, carminative and alterative and is useful in leprosy, rheumatism, syphilitic disorders, nervous and skin diseases and urinary disorders Gum resin contains 32 per cent. gum, 19.5 per cent. mineral matter and 1.45 per cent of an essential oil. The pharmacological action of this oleoresin resembles the action of copaiba and cubebs Taken internally it acts as bitter, stomachic and carminative It also acts as diaphoretic, expectorant and diuretic, It is said to be uterine tonic and regulates menstrual functions No proper chemical trials have been carried out

Coptis teeta Wall. (Vern H -Mamira, B -Tita, Bo-Mahmira) I P.L.

Grows in upper Assam. It is used as an eye salve in indigenous medicine. Active principles berberine 8 5 per cent , coptine (0.08 per cent), palmatine (traces), coptisine 0 02 per cent and jatrorrhine (0 01 per cent) It is used as a bitter tonic in the same way as Calumba.

Coscinium fenestratum Colebr. (Vern —Haligach, S.-Daru haridraka, B -Haldi-gach, M -Mara-manjal) I.P C.

Grows in the forests of Western India and is used as a bitter tonic. It contains berberine

Crocus sativus Linn (Vern.—Kesar, Safran, S -Kumkuma, H & Bo.-Jafran, M -Kungumapu) B P C , I P C & I P L

Is commonly found in Kashmir and Quetta and also cultivated in Spain It is considered to be stomachic, stimulant, antispasmodic and aphrodasiac. It is used as flavouring and colouring agent in cookery. It contains three crystalline colouring matters, α-crocetin, β-crocetin, γ-crocetin, a fatty oil, a volatile oil and a bitter substance The essential oil has the characteristic features of other essential oils No proper clinical trials have been carried out.

Curcuma longa Linn Syn -*C domestica* Valeton (Vern —S -Haridra, H.-Haldi, B -Halood, Bo -Halada, M -Munjal). B. P C. & I P C.

Grows throughout Southern parts of Asia, in India, China and East Indies. Also cultivated in almost all the States of India, particularly in Madras, Bengal and Bombay Used as a spice Hakims use it as a stomachic, tonic and blood purifier The juice of rhizome has anti-parasitic action in affections of skin Rhizome contains an essential oil 5.8 per cent., protein, mineral matters, carbohydrates, and vitamin A. The oil has slight anti-bacterical properties Taken internally it has antacid action and relaxes the intestines. Used as a stomachic and carminative.

Curcuma zedoaria Rosc. (Vern.—S.-Sati, H. & Bo.-Kachura, B -Shori, M.-Pulan-ki-zhanga).

Grows in Eastern Himalayas and Kanara, Ceylon and China. It is also cultivated in many Indian States. Rhizomes are used as aromatic; stomachic, stimulant and carminative. Externally applied on bruises

and sprains. Contains an essential oil 1,5 per cent., starch 82.6 percent., ash 1.01 per cent and moisture 13'1 per cent The essential oil is carminative and produces relaxation of the intestines

Cuscuta reflexa Roxb. (Vern —S -Amaravela, H -Akasbel, B.-Algusi, Bo -Nirmuli, M.-Sitama-purgonalu).

Grows throughout the plains of India upto an altitude of 5,000 ft Used as carminative, purgative, diuretic and purifier of blood. Seeds contain a bitter substance, a glucosidal resin and quercetin Plant also contains, cuscutin, cuscutalin, brown wax and reducing sugars. No pharmacological studies have been carried out.

Daemia extensa R Br (Vern —S.-Phala kantak, H.-Utran, B.-Chagulbanti, Bo & M -Utarni) I.P C

Grows throughout the hotter parts of India upto an altitude of 3,000 ft Used as cooling, anthelmintic, laxative and antiperiodic. Considered useful in eye troubles, uterine disorders and urinary troubles The plant contains 2 4 per cent of inorganic salts, a bitter resin, and three bitter substances Bitter principle C is the most active pharmacologically Has a stimulant action on uterus, intestines, due to direct action on the involuntary muscle Further pharmacological studies are in progress

Datura stramonium Linn (Vern —B -Sada dhutura, P -Tattu dattura, M -Umatai) B P ,B P C & I P C

Grows throughout the temperate Himalayas from Kashmir to Sikkim Used as intoxicant, emetic, digestive and as antispasmodic in asthma and whooping cough. Seeds and leaves contain hyoscyamine, atropine and hyoscine Datura possess properties similar to belladonna

Dichora febrifuga Lour. (Vern.—H -Basak, Nep -Aseru).

Grows in the temperate Himalayas from Bhutan to Khasia hills, at altitude of 4,000 to 8,000 ft , also in Upper Burma, Malayan Peninsula and China Roots are used to cure malaria in China and as a febrifuge in India. Roots and leaves contain two alkaloids, febrifugine and Isofebrifugine Clinical trials have revealed that the drug has antipyretic action like quinine (1 5 times) but no anti-parasitic action. Further work is indicated

Didymocarpus pedicellata R Br

Grows in subtropical regions from Chamba to Kumaon at altitude between 2,500 and 5,000 ft Leaves are used as a remedy for stones in kidney and bladder The plant contains pedicin 1 per cent., iso-pedicin 0.4 per cent , pedicinin 0.3 per cent. and pedicellin 1 per cent which are crystalline No detailed pharmacological and clinical trials have been carried out.

Digitalis lanata Ehrh. Grecian Foxglove B P , B.P.C. & I.P.C

Grows in Kashmir at altitudes about 7,000 ft Fresh leaves contain 3 natural glucosides lanatosides A, B & C which on hydrolysis give digitoxin, gitoxin and digoxin Digoxin produces the same cardiac effects as digitalis. It is 300 times more potent than prepared digitalis

and is of particular value for rapid digitalization. Digitalis increases the force of systolic contractions and efficiency of the decompensated heart Slows the heart rate and reduces cardiac œdema with diuresis

Digitalis purpurea Linn Foxglove. B P., B P.C. & I P. C.

Cultivated or found as an escape in hilly regions at altitudes of 5,000 – 6,000 ft Originally a native of Western Europe but now extensively grown in many parts of the world In India, it is found in Kashmir Himachal Pradesh, Darjeeling area and other parts. It is used as a myocardial stimulant in congestive heart failure Used as a heart tonic and diuretic Contains several glucosides but the leaves contain digitoxin, gitoxin and gilatin

Dregea volubilis Benth. ex Hook (Vern —S -Madhu malati, H.-Nakchikni, B -Titakunga, Bo -Dodhi, M -Kodicpalay)

Grows wild in Assam, Bengal, Deccan, Peninsula from Konkan southward to Ceylon The plant is used to cure piles, leucoderma, asthma and urinary discharges It contains a substance of glucosidic nature with low toxicity and traces of an alkaloid It stimulates all organs having cholinergic nerve supply and causes a prolonged fall of blood pressure Further work is indicated

Eletttaria cardamomum Maton (Vern —S -Ela, H. & B -Chotielachi, Bo -Elachi, M -Ellakai) B P , B P C , I P C & I P L.

Grows in Western and Southern India in the forests of Kanara, Mysore, Coorg, Wynaad, Travancore, Cochin Also found wild in Burma, Ceylon, China and Malaya Archipelago Used as a spice and masticatory and in medicine as carminative and flavouring agent. Seeds contain an oil to the extent of 2 to 8 per cent.

Entada phaseoloides (Linn) Merr Syn.-*E. pursaetha* DC., *E. scandens* Benth

Grows in Central and Eastern Himalayas at altitudes of 4,000 ft and is also found in damp forests of Eastern Bengal, Bihar, Orissa, Eastern and Western Ghats Seeds are used as soap to wash hair Locally it is applied to swellings , it is a fish poison Seeds contain two saponins, entada-saponin A & B and 8 per cent of a fixed oil. Both saponins are toxic to rabbits and guinea pigs They produce hæmolysis of red blood cells and a depressant action on circulatory and respiratory systems Movements of smooth muscles are inhibited No clinical trials have been carried out

Ephedra gerardiana Wall Syn -*E vulgaris.* and allied sp. (Vern Amsania) B P C., I P C & I P L

All species grow in Northern India, Bashahr Division, Chakrata, Kangra, Kulu, Baluchistan, Kashmir, Hazara, Kagan and Waziristan. Contains ephedrine and pseudo-ephedrine. Ephedrine is a useful remedy for asthma Tincture of Ephedra is an excellent cardiac stimulant in toxic conditions

Erythroxylum coca Lam. B P. & B P.C.

Originally a native of South America but is grown in the West Indies, India, Ceylon and Java. Leaves used for euphoric purposes and

contain cocaine to the extent of 0 15 to 0 8 per cent. and other alkaloidal substances. Cocaine is used for local anaesthetic purposes and also as a euphoric

Eucalyptus L. W Erit (Vern.—Karpura Maram) Blue gum tree. B.P., B P C., I.P C. & I P.L

Grows naturally in Australia but cultivated in other parts of the world The oil from leaves is used in soap industry and as an antiseptic and disinfectant. Leaves and terminal branches yield an essential oil

Eugenia aromatica Kuntze Syn -*Caryophyllus aromaticus* (Vern. Long) B P & I P C

Native of Malacca Islands and is largely cultivated in Zanzibar, Pemba, the Amboyna Islands, Penang, Madagascar, Seychelles, Reunion, Mauritius and Ceylon and Southern India Dried buds contain an essential oil and are used as aromatic stimulant and carminative

Euonymus tingens Wall (Vern —Barphali, H -Kungku) (Dogwood) I P.C

Grows in tropical regions of Asia, Malaya, Europe and America. Used to stimulate torpid liver producing flatulence and indigestion.

Euphorbia hirta Linn Syn -*E pilulifera* Linn (Vern —H -Dudhi, B -Bara keru, Bo -Nayeti, M .-Amumpatchaiarissi). B P.C & I.P C.

Grows all over the hotter parts of India. It is used as a remedy for respiratory troubles, dysentery, colic and worms in children It contains gallic acid, quercetin, traces of an alkaloid and an essential oil. The extract possesses depressant action on heart and respiration and produces relaxation of bronchioles It is used to relieve asthma and other spasmodic conditions of the respiratory tract

Ferula foetida Regel. (Vern —S -Hingu, H & B.-Hing, Bo -Kayam) Asafoetida B P C , I P C. & I P L

Found in Turkestan Gum resin used as a carminative, antispasmodic and as flavouring agent in spices Contains a volatile oil, resin, gum and impurities Used in medicine as carminative and antispasmodic in nervous disorders of childern and women

Foeniculum vulgare Mill (Vern —S -Madhurika, H —Bari-saunf, B -Pan-mauri, Bo -Bari sopha, M -Shombu). B P., B.P.C , I.P.C. & I P.L

Cultivated throughout India and also in cold places upto an altitude of 6,000 ft. Used as aromatic, stimulant and carminative. Fruit contains a volatile oil with pleasant aromatic odour, chief constituent of which is anethol but small quantities of substances like fenchone are also present.

Gaultheria fragrantissima Wall. (Vern.—Jar Gandapuro). Indian winter green. I.P.C & I P.L.

Grows freely in the Nilgiris, Travancore and Assam. Oil of winter green used as an external application for rheumatic affections, sciatica,

neuralgia and as a flavouring agent in tooth pastes, etc. Leaves contain an oil which mainly consists of methyl-salicylate.

Glycyrrhiza glabra Linn. (Vern —S -Yashti-madhu, H -Jethimadh, B. & Bo -Jashti madhu, M -Atimaduram) B P., B P.C. & I.P C.

Found in sub-Himalayan tracts from the Chenab westwards, Burma and the Andaman islands Used as a tonic, demulcent in catarrh of respiratory treat and genito-urinary passages and as laxative Roots contain a tribasic acid glycyrrhizic acid and potassium and calcium salts, glucose, sucrose and fat o 8 per cent. Used in medicine to cover the acrid taste of many nauseous drugs.

Gymnema sylvestre R Br (Vern —Gur mar, S -Meshasringi, H & B —Merasingi, Bo -Kavali, M -Shiru kurunja)

Grows in Central and Southern India and in Tropical Africa It is used as stomachic, diuretic, antiperiodic, antidiabetic and in urinary disorders It contains resins, a glycoside named as gymnemic acid, a bitter principle and some tartaric acid and calcium oxalate Experiments on animals showed no reduction of blood sugar The leaves stimulate heart and circulatory system and increase urine secretion Clinical trials showed no effect on blood or urine sugar of diabetic patients

Helicteres isora Linn (Vern —Kapasi) The East Indian screw tree

Is found throughout Central and Western India It is chiefly used in intestinal disorders such as colic, flatulence and diarrhoea It contains a demulcent substance and tannins The pods are used even today in some parts of India in the treatment of chronic diarrhoeas but clinical trials showed no marked action

Hemidesmus indicus R Br (Vern —S -Ananta, H -Magrabu, B. Anantamul, M.-Nannari) Indian Sarsaparilla I.P C.

Grows in Northern India, common in Bengal and in the Deccan, extending to Travancore and Ceylon and Bombay State. Used in the treatment of nutritional disorders, syphilis, chronic rheumatism skin affections and as a blood purifier Roots contain a steroptene, an essential oil, sterols, resin, tannin and small amount of glycosides Clinical trials show that its medicinal value is in no way inferior to imported Sarsaparilla.

Holarrhena antidysenterica Wall (Vern —Dhudi, S -Kutaja, H.-Karchi, B -Kurchi, Bo.-Pandhrakura, M.-Kashappuvetpalarishi) B P., B.P C , I P.C. & I P L

Grows throughout India upto an altitude of 3,500 ft and even as far south as Travancore Bark and seeds are used as a powerful antidysenteric, astringent, febrifuge, anthelmintic, carminative and aphrodasiac Bark and seeds contain kurchine, conessine, kurchicine and holarrhine Conessine is toxic to protozoa and is locally irritant The circulatory system is depressed and intestinal movements are stimulated, it has narcotic action on frogs The total alkaloids have similar action The extract from bark (containing one grain of total alkaloids in 4 c.c) given in doses of 2 drams 3 times a day has beneficial effect in amoebic dysentery especially the chronic form. It has proved effective against trichomona infestations.

Holarrhena febrifuga Klotizseh

Grows in Kenya, Tanganyika, Northern Rhodesia and Nyasaland. Used in the treatment of influenza, bilharzia, syphilis and also as substitute for quinine. Stem bark and leaves contain conessine 6.0 per cent and a subsidiary norbase isoconessiomine. Further studies indicated.

Hydnocarpus anthelmintica Pierre.

It grows in Siam, Northern Cochin and Gamboja

Hydnocarpus wightiana Blume (Vern.—Janglibadam, Kowti, Bo-Kava, M.-Yetti). B P., I P.C & I P L

Grows in western parts of the Indian Peninsula from South Konkan along the coastal range. The oil has been used in the treatment of leprosy for centuries Seeds contain Chaulmoogra oil and Palmitic acid. Chaulmoogra oil has a bacteriostatic action on acid fast bacilli It is extremely irritant to the tissues and for this reason ethyl esters such as moogrol (British) and Antileprol (German) are used by injection in the treatment of leprosy

Hyoscyamus niger Linn. (Vern —S -Parasikaya, H -Khurasani-ajvayan), B -Khorasani ajowan, Bo -Khorasani-owa, M -Khorasani-yomam) B P, B P C & I P C

Grows wild in temperate Himalayas at altitudes of 6,000 to 12,000 ft and also cultivated. Leaves contain alkaloids hyoscine and hyoscyamine

Ipomoea hederaceae Jacq & **I turpethum** Br (Vern —H , B & Bo -Kaladana, M -Jirkiviral) B.P C, I P C & I.P.L

Found throughout India ascending to altitudes of 3,000 ft The resinous substance from the root bark is an excellent substitute for Jalap which is a drastic purgative It is official in the I P L

Juniperus communis Linn. (Vern. H -Aaraar, P -Petthri) I.P C

Found in the Western Himalayas, Kumaon and Kurram valleys at an altitude of 11,000 ft above the sea level It is used as digestive and diuretic. Berries contain a voltaile oil 0 2 per cent , invert sugar about 33 per cent , resin 10 per cent , a bitter principle, organic acids and wax Beverage gin is alcohol flavoured with juniper berries or the essential oil

Luffa acutangula Roxb. (Vern. S -Koshataki, H & Bo -Torai, B.-Jhinge, M -Pikumkai)

Grows in N W India, Sikkim, Assam and East Bengal and also cultivated Leaves used locally in splenitis, haemorrhoids, leprosy and granular conjuctivitis Fruit contains a bitter substance luffein and seeds contain a fatty oil and a saponin. The latter is toxic to frogs and has a haemolytic action. It shows digitalis-like action and produces vomiting and diarrhoea. Further work is indicated

Luvunga scandens Ham. (Vern.—Jivanti, S -Lavangalata, B -Labangaphal).

Grows in Eastern Bengal, Assam, Khasia hills, Chittagong & Burma Roots and berries are used to cure tuberculosis of lungs, billiousness

and fevers. Berries contain four crystalline neutral substances. Further chemical and pharmacological investigations are indicated

Madhuca indica J. F. Gmel. Syn -*M. latifolia* Mcbrd. The Mahu tree. (Vern -Mohua). I.P.C.

Grows in forests of Madhya Pradesh and is cultivated all over India. Also found in Uttar Pradesh, Bihar, Gujrat and Bombay Its flowers, seeds and timber are of economic value The seed-oil is edible and is also used for soap manufacture. The seeds contain 50 to 55% of a fatty oil and a saponin Leaves contain a glucosidic saponin and traces of an alkaloid. Flowers contain sugar and enzymes The plant is used as an astringent It is largely employed as a lotion for chronic ulcers and bleeding spongy gums. Also considered beneficial in chronic bronchitis and chronic rheumatism. No proper clinical trials have been carried out.

Madhuca longifolia Linn. Macbd (Vern.—Mohua) Syn -*Bassia longifolia*. I.P C

Is found in forests of Western India, Kanara, Malabar and Mysore in the same way as of *Madhaca latifolia*. Seeds contain 40 per cent. of fatty oil and sapo-glucoside called 'Mowrin' Fruit contains saccharose 4 6 to 16.2 per cent. and maltose 2.39 per cent , tannins and enzymes. Acts in the same way as *M. latifolia* The sapo-glucoside acts on the heart in the same way as digitalis. Requires further investigation

Mahonia nepalensis DC. Syn.-*Berberis nepalensis* Spreng (Vern.— Chiror Anrudana, P.-Amudanda, Nep.-Chatri)

Grows in the outer Himalayas from Ravi eastward to Khasia and Naga Hills also in the Nilgiris at an altitude of 5,000 ft It contains umbellatine

Mallotus philippinensis Muell -Arg (Vern —Kambila, S -Rechanaka, H. & B.-Kamala, Bo -Shendri, M -Kapila) I P C

Grows throughout the plains of India and Ceylon, tropical parts of Asia and Australia, Bombay and Bengal It is used as anthelmintic especially against tape worms Fruit contains a brownish-red resin called rottlerin, an impure rottlerin, traces of a volatile oil, starch, sugar, tannins and oxalic acid. It has toxic action on frogs, tadpoles and worms. The drug has a marked cathartic and anthelmintic action in doses of 2 to 3 drams.

Melia azedarach Linn. (Vern.—Drek)

Grows wild in Sub-Himalayan tract at altitudes of 2,000-3,000 ft and is also cultivated Root bark, fruit, flowers and leaves are considered hot, dry and to have deobstruent and resolvent properties.[5] Juice of leaves used as anthelmintic, diuretic and emmenagogue It contains a resin and the fruit a bitter substance. No systematic phamacological studies or clinical trials have been carried out

Mentha arvensis Linn (Vern.—H.-Podina, B. & Bo.-Pudinah. M.-Pudina) The marsh mint B.P., B P.C , I.P C. & I.P.L

Grows abundantly in northern and western Himalayas and in Kashmir at altitudes of 5,000 to 10,000 ft It is used as a flavouring agent and as carminative and stimulant Yields an oil similar to peppermint oil, to the extent of 0 18 to 0 2 per cent The oil contains 4 to 14 per cent. esters as menthyl acetate and 46% of free menthol Official in I.P.L

Moringa oleifera Lam Syn -*M pterygosperma* Gaertn. (Vern — Sanjna, Saonjna). I P C.

The plant is found wild in the Sub-Himalayan tract from the Chenab to Oudh and is also cultivated throughout India The root of this tree is extensively employed in intermetent fevers, epilepsy, hysteria, palsy, chronic rheumatism, dropsy etc The root contains physiologically active bases and an antibiotic substance pterygospermin

Myristica fragrans Houth (Vern —S -Jatifalam, H , B. & Bo.-Jayphal, M -Jadikkay). The nutmeg B P , B P C , I.P.L & I.P C.

Grows in the Nilgiri hills and on the Malabar coast Oil is used in the manufacture of pharmacopoeial preparations and aperient pills to prevent griping Nutmegs yield 5 to 15 per cent of a volatile oil and 30 to 40 per cent of fats, also phytostirin, starch, amylodextrin, colouring matter and a saponin. It is largely used for flavouring and as a carminative.

Myrsine africana Linn (Vern Gugul, H. Chapra).

Grows in Afghanistan, salt range and the Himalayas from Kashmir to Nepal at altitudes from 1,000 to 8,000 ft. Berries used as anthelmintic and as a remedy for skin diseases They contain embellic acid 3 per cent. and quercitol 1 per cent No pharmacological work or clinical trials have been carried out

Oldenlandia auricularia Schum Syn -*Hedyotis auricularia* Linn. (Vern.-Gatta colla, B.-Muttia-lata)

Grows in Eastern Bengal and from Nepal, Sikkim and Khasia hills to Assam, Chittagong, Manipur and Burma. Also found in Kanara and Ceylon Used in the treatment of dysentery and diarrhoea Roots contain a fatty oil and glucosides, also reducing sugars, colouring matters, tannin, alkaloids auricularine, heydyotin and albumin No pharmacological work has been done.

Papaver somniferum Linn. (Vern —S -Ahiphena, H & B -Afim, Bo -Aphu, M -Postakatol). The Opium B.P., B P C , I P C & I P.L.

Cultivated in many parts of India especially in Utter Pradesh, Bihar and other parts of the world, China, Asia Minor, Turkey, Persia and Egypt. Dried inspissated juice obtained by incising the green fruit is opium which is used in the treatment of dysentery and diarrhoea as an anodyne sedative and euphoric Opium contains about 25 alkaloids, the important ones being morphine, codeine, thebaine, papaverine and narcotine In addition to these acetic, lactic, sulphuric and meconic acids, gummy and pectinous substances, albumin, wax, fat, resin, etc. also occur.

Opium is a soverign remedy for relieving pain and producing sleep. Opium and morphine are habit forming drugs and produce addiction

Peganum harmala Linn (Vern —H. & Bo.-Hurmal, B.-Isband, M.-Shimai-azha-vanai-virai) I.P C

Grows wild all over N.W India, Sind, Punjab and Kashmir, Agra and Western Deccan Also found in Arabia, North Africa, Hungary and Spain. Seeds are used as alterative, aphrodisiac, lactagogue and anthelmintic. Contain alkaloids harmine, harmaline and harmalol to the extent of 4% The alkaloids have some anthelmintic action and depress respiration and heart Clinical trials revealed no effect in malaria. The seeds have been used to expel tape worms

Peucedanum graveolens Benth (Vern —S -Shatapuspi, B.-Soolpha, Bo -Balunt-shep, M -Satakuppi)

Grows in central and southern Europe and in tropical and subtropical countries. In India, it is cultivated as a cold weather crop. Used as a condiment, carminative, digestive and flavouring agent Largely used in soap manufacture as a perfume Yields 0.06 per cent. of an essential oil

Picroena quassioides Benth. (Vern Bharangi) I P C & I P L.

Found in the outer Himalayas from Kashmir to Nepal and also in China. Bark and leaves used as febrifuge and insecticide. Contains bitter principles quassin and neo-quassin and a bitter alkaloid 0.05 per cent. Quassia is a popular bitter

Picrorhiza kurrooa Benth (Vern —S -Katuka, H & B -Katki, Bo -Kalikutki, M -Katuka-rogani). B P.C , I P C & I P.L

Grows in the North West Himalayas from Kashmir to Sikkim at altitudes of 5,000 to 11,000 ft. Used as a bitter tonic and stomachic. Contains 26 6 per cent. of a bitter substance and a glucoside Can be used in medicine as a bitter if properly standardised.

Pimpinella anisum Linn. (Vern.—S -Shetapuspa, H.-Saonf, B.-Muhuri, Bo.-Sonf, M.-Shombu). The Anise B P , B P.C. & I.P C.

Grows in Europe, Russia and Middle East and many other parts of the world In India, found in Uttar Pradesh, the Punjab and Orissa. Used to prevent flatulence and intestinal colic and as an aromatic. It contains an essential oil which is used in medicine to prevent flatulence, intestinal colic and as a flavouring agent.

Pinus roxburghii Sargent Syn.-*P longifolia* Roxb (Vern -Chir, S -Sarala) Chir Pine. B.P , I.P C. & I.P L

Grows in abundance in U S.A. and France. In India, it is found in the temperate Himalayas from Kashmir to Burma at altitude from 2,000 ft. to 6,000 ft. The oil is used in the manufacture of paints and varnishes and colophony for plasters in medicine The oleoresin yields about 20 per cent. of turpentine oil and about 80 per cent. of residue named as colphony which is used in medicine for preparation of plasters and ointments. Oil of turpentine is used externally as a counter-irritant

and rubefacient. Small doses of oil are given internally in bronchitis and larger doses as an anthelmintic.

Piper betle Linn. Betel leaf. (Vern.-S.-tambula, H., B. & Bo.-Pan, M.-Vettilai) I P C.

It is cultivated in Madras, Central Provinces, Bengal, Orissa, Bombay, U. P. and Burma. Leaves are used as an aromatic, stimulant, carminative, astringent and aphrodasiac Contain an essential oil, starch, sugar, tannins and diastases No pharmacological studies have been carried out Leaves with other substances such as betel nut, lime, catechu, etc. are widely chewed in India They act as mental stimulant and digestive

Piper cubeba (Vern.—S -Sugandha muricha, H., B. & Bo.-Kababchini, M -Val milaku) Cubebs. I P.C & B P.C

Found in Java, Sumatra and Malaya Archipelago and cultivated to a small extent in India in Mysore Fruit is used as a condiment and in the treatment of genito-urinary diseases, cystits, gonorrhoea and gleet Contains an essential oil upto 10 15 per cent.

Pistacia integerrima Stew (Vern —Kakra, S -Karkata sringi, H & B.-Kakra shingi, M -Kakkata shingi). I P C

Is found in the sub-alpine Himalayas. It is used as a remedy for cough, phthisis and asthma in doses of 20 grs combined with aromatics Galls contain an essential oil 1 21 per cent a crystalline hydrocarbon 3.4 per cent tannins and gum mastic 5 0 per cent No systematic investigations have been carried out Further clinical trials are needed

Plantago ovata Forsk. Ispaghula (Vern.—H -Ispaghul, B -Isabgul, Bo.-Isabghol, M.-Ishappukol-virai). B P C , I P.C. & I P L.

Grows in the plains of the Punjab and Sind and low hills from Sutlej westwards It is also cultivated in Bengal, Mysore and Coromandel Coast Seeds are cooling and demulcent and are used in dysenteries, diarrhoea and other inflammatory conditions of digestive and urinery tract. Contain a fatty oil, albuminous matter and mucilage Some species contain a glucoside named aucubin, which is pharmacologically inactive. The mucilage from seeds inhibits growth of common intestinal organisms in the gut The colloidal nature of the mucilage has a remarkable power of adsorbing bacteria and toxins and is soothing. The seeds are prescribed in chronic bacillary and amoebic dysentery, chronic constipation and chronic diarrhoea in doses of 1—2 ounces soaked in water. The pericarp of seeds separated is more convenient for use and is taken in doses of 2 to 3 teaspoonsful once or twice daily.

Plumbago indica Linn Syn.-*P. rosea* Linn. (Vern.-S.-Chitraka, H., B. & Bo.-Lal chitra).

Commonly cultivated in gardens of India. Root is used as abortifacient and vesicant in rheumatic affections of joint and paralytic conditions Contains, plumbagin, the active principle, amorphous brown pigment and a reducing sugar. No systematic pharmacological investigations have been carried out.

Plumbago zeylanica Linn. (Vern.—S.-Chitraka, H. & B.-Chita, Bo.-Chitaro, M.-Chittira)

Grows throughout India, particularly in Uttar Pradesh, Bengal and Southern India. Root is used in digestive disorders, piles, anasarca and skin diseases and as abortifacient It contains an active principle named as plumbagin, which stimulates the central nervous system, heart and plain muscles in small doses Larger doses have a depressant action. Systematic clinical trials are needed to establish its therapeutic value.

Podophyllum hexandrum Royle. (Vern.—H.-Papra). B.P., B.P.C., & I.P.C.

Grows in the temperate forests of the Himalayas from Sikkim to Kashmir at a height of 7,000 ft Used as a drastic purgative and as a cholagogue Contains a resin 8 to 13 per cent, flavonol, quercetin, starch and calcium oxalate. Podophyllum is irritant to the eyes and mucuous membrane. It is an active purgative and is administered in average doses of 0.01 gm.

P. paltatum contains podophylotoxin and α & β paltatins, the farmer has been shown to be present in *P. hexauclum*, but not paltatins. The resin is effective in destroying soft warts or condylomata.

Pongamia pinnata Merr. Syn -*P. glabra* Vent (Vern —Kanga, S., H. & Bo -Karanja, B.-Dahar karanja)

Grows all over India especially near the Coast, from Central and Eastern Himalayas to Ceylon The seeds and leaves are used in treatment of skin diseases and rheumatism Internally the seed oil is used as stomachic, cholagogue, febrifuge and expectorant It is bitter fatty oil forming 27 to 36 4 per cent. of seeds containing traces of an essential oil It also contains two crystalline compounds named as karanjin and pangamol No systematic pharmacological investigations or clinical trials have been carried out

Premna integrifolia Linn. (Vern.—S.-Ganikarika, H. & Bo.- Arni, B.-Ganiari, M -Munni-vayz)

Grows near the sea from Bombay to Malacca and in Ceylon. Root is used as laxative, carminative and stomachic. The decoction of whole plant is used in the treatment of fevers and rheumatism and neuralgia. Stem bark contains three alkaloids, premine, ganiarine, ganikarine and some unsaturated aromatic hydrocarbons The alkaloids ganiarine and premnine have a sympathomimetic action Further work is indicated.

Pristimera indica Willd Syn.-*Hippocratea indica* Willd.

Grows wild in forests from Konkan to Madras in South Bengal and Assam. Paste from roots used as a remedy for respiratory troubles Root contains dulcitol and an antibiotic pristimerin which is effective against gram positive organisms Preliminary clinical trials revealed that pristi merin is effective against throat affections Further investigations are in progress.

Psoralea corylifolia Linn. (Vern —S -Vakuchi, H. & B.-Babachi, Bo -Bobawachi, M.-Karpo-Karishi). I P.C & I P L.

Grows all over the plains of India. The seeds are considered anthelmintic, diuretic, diaphoretic and a remedy against skin diseases and leprosy. They contain an essential oil a fixed oil, a resin and traces of an alkaloidal substance. The oil has an irritant action on skin. It is

toxic to paramoecia and bacteria Plain muscle is stimulated and areterioles are dilated An oleoresinous extract made from seeds has beneficial effect in leucoderma when applied locally. Seeds are also taken internally in this affection.

Randia dumetorum Lamk. (Vern.—S.-Madan, H.-Manphal, B -Menphal, Bo.-Gelaphal, Maruk-kallan-kai).

Grows throughout India. Fruit is used as an emetic and aperient and as a substitute for ipecacuanha Contains a small quantity of an alkaloidal substance, Randia saponin and a glucoside. Randia tannic acid and Randia fat are also present. The drug produces irritation of the mucous membrane and haemolysis both in vitro and in vivo. Frog's heart is arrested when perfused with drug solution in concentration of 1 in 50,000. No clinical trials have been carried out.

Rauwolfia canescens Linn

Grows wild in Howrah district near Calcutta in Bengal. Roots used to adulterate those of *R serpentina* Root bark stem bark, and leaves contain an alkaloidal body named as rauwolscine The alkaloid is depressant to the cardio-vascular system It is also sympatholytic in action and abolishes the pressor effects of adrenaline Further work is in progress

Rauwolfia serpentina Benth. (Vern —S.-Sarpagandha, H -Chota chand, B. & Bo -Chandra, M.-Covannamilpori) B.P.C , I P.C. & I.P L

Grows near the foot of hills from Moradabad to Sikkim, in Assam, Pegu, in Southern India along the Ghats , also occurs in Java and Malaya. Root is used as an antidote in snake bites and as a remedy for insomnia, hypochondria and insanity It contains alkaloids, ajmaline, ajmalinine, serpentine and serpentinine Extracts containing the total alkaloids, including serpentine, lower blood pressure in animals , ajmaline and serpentinine raise it Recently a new alkaloid, Reserpine, has been isolated which has a marked sedative & hypotensive action. Doses of 20 to 30 grains of powdered root produce sedative effect and lowers blood pressure especially the diastolic. Considerable work is being carried out on this drug in Europe and America on account of blood pressure lowering properties and sedative effects

Rheum emodi Wall (Vern —H & B -Revandchini, Bo -Iadakirevanda chini, M -Nattu-ireval-chinni) Indian Rubarb I P.C. & I.P L.

Grows wild in the Himalayas in Nepal and Sikkim at altitudes of 4,000 to 12,000 ft Used in ailments of children and as a purgative Contains derivatives of anthraquinone to the extent of 2.0 to 4.5 per cent gallic acid, tannin, starch, fat, dextrose and levalose, pectin and calcium oxalate

Ricinus communis Linn (Vern.—S.-Eranda, H.-Arand, B -Verenda, Bo.-Erendi, M -Amanakham chedi) Castor seeds. B.P., B.P.C. & I.P.C.

Cultivated throughout India particularly in Madras, Bombay and Bengal Seeds contain a fixed oil which consists of ricinoleate of glycerol with a small quantity of palmitin and stearin and a toxic substance ricin. Oil from seeds is used as a purgative

Rosa damascena Mill. (Vern. –H.- Gulab ka phul, S.- Satapatri, B.- Golapphul, Bo.-Gul, M.-Golappu). The Rose. B.P. & B P.C.

Chiefly cultivated in Bulgaria In India it is cultivated in the Punjab, United Provinces, Bihar and Orissa Rose water is used for eye lotions and the oil as a flavouring agent Flowers contain a pleasant smelling essential oil used as perfume, and fruits contain Vit. C.

Santalum album Linn. (Vern —S -Swet-chandan, H.-Safedchandan, B.-Sadachandan, S -Mhandanak-kattai). B P., B. P. C & I. P. C.

Grows wild and is cultivated in Mysore State, Coorg, Coimbatore and Southern parts of Mysore. Wood is used as bitter, cooling, astringent and useful in biliousness, fever and thirst. Paste made from wood as a cooling application for skin affections. The heart wood of the tree yields an essential oil 2 5 to 6 per cent which is used for flavouring purposes and as urinary disinfectant

Saraca indica Linn. (The Asoka tree) Vern —S -Asoka, H , B. & B.-Asok, M -Asek) I P C.

Grows in Bengal, South India, Arakan, Tenasserim and is also cultivated in many parts Bark is used as astringent, uterine sedative in menorrhagia and as a remedy for piles and dysentery It contains tannins and catechol. Various fractions isolated from the bark produced no marked action on uterus but faith in its beneficial effects in uterine disorders persists.

Saussurea lappa Clarke Comp The Costus (Vern – S -Kushtha, H.-Kut, B.-Pachak, Bo -Ouplate, M -Goshtam). I.P C & I.P.L.

Grows in N W Himalayas especially on the slopes of mountains round the Kashmir valley upto an altitude of 13,000 ft. Root is used as hairwash to kill lice and as insect repellent, carminative, stimulant and antiseptic Root contains an essential oil, 1 5 per cent., an alkaloid named as sausserine, resin 6 0 per cent , traces of a bitter substance, tannin, inulin 18 per cent, a fixed oil and potassium nitrate Leaves contain only the alkaloid The essential oil has strong antiseptic and disinfectant properties. The circulatory system is stimulated and smooth muscle of bronchi and intestines is relaxed The alkaloid produces relaxation of bronchioles and has a general depressant action in involuntary muscles The heart is stimulated An alcoholic extract of root in 1/2 to 2 dram doses relieves asthmatic attacks

Scilla indica Baker (Vern.—H. & B -Suphadie khus, Bo.-Bhui-kanda, M -Shirunari-vengayam).

Grows frequently in sandy places near the sea in the Deccan Peninsula from the Concan and Nagpur southwards Used as expectorant, cardiac stimulant and diuretic. Squill contains a number of cardiac glycosides Clinical trials have revealed that it has beneficial effect on patients suffering from bronchial catarrh and chronic bronchitis

Semecarpus anacardium Linn. The marking nut tree (Vern —S.-Bhallatamu, H & B.-Bhela, Bo.-Biba, M.-Shayrang)

Grows in Sub-Himalayan tract from Sutlej eastwards down to Assam at an altitude of 3,500 ft. Juice is used in preparation of a varnish, fruit is used as a substitute for marking, counter irritant in rheumatism, sprains

and leprotic nodules Internally juice is used in syphilis, scrofulous affections, piles and nervous debility. The pericarp of the fruit contains 20 per cent. of an oil, which gives bhilawanol and other compounds No systematic investigations have been carried out to determine its beneficial effects in treatment of disease.

Sida cordifolia Linn. (Vern —Khareti, S -Bala, H -Kungyi, B.-Brela, Bo -Chikana, M.-Chiribenda). I P C.

Grows throughout tropical and sub-tropical India and Ceylon. Root is used as cooling, stomachic, tonic, febrifuge and diuretic The seeds are considered aphrodisiac and are also used in the treatment of cystitis, rheumatism and spermatorrhoea. The seeds contain fatty oil, and an alkaloid (0 085 per cent) recognised as ephedrine, which probably is responsible for its cardiac stimulant effects.

Skimmia laureola Sich & Zucc (Vern -Bassu, Nep.-Chumlani, P.-Ner)

Grows throughout temperate Himalayas from Kashmir to Khasia hills The leaves are used as an incense and burnt near smallpox patients Leaves contain an essential oil with pleasant smell and an alkaloidal substance. No systematic pharmacological studies have been carried out. The essential oil could be used as perfume in soap and cosmetic industry.

Sonneratia acida Linn (Vern -B -Archaka)

Grows in Bengal and the Deccan Peninsula. Used as a poultice in sprains and swellings Juice is used for stopping haemorrhage Contains colouring matter, archin and archinin and a crystalline compound archicine. No pharmacological studies have been carried out

Stephania glabra Roxb. Miers (Vern -Pusha)

Grows in tropical and temperate Himalayas ascending to an altitude of 7,000 ft from Sindh eastwards to Khasia hills and Pegu Root is used as a remedy against phthisis, asthma, dysentery, fevers and intestinal complaints. Tubers contain three alkaloids, gindarine, gindaricine and gindarinine. No pharmacological and clinical studies have been carried out.

Strychnos nux-vomica Linn. (Vern —S -Visha-mushti, H -Kuchla, B -Kuchila, Bo -Kajra, M.-Yetti) B P., B P.C , I P C & I P L

Grows wild throughout tropical India upto an altitude of 4,000 ft in Orissa, Madras, Cochin, Travancore and the Coromandal Coast Seeds are used in the treatment of dyspepsia and diseases of the nervous system Fruit, bark, leaves and wood contain strychnine, brucine, vomicine, etc Seeds contain 1.53 to 3.42 per cent. of the total alkaloids of which half is strychnine

Swertia chirata Ham. (Vern.– S -Kirata, H -Charayatah, B -Chireta, Bo.-Chiraita, M.-Nila-Vembu) Chiretta. B.P C., I.P C & I.P.L.

Grows abundantly in the temperate Himalayas from Kashmir to Bhutan and Khasia Range between altitudes of 4,000 to 10,000 ft. Used as a bitter tonic, stomachic febrifuge and anthelmintic. Contains a bitter glucoside chiratin, resins, tannin and ash 4 to 8%.

Symplocos racemosa Roxb. (The Lodh tree) Vern —S -Lodhra, H. B. & Bo -Lodh, M.-Ludduga). I.P.C.

Grows in plains and lower hills of Bengal, Assam and Burma. It is also found in dry forests of Chhota Nagpur plateau Bark and leaves are used for dyeing purposes. The bark is used as astringent in bowel complaints, eye diseases and ulcers Bark contains three alkaloids, loturine 0.24 per cent. collutrine 0 02 per cent and loturdine 0 06 per cent No systematic pharmacological investigations or clinical trials have been carried out.

Terminalia arjuna W. and A. (Vern.—S & Bo -Arjuna, H & B -Arjun, M.-Vellai-maruda-maram). I P C.

Grows throughout the Sub-Himalayan tracts of United Provinces and in the Deccan, Southern Bihar, Chhota Nagpur, Burma and Ceylon Bark is used as a cardiac tonic in heart diseases and contains large quantities of calcium salts, about 12 per cent. tannins, an organic acid, an ester and some colouring matters and sugar. It is reputed to have a stimulant action on heart Clinical trials however showed no beneficial effects in heart disease. Further investigations needed

Thalictrum foliolosum D C. (Vern —H -Pinjari, P -Gurbiani, Bo.-Mamiran).

Grows in Khasia hills between altitudes of 4,000 ft to 6,000 ft. It is used as a bitter tonic and laxative. It is a cheap but valuable substitute of Mamira used in preparation of eye solves It contains two alkaloids berberine and thalictrine No proper investigations have been carried to study its efficacy in conjunctive affections

Thevetia peruviana Schum Syn -*T neriifolia* Steud See poisonous plants of India (Vern -H & Bo -Pila-kaner, B -Kolka-phul M -Pachchai-alari)

Grows in plains all over India, but originally a native of west Indies. Seeds are used as abortifacient and as poison for man and cattle They contain 57 per cent of a colourless oil, which yields glucoside thevetin which is not toxic to unicellular organisms The drug is toxic to higher animals and is not likely to be of therapeutic value because of powerful toxic effect on heart muscle

Tinospora cordifolia Miers (Vern — Giloe, S -Guduchi, H & B -Gulancha, Bo -Gulwail, Bo -Gulwail, M -Sindilkodi) I P C & I P L

Grows in Uttar Pradesh It is used in treatment of rheumatism hyperacidity and dyspepsia and as a tonic, antiperiodic and diuretic. Stems contain glucoside giloin and non-glycosidic bitter, gilenin and gilosterol No Pharmacological investigations or clinical trials have been carried out

Toddalia asiatica Lam. Syn -*T aculeata* Pers. (Vern —S -Kanchana, H -Kanj, B.-Koda-todali, Bo.-Jun-li-kali-mirchi, M Milkaranai)

Grows in the Nilgiris and sub-tropical Himalayas from Kumaon to Bhutan upto an altitude of 3,000 ft. Root bark is considered to be antimalarial. The plant contains two alkaloids, todaline and todalinine. Leaves contain an essential oil and root bark contains an essential oil;

resin, bitter substance, citric acid, pectin and starch. A freshly prepared infusion of the plant is very feebly toxic. The alkaloid todaline has no antipyretic action The root bark has no antimalarial properties.

Trachyspermum ammi (Linn) Sprague, Syn.-*Carum copticum* Benth (Vern -S -Yamani, H. & Bo.-Ajowan, B -Jowan, M.-Oman) B.P. & I.P.C.

Widely cultivated all over India. Seeds are used in diarrhoea, dyspepsia, cholera, colic, flatulence and indigestion and as tonic. They contain thymol

Tribulus terrestris Linn. (Vern.—Gokhru, S.-Gokshura, H.-Chota-gokhru, B -Gokhuri, Bo.-Lahana-gokhru, M -Nirunji) Small Caltrops I.P.C

Grows throughout India and Ceylon. Fruit is used as diuretic, tonic, aphrodisiac and in the treatment of urinary disorders and impotency. The fruit contains an alkaloid in traces, a fixed oil, an essential oil, resins and nitrates Preliminary investigations revealed that the alkaloid produces slight rise of blood pressure and appreciable rise in kidney volume. Clinical trials confirm its diuretic action. Further investigations are needed.

Tylophora indica Merr Syn -*T asthmatica* W. & A (Vern.—H. & B.-Antamul, Bo.-Anthalmul M -Nay-palai) I P C

Grows in North and East Bengal, Assam, Kachar, Chittagong and the Deccan Peninsula upto a height of 3,000 ft. Root and leaves have been used as a substitute for Ipecacuanha in dysentery catarrh and other affections. Root contains two alkaloids tylophorine and tylophonine The drug produces dermatitis on contact The alkaloid is toxic to protozoa and has a depressent action on the heart No clinical trials have been carried out Deserves further study

Urginea indica Kunth. (Vern —S -Vana—palandam, H & B.-Jangli piyaz, Bo -Jangali-kanda, M -Nari-vengayam) B.P.C & I.P.C.

Grows in the Western Himalayas upto 7,000 ft., shores of Coromandal, Konkon and Western Ghats. Contains glycosides scillaren-A and scillaren-B, musilage, carbohydrates, a phytosterol and calcium oxalate. Used as cardiac stimulant and expectorant.

Valariana wallichii D C (Vern —S -Tagara, H. & B.-Tagar, Bo.-Tagar-ganthoda) B.P C & I P.C.

Grows in the temperate Himalayas from Kashmir to Bhutan at altitudes ranging from 4,000 to 12,000 ft Used in the treatment of hysteria and nervous troubles of women Contains 0 3 to 1 0 per cent of volatile oil containing esters of iso-valerianic acid and formic acid.

Vanda roxburghii R. Br (Vern.—S , H., B & Bo.-Rasna, M.-Knapa-chettu).

Grows in Bengal, Chota Nagpur, Madhya Pradesh, Cochin and Travancore. Roots considered beneficial in rheumatism, diseases of nervous system and in secondary syphilis. It contains a glycoside tannin, saponins, sterols, fatty oil and resins The glycoside, has low toxicity and stimulates all organs having cholingeric nerve supply. The heart is slowed and cardiac output is diminished. No clinical trials have been carried out.

Vitex peduncularis Wall. (Vern —H.-Nagbail, B.-Goda, M.-Navaladi). I.P.C.

Grows in Bihar, Eastern Bengal and Madhya Pradesh. The leaves and roots are used in the treatment of malarial and black water fevers Leaves contain a glycoside. Clinical trials with infusion of leaves showed no beneficial effect in malaria or black water fever.

Withania somnifera Dunal (Vern —S., B. & Bo -Ashwagandha, H -Asgandh, M -Amku lang-kalang) I.P.C.

Grows throughout the drier parts of India, Baluchistan and Ceylon. Leaves and roots are used in the treatment of tumours tuberculus glands, bronchitis, rheumatism and dyspepsia The plant contains pot nitrate, tannin, colouring matter, glucose and an alkaloid. It has a mild sedative effect on animals Deserves further study

Xanthium strumarium Linn (Vern —S -Aristha, H Chhota-gokru, B -Bon-okra, Bo -Shankeshvara, M -Marlu-mutta) I P C

Grows throughout the hotter parts of India upto an altitude of 7,000 ft. The whole plant is used as diaphoretic and sedative Root is used as bitter tonic and in the treatment of cancer and strumous diseases. It contains fat 38.6 per cent., albuminoids 36 6 per cent and a glycoside Xanthostrumanin 1 3 per cent The glucoside is physiologically inactive. No clinical trials have been carried out.

Zingiber officinale Rosc. (Vern —S.-Adrakam, H -Adrak, B -Ada, M.-Inji) B.P & I.P C

Cultivated in Cochin, Bengal, Bombay, Kumaon and many other parts of India Used as a condiment in food and as carminative and stimulant of the gastro-intestinal tract. Contains 0 25 per cent. of a volatile oil of light yellow colour, resinous matter, starch and mucilage.

CHAPTER VII

I. TOXICOLOGICAL ASPECTS OF INDIAN PLANTS

The second line of basic research has been in connection with the Poisonous Plants of India including the group of insecticides, insect repellent etc Here we will briefly refer to the present position with regard to the toxicological aspects of Indian Plants.

(1) CRYPTOGAMS OR THE FLOWERLESS PLANTS

The toxicological aspects of the Cryptogams are little known so far as India is concerned

(a) **Bacteria.** The bacteria are among the simplest form of plant life and are met with universally. The majority of them are harmless, but some are injurious to man and animals They produce deleterious effects in two ways —Firstly as parasites, when they derive nourishment from living animals and many of them produce, within the body, toxins which are harmful Secondly many saprophytic bacteria produce poisonous substances, especially such as those occuring in putrid flesh, fish and other decaying organic matter It is not our intention to give details of toxic properties of this group These are fully discussed in books on bacteriology Bacteria are a class by themselves and although they belong to vegetable kingdom do not come under our purview here.

On the other hand a number of organisms of this group produce antibiotic substances which are of very great value in the treatment of disease Streptomycin has been derived from *Streptomyces griseus* (*Actinomyces griseus*), Aureomycin from *Streptomyces aureofaciens*, Terramycin from *Streptomyces rimosus*, Chloromycetin from *Streptomyces veneznelae*, Bacitracin from strains of *B. subtilis*, etc.

A large amount of work is now being done in this country on the isolation of antibiotic substances from various soil and other micro organisms found in this country.

(b) **Algae** The algæ that cause poisoning are mostly those which are found in stagnant waters. The normally offensive odour may be sufficient to indicate their presence, but only a microscopic examination can determine just what the forms of algæ present may be. Blue-green algæ, as a group, are perhaps the most pronounced in their toxic effect. Parker and other workers have shown that when odours in water are pronounced, these microscopic organisms are present in considerable numbers He claims that of the organisms which produce objectionable and deleterious qualities in water, the microscopic ones are the more important and very few cases have been observed in which really serious trouble in water supplies could be attributed directly to the growth of larger plants In any study of the algæ from this point of view, however, account must be taken of the products of decomposition by the associated bacteria, since poisoning may be produced by the toxins produced by bacteria rather than by the algæ.

Certain algæ, such as *Microcystis flos-aquae* (Wittr.) Kirch, *Aphanizomenon flos-aquae* (Linn.) Ralfs. and species of Anabœna, etc. form on the surface of water what is generally called water bloom. The presence of water bloom on the surface of lakes, ponds, and other open sheets of water is distasteful to bathers and obnoxious to those living in the vicinity. Livestock compelled to drink water containing water bloom are reported to have suffered from poisoning. In Minnesotta, (U.S.A.) during recent years, horses, cattle, sheep and donkeys have died in large numbers on the shores of lakes where water bloom is present. All the above mentioned algæ forming water bloom have been recorded in various parts of India but no work has been done in connection with their toxic effects. According to Dr Bhardawaja of the Benaras Hindu University, water blooms containing these species occur commonly on the surface of many temple tanks in different parts of India. Of the other possibly harmful algæ may be mentioned species of *Nodularia, Clathrocystis, Nostoc, Oscillatoria, Pandorina,* and *Volvox* when present in large numbers.

The question of growth of algæ in water reservoirs leads us to a very important public health problem. Although in India very little information is available about the contamination of the water supplies with the group of toxic algæ, the important question of checking their growth in the reservoirs of water supplies needs study. One of the essentials of the algal growth is light. This growth may, therefore, be prevented, or at any rate considerably reduced, by covering up the reservoirs and cutting off sunlight. Most of the reservoirs for the supply of water to both animals and man, however, in India are generally not covered and are often largely contaminated with algal growth. The removal of organic matter by keeping the source of water supply in as pure a state as possible will not doubt keep down the algal growth. It must be understood, however, that nearly all water contains sufficient organic matter for the growth of algæ, especially the water coming from water-sheds. Growth of algæ can also be successfully prevented by additions of copper sulphate in dilutions of one in five millions or even more. Such dilutions do not render the water deleterious to man and animals. The problem of toxic algæ is important and deserves the attention of workers in this field. It is also possible that some of these algæ may contain active principles having therapeutic properties in the treatment of disease. This aspect has yet to be investigated.

(c) **Fungi.** (i) Some fungi live on the skin and mucous membranes of man and animals and cause various, diseases, e.g. ringworm, thrush, etc.

(ii) There are others which attack foodstuffs and among these may be mentioned —

(1) **Smuts.** Many of these are destructive parasites which invade plants of vital economic importance on account of their food value, such as oats, millet and other cereals. Some are supposed to be piosonous if taken in large quantities, and others are said to produce irritation of the mucous membranes. There is difference of opinion with regard to the injurious effects produced by particular kinds of smut and hardly any authentic information is available regarding those occuring in India. The subject deserves the attention of mycologists.

(2) **Rusts.** Annual reccurrence of the outbreaks of rust attacks of cereals in India, especially those attacking wheat, is of great economic importance to the country. These, especially the uredo stage, produce

inflammation of the mucous membrane of the mouth and nose. The dust coming from the infested straw when the grain is thrashed is stated to cause serious disturbances of the respiratory tract of man and animals. Very little information is available about the Indian strains.

(3) **Ergot**, which grows on rye, is a well known example of a fungus which produces highly poisonous substances. Some species of this fungus grow on wild grasses in India and Ergot itself has been grown in this country for medicinal purposes.

(4) The poisonous nature of the seeds of darnel (*Lolium temulentum* Linn.), a grass and annual weed cultivated especially in northern India, is believed to be due to a fungus, and cases of poisoning due to admixture of the seeds with wheat grains are not infrequently reported in India and abroad. Cases of death among livestock have also been reported.

(5) **Moulds**. Very variable data are available as regards the poisonous effects of mouldy food stuffs in India, but there appears to be little doubt that the presence of certain species may occasionally produce harmful effects in man and animals Species of *Mucor*, *Aspergillus*, *Penicillium* and *Fusarium*, etc. deserve special investigation in this connection. It appears, however, that there is an appreciable difference in the susceptibility of different species of animals to the effects of mouldy foodstuffs. In general it has been stated that horses, dogs and pigs are more susceptible than ruminants and poultry, while in other animals the case may be the reverse. Very little information is available about the toxicity of moulds occurring in India and the problem requires a thorough investigation because of its great economic importance In the meantime it would be safer to consider all fungus-infected food-stuffs as deleterious Acute poisoning with the moulds is rarely met with and if they are taken in small quantities there is hardly any danger Mouldy food should however be avoided It is worthy of note that some of these fungi are the source of powerful antibiotic substances such as penicillin from *Penicillium notatum*, fumigaen from *Aspergillus fumigattus*, calvicin from *Aspergillus clavatus*, etc. A large amount of work is now being done in this country in connection with antibiotics from local strains of these organisms.

(iii) **Mushrooms or fleshy fungi** The third group of the poisonous fungi belongs to the mushroom class A number of these are edible and many occurring in India are indiscriminately eaten by people and if properly cooked few produce serious toxic effects. Cases of fungus poisoning, however, are not infrequently met with, particularly in the hills. Unfortunately very little information is available about the poisonous fungi growing in this country and inspite of numerous cases of poisoning, little attention has been paid to the subject.

Stropharia semiglobata (Batsch) Quel. from Khasia hills, *Hypholoma fasiulare* (Huds) Fr. from Darjeeling and Simla and *Lactarius vellereus* Fr. from Sikkim are regarded as poisonous. There is also evidence on record that there exists in Bengal a fungus which closely resembles an edible variety but which contains amanitine or muscarine, the poisonous principle of the foreign *Amanita muscaria* Pers. There are probably many more poisonous species that have really been incriminated as poisonous, but on the whole their number may be small and indeed if properly cooked

only a few are dangerous. If washed in water and macerated in vinegar before cooking, and if eaten with plenty of bread there is almost no danger in most cases. The safest method, however is to learn to recognize the edible species and never to eat a fungus until its identity is certain.

Some of the foreign poisonous fungi, e g *Lepiota cristata* Quel. *Volvaria gloiocephala* Gill , *Amanita muscaria* Pers. and *Amanita phalloides* Secr. are well known. The last-mentioned is responsible for perhaps 90 per cent. of the deaths caused by fungus poisoning in Europe, Great Britain and U.S A. During the World Wars, when food scarcity became acute in Germany and Austria, poisoning from fungi appreciably increased.

According to Ford there are four main types of mushroom intoxication —(1) Gastro-intestinal type in which the attack ceases when the stomach is emptied (2) General catharsis which is painless. (3) Violent vomiting and pain but no involvement outside the gastro-intestinal tract. (4) Choleriform type producing widespread degeneration of cells With regard to edible fungi it may be stated here that the nutrition value of mushrooms is small, their chief value being their flavour and the feeling of satiation they produce The following is a list of mushrooms met with in India :—

Agaricus compestris Linn.—The Mushroom; (Sans.—Chattrak; Kash —Manskhel, Beng.-Banger chhata, Sant —Ot : Bomb.—Alombe). Generally in damp debris throughout India during rainy season; universally eaten fresh or dried.

Amito pers. Kurrum.

A. ostreatus Jacq (Cutch & Bomb.—Phanasa-alambe, or vulgarly phansamba) Grows upon stumps of old jack-trees (phanas).

Cantharellus cibarius Fr Kashmir, Peshawar, Mussoorie.

Collybia albuminosa (Berk) Betch. (Syn -Lepiota albuminosa Berk). (Beng —Durga chhata) Bengal, C P , and Berar Grows from inside the termites, nests, eaten with relish.

Coprinus comatus (Battara) Fr The Mushroom; Hindi. & Punj.-Khumbi, Khumb) Punjab, Uttar Pradesh and several other parts of India. Eaten fresh or dried Collected during rainy season.

Entoloma microcarpum Berk & Broome. (Beng -Wee-chhata). Bengal. Grows on the surface of outer crust of termites' nests, commonly eaten by villagers.

Fistulina hepatica Fr. Darjeeling.

Helvella crispa Fr. Common in Afghanistan

Hirneola polytricha Mont. (Syn.-*Exidea polytricha* Mont.). Belgium, Poona, Dharwar, Nidungayam, Malabar and Burma.

Hydnum caralloides Scop Darjeeling 7,500 ft. , Chitral, N.W.F.P. (common) and Afghanistan. In crevices of old tree-trunks collected during August; dried in the sun and largely used.

H. repandum Linn. Mussoorie, Uttar Pradesh

Lactarius deliciosus Fr. Sikkim.

Lentinus subnudus Berk Common in Bengal, Kadala and Bombay. On dead branches of logs Eaten by Kholes fresh and young.

Lepiota mastoidea Fr. Bengal.

L. procera (Scop) Sacc Saharanpur.

Lycoperdon sp. Puff-balls Bengal, Kashmir and many parts of Western Himalayas.

Malanogaster durissiums Cooke Truffle. Simla (abundant), Kangra. Occasionally eaten.

Morchella esculenta Pers. The Morell, (Punj -Guchhian) Plains, kana kach hills. Fleshy fungus abundant in Kashmir, Chamba and many parts of Northern Punjab It appears on hills as snow melts in early spring. It is dried and is eaten with much relish.

Pleurotus cretaceus Massee (Vern -Dhingri). Peshawar and Madhya Pradesh. On wood.

P. fimbriatus Bolt C P. and Berar

Polyporus squamosus (Huds) Fr Darjeeling, 7,500 ft., Pangi, N.W. Himalayas. On dead wood.

Truffles. Stewart describes some being found in Kashmir Badhwar has recently collected some blackish-brown ones from the Kagan valley locally known as 'usri'. They are highly flavoured and their presence in the soil is discovered by the villagers by smell in September-October when they are said to develop the flavour best Goats are also said to dig out some during grazing and eat them. (Also see *Melanogaster durissiums* above).

Volvaria diplasia Berk & Broome (Beng -Pawal-chhatta) Bengal, Burma.

V. terrestria Berk & Broome. (Beng. - Poal-chhattea) Bengal Grows on heaps of waste paddy straw

In addition Stewart mentions another species as being freely eaten in the Punjab, which is known as 'shirian' in the Jhelum and 'batbakri' in the Kair valley It is a thin, flat ragged-looking fungus, yellow above and with white gills below, which is found on dead trees in various parts of the Punjab Himalayas at altitudes of 8,000 to 8,500 ft. He also mentions an 'underground mushroom' of doubtful species found near Multan called 'boinphal'.

So far as the Poisonous Mushrooms are concerned little information is available. The following is a list of mushrooms which are known to be poisonous :-

POISONOUS FUNGI

Very little information is available regarding the Indian poisonous fungi From time to time cases of fungus poisoning are reported but, it is to be regretted that little or no attention has been paid to the subject

Stropharia semiglobata (Batsch) Quel from Khasia Hills

Hypholoma fasciculare (Huds). Fr. from Darjeeling and Simla and *Lactarius vellerens* Fr from Sikkim, are regarded as poisonous. There is also evidence on record that there exists in Bengal a fungus which closely resembles an edible form but which contains amanitine or muscarine, the poisonous principle of *Amanita muscaria* by eating which, symptoms closely resembling those of intoxication rapidly ensure. Furthermore, mucor has been regarded as a harmful in India since ages, and the pickles and all edible stuff attacked by it are not thought fit for eating

There are, however, some foreign fungi which are definitely reported to be poisonous

Amanita phalloides The death cap is responsible for perhaps 90 per cent of the deaths caused by fungus poisoning in Europe, England and U S A It is the most dangerous fungus known and very small quantities will cause intense suffering and often death. There are indeed, several other species of the genus that are very poisonous, e g , *Amanita muscaria* -fly agaric and *A. pantheriana* - warted agaric, etc which are intensely poisonous

Lepiota cristata - crested agaric, and several other small species of Lepiota are regarded with suspicion and should be avoided. *Volvaria gloiocephala* - glutinous agaric and its allied mushrooms have always been regarded as poisonous, but there is recent evidence that they may be eaten without ill effects.

Psalliota xanthoderma - yellow staining mushroom has caused illness in some cases.

(d) **Lichens.** Very little is known about these symbiotic organisms which consist of algal cells enveloped by the mycelium of the fungus forming a felted mass Although this group is not to be regarded as a serious menace to livestock, cases of poisoning due to *Parmelia* and *Cretaria* species, etc are mentioned in foreign literature *Parmelia molliuscula* has been said to effect sheep and cattle, producing lack of coordination of the hind limbs In more severe cases the animal lies down and is unable to move either its front or hind limbs. Little or no information is available about lichens in India and even their systematic botany has not been sufficiently worked out.

(e) **Bryophyta** (Liverworts and Mosses). This is the least-known group of plants from the view-point of poisoning and there is little to be said about it.

(f) **Pteridophyta** (Vascular Cryptogams). This group includes ferns and allied plants but unfortunately little or no work has been done in India with regard to their toxicity. Greshoff and others have reported

the presence of hydrocyanic acid in a number of ferns, especially when young. References to the supposed poisonous properties of the bracken fern (*Pteris aquilina*) have appeared in the literature for a long time, and Stockman in great Britain showed that it is poisonous to cattle when eaten in considerable quantities The plant is found in India *Aspidium filix-mas*, the male fern, is suspected of being poisonous. The roots are used in medicine and large quantities of it produce haemorrhagic gastro-enteritis, tremors, weakness, stupor coma, acute nephritis, and cystitis Six drachms of the oleoresin have proved fatal in man and three ounces in the cow. This fern is not found in India, but there are several other foreign species of Aspidium which are also suspected of being poisonous. The present authors have examined Indian representatives of these plants. Some foreign species of Osmunda, Davallia and Adiantum are also suspected of being poisonous and medicinally active. The following ferns growing in the North Western Himalayas have been examined and found to be active as anthelmintics.

Name of Ferns	*Locality*	*Active principles*		
Dryopteris odontoloma	Kashmir & Mussoorie	Up to B P	& U S P	standard
D. barbigera	Kashmir	do	do	do
D. marginata	Kashmir	do	do	do
D. blanfordii	Chamba	do	do	do
D. schimperianum	Mussoorie	do	do	do
D. calcarata	Mussoorie	Not up to standard		
D. marginata	Mussoorie	Up to standard		

Some of the foreign species of *Equisetum* (horsetail) have long been recognized in foreign countries as injurious to cattle and horses. They produce an intoxication in which the animals stagger about and wander aimlessly There is no information available in India with regard to the Indian horsetail *Equisetum arvense*, but several European and American workers are convinced that it is definitely poisonous to horses, while others hold a contrary opinion This plant grows commonly in certain places in India where it might be a menace to livestock.

CHAPTER VIII

II. TOXICOLOGICAL ASPECTS OF PHANEROGAMS (THE FLOWERING PLANTS)

After having given a very brief survey regarding the toxicological aspects of the Cryptogamic flora we will now take up the Phanerogams or the flowering plants Economically this is the most important group both for man and animals from the point of view of everyday necessities of life, e g, food, medicines, etc It is probably on account of this that more information is available with regard to this group.

From a toxicological point of view the Phanerogams may be divided into two main groups.

I PLANTS POISONOUS TO MAN AND LIVESTOCK.

(a) **Medicinal Plants** —Primitive man in the quest for food must have come across plants containing poisonous principles by accident and by experience must soon have learned to avoid them. He even made use of them for the purpose of fighting against his enemies and for procuring his food by killing animals with them. On the whole, our knowledge is fairly well advanced so far as the relationship of poisonous plants have been used for criminal purposes from ancient times but the majority of them are used as medicinal agents for the amelioration of human suffering. It is well known that many plants that are harmful to life in large quantities, produce remarkably beneficial effects in small regulated doses in disease conditions. From the economic point of view, the abundance of this group of plants in our midst is of very great importance in as much as it provides us with medicinal agents of every description, not only sufficient for our own use but also for purposes of export. We have already given brief review of the work done on this group.

(b) **Poisonous to livestock**. The second important aspect of these plants is in connection with other countries, our knowledge is very meagre. In India, there are hundreds of plants that are intimately connected with the food supply of the bovine population such an essential factor in connection with agriculture and food supply. The fodder required for this livestock amounts to about fifty million maunds daily (excluding the concentrates that are in use). Even in its present unsatisfactory condition, the cattle industry contributes very largely to the annual agricultural income of this vast country. Its welfare is therefore of the utmost importance. There is no doubt that the mortality among cattle through poisoning in India is very large. Unfortunately no figures are available of the loss suffered on this account but it must be enormous.

Even in a country like the United States of America where knowledge with regard to Poisonous Plants is well advanced as compared to India, great losses are incurred every year through the poisoning of cattle. The loss occuring in this sub-continent can thus be imagined.

Though the number of plants which are known to have markedly poisonous properties is perhaps small as compared with the total species included in the Indian floras, there are many which are of common occurrence and which no doubt produce serious losses by death or illness among sheep, cattle and other domestic animals The toxic effects produced may be indicated by reduction in the yield of milk, the milk may become unpalatable through excretion in it of toxic products, or it may even become unfit for consumption Some time even the meat of these animals becomes poisonous

It may be stated here that animals do not instinctively select toxic plants as forage, that all classes of livestock are not necessarily equally susceptible to the same poisonous plants, that not all poisonous plants are dangerous from their initial appearance up to maturity and that in some the animals do acquire a depraved appetite for harmful plants, especially when the fodder supply is scarce, a condition which is of frequent occurrence in many parts of India The losses in many cases may be avoided by increasing our knowledge about these plants.

Prevention. The question arises as to what should be done to prevent poisoning by plants. The adage 'prevention is better than cure' is applicable to the problem of plant poisoning with just as much force as in other spheres Often cases are brought to notice when the symptoms have developed and the poison has already circulated in the blood stream and has done irreparable damage to the system Increased knowledge of the poisonous plants is the first step in this direction and this is sure to have an effect in decreasing fatalities among human beings and livestock. It is for this reason that Indian Council of Agricultural Research and Indian Council of Medical Research have encouraged research on this group Keeping the animals always as far as possible from dangerous areas and exercising special care during periods of drought are likely to decrease the mortality amongst livestock Eradication of poisonous plants is a difficult matter, involving an enormous amount of labour and capital, but wherever and whenever possible it should be resorted to This depends upon the habits of the particular plant. Such plants may be annual, biennial or perennial herbs, or shrubs or trees Annuals should be pulled out or dug out before seeding and biennials may be dealt with as the annuals. Perennial are propagated both by the seeds and by the underground organs, such as tubers, rootstocks, bulbs, etc and may be dug out if not deeply rooted Shrubs are woody perennials and should be cut down or dug out Cutting down of lower branches of trees within the reach of animals or children is advocated

The indiscriminate importation of ornamental plants has recently increased the number of poisonous plants in India. Some of these do not find much competition in their adopted home and are likely to spread in this country at an enormously rapid pace. The time is now ripe to agitate for a law prohibiting the importation of poisonous plants for gardens or to take measures to forbid the cultivation of those already introduced. The foodstuff dealers should make sure that adulteration is not practised either with poisonous plants or with plants whose properties are doubtful. Recent work in connection with the causation of epidemic dropsy has shown that in some epidemics, mustard oil adulterated with 'katakar' oil from the seeds of *Argemone mexicana* Linn , the mexican poppy or 'Shialkata', was the cause of the outbreak.

Food Poisons. There is a number of common articles of food which produce poisoning in man and animals and a brief reference may be made to these.

(1) **Khesari dal, Lathyrus sativus** Linn , an important article of diet in man and animals, has been responsible for a large number of cases of poisoning under certain conditions in man, cattle, sheep, pigs, horses, pigions, ducks etc. Examples of lathyrism in man in the form of spastic paralysis are commonly seen every day in the streets of Calcutta and towns in Bihar and its toxic effects in horses and cattle are well known Moderate amounts of this pulse can be taken with impunity, but large amounts, especially to the exclusion of other fodders or foods, produce poisoning.

(2) Grasses (Gramineae) form an important part of the food of animals Some of these develop dangerously large quantities of hydrocyanic acid under certain climatic and soil conditions especially at times of drought or when the plants are wilting, stunted or young Unfortunately our knowledge of Indian grasses in this connection is meagre and it is not possible to estimate the losses in livestock from this source

The common Jowar (*Sorghum vulgare* Pers), the Indian millet, is largely cultivated as fodder for cattle and also for human food It has caused serious outbreaks of poisoning among livestock when wilted or stunted under drought conditions *Sorghum halepense* Pers , a tall perennial grass, with creeping rhizomes and numerous suckers, known as Johnson grass, grows all over India under the name of 'baru' in Hindi and 'Kala-mucha' in Bengali. It has been responsible for serious losses among livestock during recent years in the North-Western parts of India It has been stated that the amount of hydrocyanic acid in these plants decreases with the age of the plant but never entirely disappears The points to be remembered about these grasses are that they are dangerous during wilting and under conditions of draught, that the younger and more succulent ones are often more likely to contain lethal doses of hydrocyanic acid and that if well dried, these plants are generally not dangerous. The toxicity in the case of cyanogenetic compounds depends on the quantities of hydrocyanic acid liberated, and according to the amount and speed at which they are eaten Often such large quantities are given that the animal will die before any veterinary aid can be given. The only remedy is prevention The problem of poisonous grasses is of great economic importance in certain parts of India where rains often fail and drought conditions prevail.

The workers of the Drug Research Laboratory have been engaged in carrying a survey of grasses occurring in the North Western Himalayan Region. About 150 different species have been collected and are now being studied with regard to their nutritive values and possible poisonous properties A number of poisonous grasses have been discovered which the cattle of these parts instinctively avoid, but new-comer gets poisoned *Stipa siberica* commonly growing on hills round Kashmir Valley is an example and there are others.

(3) The linseed plant, *Linum usitatissimum* Linn., contains a cyanogenetic glucoside, the maximum amount of which is reached very early in the development of the plant and finally disappears, except in the seed, which

still contains small quantities. An oil is expressed from the seeds and the remaining cakes are used to feed livestock. Cases of poisoning have been frequently reported amongst animals feeding on this plant. It is unsafe to feed the cattle on the immature plant, especially when it is wilted. The cake after extraction of the oil should be treated with boiling water to destroy the enzyme responsible for liberating hydrocyanic acid from the glucoside, and should not be soaked in cold water overnight. It should be given only in small quantities at a time.

(4) The *mustard cake* which is fed to cattle after the extraction of oil may produce chronic irritant poisoning, colic, lassitude, etc , if fed in large amounts and over prolonged periods on account of the liberation of an essential oil by the action of an enzyme on the glucoside contained therein

(5) Several members of the cucumber family (*Cucurbitaceae*) are edible but bitter varieties are occasionally met with. The latter have a strong purgative action and should be discarded. Incidentally it may be remarked that most of the wild members of the family are toxic Colocynth which is a powerful intestinal irritant is a familiar example The bitter members of this family have more or less a similar action.

(6) The potato, *Solanum tuberosum* Linn., when sprouting, produces dangerously large quantities of the toxic alkaloid, solanine, and must be thrown away Certain plants, such as buck-wheat (*Fagopyrum esculentum* Moench) is largely cultivated for human and animal consumption, under certain conditions not yet fully understood, become toxic and give rise to inflammatory swellings of the face, eyelids and ears.

Reputed Poisonous Plants of India including Insecticidal, Piscicidal and Abortifacient

Ranunculaceae

(Buttercup Family)

Anemonin, aconitine, indaconitine, pseudaconitine, adonidin, delphinine, staphysagroine, cyanogenetic glucosides, essential oils, saponins, etc.

1 **Aconitum balfourii** Stapf (Vern - Nep - Gobari), *A chasmanthum* Stapf ex Holmes (Vern - Kash -Banbalnag), *A deinorrhizum* Stapf. (Vern - Mohra, Maurabikh), *A elwesii* Stapf *A falconeri* Stapf (Vern - Bis, Bikh, Meetha-tellia), *A. ferox* Wall ex Seringe, (Vern - S - Visha, H & B.-Bish, Bo.-Vachnag, M -Vashanavi), *A. lacinatum* Stapf (Vern - Kalo bikhmo), *A. laeve* Royle, *A lethale* Griff , *A luridum* Hk f. & T , *A moschatum* Stapf, *A. soongaricum* Stapf *A spicatum* Stapf (Vern - Bikh, Kalo bikhoma donghi), *A violaceum* Jacq

These are all cardiac depressant and nerve poison. Cause deaths among livestock and are also used as arrow poison. Used in Hindu Medicine after mitigation as cardiac tonics Mitigation or correction generally consists in boiling the aconite roots with cow's urine

2. **Actaea spicata** Linn.

It is acrid and poisonous and deaths among horses are reported

3. **Adonis aestivalis** Linn , *A. chrysocyathus* H. f & T.
These are poisonous to animals and act as poison to heart.

4. **Anemone obtusiloba** D. Don. (Vern - P.-Rattanjog)

It is a vesicant and when taken internally produces vomiting and purging, drying alters properties.

5 **Aquilegia vulgaris** Linn
It is poisonous to animals.

6 **Caltha palustris** Linn.

It is acrid and poisonous and deaths among horses are reported.

7. **Cimicifuga foetida** Linn (Vern - P -Jiunti)
It is a heart depressant and acts as insect repellent.

8. **Clematis gouriana** Roxb , *C graveolens* Lindl , *C napaulensis* DC. (P.-Oandak), *C orientalis* Linn , *C triloba* Heyne (Vern - S.- Laghukarni, H & Bo- Moravela) *C wightiana* Wall.

These are irritant and produce blisters, such properties are altered by drying

9. **Delphinium brunonianum** Royle (Vern - P -Laskar), *D. caeruleum Jacq* (Vern - P.-Dhakangu), *D. elatum* Linn , *D. vestitum* Wall

These are cardiac and respiratory depressants, acrid in taste, insecticidal and poisonous to animals

10. **Nigella sativa** Linn. (S -Krishnajiraka, H & B.-Kalajira, Bo - Kalenjire, M -Karun-shiragam)

Produces abortion in larger doses.

11. **Paeonia emodi** Wall Vern.- H.-Udsalap, P - Mamekh).

It is said to have narcotic properties

12. **Ranunculus arvensis** Linn (Vern - P - Chambul), *R. falcatus* Linn , *R. laetus* Wall , *R lingua* Linn , *R pensylvanicus* Linn. f , *R. sceleratus* Linn (Vern - Pers.- Kabiraj).

These are vesicant and poisonous to livestock when fresh, drying alters these properties.

Magnoliaceae

(Magnolia and Champa Family)

Shikimin, illicin, essential oils.

13. **Illicium griffithii** Hk. f & T., *I anisatum* Linn., (*-I religiosum* Sieb. & Zucc.) (Vern.- H -Anasphal, Bo.-Badian, M.-Anashuppu)

This is the star anise of China (*I verum* Hook.f.) which was imported into India sometimes adulterated with *I religiosum*, it has produced deaths. The latter is respiratory and cardiac poison. Indian *I. griffithii* is also referred to as being poisonous

Annonaceae

(Custard apple Family)

Resin, alkaloid, etc.

14. **Annona reticulata** Linn (Vern - H.-Lona, B -Nona, Bo.-Ramphal, M -Ramsita), *A squamosa* Linn (Vern - S.- Gandhagatra, H - Sitaphal, B -Ata, M - Sitapalam).

The seeds are intensely irritant to the conjunctiva; applied locally they act as abortifacient They have insecticidal properties. Roots act as drastic purgative.

Menispermaceae

(Moonseed Family)

15. **Anamirta cocculus** (Linn) W. & A (Vern - S -Kakaphala, H. & B - Kakmari, M -Kakkay-kolli-virai)

It is a convulsant poison to animals. It acts as an insecticide and it is also used to poison fish and cattle.

16 **Pachygone ovata** (Poir) Miers

It is an insecticide and piscicide

Berberidaceae

(Berberry Family)

Berberine, podophyllum resin

17 **Berberis aristata** DC (Vern -S -Daru haridra, H -Dar-hald (and probably few more species)

It is poisonous to lower animals and is a fish poison

18 **Podophyllum hexandrum** Royle (*P emodi* Wall. ex Hk f et T)

It contains a resin which is a drastic purgative and is irritant to the mucous membranes

Papaveraceae

(Poppy Family)

Morphine, codeine, papaverine, berberine, argemone oil, etc

19 **Argemone mexicana** Linn (Vern - S -Srigala-kantaka, H & B - Sialkanta, M -Birama-dandu)

The oil expressed from its seeds is occasionally mixed with mustard oil which has been held responsible for producing symptoms resembling epidemic dropsy

20 **Meconopsis aculeata** Royle (Vern - Simla - Kanta), *M. napaulensis* DC.

The roots are considerd to have narcotic properties.

21. **Papaver dubium** Linn., *P. nudicaule* Linn , *P. rhoeas* Linn. (Vern - S -Rakra-posta, H -Lalpost, Bo.-Janglimudrika, M.- Shivappu-postaka chedi), *P. somniferum* Linn (Vern.- S.-Ahiphena, H. & B.-Afim, Bo -Aphu, M.-Postakatol)

All species yield opium more or less but *P. somniferum* is the chief source Opium has been used for suicidal purposes

Cruciferae

(Mustard Family)

Glucocides on contact with water produce vesicant active essential oils

22 **Brassica cernua** (Thunb) Forbes et Hemsl , *B integrifolia* (West) O E Schulz, *B juncea* (Linn) Czernjaew et cosson (rai) , *B napus* Linn with four varieties (Toria, sarson), *B nigra* (Linn.) Koch (black mustard).

Seeds of these are vesicant, and mustard cakes if fed in large quantities and over prolonged periods is harmful to cattle Sarson cake is safest, mixture with rai or black or white mustard dangerous

23 **Lepidium draba** Linn (Vern - Afgh -Bijindak)

It is a fish poison

24 **Sinapis alba** Linn (white mustard)

It is discussed under Brassica

Capparidaceae

(Caper Family)

Essential Oils

25 **Capparis decidua** Edg (*C aphylla* Roth) (Vern - S -Karira, H -Karer, P Karia, Bo -Kari, M Karyal)

It acts as a vesicant

26 **Cleome felina** Linn f *C viscosa* Linn , (Vern -S -Swarnakshira)

It acts as a vesicant

27 **Gynandropsis gynandra** (Linn) Merr (*G pentaphylla* DC)

It is an insecticide and a piscicide It has vesicant properties

Bixaceae

(Chaulmoogra family)

Cyanogenetic glucoside, chaulmoogra oil.

28 **Gynocardia odorata** R Br (Vern -H., B. & Bo -Chaulmoogra)

The fruit acts as a piscicide.

29 **Hydnocarpus kurzii** (King) Warb. (*-Taraktogenos kurzii* King), *H. laurifolia* (Dennst.) Sleumer (*-H. wightiana* Bl)

The fruit acts as a piscicide and the seed oil is gastro-intestinal irritant.

Polygalaceae
(Milkwort Family)

Saponins.

30. **Polygala chinensis** Linn. Vern -H -Meradu, Bo -Negli), *P. crotalarioides* Buch -Ham (Vern -Santh -Lil kathi), *P telephioides* Willd.

It has an acrid taste and is an emetic It has expectorant properties.

Caryophyllaceae
(Carnation Family)

Saponins.

31. **Saponaria vaccaria** Linn (Vern - H.-Musna, B.-Sabuni) and probably some others of the family.

It is acrid and its toxicity is partially removed by boiling

Hypericaceae
(St John's wort Family)

Balsamic resinous juice

32 **Hypericum perforatum** Linn (Vern -H & P -Basant)

It is poisonous to animals, especially horses if taken in excess, usually it is not eaten

Guttiferae
(Gamboge Family)

Gum resins

33 **Calophyllum inophyllum** Linn (Vern -S -Punnaga, H. Sultana, champa, B -Punnag, Bo -Undi, M -Punnagam).

It is a fish poison.

34 **Garcinia morella** Desrouss, and probably others

It yields a gum resin which is a violent gastro-intestinal irritant

Ternstroemiaceae
(Tea Family)

Caffeine, theophylline

35 **Thea sinensis** Linn

Its excessive indulgence only is harmful.

Malvaceae
(Cotton family)

Gossypol, resin, ephedrine, pseudo-ephedrine.

36. **Gossypium** species.

The root bark is emmenagogue and used as abortifacient; occasionally harmful effects of cotton seed cakes on animals have been reported.

37. **Malva parviflora** Linn. (Vern -H. Panirak)

It is reported to have produced narcotic poisoning in animals.

38. **Sida rhombifolia** Linn (Vern - S -Atibala, H & B -Swet-berela, M -Athiballachetu)

The ripe capsules are reported to be fatal to fowls.

Linaceae

(Flax Family)

Cyanogenetic compounds, cocaine

39. **Erythoroxylum coca** Lam

It is a central nervous stimulant, sensory nerve endings-paralysant, addiction to it is harmful

40 **Linum usitatissimum** Linn (Vern - S -Atasi, H & B -Tisi, Bo -Alasi, M -Alshiviral).

The young plants are known to produce deaths in animals, sometimes seed cakes also harmful

Zygophyllaceae

(Bean-caper and Guaicum Family)

Harmine, harmaline, harmalol, peganine, essential oils, saponins, resins.

41 **Peganum harmala** Linn. (Vern.-H & Bo -Hurmal, B -Isband, M -Shimai-azha-vanai-virai)

It is an insecticide, narcotic, nauseant and emetic It is used as abortifacient It is a protoplasmic poison and paralyses skeletal and cardiac muscles of frog

42. **Tribulus terrestris** Linn. (Vern -S -Gokshura, H -Chotagokhru, B -Gokhuri, Bo -Lahana-gokhru, M.-Nirunji)

It causes the disease goeldikkop (dikgeel) in South Africa in small stock which is characterized by oedema of head, fever and jaundice.

Rutaceae

(Rue Family)

Essential oils, rutin, skimmianine, saponins, resins, etc.

43. **Acronychia pedunculata** (Linn) Miq (*-A laurifolia Bl*)
It is a fish poison.

44. **Ruta graveolens** Linn var *angustifolia* Hk f., (Vern - S.-Somalata, H.-Sadab, B.-Ermul, B -Satap, M.-Arvada), *R. tuberculata* Forsk.

These are acro-narcotic poisons which are rubefacient. The oil and herb frequently used to produce criminal abortion.

45 **Skimmia laureola** Sieb & Zucc ex-Walp (Vern - Nep -Chumlani, P -Ner).

It is reported to be poisonous to sheep and goats.

46. **Zanthoxylum alatum** Roxb (Vern - S -Tumburu, H.-Tejmal, B.-Nepali dhania) (Probably some more species)

It is a fish poison.

Simaroubaceae

(Bitter-bark Family)

Essential oils, saponins, resins, bitter substances

47 **Ailanthus altissima** (Mill) Swingle (—*A glandulosa* Desf)

It is nauseant and nervous system depressant Accumulation of its leaves in well water is reported to produce chronic gastritis

48. **Balanites aegyptiaca** (Linn) Delile (—*B roxburghii* Planch) (Vern -S -Ingudi, H & B -Hingan, Bo -Hinger, M.-Najunda)

It is believed to be a fish poison and a purgative

49 **Brucea amarissima** (Lour) Merr - (*B sumatrana* Roxb)

Its seeds produce nausea, vomiting, abdominal pain and purging

50 **Picrasma napalensis** Benn

It is stated to be used as larvicide in Sikkim

Meliaceae

(Neem and mahogany Family)

Bitter substances, bitter oil, saponins

51. **Azadirachta indica** A Juss

It is parasiticidal and the leaves are used as insect repellent

52. **Melia azedarach** Linn

Its berries are especially poisonous to man and animals, they are narcotic and gastro-intestinal irritant

53 **Walsura piscidia** Roxb (Bo & M -Walsura).

It is a dangerous emmenagogue and a violent emetic. It is largely used as a fish poison.

Celastraceae

(Spindle-tree Family)

Alkaloid, essential oil, resin.

54. **Elaeodendron glaucum** Pers. (Vern -H.-Bakra, Bo.-Bhuta-pala, M -Selupa.)

It is emetic, overdoses fatal.

Sapindaceae
(Soap-nut Family)

Saponins, cyanogenetic compounds

55 **Cardiospermum halicacabum** Linn (Vern -S -Karnaspota, H.-Kanaphata, M -Mooda cotton).

Its leaves are emetic and rubefacient

56 **Dodonaea viscosa** Linn (Vern -H -Aliar, Bo -Bandurgi, M - Virali)

It is fish poison, it is deleterious to camels

57 **Harpullia cupanioides** Roxb.

It is a fish poison

58 **Melianthus major** Linn.

It produces acute diarrhoea, salivation and colic, honey from flowers stated to be poisonous.

59 **Sapindus mukorossi** Gaertn. (Vern -S -Phenila, H B & Bo -Ritha), *S trifoliatus* Linn (Vern -S -Phenila, H B. & Bo -Ritha, M -Ponnan-kottai)

Both are fish poisons, emetic, purgative and are used for procuring abortion

60 **Schleichera oleosa** (Lour) Merr. (*S trijuga* Willd.) (Vern -H - Kosum, M -Pu-maram).

The oil is occasionally mixed with mustard oil or ghee It produces irritant poisoning and the seeds are used as insecticide.

Anacardiaceae
(Cashew and mango Family)

Toxic phenolic compounds, toxic resin.

61 **Anacardium occidentale** Linn. (Vern -H. & Bo.-Kaju, B.-Hijli badam, M -Mundiri-kai).

The pericarp contains powerfully vesicant juice, used to preserve floors, wood, books, etc. from white ants, tar from bark also vesicant.

62. **Holigarna arnotiana** Hook. f. (Vern -Bo -Bibu), *H. ferruginea* March, *H grahamii* (Wight) Hook f , *H longifolia* Buch. Ham ex-Roxb. (Vern- B.-Barola, Bo -Hulugiri).

The juice is vesicant although not equally powerful in all species.

63. **Rhus insignis** Hook f., *R. punjabensis* J. L. Stewart, *R. succedanea* (Vern -S -Karkata singi, H & B.-Kakrasingi, Bo -Takadasingi), *R. wallichii* Hook. f. (Vern -Nep.-Ghosi, H -Akoria.)

Dreaded by local people; even smoke from burning wood is considered poisonous, its juice is vesicant

64 **Semecarpus anacardium** Linn f (Vern -S -Bhallatamu, H. & B -Bhela, Bo -Biba, M -Shayrang), *S travancoricus* Bedd

The pericarp contains vesicant juice It is used in some parts as abortifacient.

Coriariaceae

(Coriaria Family)

Coriamyrtin, tutin in foreign species.

65. **Coriaria nepalensis** Wall

It is stated to be narcotic, the foreign species are very poisonous acting like picrotoxin and producing convulsions.

Moringaceae

(Horse-radish Family)

Essential oils, alkaloid, moringine, moringinine

66 **Moringa oleifera** Lamk (—*M pterygosperma* Gaertn) (Vern -S - Sobhanjana, H -Sajnah, B -Sojna, Bo-Sujna, M -Murungai)

The fresh root bark is said to be vesicant and is used to procure abortion Moringinine acts on sympathetic nervou, system.

Leguminosae

(Pea Family)

Alkaloids, glucosides, saponins, cyanogenetic compounds, rotenone, toxic albumin, bitter substances ; globulins

67. **Abrus precatorius** Linn (Vern -S & Bo -Gunja H -Gaungchi B.-Kunch, M -Gundumani)

It is specially a blood poison and is used to poison cattle and to procure abortion.

68 **Acacia pennata** Willd (H.-Biswul, Kumaon -Agla, Nep -Arfu) It is a fish poison

69 **Albizzia procera** Benth (Vern -H -Safed siris, B -Kori, Bo - Kinai tihiri, M -Konda vaghe)

It is a fish poison

70 **Butea monosperma** (Lam) O Ktze (—*B frondosa* Koen ex-Roxb) (Vern - S -Kinsuk, H & B -Palas)

Its seeds are insecticide and painful if taken internally

71 **Caesalpinia nuga** Ait (Vern -M -Kakumullu) It is a fish poison.

72. **Canavalia virosa** W. & A. (*C. ensiformis DC.* var *virosa* Baker. (Vern .-M -Kattuvalari).

Its fruit is stated to be poisonous.

73 **Cassia absus** Linn. (Vern -H -Chaksu, M -Karun Kanam), *C acutifolia* Delile, *C alata* Linn , *C. angustifolia* Vahl, *C. fistula* Linn , *C obovata* Collad (Vern —Surati sonnamukai).

These are all purgatives which are irritant in large doses *C absus* seeds are irritant application to eyes, and *C alata* is a fish poison

74 **Clitoria ternatea** Linn (Vern -S -Aparajita, H. & B -Aparajit, M -Kakkanan)

The roots are powerful cathartic like Jalap, not a safe medicine

75 **Cytisus scoparius** Linn

The plant is not eaten by cattle It is an emetic and a cathartic

76 **Dalbergia stipulacea** Roxb.

It is a fish poison.

77 **Derris elliptica** Benth (Vern -Malay -Tubah), *D. scandens* Benth (Vern -B -Noalata, P -Gunj, M -Nala tige), *D uliginosa* Benth. (Vern -B -Panlata, Bo -Kirtana), (possibly *D ferruginea* Benth)

These are all fish poisons and *D elliptica* is a powerful insecticide.

78. **Entada phaseoloides** (Linn) Merr (—*E. scandens* Benth) (Vern -H -Chian, B -Gilagach, Bo -Gardal).

It is a fish poison

79 **Lathyrus aphaca** Linn , *L sativus* Linn (Vern -S -Triputi, H. & B -Khesari, Bo -Lakh)

It is a food and fodder *L sativus* if taken in large amounts and over prolonged period produces lathyrism in man and animals Ripe seeds of *L aphaca* are stated to be narcotic if taken in excess

80 **Melilotus alba** Desr.

It is stated to be poisonous to cattle.

81 **Milletia auriculata** Baker, *M pachycarpa* Benth., *M. piscidia* Wight.

These are fish poisons *M. auriculata* is an insecticide.

82 **Mundulea suberosa** Benth.

It is a fish poison.

83 **Ougenia dalbergioides** Benth. (Vern - S.-Tinisasegandun, H.-Sandan, B.-Tinis, Bo Tiwas, M-Tella-motuku).

It is a fish poison.

84. **Phaseolus lunatus** Linn.

The coloured variety sometimes exhibits poisonous properties if eaten.

85. **Pithecellobium bigeminum** Mart (Vern -H. & Bo -Kachlora)

It is a fish poison The seeds are stated to be eaten in Burma but sometimes produce severe poisoning

86 **Pongamia pinnata** (Linn) Merr (—*P glabra* Vent) (Vern - S., H. & Bo -Karanja, B Dahar karanja, M -Pungammaram)

It is piscicide and insecticide.

87 **Sophora mollis** R. grah var. *Hydaspidis* Baker, *S tomentosa* Linn.

The Seeds of *S. mollis* have insecticidal properties and the leaves of *S. tomentosa* are powerfully emetic and cathartic in large doses.

88 **Tephrosia candida** Linn , *T. purpurea* Pers (Vern - S -Sarapunkha, B -Bon-nil, H & Bo -Sarphankha, M -Kolluk-kay-velai)

These are fish poisons Some foreign species are insecticides Some species of Tephrosia in India are likely to prove of value as insecticides

89. **Trifolium repens** Linn

It is highly prized fodder in Europe In the Himalayas poisoning has been reported in horses

90 **Vicia sativa** Linn. (Vern -H -Ankra, B.-Ankari)

It is suspected to be the cause of lathyrism.

Rosaceae

(Rose family)

Cyanogenetic glucosides, phloridzin

91 **Prunus amygdalus** Batsch. (bitter variety). (Vern -H , B. & Bo - Badam, M -Vadam-Kottai), *P armeniaca* Linn. (Vern -H -Khubani, P -Gurdlu), *P. avium* Linn , *P. cerasus* Linn , *P. mahaleb* Linn (Vern .-S , Priyangu), *P padus* Linn. (Vern -H -Jamana, P -Jamma), *P. persica* Stokes, (Vern -H -Aru), *P. puddum* Roxb , (Vern -S.-Padmaka, H -Paddam, Bo -Padma-kasta), *P. undulata* Buch -Ham

The seeds are poisonous and the leaves of many species are said to be dangerous to live-stock when wilted; harmless when on the plant, suspicious when dried.

92. **Pygeum gardneri** Hook. f.

The seeds are fish poison.

93. **Pyrus aucuparia** Linn., *P. malus* Linn.

The bark of *P. aucuparia* is irritant to the alimentary tract and wilting leaves of other species are occasionally poisonous to animals browsing upon them.

94. **Rubus moluccanus** Gaertn (Vern-Kumaon.-Katson)

The leaves are reported as powerful emmenagogue and abortifacient.

Crassulaceae

(Life-Plant-Family)

Glucosides-in foreign species.

95 **Kalanchoe spathulata** DC. (Vern:-H.-Tatara).

The expressed juice of bitter variety is a drastic purgative and is poisonous to goats. It is not eaten by cattle. Leaves are said to be insecticidal.

Droseraceae

(Sundew Family)

Organic acids, proteolytic enzymes

96 **Drosera peltata** Sm var. *lunata* Clarke, (Vern - H -Mukajali, P Chitra), *D. spathulata* Labill. (*D. burmanni* Vahl)

These are rubefacient Some Australian species are reported to be injurious to sheep.

Combretaceae

(Myrobolan Family)

Tannins

97 **Terminalia bellerica** Roxb (Vern -S -Bahira, H. & B.-Bahera, Bo -Behaira, M.- Vallaimurdu), *T chebula* Retz. (Vern - S & B - Haritaki, H -Harara, Bo - Hirda, M - Kadukkay-pu)

T bellerica is reported to be fish poison. The kernel is stated to be poisonous and cases are reported where narcotism followed nausea and vomiting Some varieties of *T chebula* are drastic purgatives.

Myrtaceae

(Myrtle and jamun Family)

Saponins, essential oils, tannins.

98 **Barringtonia acutangula** Gaertn (Vern - S. - Dhatriphal, B - Hijal, H -Hijjal, Bo -Samudraphala, M -Samutrapullam), *B asiatica* Kurz (*B speciosa* Forst), *B racemosa* Bl (Vern - S - Samudrapad, H -Norvishee, B -Samudraphal, M -Samudra).

These are believed fish poisons

99 **Careya arborea** Roxb. (Vern -S , H & B.-Kumbhi).

It is a fish poison The inner bark rubbed on shoes keeps off leeches.

100 **Eucalyptus globulus** Labill (Vern - M -Karpura maram)

The essential oil contained is an important ingredient of insecticides, internally gastro-intestinal irritant.

101 **Melaleuca leucadendron** Linn. (Vern - H., B & Bo.-Nim, M - Vembu)

Its essential oil is an irritant and a mosquito repellent.

Lythraceae

(Henna and pomegranate Family)

Acrid principle

102 **Ammannia baccifera** Linn (Vern - S - Agnigarva, H - Janglimendi, M - Nirumel-neruppu) *A. senegalensis* Lamk (Vern - P.-Fauglimehandi)

These are vesicant, taken internally they cause great pain.

103 **Lagerstroemia indica** Linn, *L speciosa* (Linn) Pers (*-L flosregineae* Retz) (Vern - S - Arjuna, H & B -Jarul, B - Taman, M Kodali)

The bark and leaves are purgative, seeds of the former species are narcotic

Samydaceae

(Casearia Family)

104 **Casearia graveolens** Dalz (Vern - H - Chilli, Bo - Naro), *C. tomentosa* Roxb (Vern - H - Chillara)

The pounded fruit is used as a fish poison

Caricaceae

(Papaw Family)

Carpaine, carposide, caricin in seeds yielding essential oil on hydrolysis, papain

105 **Carica papaya** Linn (Vern - H - papaya, B -Papey, Bo.-Papai, M.- Pappayi)

The seeds are believed to have powerful emmenagogue properties and are used as abortifacient The juice of unripe fruit is acrid or even vesicant

Passifloraceae

(Passion-flower Family)

Hydrocyanic acid, saponins

106. **Adenia (modecca) palmata** Engl., *A wightiana* Engl.

The roots and fruits are poisonous. Deaths from eating fruits of *A palmata* are reported

Cucurbitaceae

(Cucumber Family)

Bitter substances, such as colocynthin, alkaloids, glucosides.

107. **Citrullus colocynthis** Schrad, (Vern - S -Indra varuni, H. & Bo.-Indrayan, B.-Makhal, M -Peyt-tumatti) *C vulgaris* Schrad. (Bitter variety).

The fruit is purgative, *C colocynthis* is a drastic purgative and has produced fatal results, the dust when dry is very irritating to eyes and nostrils.

108. **Corallocarpus epigaeus** Benth. & Hook f.

Its fruit is a drastic purgative

109. **Cucumis sativus** Linn (Bitter variety) (Vern -S -Sukasa, H -Khira, B -Sasa, Bo -Kankri, M -Mulluvellari), *C trigonus* Roxb (Vern -S -Vishala, H -Bislambhi, M -Hattut-tumatti)

The fruit is a purgative, *C trigonus* excessively so

110 **Lagenaria siceraria** Standl (*L vulgaris* Seringe) (Wild variety)

It is drastic purgative Cases have been reported where beer kept in bottle gourd produced poisoning

111 **Luffa acutangula** Roxb. var *amara* C B Clarke (Vern -S -Koshataki, H & Bo -Torai, B -Jhinge, M -Pikumkai), *L aegyptiaca* Mill. ex. Hook f (Vern -S -Rajkoshataki, H -Ghiatarui, B -Dhundul, Bo -Ghosali, M -Guttibira) (Wild variety), *L echinata* Roxb (Vern -S -Koshataki, H -Kukarlata, B -Ghosalata, Bo -Kukarwele, M -Panibira).

The fruit of *L. acutangula* var *amara* is violently emetic and purgative and is not eaten. Others are also purgative

112 **Momordica balsamina** Linn (Vern -Bo -Kurelo-jangro), *M charantia* Linn (Vern -S -Sushavi H -Karela, B -Karala, Bo -Karla, M -Pavakkachedi), *M tuberosa* Cogn (—*M cymbalaria* Fenzl) (Vern -Bo -Kadavanchi)

Fruit of *M balsamina* is fatal to dogs, death occurs from violent vomiting and purging produced by the juice of this plant. *M charantia*, roots are used as abortifacient Decoction of roots of *M. tuberosa* are used as abortifacient

113. **Trichosanthes bracteata** Voigt (—*T palmata* Roxb) (Vern -S -Mahakal, H -Lal-indrayan, B.-Makal, Bo -Kaundal, M -Korattai) *T. cucumerina* Linn. (H -Jangli-chichonda, S -Patola, B -Bonpatol, Bo -Ranparul, M.-Pudel), *T dioica* Roxb (Vern -S -Patola, H.-Parvar, B.-Potal, Bo -Potala, M -Kombupudalai)

The roots are powerful cathartic Fruit of *T cucumerina* is never eaten, because of powerful cathartic action Fruit of *T. bracteata* is used as cattle poison and to destroy crows

114 **Zanonia indica** Linn. (Vern -S.-Dirghapattra, H -Chirpoti, Bo.-Chiraputi, M.-Penarvalli).

The fruit is very acrid and a powerful cathartic

Begoniaceae

(Begonia Family)

111 **Begonia rex** Putzeys

Its juice is poisonous to leeches.

Ficoideae

116. **Trianthema portulacastrum** Linn. (*T. monogyna* Linn), *T pentandra* Linn (Vern -P. & Bo -Bishkapra).

The roots are irritant and cathartic The leaves and stems are used as pot herb occasionally said to produce paralysis and diarrhoea

Umbelliferae

(Carrot and coriander Family)

Essential oils, cicutoxin, cicutoximin, vellerin

117. **Apium graveolens** Linn (Vern -S & H.-Ajmoda, B.-Chanu).

The seeds are irritant and poisonous in large overdoses.

118. **Centella asiatcia** (Linn) Urb (*-Hydrocotyle asiatica* Linn)

It is stupefying narcotic in large doses and a cumulative poison

119. **Cicuta virosa** Linn

The plant is the cause of extensive poisoning in Europe. The active principle belongs to picrotoxin groups of poisons which are convulsant.

120 **Daucus carota** Linn (Vern:-S -Shikha-mulam, H., B. & P.-Gajar, M -Gajjara kelangu)

The seeds are used for procuring abortion and the tuberous roots are eaten

121. **Hydrocotyle javanica** Thunb.

It is stated to be a fish poison.

Araliaceae

(Ivy and Panax Family)

Resin, a-hederin saponin.

122. **Hedera helix** Linn (Vern.-H.-Lablab, P.-Banda, Kash.-Karmora).

The decoction made from leaves is used to kill lice and other poisonous properties are also assigned to the plant.

Caprifoliaceae

(Honey-suckle family)

Sambucine, cyanogenetic glucoside, sambunigrin, bitter substances, resin (cathartic).

123. **Sambucus ebulus** Linn , (Vern -P —Mushkiara), *S. nigra* Linn

Both are strongly purgative *S ebulus* has foetid smell when bruised and is not eaten by cattle, poisoning amongst boys and fowls reported

Rubiaceae

(Madder and coffee Family)

Quinine, quinidine, cinchonine, cinchonidine, caffeine, emetine, cephaeline, ipecacuanhin, essential oils, saponins.

124 **Adina cordifolia** Benth & Hook f (Vern —S.-Dharakadamba, H -Hardu, B -Keli kadam, M -Manja kadambe)

Its juice is used as insecticide

125 **Cinchona calisaya** Wedd and var. *ledgeriana* Howard, *C officinalis* Linn f , *C succirubra* Pavon

The cinchona alkaloids are general protoplasmic poisons and parasiticide The plants are believed fish poisons.

126 **Coffea arabica** Linn (Vern —H -Coffee, B.-Kafi)

Its excessive indulgence produces harmful effects and chronic poisoning

127 **Cephaelis ipecacuanha** Stokes

It is emetic and irritant and a powerful cardiac depressant

128. **Randia dumetorun** Lamk , (Vern —S -Madan, H -Mainphal, B -Menphal, Bo -Gelaphal, M -Maruk-kallan-kai), *R uliginosa* DC (Vern —S -Pindaluka, H -Pindalu, B -Piralu, Bo -Pendari, M - Wagata)

These are believed fish poisons *R dumetorum* is used to preserve grain from attacks of insects, it is also used as abortifacient.

Compositae

(Sun-flower Family)

Essential oils, artemisin, santonin, bitter substances (absinthin lactucin, etc), saponins, resin, senecio alkaloids, xanthostrumarin, pyrethrins.

129. **Anthemis cotula** Linn.

It is an undesirable food for livestock, it is acrid and vesicant.

130. **Artemisia absinthium** Linn. (Vern —H. & Dec -Vilayati afsantin), *A maritima* Linn (Vern —S -Gadadhar, H -Kirmala, Bo - Kiramani owa), *A. vulgaris* Linn (S -Nagadamani, H -Nagadouna, B -Nagdona).

The essential oil from *A absinthium* is a violent narcotic poison producing convulsions *A maritima* is an irritant poison when given in large doses, and fatal cases are reported *A vulgaris* produces epileptiform spasms in man and is also reported as fish poison.

131. **Centratherum anthelminticum** O. Ktze.(-*Vernonia anthelmintica* Willd).

It is used as insecticide and insect repellent.

132. **Chrysanthemum cinerariæfolium** Vis., *C. coccineum* Willd., *C. roseum* Adam.

All are reputed insecticides

133. **Erigeron canadensis** Linn.

It is an irritant.

134 **Eupatorium odoratum** Linn.

It is stated to be a fish poison. *E urticifolium* L. f. of foreign countries produces acidosis and trembles in sheep and cattle

135. **Gnaphalium luteo-album** Linn (Vern –P -Balraksha).

It is suspected to produce livestock-poisoning in South Africa.

136. **Inula graveolens** Desf.

It is suspected to be poisonous to livestock.

137. **Lactuca tatarica** C A Meyer var *tibetica* C B Clarke

It is occasionally browsed by sheep, sometimes injurious

138. **Saussurea lappa** C.B Clarke (Vern —S.-Kushtha, H -Kut, B.-Pachak, Bo.-Ouplate, M.-Goshtam).

The roots are used to protect woollen fabrics against insects.

139. **Senecio** *species* (*S vulgaris* Linn introduced plant)

The Indian species should be studied in Ragwort poisoning due to several species of this plant is well known in foreign countries, some species produce hepatic cirrhosis

140. **Sphaeranthus indicus** Linn (Vern —S.-Munditika, H & Bo -Gorakmundi, B.-Murmuria, M -Kottak).

It is believed to be a fish poison.

141. **Xanthium strumarium** Linn. (Vern —S.-Arishta, H.-Chhotagokru, B -Bon-okra, Bo.-Shankesh vara, M -Marlu-mutta)

It is reported to be poisonous to cattle and pigs in America and Australia.

Campanulaceae

(Bell-flower Family)

Alkaloids.

142. **Lobelia excelsa** Leschen, *L. nicotianifolia* Heyne. (Vern—Bo.-Dhavala, M.-Rattu papillay).

It is irritant to nose and deaths have been reported in man. Its action resembles nicotine, except that more burning pain in the stomach is produced. It is used as substitute for datura.

Ericaceae

(Rhododendron Family)

Andromedotoxin, ericolin, Essential oils

143. **Gaultheria fragrantissima** Wall.

It is an irritant poison and deaths are reported from its use as abortifacient

144 **Pieris ovalifolia** D. Don.

It is poisonous to goats and is an insecticide.

145 **Rhododendron anthopogon** D Don , (Vern —Kash.-Tazak-tsun), *R arboreum* Sm (Vern -P —Ardawal), *R. barbatum* Wall (Vern — Nep -Guras), *R campanulatum* D Don (Vern -H -Cheraelu Kash -Gaggar), *R cinnabarinum* Hook f (Vern.--Nep -Bulu), *R falconeri* Hook f (Vern —Nep -Kurlinga), *R. setosum* D. Don (Vern - Bhutia -Tsallu).

These are probably all poisonous to live-stock; some are reported to be fish poisons and honey from some is reported to be poisonous.

Plumbaginaceae

(Plumbago Family)

Plumbagin

146 **Plumbago indica** Linn (*-P. zeylanica* Linn) (Vern:-S —Chitraka H & B -Chita, Bo -Chitaro, M -Chittira), *P rosea* Linn. (Vern — S -Chitraka, H., B. & Bo -Lal chitra, M.-Chittur-mol).

These are strong irritants externally and internally and are used to procure abortion.

Primulaceae

(Prime-rose Family)

Saponins

147. **Anagallis arvensis** Linn. (Vern —H.-Jonkhmari)

It produces gastro-enteritis in dogs and horses. It is used to poison fish and expel leeches from nostrils of animals.

148 **Cyclamen persicum** Miller (Vern --Ind. Baz.-Bankhur-i-Miryam).

It is believed to be a fish poison.

149. **Primula reticulata** Wall. (Vern -Kumaon.—Bishcopra).

It is stated to be poisonous to cattle.

Myrsinaceae

(Ardisia Family)

Saponins.

150 **Maesa indica** Wall. (Vern —M -Kirithi) Leaves are stated to be fish poison.

Sapotaceae

(Sapodilla and mohwa Family)

Saponins

151 **Madhuca indica** J F Gmel *M* (*Bassia*) *latifolia* (Roxb.) Macbride, *M. longifolia* (Linn) Macbride.

The residual cake is used as fish poison and is said to have insecticidal properties *Mohwa meal* is used to kill worms on lawns

Ebenaceae

(Ebony Family)

Bitter substances

152 **Diospyros ebenum** Koenig (Vern – H & Bo -Tendu, M -Acha), *D. montana* Roxb (Vern -S --Tumala, H -Lohari, B -Ban-gal, Bo Kundu, M -Muchi-tanki), *D paniculata* Dalz (Vern —S -Thinduka, M -Karinthuvari)

All these are believed to be fish poisons.

Salvadoraceae

(Salvadora Family)

153 **Salvadora oleoides** Dcne (Vern —H , S. & B -Pilu, Bo.-Kankhina, M -Ughaiputtai), *S persica* Linn (Vern —S -Pilu, H & B -Chota-pilu, Bo -Pilvu, M -Ughaiputtai)

The root bark is vesicant

Apocynaceae

(Dog-bane and Oleander Family)

Kurchi and rauwolfia alkaloids, glucosides, e g cerberin, karabin, nerin, neriodorin, oleandrin, I-strophanthin, thevetin, etc., bitter substances

154 **Allamanda cathartica** Linn. (Vern —Bo -Jahari sontakka) It is a hydragogue cathartic

155. **Cerbera manghas** Linn (*-C odollam* Gaertn) (Vern.—B.-Dhakur, M -Katarali)

The green fruit is used to poison dogs. The seeds are irritant poison and the plant is a fish poison

156 **Ervatamia dichotoma** (Roxb) Blatter (*-Tabernaemontana dichotoma* Roxb)

The seeds are powerfully narcotic and poisonous.

157 **Holarrhena antidysenterica** Wall (Vern – S -Kutaja, H Karchi, B -Kurchi, Bo -Pandhrakura, M -Kashappu-vetpalarishi)

It is not browsed by cattle and goats, it is an anthelmintic, kurchicine is a general protoplasmic poison.

158. **Lochnera pusilla** K Schum (-*Vinca pusilla* Murr.), *L rosea* (Linn) Reichb (-*Vinca rosea* Linn)

These are cardiac poisons and *L pusilla* is regarded as poisonous to cattle.

159 **Melodinus monogynous** Roxb (Vern —B -Sadul kou)
It is a fish poison

160 **Nerium-indicum** Mill. (*N odorum* Soland) (Vern —S -Karavi, H -Karber, B -Karabi, Bo -Kanhera, M Alari)

It is very poisonous and is used for suicidal purposes and to procure abortion, depresses nervous system and heart

161 **Plumeria acuminata** Ait (-*P acutifolia* Poir) (Vern —S -Kshira champa, H & Bo -Khair Champa, B -Gobar champa, M -Vadaganneru)

Its milk is rubefacient and is used to procure abortion, internally it acts as purgative and is poisonous

162 **Rauwolfia serpentina** Benth ex Kurz (Vern —S -Sarpagandha, H -Chota chand, B & Bo -Chandra, M -Covannamilpori)

It is hypnotic and a fish poison

163 **Thevetia peruviana** (Pers) Merr (-*T neriifolia* Juss) (Vern —H & Bo-Pila-kaner, B -Kolkaphul, M -Pachch-ai-alari)

All parts especially seeds are very poisonous It is used to poison cattle and produces violent vomitting and purging Action on heart like digitalis It is also a fish poison

Asclepiadaceae

(Milk-weed Family)

Tylopharine, glucosides, bitter substances, resins, saponins, etc

164 **Asclepias curassavica** Linn (Vern —H -Kakatundi, Bo-Karki)

It is a fish poison, emetic and cathartic

165 **Calotropis gigantea** R Br (Vern —S -Arka, H -Ak, B & Bo -Akanda, M -Erukku), *C procera* R Br (Vern - S Alarka, H -Madar, P -Shakar-al-lighal, Bo -Mandara, M -Vellerku)

Milk of these plants is drastic purgative and caustic Stated to be used for suicidal and homicidal purposes and as an abortifacient and cattle poison

166 **Cryptostegia grandiflora** R Br (Vern —Bo -Vilayati-vakhandi, M.-Palai).

A fatal case from leaves is reported in which persistent vomiting was observed

167. **Cynanchum arnottianum** Wight, *C. vincetoxicum* Pers.

C. arnottianum is used as insecticide. *C. vincetoxicum* is not eaten by cattle and is regarded poisonous Its root is emetic.

168 **Sarcostemma acidum** (Roxb.) Voigt (*-S. brevistigma* W. & A.) (Vern —S & Bo -Soma, H. & B -Somlata, M.-Kondapala).

Stated to have insecticidal properties.

169 **Secamone emetica** R. Br. (Vern —B -Shada-buri).

The roots are acrid and the plant is powerfully emetic.

170 **Tylophora indica** (Burm f) Merr (-T *asthmatica* Wight and Arn), (Vern -H. & B -Antamul, Bo -Anthamul, M.-Nay-palai), *T. fasiculata* Buch.-Ham (Vern -Bo -Bhuidari)

Fatal cases are reported in man, it is emetic and *T. fasciculata* is used as rat poison.

Loganiaceae

(Nux-vomica Family)

Strychnine, brucine, etc.

171 **Strychnos colubrina** Linn (Vern —H & B -Kuchila-lata, Bo -Goagarilakei, M -Nagamusadi), *S nuxvomica* Linn. (Vern —S -Visha-mushti, H -Kuchla, B -Kuchila, Bo.-Kajra, M -Yetti)

These are poisonous *S nux-vomica* seeds are used as fish poison and are source of strychnine, one of the deadliest poisons known Suicidal and homicidal cases are recorded. It is employed to kill dogs, rodent, etc

Boraginaceae

(Borage and Sebestan Family)

Alkaloids.

172 **Heliotropium eichwaldii** Steud (Vern —H & P -Nilkattei, Kash.-Chirghas), *H indicum* Linn (Vern —S -Hastisunda, H. & B -Hatisura, Bo -Burundi, M -Tel-kodukki)

These are suspected to be poisonous

Convolvulaceae

(Convolvulin, pharbitin, terpithin, terpethein, cucutalin, resin.

173 **Calonyction muricatum** (Linn.) G Don (*-Ipomoea muricata* Jacq)

See Ipomoea.

174 **Convolvulus arvensis** Linn (Vern —H -Hiranpadi, Bo -Hiran-pag, M -Naranji), *C scammonia* Linn.

The roots are strongly purgative

175. **Cuscuta reflexa** Roxb. (Vern —S Amaravela, H.-Akasbel, B -Algusi, Bo.-Nirmuli, M -Sitamapurgonalu).

It is nauseant and emetic, it is used to procure abortion.

176 **Ipomoea reptans** (Linn) Poir (*-I aquatica* Forsk) (Vern — S -Kalambi, B -Kalmisak, Bo.-Nalichi baji, M -Sarkarei-valli), *I. nil* Roth (—*I. hederacea* Jacq.) (Vern —H., B. & Bo -Kaladana, M -Jirkivirai), *I purga* Heyne,

These are strongly purgative and irritant poisons in overdoses.

177. **Operculina turpethum** (Linn) Manso (—*Ipomoea turpethum* R. Br.)

See Ipomoea

Solanaceae—

(Datura and nightshade Family)

Hyoscyamine, hyoscine, atropine

178 **Atropa belladonna** Linn. (Vern —H -Sag-angur, B -Yebruj, Bo.-Girbuti)

Fatal cases of poisoning are reported, dryness of mouth and throat, dilation of pupils and delirium characteristic features

179 **Capsicum annuum** Linn. (Vern —H. & P -Mirch), *C frutescens* Linn., *C. minimum* Roxb.

The seeds are gastro-intestinal irritant and are used for torturing purposes

180. **Datura fastuosa** Linn. (Vern —S -Krishna dhatura, H & B -Kala Dhatura, M -Karu umattai), *D metel* Linn , *D. stramonium* Linn (Vern —B.-Sada dhutura, P -Tattu dattura, M -Umatai).

These are commonly used by criminals for stupefying their victims , symptoms resemble those of Atropa

181. **Hyoscyamus muticus** Linn , *H. niger* Linn (Vern —S -Parasikaya, H -Khurasani ajvayan, B.-Khorasani ajowan, Bo -Khorasani-owa, M -Khorasani-yomam), *H. pusillus* Linn , *H. reticulatus* Linn.

Cases of livestock and children poisoning are on record, action like Atropa.

182 **Lycium barbarum** Linn. (Vern —Baluchi -Koh-tor)

It is reported to be poisonous to livestock

183 **Mandragora caulescens** Clarke

It is suspected to be poisonous

184 **Nicandra physaloides** Gærtn.

It is considered to be insecticide

185 **Nicotiana rustica** Linn. (Vern —H & B -Vilayeti tamaku, P -Kakkar tamaku), *N. tabacum* Linn (Vern —H -Tamaku, B -Tamak., Bo -Tambaku, M.-Pugai-ilai)

These are insecticide and are also used to ward off leeches fatal cases reported among human beings and livestock.

186. **Physochlaina praealta** Miers (Vern —P -Nandru)

It is reported to be poisonous.

187. **Scopolia anomala** (Linn et Otto) Airy-Shaw. (*S. lurida* Dunal).

It is poisonous and action is like Atropa.

188 **Solanum dulcamara** Linn (Vern —S Kakmachi, P -Ruba-barik) *S. incanum* Linn , (-*S coagulans* Forsk), *S. nigrum* Linn (unripe berries) (Vern —S & B -Kakmachi, H -Makoi, Bo -Mako, M -Manattak kali), *S spirale* Roxb (Vern —H -Mungas kajur, Bagua), *S. tuberosum* Linn (Sprouting).

The cases of poisoning among human beings and animals are reported, some fatal These are gastro-intestinal irritant , occasionally associated with atropa-like symptoms

189 **Withania somnifera** Dunal (Vern —S , B & Bo -Ashwagandha, H -Asgandh, M -Amku-lang-kalang)

It is reported to be used as abortifacient and as an insecticide, it is also stated to be hypnotic

Scrophulariaceae

(Mimulus and Digitalis Family)

Digitalin, digitonin, digitoxin, gitalin, gitonin, etc , saponin, bitter substance

190 **Digitalis purpurea** Linn

It is a cardiac poison and fatal cases due to eating of plant are reported in India

191 **Verbascum thapsus** Linn

It is a fish poison and the seeds are narcotic

Bignoniaceae

(Bignonia Family)

192 **Dolichandrone falcata** Seem. (Vern.-H —Hawar, Bo -Manchingi, M -Kadatathie)

It is fish poison reputed to be abortifacient.

Pedaliaceae

(Sesamum Family)

Sesamol (a phenolic substance), seasmolin

193 **Sesamum indicum** Linn (-*S orrientale* Linn) (Vern.—S , H , B , & Bo -Til, M -Yellucheddie

Seed cakes are commonly fed to cattle in India but it is stated to be toxic to livestock in Europe producing colic, tremors, dyspnoea and distantion.

Verbenaceae

(Verbena and teak Family)

194. **Callicarpa longifolia** Lamk var *lanceolaria* C. B. Clarke.
It is a fish poison

195 **Duranta repens** Linn (*D plumieri* Jacq)

It is very bitter and believed to be poisonous to livestock, but generally refused

196. **Lantana aculeata** Linn (*-L camara* Linn) (Vern -Bo.—Vhaneri, M -Aripu)

The reports about being poisonous to livestock received from the Punjab and Assam Government Departments described it to be abortifacient.

196A **Stachytarpheta jamaicensis** (Linn) Vahl var. *indica* Lam (*-S indica* Vahl.)

It is described as abortifacient

197 **Verbena officinalis** Linn (Vern.—P -Pamukh)

It is stated to be irritant poison

Labiateae

(Mint and sage Family)

Essential oils, saponins

198. **Eremostachys acanthocalyx** Boiss , *E vicaryi* Benth (Vern —P.-Gurgunna)

E acanthocalyx is stated to be poisonous and *E vicaryi* is used as a fish poison

199 **Lamium amplexicaule** Linn

It is regarded as injurious in America

200 **Pogostemon heyneanus** Benth (*P patchouli* F B I , non Pelletier) (Vern --H -Pacholi)

The leaves are used as insecticide.

Chenopodiaceae

(Spinach and beet Family)

Essential oils, saponins, salsoline, oxalic acid

201 **Chenopodium ambrosioides** Linn, *C botrys* Linn

Anthelmintic against hook-worm and round worm Fatal poisoning is on record

202 **Haloxylon recurvum** Benge ex Boiss , *H salicornicum* Bunge ex Boiss.

These are stated to be poisonous but *H. recurvum* is a favourite food of camels

203 **Salicornid brachiata** Roxb (Oomarie keeray) Ash is stated to be abortifacient

204. **Salsola kali** Linn.

It is suspected to be poisonous but a feeding test with half dried plants in flowering stage was negative

205 **Suaeda fruticosa** Forsk

It is stated to be poisonous.

Phytolaccaceae

(Phytolacca Family)

Bitter substances.

206 **Phytolacca latbenia** (Buch-Ham) H Walt. (*P. acinosa* Hook f., B I non-Roxb.) (Vern —H.-Matazor)

It is stated to be poisonous if eaten raw, but it is edible when cooked.

Polygonaceae

(Buck-wheat and rhubarb Family)

Rutin, essential oils, anthra-quinone derivatives, oxalic acid, oxalates.

207. **Fagopyrum esculentum** Moench, *F. tataricum* Gaertn.

It is commonly eaten but under certain conditions, not properly understood at present, produces eruptions and urticaria.

208 **Polygonum aviculare** Linn. (Vern —S -Nisomali, H -Bannatia, B -Machutie), *P flaccidum* Meissn , *P hydropiper* Linn. (B -Packur-mul), *P orientale* Linn., *P persicaria* Linn., *P. tomentosum* Willd

P hydropiper is biting to a degree that no animal will eat it. Acrid, emetic, vesicant, insecticidal and piscicidal properties to varying degree are strongly suspected.

209 **Rheum emodi** Wall (Vern —H. & B -Revandchini, Bo.-Ladaki-revanda chini, M -Nattu-ireval-chinni) and probably some other species

Its leaves and petioles are edible but the latter is responsible for occasional poisoning

210 **Rumex acetosa** Linn , *R acetosella* Linn (Vern —S. Chutrika, B -Chukapalam)

Oxalic acid poisoning is produced if eaten in excess.

Aristolochiaceae

(Birth-wort Family)

Aristolochin, glucoside, essential oils, bitter substance.

211. **Aristolochia bracteata** Retz. (Vern —S.-Dhumrapatra, H.-Kirmar, M -Adutina-palai), *A. indica* Linn (Vern —S -Rudra-jata, H. & B -Isharmul)

These are nauseous and bitter, emmenagogue and abortifacient. *A bracteata* is insecticide

Piperaceae

(Pepper Family)

Essential oils, piperine, piperovatine

212. **Piper sp.**

Harmful effects of *P. betle* Linn. and *P. nigrum* Linn. are well known.

Myristicaceae

(Nutmeg Family)

Essential oil (with myristicin), saponins.

213 **Myristica fragrans** Houtt., *M malabarica* Lamk., (Vern —Bo.-Ramphal), possibly some others also.

These are narcotic and occasional cases of poisoning are reported

Lauraceae

(Laurel Family)

Essential oils

214. **Cassytha filiformis** Linn

It is stated to be used as insecticide

215 **Cinnamomum camphora** F Nees (imported).

It is protective against moths, counterirritant, systemically stimulates then depresses and paralyses central nervous system.

Thymeliaceae

(Mezereum Family)

Saponins

216 **Daphne cannabina** Wall , *D oleoides* Schreb

These are severe gastro-intestinal irritants. Camels do not eat *D oleoides*

217. **Edgeworthia gardneri** Meissn.

It is a fish poison

218. **Lasiosiphon eriocephalus** Decne

Dust from dried plant very irritant and is not eaten by livestock It is a fish poison

219. **Wikstroemia viridiflora** Meissn. (*W. indica* C. A Mey, var. viridiflora Hook. f)

It is a fish poison.

Loranthaceae

(Mistletoe Family)

220. **Viscucm album** Linn and possibly others.

The poisonous properties are probably acquired if growing on poisonous hosts, e g. *Strychnos nux-vomica.*

Euphorbiaceae

(Croton and eastor oil Family)

Cyanogenetic compounds, saponins, crotononside, ricinine, essential oils, euphorbon, phenolic substance, resins, toxalbumins

221. **Andrachne cordifolia** Muell. Arg. (Vern·—P.Gurguli).

It is reported to be cattle poison and African species are used as insecticide.

222 **Baliospermum montanum** Muell , Arg (-*B axillare* Blume) (Vern —S , H , & B -Dantı, Bo -Dantımul, M.-Naga-danti).

Its seeds and oil are drastic purgative and the seeds in overdoses are acro-narcotic poison

223 **Buxus sempervirens** Linn. (Vern —Kash -Chikri, P -Papri)

It is stated to be fatal to camels, cattle and goats are probably immune

224 **Chrozophora rottleri** A Juss ex Spreng (-*C tinctoria* Hook f in part) (Vern —H -Subali, P -Kukronda)

It is emetic and cathartic and animals avoid it

225 **Cleistanthus collinus** Benth. & Hook f (Vern —M -Nachuta)

It is used as fish poison and occasionally as human poison An extract of it is a voilent gastro intestinal irritant.

226 **Croton oblongifolius** Roxb , *C tiglium* Linn (Vern —S - Kanakaphala, H -Jamalgota, B-Joypal, Bo.-Geyapal, M - Nervalam)

The seeds especially and the oil also is a drastic purgative and is reported to be poisonous The seeds are stated to be used as insectide and piscicide

227 **Euphorbia acaulis** Roxb , *E antiquorum* Linn (Vern —S - Vajrakantaka, H -Tridhara-sehund, B -Tekata sij, Bo -Naraseja, M -Shadhurakkalli), *E cattimandoo* W Elliot, *E heliioscopia* Linn , *E hirta* Linn , *E hypericifolia*, Linn (Vern —Bo -Nayeti, P -Hazardana), *E nerifolia* Linn (Vern —S -Snuhi, H -Sehund, B -Mansa-sij, Bo.-Minguta, M -Ilaikal, *E. nivulea* Buch Ham , (S patta karie, B -Sij, Bo -Newrang, M -Aku-jemudu), *E peplus* Linn *E pilosa* Linn., *E rothiana* Spreng , *E royleana* Boiss (Vern — H. & P -Shakar pitan), *E thomsoniana* Boiss , *E thymifolia* Linn (Vern —S -Racta-vinda-chada, H -Chhoti dudhi, B - Dudiya, Bo -Nayeti, M -Sittrapaladi), *E tirucalli* Linn (Vern:— H.-Sehud, B -Lankasij, Bo -Shera, M -Kombu-Kalli), *E trigona* Haw.

Acrid and vesicant juice is found in most species and some are used as abortifacient when applied locally *E antiquorum*, *E. nerifolia*. *E royleana*, *E tirucalli* are fish poisons and *E. antiquorum*, and *E thymifolia* are stated to be used as insecticides, some are poisonous to livestock

228. **Excoecaria agallocha** Linn (Vern —B -Gangwa, B -Geva, M - Chilla)

Its fresh sap is extremely acrid and causes intolerable pain if it gets into the eye The woodcutters have suffered and call it blinding tree It is also fish poison

229 **Fluggea leucopyrus** Willd , *F virosa* Baill (*F microcarpa* Bl) (Vern —H.-Dalme, B Pandharphali)

It is a fish poison and is used to destroy worms in sores.

230. **Hura crepitans** Linn.

Its seeds and oil are violent purgative, milky juice very irritant

231 **Jatropha curcas** Linn , *J glandulifera* Roxb , *J gossypifolia* Linn , *J multifida* Linn

These are violent purgative like *Croton* sp and *J curcas* is a fish poison

232 **Manihot esculenta** Crantz (*M utilissima* Pohl.) (Vern —Baz - Cassarva M -Maravuli

Its fresh tubers are extremely poisonous and cassava or tapioca meal is specially prepared

233 **Phyllanthus urinaria** Linn (Vern —S -Tamara valli, H & B - Hazarmani, M -Shiyappunelli)

It is stated to be a fish poison

234 **Ricinus communis** Linn (Vern —S -Eranda, H -Arand, B.- Verenda, Bo -Erendi, M -Amanakham chedi)

The seeds produce violent grastro-enteritis, subcutaneously very poisonous The oil is stated to be an active poison for flies

Plant is a fish poison

235 **Sapium indicum** Willd (Vern —B -Hurua, Bo -Hurna), *S - insigne* Trimen (Vern —H -Khinna, Bo -Dudla)

S indicum juice is narcotic poison The fruit is extremely nauseous and seeds are fish poison *S insigne* juice is vesicant

236 **Tragia bicolor** Miq , *T involucrata* Linn (with varieties) (Vern — S -Vrischikali, H -Barhanta, B -Bichuti, Bo -Kanchkuri, M - Kanchuri-vayr)

These are stinging nettles

Urticaceae

(Nettle, hemp and mulberry Family)

α-β & γ-antiarin, saponin, resin containing cannabindol (toxic), formic acid

237 **Antiaris toxicaria** Lesch O (Vern —Bo -Chandla, M.- Nettavil, Sing -Riti, Burma -Hmyaseik)

Its sap is used as an arrow poison, it is a powerful heart poison

238 **Cannabis sativa** Linn. (Vern —S -Ganjika, H , B & Bo-Ganja, M -Bhangi)

The preparations of bhang, charas and ganja are well known in India, excessive indulgence produces physical and mental injury. Stated to be used as a fish poison in Bengal It is spread on beds to drive away bugs.

239. **Ficus sp.**

Some species contain acrid juice, according to Watt fruit of *F bengalensis* is poisonous to horses.

240. **Fleurya interrypta** Gaud.

Stings.

241. **Girardinia leschenaultiana** Dcne , *G zeylanica* Dcne.

These are stinging nettles.

242. **Laportea crenulata** Gaud., *L terminalis* Wight

These are stinging nettles

243 **Urtica dioica** Linn , (Vern —H & P -Bichu), *U. hyperborea* Jacq., *U parviflora* Roxb., *U. pilulifera* Linn.

These are stinging nettles.

Juglandaceae

(Walnut Family)

244. **Juglans regia** Linn (Vern —S -Akshota, H. & B -Akhroot, Bo -Akroda, M -Akrottu).

The rind of unripe fruit is stated to be fish poison in Jaunsar and Tehri Garhwal

Myricaceae

(Sweet-gale Family)

Essential oils, myricelin).

245. **Myrica nagi** Thunb. (Vern—S.-Katphala, H., B & Bo -Kaiphal, M.-Marudam-pattai).

Bark is stated to be used as fish poison in Khasia hills.

Gnetaceae

(Gnetum Family)

Saponins, bitter substance

246 **Gnetum scandens** Roxb.

It is a fish poison

Coniferae

(Pine Family)

Essential oils, taxine, taxicatin.

247 Several members especially *Taxus baccata* Linn.

Most members possess toxic essential oils and poisoning due to the use of Juniper oil as abortifacient is reported. Deaths in man and animals due to eating the berries and leaves of *T. baccata* are reported and the seeds are very poisonous.

Iridaceae

(Iris Family).

Saponins, picrocrin (bitter substance), essential oils.

248. **Crocus sativus** Linn. (Vern·—S -Kumkuma, H., B & Bo -Jafran, M · Kungumapu)

Bulbs are toxic to young animals and stigmas in overdoses are narcotic poison. It is used as abortifacient

Amaryllidaceae

(Amaryllis and Agave Family)

Saponin, Lycorine, tazetitne.

249 **Agave americana** Linn (Vern —H -Kantala, B -Jungli anarash)

It is stated as fish poison and it is also stated to be toxic to livestock under field conditions The wall paper impregnated with expressed juice is said to be proof against white-ants

250 **Cirnum asiaticum** Linn Vern —S -Vishamandala, H Pindar B -Bara-kanur, Bo -Nagdowan, M -Vishomangil), *C. lotifolium* Linn (Vern —H & B Sukh-darsan)

251 **Narcissus tazetta** Linn (Vern —P Nargis)

Its bulbous roots are emetic and purgative, and produce irritant poisoning in overdoses

Taccaceae

252. **Tacca pinnatifida** Forst (Vern —Bo -Diva, M -Karachunai)

The tuber is intensely bitter, acrid and poisonous when fresh, yields nutritious starch by maceration and repeated washing.

Bromeliaceae

(Pine-apple Family)

253 **Ananas comosus** Merr (--*A sativus* Schult) (Vern —H.-Anannas, B.-Anaras, M -Anasha pazham).

The juice of leaves and unripe fruit is purgative and is sometimes used as abortifacient.

Dioscoreaceae

(Yam Family)

Dioscorine, glucoside (toxic).

254 **Dioscorea bulbifera** Linn. *D hispida* Dennst, (—*D. daemona* Roxb), *D prazeri* Prain & Burk (—*D deltoidea* Wall.).

Tubers are very acrid but in most cases boiling makes them edible

Liliaceae

(Lily Family)

Imperialine, colchicine, methyl-colchicine, saponine, borbaloin emodin, sicaloin, resin, essential oils, etc.

255. **Allium sativum** Linn (Vern —S -Lasuna, H. & Bo. Lasan, B.-Rasun, M.-Vallaipundu)

Its essential oil is very irritant and pungent, produces irritant poisoning in excess. It is also stimulant, narcotic and anthelmintic.

256 **Aloe species.**

Insipisated juice 'Mushabbar' of commerce is a powerful drastic purgative and fatal cases are reported from its use. It has been used to procure abortion

257. **Colchicum luteum** Baker (Vern —Surinjan)

Resembles closely the foreign *C autumnale* which is poisonous and produces gastro-intestinal irritation The Indian species is also poisonous

258 **Fritillaria imperialis** Linn.

The bulbs are toxic when fresh and are said to act as heart poison.

259 **Gloriosa superba** Linn (Vern —S -Sukra puspita, H -Kalihari, B -Bishlanguli, Bo.-Karianag, M -Agnişikha)

The root is stated to be sometimes used for suicidal purposes and as abortifacient, it is an acronarcotic poison The juice of leaves is stated to be used to destroy lice in hair

260 **Scilla iddica** Baker

.. ..

261. **Urginia coromandeliana** Hook. f., *U indica* Kunth (Vern —S.-Vana-palandam, H & B.-Jangli piyaz, Bo.-Jangli-kanda, M -Nari-vengayam)

The bulbs are irritant poison The foreign species *U. scilla* is a fish poison, as also is the Indian representative

Juncaceae

(Rush Family)

262 **Juncus effusus** Linn

It is suspected to be poisonous to livestock in South Africa This and other species in India are worth investigating.

Palmaceae

(Palm Family)

Arecaine, arecolidine, arecoline, guvacine, guvacoline, saponins

263 **Areca catechu** Linn (Vern —S -Gubak, H & B -Supari, Bo -Sopari, M Kamugu)

Young and undried nut when chewed in excess gives rise to temporary giddiness It also produces gripping and strong intestinal irritation, resulting in loose motions.

264. **Arenga obtusifolia** Mart.

Juice from fruit is used in Malaya to poison enemies and *A obtusifolia* is stated to be used as fish poison.

265 **Corypha umbraculifera** Linn. (Vern —S.-Alpayushi, M.-Talipanai)

The fruit is stated to be used as fish poison.

266 **Wallichia disticha** T Anders.

Watt states that berries and perhaps the leaves are irritant to skin

Araceae

(Aroid Family)

Calcium oxalate (acicular crystals), bitter substance, sharp acrid substance, essential oil (alkaloid and saponin in foreign plant)

267 **Acorus calamus** Linn (Vern —H & B.-Bach, Bo -Vaj, M Vashambu), *A gramineus* Soland

Roots are stated to be used as effective isecticides and insectifuge Doubtful case reported when *A calamus* proved poisonous to camels during the Afghan Campaign The rhizome is used in medicine but in overdoses produces violent and persistent emesis.

268 **Alocasia indica** Schott (Vern —S.-Manaka, H -Mankenda, B -Mankachu), *A montana* Schott, *A odora* (Roxb) C Koch (-*A macrorhiza* Schott)

Fresh tubers are acrid and irritant

269 **Amorphonphallus campanulatus** (Roxb) Bl *A lyratus* Engl, *A sylvaticus* (Roxb) Kunth (*Sunantherias sylvatica* Schott)

Fresh tubers are acrid and irritant and the seeds are intensely acrid Seeds of *A sylvaticus*, like *Plesmonium*, and fruit are intensely acrid

270 **Arisaema speciosum** Mart (Vern —P -Kiralu), *A tortuosum* Schott (Vern —P -Samp-ki-kumb)

The tubers are poisonous and have insecticidal properties. The fruit is also probably poisonous

271 **Homalomena rubescens** Kunth

It is stated to be poisonous

272 **Lagenandra ovata** (Linn) Thw (-*L toxicaria* Dalz) (Vern:—Bo -Rukh-alu, M.-Maravara Tsjembu)

It is stated to be very poisonous and has insecticidal properties.

273. **Plesmonium margaritiferum** Schott

Its crushed seeds produce local anaesthesia and it is used as a cure for toothache

274 **Sauromatum guttatum** Schott.

Tubers are regarded as very poisonous.

275 **Steudnera virosa** (Kunth) Prain (-*Colocasia virosa* Kunth.).

It is believed to be poisonous.

276 **Thomsonia nepalensis** Wall

It is acrid when fresh.

277. **Typhonium trilobatum** Linn. Schott. (Vern.—B -Ghet-kachu, M.-Karunaikkizhangu).

Fresh tubers are exceedingly acrid

Cyperaceae

(Sedge Family)

Essential oil.

278 **Carex cernua** Boott

It is said to be one of the causes of 'vlei' poisoning in cattle in South Africa.

279. **Cyperus longus** Linn

It is regarded as poisonous in South Africa.

280. **Scirpus corymbosus** Heyne

See *Carex cernua.*

Gramineae

(Grass Family)

Cyanogenetic glucosides, hydrocyanic acid, temuline, saponins, oxalic-acid, selenium protein (toxic)

281 **Avena fatua** Linn (Vern —H -Kuljud), *A sativa* Linn

It is good fodder but occasionally deleterious probably on account of 'hair balls' that are developed in the stomach

282 **Bambusa bambos** Druce (-*B. arundinacea* Willd) (Vern:—S -Vansa, B & H -Bans, Bo.-Mandgay, M -Mangal).

Fresh young shoots are stated to have insecticidal properties

283. **Dendrocalamus strictus** Roxb.) Nees (Vern:—H -Bans kaban, B -Karail, Bo -Bas, M.-Kanka).

The leaves are stated to be used to procure abortion

284 **Lolium perenne** Linn., *L. temulentum* Linn (H -Machni)

Several cases of poisoning, mostly non-fatal in man and animals, from eating the seeds of *L temulentum* have been recorded Gastro-intestinal irritation and severe nervous symptoms are reported.

285. **Panicum maximum** Jacq

It is suspected to be responsible for the production of 'Dikoor', a disease affecting young sheep in Africa.

286. **Paspalum scrobiculatum** Linn (Vern —S.-Kodrava, H.-Kodo, B -Kodoa, dhan, Bo -Kodra, M -Kiraruga).

The 'kodra' poisoning is very similar to *L temulentum* poisoning, the animals suffer much more than men The animals should be prevented from grazing the crop when ripening

287 **Sorghum halepense** (Linn) Pers (Vern —H -Baru, B -Kalamucha), *S saccharatum* Pers (Vern —H & Bo -Deo-dhan, M -Tellajonna), *S vulgare* Pers (Vern —S Javanala, H , B & Bo -Jowar, M -Cholam)

These are good fodder but occasional poisoning is reported with stunted growth, under drought conditions. The frosted leaves, or second growth is dangerous

288 **Stipa sp** (some)

It is believed to be poisonous and mechanical action of 'seeds' may not be overlooked

289. **Triticum aestivum** Linn (Vern —S -Godhum, H -Gehun, Bo -Gam, M -Godumai)

Under certain conditions it becomes deleterious fodder.

290 **Zea mays** Linn (Vern —S -Yavanala, H. & B -Bhutta, Bo.-Makai, M -Makka-scholam)

The pollen from it is stated to be a possible cause of hay fever It is said to be occasionally responsible for deleterious effects, as yet not fully understood

CHAPTER IX

We have studied the literature on Indigenous Systems of Medicine as practiced in India with a view to find out the uses to which different medicinal plants have been put in the treatment of various important diseases For instance certain plants have the reputation of being effective in the treatment of bowel diseases particularly dysentery and cholera Others are said to be useful in the treatment of prolonged fevers such as the group of enteric fever Certain other plants are alleged to be effective in the treatment of tuberculosis (pulmonary) Yet another group is considered to have action on genital organs and emenagogue and abortifacient properties are attributed to them Investigation on some of these plants has been carried out and brief summaries of the results achieved are given Others marked with an asterisk have not been investigated so far and will form a field for research for the investigators

Besides these certain plants are considered to have insecticidal and insect-repellent properties and another allied group is toxic to fishes These, as has been already indicated, are a very important group from point of view of control of insect and other pests which do incalculable harm to man, livestock and agricultural crops

The object of giving these lists is to bring these to the notice of those who are interested in indigenous drugs To the lay public these will be of general interest, as they will find many names of common plants growing at their very door which have been considered effective in the treatment of common ailments and are household remedies For the research worker these lists will open up a field for investigation and will give an idea as to the avenues in which investigations might be directed

Indian Medicinal Plants Alleged to be useful in Tuberculosis in the Indian Indigenous Medicine.

1 ABRUS PRECATORIUS Linn

2 ADHATODA VASICA Nees

3. AEGLE MARMELOS Corr

4 ALANGIUM SALVIFOLIUM (Linn) Wang Syn -*A lamarckii* Thwaites.

*5 ALBIZZIA JULIBRISSIN Durazz , (B -Kalkora, H -Lal siris) , outer Himalaya from the Indus to Sikkim, ascending upto 6 000 to 7,000 ft

*6 ALBIZZIA LEBBECK Benth , (S -Shirisha, H , B & Bo -Siris) , throughout India, ascends to 4,000 ft in the Himalayas usually planted

*7 ALLIUM CEPA Linn , (S -Palandu, H -Piyaz, B -Piyaj, Bo.-Kanda), extensively cultivated all over India.

8 ALLIUM SATIVUM Linn.

*9. ALLIUM SCHOENOPRASUM Linn., Kashmir to Kumaon 8,000 ft to 11,000 ft, grown as a garden crop

10. ALPINIA GALANGA Swartz

11. ASCLEPIAS CURASSAVICA Linn

*12 ASPARAGUS RACEMOSUS Willd, (S & B -Shatamuli, H -Shatawar, Bo -Satavari) Himalayas, from Kashmir eastwards

13. AZADIRACHTA INDICA A Juss, Syn *Melia azadirachta* Linn

*14 BAUHINIA RACEMOSA Lam, (S -Svetakanchan, H -Kanchnal), throughout India.

*15 BAUHINIA VARIEGATA Linn, (S -Kovidara, H -Kachnar, B -Rakta kanchan), sub-Himalayan tract from Indus eastwards, dry forest of Eastern, Central & South India

16 BUTEA MONOSPERMA (Lam.) Kuntze

*17 CANSCORA DECUSSATA Schult, (S -Sankhapuspi, H -Sankhaphuli), throughout India upto 4,000 ft, grows in moist situations

*18 CAPPARIS SPINOSA Linn, (S -Kakadani, H & P -Kabra), plains between the Indus & Jhelum, Salt Range, Chamba, Kumaon, Nepal, Deccan, W Ghats, Baluchistan, Wazirstan

19 CASSIA FISTULA Linn

*20 CASSIA TORA Linn Syn -*C obtusifolia* Linn, (S -Chakramards, H & Bo -Chakunda), throughout India

*21 CHIONACHNE KOENIGII (Spreng) Thw *Polytoca barbata* Stapf (B -Gurgur, H -Kansa, S -Kanda), throughout India in hot and damp parts

22 CIMICIFUGA FOETIDA Linn, (P -Jiunti)

23 CITRULLUS COLOCYNTHIS Schrad

*24 CLERODENDRON SERRATUM (Linn) Moon, (H -Barnagi, S & Bo -Bharangi), more or less throughout India

25 CLITORIA TERNATEA Linn.

26 COCCINIA CORDIFOLIA Cogn. Syn -*C indica* W & A, *Cephalandra indica* Naud

*27. COCCULUS HIRSUTUS Diels, (B -Hayer, Bo -Vasanvel, H.-Jamti-ki-bel), from the foot of the Himalayas to S India

*28 COCOS NUCIFERA Linn, (H -Nariyal, Bo -Narel, M -Tanba), cultivated in the hot damp regions of India especially near the sea

*29 COMMIPHORA AGALLOCHA Engler (B, Bo & H -Guggul), Assam, Sylhet, Bengal, Madhya Pradesh

30. COMMIPHORA MUKUL Engl, Syn.-*Balsamodendron mukul* Hook. ex Stock, Bellary, Mysore, Khandesh, Kathiawar, Rajasthan, Sind, Baluchistan.

*31. CORIANDRUM SATIVUM Linn, (S.-Dhanyaka, H. & B -Dhania); extensively cultivated throughout India.

*32. CRATAEVA NURVALA Ham ; (H -Barun, S -Varuna) almost all over India, wild or cultivated, often found along streams

*33 CRESSA CRETICA Linn , (H & B -Rudranti, Bo -Khardi), throughout India

*34. CUMINUM CYMINUM Linn , (S Jiraka, H -Jira, Zira, Tam.-Shiragam); cultivated throughout India (except Bengal & Assam),

*35 CURCUMA ANGUSTIFOLIA Roxb , (H & B -Tikhur Tam -Kua), outer ranges of Central Himalayas, Bihar, Bengal

*36. CURCUMA ZEDOARIA Bose

*37. DIOSPYROS MELANOXYLON Roxb , (H -Kendu, B Kend, Bo -Temru), deciduous forests of the Madhya Pradesh, Bihar, W. Peninsula, Chota Nagpur.

38 DRYNARIA QUERCIFOLIA J Sm

39. ELETTARIA CARDAMOMUM Maton

*40 EMBELIA TSJERIAM-COTTAM A Dc (H -Bayabirang, Bo -Barbatti)' Ceylon, Malabar Coast, Sylhet and Assam

*41. EULOPHIA NUDA Lindl , (S -Manya, H -Goruma, H -Badbar) Tropical Himalaya, Nepal to Sikkim, Assam, Khasia Hills, Chota, Nagpur, Manipur, W Peninsula

*42 FRITILLARIA CIRRHOSA Don Prodr , Central & E Himalayas, Sikkim, 11,000 to 16,000 ft

*43 FRITILLARIA ROYLEI Hook , Kashmir to Kumaon, 8,000 to 13,000 ft

*44 GERANIUM ROBERTIANUM Linn , Kashmir to Garhwal

45 GLYCYRRHIZA GLABRA Linn

*46. GMELINA ARBOREA Linn , (S -Gumbhari, H -Kambari), throughout India

*47 GOSSYPIUM ARBOREUM Linn , (H -Nurma, P -Papas), Indian gardens.

*48 GRAPTOPHYLLUM PICTUM (L) Griff , (M.-Ysjudemaram), Indian gardens

*49 GREWIA ASIATICA Linn (S -Parusha, H. & B.-Phalsa) , extensively cultivated throughout India

*50. HAUCLEA SESSILIFOLIA Roxb

*51 HYSSOPUS OFFICINALIS Linn ; '(Zufah-Yabis), Kashmir to Kumaon, 9,000 to 11,000 ft

*52 INDIGOFERA TINCTORIA Linn , (S -Nilika, H & Bo.-Nil) , widely cultivated in many parts of India

*53. JASMINUM AURICULATUM Vahl , (S. & Tel , Magadhi) ; Deccan, Carnatic, W. Peninsula.

*54 LACTUCA SERRIOLA Linn. Syn *L scariola* Linn. (H -Kahoo, B.-Salad); W Himalaya, 6,000-12,000 ft.

*55 LEEA AEQUATA Linn , (S , H. & B -Kakajangha), Sikkim, Himalaya, Assam, East Bengal, Sylhet, Andamans.

56. LINUM USITATISSIMUM Linn

*57. LITSEA CHINENSIS Lam (H -Garbijaur, B -Kukarchita), throughout the hotter parts of India

*58. LORANTHUS ASPER Lour

59. LUFFA ACUTANGULA var *amara* Clarke

60 LUFFA ECHINATA Roxb Host Beng

61 LUVUNGA SCANDENS Ham.

*62 MACHILUS MACRANTHA Nees. , (Tam -Kolamavu, Mal -Uravu) , W Peninsula

*63 MARTYNIA ANNUA Linn , (H -Bichu, B -Baghnoki, Tel -Garudamukku) , naturalized in India

*64 MELISSA PARVIFLORA Benth , (H -Bililotan), Garhwal to Sikkim & Mishmi, Khasia hills

*65 MIMUSOPS HEXANDRA Roxb , (S -Rajadani, H -Khirni, M -Palla) , upper Gangetic plain, Madhya Bharat, S India

*66 MOMORDICA DIOICA Roxb , (S -Vahisa, M -Palupaghel kalung) , throughout India, ascending to 5,000 ft in Himalayas

67 MORINGA OLEIFERA Lam

68 MUSA SAPIENTUM O Kuntze.

*69 OCHNA PUMILA Ham ex D Don , (Santh -Champabaha), Kumaon to Sikkim, Bihar, Chota Nagpur

*70. OLDENLANDIA UMBELLATA Linn , (M.-Saya, H -Chirval) , Orissa, Bengal, Deccan, Circars, Carnatic

71. OPERUCLINA TURPETHUM (Linn) Silva Manso.

*72 PHASEOLUS ACONITIFOLIUS Jacq (H -Mat, Tam -Tulkapyre), Himalayas to Ceylon

*73. PHASEOLUS TRILOBUS Art , (H. & B.-Mugani) , Himalayas to Ceylon and Burma

*74 PIPER CHABA Hunter (H B & Bo.-Pan) , cultivated in various parts of India

*75 PIPER LONGUM Linn., (S -Pippali, H.-Pipal), hotter states of India.

76. PINUS ROXBURGHII Sargent, Syn.-*P longifolia* Roxb.

77 PISTACIA INTEGERRIMA Stew.

*78 PISTIA STRATIOTES Linn , (H -Jalqumbhi, Bo -Prashni), throughout India in still sweet waters

*79 PLANTAGO MAJOR Linn , (H -Lahuriya, Bo Bartang), Assam, Khasia hills, W Ghats, Konkan, Nilgiris, Pulneys, Baluchistan

80. PLUMBAGO ZEYLANICA Linn.

*81 RHODODENDRON CAMPANULATUM Wall. ex G Don., (H.-Cherailu, Kash -Gaggar), alpine regions from Kashmir to Bhutan

82. RHUS SUCCEDANEA Linn

*83 SANSEVIERIA ROXBURGHIANA Schult , (B.-Murba, Bo -Morwa), Coromandel coast

*84 SANTALOIDES MINUS Schellenb. Syn -*Rourea santaloides* W & A , (B -Vitaraka, Bo -Vardara), Konkan, S M Country, Kanara to Travancore

*85 SAPINDUS TRIFOLIATUS Linn (S -Phemila H , B & Bo -Ritha).

*86. SESBANIA SESBAN (Linn) Merr Syn -*S aegyptiaca* Pers , (H & B -Jayanti), plains from the Himalayas to Ceylon, ascending upto 4,000 ft in north-west

87. SIDA CORDIFOLIA Linn

88 SIDA RHOMBIFOLIA Linn

*89 SOLANUM TRILOBATUM Linn , (S -Alarka, M -Tudovullay), Gujerat, Deccan, S M Country, N Circars, Carnatic

*90. SPHAERANTHUS INDICUS Linn , (S -Munditika, H & Bo -Gorakmundi), throughout India, ascending the Himalayas upto 5,000 ft. from Kumaon to Sikkim

*91. SPONDIAS PINNATA Kurz , (S -Amrataka, H , B & Bo -Amra), from the Chenab eastwards, Salt Range, Andamans, W Indian Peninsula

92 STEPHANIA GLABRA Roxb

*93. STREBLUS ASPER Lour , (H. Siora, B -Sheora), drier parts of India, Travancore, Andamans

*94. TERAMNUS LABIALIS Spreng , (S -Masha-Parui, H -Mashparui), throughout India

*95. URTICA DIOICA Linn., (H & P -Bichu), N W. Himalaya, from Kashmir and the Salt Range to Simla, 8,000-10,700 ft.

*96 VANDA SPATHULATA Spreng , (Mal -Ponnam-penmaraiva), W Peninsula from Malabar to Travancore

*97. VATERIA INDICA Linn , (S -Ajakarua, H -Safed damar), Western India, from N. Kanara to Travancore, chiefly in evergreen forests, Coorg.

98 VERBASCUM THAPSUS Linn

*99 VERNONIA CINEREA Less., (S & H -Sahadevi, B -Kukseem), throughout India, ascending to 8,000 ft in the Himalaya Khasia and Peninsula mountains

*100. VERNONIA ROXBURGHII Less, Upper Gangetic Plain, Kumaon, Bengal, Central & W. India.

101 WITHANIA SOMNIFERA Dunal.

*102. ZIZYPHUS JUJUBA Lam., (S -Badari, H.-Baer, B -Kul), indigenous and naturalized throughout India, in the Outer Himalaya upto 4,500 ft.

Indian Plants Considered to be useful in Dysentery in Indian Indigenous Medicine.

1 ABRUS PRECATORIUS Linn.

*2 ABUTILON INDICUM Sw , (H -Kanghi, B.-Potari), throughout the hotter parts of India

*3 ABUTILON THEOPHARSTIC Medic, (S -Jaya, Bo -Nahani khapat), N W India, Sind, Kashmir, Bengal

*4 ACACIA ARABICA Willd , (H -Kikar, B -Babla), naturalized in all parts of India, indigenous to Sind and the Deccan.

*5 ACACIA CATECHU Willd , (H -Khadira, H -Khair), Punjab, N. W. Himalaya, Central India, Bihar, Ganjan, Konkan, Deccan

*6 ACACIA FERRUGINEA DC , (Nep -Khour, Bo -Ker), Gujerat, Berar, Circars, Deccan, Konkan, Carnatic, W Ghats

7 ACORUS CALAMUS Linn

*8 ADENSONIA DIGITATA Linn , (H & Bo -Gorakh-amli,), occasionally cultivated in some parts of Uttar Pradesh, Bihar, Bombay & Madras

9 ADHATODA VASICA Nees

*10 ADIANTUM LUNULATUM Burm , (H & B -Kali-Jhant, Bo -Hansraj), throughout N India in moist places, South India

11 ADINA CORDIFOLIA Benth

12 AEGLE MARMELOS Corr.

*13 AGARICUS OSTREATUS (Jacq) Fries., (Bo -Phanasa-alambe)

*14 AGERATUM CONYZOIDES Linn ; (B -Dochunty, Bo -Osari), throughout India upto 5,000 ft

15 AILANTHUS ALTISSIMA (Mill) Swingle, Syn —*A glandulosa* Desf ; (English-Ailanto).

*16 AILANTHUS EXCELSA Roxb , (H -Mahanimb, Tam -Peruppi), Bihar, Chota Nagpur, Madhya Pradesh, Ganjam, Vizagapatnam, Deccan

*17 AILANTHUS MALABARICA DC , (Bo -Guggula-dhup Tel -Maddi-palu), Konkan, Deccan, W Ghats.

18 ALBIZZIA LEBBECK Benth

19 ALSTONIA SCHOLARIS R Br.

*20 ALTHAEA ROSEA Cav , Indian Gardens

*21 AMARANTHUS TRICOLOR Linn Syn —*A gangeticus* Linn.; (H -Lal-sag, S -Marisha), cultivated throughout India

*22 AMOMUM XANTHIOIDES Wall., (H -Ilayechi, Tam.-Elam), Tavoy, Tenasserim.

23 ANDROGRAPHIS PANICULATA Nees, (S.—Bhunimba, H -Kiryat).

*24. ANNONA MURICATA Linn ; (S -Mullanjakka, Tam.-Mullu-chitta); grown to a small extent in Assam.

25 ANNONA RETICULATA Linn

*26. ANTHOCEPHALUS INDICUS A. Rich Syn -*A cadamba* Miq S. & H.-Kadamba), sub-Himalayan tract from Nepal to Burma, Circars and W. Ghats.

27. ANTIARIS TOXICARIA Leschen

*28. ASPARAGUS ADSCENDENS Roxb , (H -Safed musli, Gharwal-Jhirna), Punjab & the Himalayas upto 5,300 ft

29. ASPARAGUS RACEMOSUS Willd

*30. ASTERACANTHA LONGIFOLIA Nees , (H -Tal-makhana, S -Kakil-akshya) throughout India in moist places

*31. AVERRHOA CARAMBOLA Linn , (H -Karmal, M -Tamarta), gardens in hotter parts of India as far north as Lahore

32 BALANITES AEGYPTIACA Del Syn -*B roxburghii* Planch.

*33. BARRINGTONIA ACUTANGULA (Linn) Gaertn (B -Hijal, H -Hijjal), common in the sub-Himalayan tracts east of the Madhya Pradesh, Jumna, Bihar, Orissa, Bengal, Assam, S. India

*34 BASELLA RUBRA Linn , (S -Potaki, H Lalbachlu), throughout India, wild & cultivated

35. BAUHINIA RACEMOSA Lam

*36. BAUHINIA TOMENTOSA Linn , (S -Aswamantaka, H -Kachnar, Bo -Asundro, M -Kanchnine), N W States, Circars, Carnatic, often cultivated in other parts of India

37 BAUHINIA VARIEGATA Linn

*38 BERGENIA LIGULATA (Wall) Engl. (B -Patharchuri, Bo Pashan-bheda), temperate Himalayas, from Kashmir to Bhutan, between 7,000-10,000 ft Khasia hills

*39 BIDENS TRIPARTITA Linn , Central & W Himalaya from Nepal to Kashmir, W Tibet

*40 BOSWELLIA SERRATA Roxb , (S -Shallaki, H & B -Luban), Madhya Pradesh, Deccan, Bihar, Orissa, Rajasthan, Madhya Bharat, Gujerat

*41. BOTRYCHIUM LUNARIA Sew , (English-Moonwort), common from Kashmir to Sikkim ascending upto 13,000 ft , extending to Korakaram Range

42. BRASSICA CERNUA (Thunb) Forbes & Hemsley

43 BRUCEA AMARISSIMA (Lour) Merr Syn -*B sumatrana* Roxb.

44. BUTEA MONOSPERMA O Kuntze.

*45. CAESALPINIA JAYABO Maza; (Arab -Bunduk, Tam.-Kalarisikkodi), Indian Peninsular region

*46 CALAMUS ROTANG Linn , (S.-Vetasa, H., B & Bo.-Bet), Madhya Pradesh, Deccan, Carnatic.

47. CALOTROPIS GIGANTEA R. Br.

48. CARAPA MOLUCCENSIS Lam.

49. CARUM BULBOCASTANUM Koch (H.-Kalajira, Kash.-Gumiyan); Kashmir, Baluchistan

50 CARUM CARVI Linn

*51 CASSIA AURICULATA Linn., (H & B.-Taruar, Tam.-Avaram), wild in the dry regions of Madhya Pradesh, W Indian Peninsula, Rajasthan, cultivated in other parts of India.

52 CASSIA FISTULA Linn

53 CASSIA TORA Linn

54 CASSYTHA FILIFORMIS Linn

*55 CASUARINA EQUISETIFOLIA Linn ; (H.-Jang'isaru, B.-Belatijau), from Chittagong Southwards

56 CEDRELA TOONA Roxb

*57 CELSIA COROMANDELIANA Vahl (S.-Kulahala, Bo.-Kolhal), throughout India

58 CENTELLA ASIATICA (Linn.) Urban Syn —*Hydrocotyle asiatica* Linn.

*59 CEROPEGIA TUBEROSA Roxb.; (P.-Galot, Bo.-Khappar kadu, M.-Manda), Konkan, Deccan, S. M. Country Circars.

*60. CINNAMOMUM INERS Reinw , (Bo.-Tikhi, H.-Jangli darchini), evergreen forests of W Ghats, Carnatic, Sheorov & Kollimalai

61 CLITORIA TERNATEA Linn

*62 CORALLOCARPUS EPIGAEUS Benth ex Hook f , (S.-Patalagaruda), Punjab, Sind, Gujerat, Rajputana, Deccan, Carnatic.

*63 CORCHORUS CAPSULARIS Linn , (S.-Kalasaka, H. & B.-Narchi), throughout the hotter parts of India

*64 CORCHORUS FASCICULARIS Lam , (B.-Bilnalita, Bo.-Hirankhori), throughout the hotter parts of India

65. CUMINUM CYMINUM Linn

*66 CYDONIA OBLONGA Mill Syn -*C vulgaris* Pers., (H.-Bihi, S.-Amritphala), cultivated in Punjab, Kashmir, Nilgiris

*67. CYLISTA SCARIOSA Roxb. (Bo.-Ranghevada), Madhya Pradesh, West & South India.

68 CYNODON DACTYLON Pers (H., B & Bo.-Dhub), throughout India ascending to 8,000 ft. in the Himalaya

*69 CYPERUS ROTUNDUS Linn , (S & Bo.-Musta, B & H.-Mutha), throughout India, common in waste grounds, gardens, road sides, in open spots & upto an elevation of 6,000 ft

*70. CYPERUS SCARIOSUS R. Br ; (S.-Nagar mustaka, H & B.-Nagar Motha) damp places in Bengal, Uttar Pradesh, eastern & southern parts of India.

71 DAUCUS CAROTA Linn

*72 DESCURAINIA SOPHIA Linn , (H.-Khubkallana), Kashmir to Kumaon upto 14,000 ft , E Himalaya, Salt Range Peshawar, Baluchistan

*73 DESMODIUM GANGETICUM DC , Outer Himalaya upto 5,000 ft. & throughout India

74. DIOSPYROS MELANOXYLON Roxb

*75 DIOSPYROS PEREGRINA Gurkein , (S -Tinduka, H & B -Gab, Bo - Tendu) , throughout India

*76. ELAEOCARPUS SERRATUS Linn ; (B -Julpai, M -Olang-karai) W Indian Peninsula

77 ERIGERON CANADENSIS Linn

78 EUPHORBIA ANTIQUORUM Linn

79 EUPHORBIA HIRTA Linn

80. EUPHORBIA HYPERICIFOLIA Linn

81. EUPHORBIA THYMIFOLIA Linn

*82 EVOLVULUS ALSINOIDES Linn , (S -Vishnugandhi, H -Sankha pushpi) Tropical and sub-tropical regions of India

83. FAGONIA CRETICA Linn

*84 FERONIA LIMONIA (Linn) Swingle, Syn —*F elephantum* Corr , (Kapittha, H -Kavitha) ,indigenous in S India, cultivated in many parts of India.

*85 FICUS BENGALENSIS Linn , (S -Vata, H -Bor, B -Bar) , sub-Himalayan tract, W Peninsula, planted elsewhere

*86. FICUS HETEROPHYLLA Linn , (S -Trayamana, B -Bhui-dumur) , throughout hotter parts of India

*87. FICUS HISPIDA Linn , (S -Kakadumbura, H -Konea-dumbar) more or less throughout India

*88. FICUS RACEMOSA Linn Syn —*F glomerata* Roxb , (S -Udumbara, H -Gular) , throughout India

*89. FIMBRISTYLIS JUNCIFORMIS Kunth , (Santh -Bindimuthi), Madhya Pradesh, W Peninsula.

90. FLEMINGIA TUBEROSA Dalz (Bo -Birmova), Konkan

91. FOENICULUM CAPILLACEUM Gilb , Syn —*F vulgare* Gærtn

92. GARCINIA MANGOSTANA Linn , (H., B & Bo -Mangustan), cultivated in Madras State, Nilgiris

93 GASTROCHILUS PANDURATA Ridleys , Konkan, Andamans

94. GOSSYPIUM HERBACEUM Linn. , (H., B & Bo -Kapas, S -Karpas), cultivated in N. W Frontier region of Pakistan, Baluchistan

95. GREWIA HIRUSTA Vahl.

96. GREWIA MICROCOS Linn.

97 HELICTERES ISORA Linn

98 HEMIDESMUS INDICUS R Br.

*99 HIBISCUS CANNABINUS Linn., (S -Nali, H -Patsan, Bo -Ambari); generally cultivated.

100 HOLARRHENA ANTIDYSENTERICA Wall.

101. HOLARRHENA MITIS R Br (Sinhalese-kiriwolla), endemic in Ceylon

102. *HYDROCOTYLE JAVANICA Thunb

*103 HYPERICUM JAPONICUM Thunb , Khasia Hills, Assam, E & W. Peninsula

*104 INDIGOFERA OBLONGIFOLIA Forsk., (S -Jhilla), throughout the Indian plains, Baluchistan

*105 IXORA NIGRICANS Br , (Tam -Mashagani), E. & W. Indian Peninsula

*106 JATEORHIZA PALMATA (Lam) Miers, (Bo -Colombo, Tam -Kolumbu) cultivated in some parts of India

107 JATROPHA CURCAS Linn

108 JUGLANS REGIA Linn

*109 JUSSIAEA SUFFRUTICOSA Linn (S -Bhallava-anga, H -Banlaunga); throughout India

*110 JUSTICIA GENDARUSSA Burm (S -Nila-nirgundi, H -Nili-nargandi), cultivated throughout India

*111 LEEA INDICA Merrill, (H & B -Kakurjiwan, Bo -Karkani), throughout India, Andamans

*112 LENS CULINARIS Medic Syn.—*L esculenta* Moench , (B -Masuri, H. & S -Masur), cold weather crop throughout India

113 LITSEA CHINENSIS Lam

114 LUFFA ACUTANGULA var *amara* Clarke

*115 MALVA ROTUNDIFOLIA Linn , (H & Bo -Khubazi), Sind, Baluchistan, Wazirstan, plains of N India, ascending to 10,000 ft, Kumaon.

*116 MANGIFERA INDICA Linn , (H B & Bo -Amb, S -Amva), indigenous in Sikim, Assam, Khasia hills, Khandesh, along the W. Ghats, cultivated in the tropical regions

*117 MELASTOMA MALABATHRICUM Linn , Burma Myetpye, Tel -Pattudu), throughout India except the Desert

*118. MENTHA LONGIFOLIA Hunds Syn – *M -sylvestris* Linn. (Bo , H & S -Pudina), W. Himalaya, 4,000 to 12,000 ft

*119 MESUA FERREA Linn (S , H. & B -Nagkeshar, Bo -Nagchampa), Mountains of E Himalaya & E Bengal, Assam, ever green rain forests of N. Kanara & S. Konkan, forests of W Ghats, Andamans.

*120. MIMOSA PUDICA Linn.; (S -Lajja, H. & Bo.-Lajalu); naturalised more or less throughout India

*121. MIMUSSOPS ELENGI Linn., (S , H & B -Bakul Bo -Borasali), Andamans, W Peninsula, Khandala Ghats, Circars

*122 MORINDA CITRIFOLIA Linn , (H & B -Ach, Bo-Aal), indigenous in the Darjeeling Terai and Andamans and along the Konkan coast, cultivated largely in India

*123 MORINDA UMBELLATA Linn., (Bo -Al, S -Pitadaru), Khasia, hills, Madras State, E Ghats , W Ghats

124 MUSA SAPIENTUM O Kuntze

*125 MUCUNA PRURITA Hook , (S -Atmagupta, H -Kiwach), Punjab plain, from base of Himalayas to Ceylon & Burma.

*126 MURRAYA KOENIGII Spreng , (S -Sourabhi nimba, H -Katnim), Konkan, W Ghats, Deccan, S M Country, Madras State, along the foot of Himalaya from Kumaon to Sikkim, upto 5,000 ft., Bengal

*127 MURRAYA PANICULATA Jack, (H Marchula, B -Kamini), outer Himalaya from the Jumna eastwards, ascending to 4,500 ft , Assam Satpura Range, Peninsula

128. MYRICA NAGI Thunb , (S -Katphala, H , B & Bo -Kaiphal)

*129 MYRTUS COMMUNIS Linn , (H -Vilayiti mehdi, B -Sutr-sowa), indigenous from the Mediterranean to N W Himalaya, gardens, throughout India

*130 NANNORHOPS RITCHIEANA H Wendl , (H Mazri), Punjab, Sind, Waziristan, Baluchistan

*131 NEPETA ELLIPTICA Rovle ex Benth , (P -Tukhmalanga), W. temperate Himalaya, from Kashmir to Garhwal, 6,000 to 8,000 ft.

*132 NYMPHAEA ALBA Linn , (Kash -Brimposh, Bo -Pandharen-kamal), Kashmir lakes

*133 NYMPHAEA PUBESCENS Willd , (M -Alli), all over India in the warmer parts

*134 OCIMUM AMERICANUM Linn Syn -*O canum* Sims , (H. & B.-Kala tulshi), plains and lower hills of India

*135 OCIMUM BASILICUM Linn , (S -Munjariki), indigenous on the lower hills of the Punjab, cultivated throughout the greater parts of India

136 OLDEN LANDIA AURICULARIA K Schum.

*137. OROXYLUM INDICUM Vent , (S.-Syonaka, H -Arlu, B - Sona), throughout India, except in the Western drier area

*138. OXALIS CORNICULATA Linn., (S -Amlika, H. & B -Amrul); nearly all regions throughout the warmer parts of India, in the Himalayas upto 8,000 ft.

*139. PAVONIA ODORATA Willd., (S -Harivera, B -Bala, Bo.-Kala vala) N W. India Bundelkhand, Sind, Baluchistan, Rajasthan, Bengal, Konkan, S M Country, Deccan

140 PEUCEDANUM GRAVEOLENS Benth

141. PHASEOLUS TRILOBUS Ait

*142. PHYLLANTHUS EMBLICA Linn , (S -Dhatriphala, H -Aoula, Bo Amla), throughout tropical India, wild or planted

*143. PHYLLANTHUS NIRURI Linn , (S -Bhumya-amalaki, H.-Jar-amla), throughout the hotter parts of India from the Punjab to Assam and southwards to Travancore

144 PHYLLANTHUS URINARIA Linn

145 PIPER LONGUM Linn

*146. PIPER NIGRUM Linn , (S -Maricha, H -Golmirch, B -Golmarich), cultivated in hot damp parts of India

147 PISTACIA INTEGERRIMA Stew

148 PISTIA STRATIOTES Linn

*149 PLANTAGO CILIATA Desf , Punjab hills, Sind, Baluchistan

150 PLANTAGO MAJOR Linn.

151. PLANTAGO OVATA Forsk

*152 PLATANUS ORIENTALIS Linn (P & Kash -Buin), N W Himalaya, 5,000 to 8,000 ft., cultivated only

153 PLUMBAGO ZEYLANICA Linn

154 PSIDIUM GUAJAVA Linn

*155. PULICARIA DYSENTERICA Gaertn Kashmir at 5,000-6,000 ft

*156 PUNICA GRANATUM Linn , (S -Darimba, H -Anar) wild in Salt range and in the Himalayas from 3 000 to 6,000 ft , also cultivated in many parts of India.

157 RHEUM EMODI Wall

*158 RUBIA CORDIFOLIA Linn , (S & B -Manjistha, H.-Manith, Bo -Manjit), throughout India in hilly districts

*159 RUMEX SCUTATUS Linn., (English-French sorrel), W Himalaya

*160. RUMEX VASICARIUS Linn , (S -Chukra, H , B & Bo -Chuka), indigenous to W Punjab, the Salt Range & Trans-Indus Hills, also cultivated.

*161. SALIX ALBA Linn , (P -Bis, Kash -Vuir), cultivated in N. W. Himalaya

162 SAPINDUS TRIFOLIATUS Linn.

163. SARACA INDICA Linn.

*164. SCINDAPUS OFFICINALIS Schott., (H & B.-Gajapipal. Bo -Thorapimple), from Sikkim eastwards, Bengal, Chittagong, Andamans.

165. SEMECARPUS ANACARDIUM Linn f

*166. SHOREA ROBUSTA Gaertn f, (S, H, B & Bo Sal), Punjab, along the Sub-Himalayan tract to Assam, Garo Hills, Orissa, Khasia hills, Jaintia hills, Madhya Pradesh

*167. SIDA SPINOSA Linn.; (S -Nagabala, H -Gulsakari, B -Bonmethi), throughout the hotter parts of India from N W. India to Ceylon.

*168. SMILAX PROLIFERA Roxb., (H -Ram dataum), Kumaon, Nepal Sylhet, Bengal, Bihar, Deccan, Peninsula

*169. SOLANUM NIGRUM Linn, (S & B -Kakmachi, Bo -Mako), throughout India, upto 9,000 ft in the W Himalayas

*170. SOYMIDA FEBRIFUGA A Juss, (S Rohuna, H, B & Bo -Rohan), dry forests of W Peninsula extending to Merwar & Chota Nagpur

*171 SPILANTHES ACMELLA Murr, (Bo & P -Akarkara), throughout India, upto 5,000 ft in the Himalayas & other mountains

*172. STACHYTARPHETA JAMAICENSIS Vahl var *indica* Lam Syn —*S indica* Vahl, (M -Simainayuruvi), tropical India from the Punjab & Sylhet to Travancore

173 STRYCHNOS NUX-VOMICA Linn

174 SYMPLOCOS RACEMOSA Roxb

*175 SYZYGIUM CUMINI (Linn) Skeels Syn —*Eugenia jambolana* Lam, (H & B -Jam), throughout India

*176 SYZYGIUM JAMBOS (Linn) Alst Syn —*Eugenia jambos* Linn, (H & Bo -Gulabjaman), Sikkim Terai

*177 SYZYGIUM OPERCULATUM Gamble, Syn —*Eugenia operculata* Roxb, (H -Rai-jaman, S -Bhumbi jambu), sub-Himalayan tract, common in Savannahs in Sal forests, Bihar, Assam, Sylhet, Chittagong

*178 TECTONA GRANDIS Linn f, (S -Saka, H & B -Segun), Konkan, W Ghats of Bombay & Madras States, Deccan, Carnatic, Madhya Bharat, Circars.

179. TERMINALIA ARJUNA W & A

180 TERMINALIA CHEBULA Retz

181. TERMINALIA CITRINA Fleming

*182 TRIUMPHETA BARTRAMIA Linn, (H -Chikti, B -Bun-akora), throughout tropical & sub-tropical India, ascending to 4,000, ft in the Himalaya

183 TYLOPHORA ASTHMATICA Wight & Arn.

*184. TYPHA ELEPHANTINA Roxb., (B -Hegla, Bo -Ramban, Kash -Pitz), marshes from N. W. India to Assam & Southwards, Indus Delta.

*185 URARIA LAGOPOIDES DC, (Bo.-Dowla, H. Pithvan); Tropical zone, Nepal, Chota Nagpur Bengal to Ava.

186 VATERIA INDICA Linn

187 WOODFORDIA FRUTICOSA Kurz, (S -Dhataki, H & B -Dhai), throughout India, Baluchistan

188 WRIGHTIA TINCTORIA R Br, (S -Asita kutanja, H -Mitha indarjou, Bo -Indrajau), Rajasthan, Madhya Pradesh, Deccan, Konkan, S M Country, Circars, W Ghats of Madras State

Indian Plants considered to be Useful in Cholera and Prolonged Fevers such as the Enteric Group

1 ACONITUM FEROX Wall

*2 ARTABOTRYS SUAVEOLENS Blume, Chittagong, Sylhet

*3. BLUMEA LACERA DC, (B —Kukursunga, Bo —Nimrudi), throughout plains of India, upto 2,000 ft

*4 CAPPARIS ZEYLANICA Linn, (H -Ardanda, B -Kalu-kera), Indian gardens

*5 CAPSICUM FRUTESCENS Linn Syn.-*C minimum* Roxb, cultivated, but not extensively

*6 CARAPA MOLUCCENSIS Lam, (B -Pussar), Littoral forests of Bengal and Andamans

7 CASSIA ANGUSTIFOLIA Vahl

*8. DESMODIUM GANGETICUM DC (S & Bo -Shal Parni, H -Sarivan), outer himalayas and throughout India

*9 DRYNARIA QUERCIFOLIA J Sm (Bo -Basingh, S -Ashvakatri), throughout plains of India, on trees or rocks

*10. ELAEOCARPUS TUBERCULATUS Roxb (S Rudraksha, M -Rutthrak-sham), Western Indian Peninsula

*11. ERYCIBE PANICULATA Roxb, (Santh -Kari, Tam -Unankodi), throughout India

*12. ERYTHROXYLUM MONOGYNUM Roxb (M Devadarum), Madras, N Circars dry evergreen forests of Deccan and Carnatic, W Ghats, dry hill forests of Travancore

13 EUPHORBIA HELIOSCOPIA Linn

*14. FAGONIA CRETICA Linn. (H -Damahan, P -Dama); Deccan, W Khandesh, Cutch, Sind, Baluchistan, Waziristan, W Rajasthan, Upper gangetic plain, Punjab

*15. FLACOURTIA INDICA Merr Syn -*F ramontchi* L' Herit, (B Bincha); Sub-Himalayan tract, Upper Gangetic Plain, common in Peninsula, W. Ghats, Forests of the N Circars and Deccan upto 3,000 ft

*16. GREWIA HIRUSTA Vahl. Symb (Bo -Gowali H Kukarbicha), Sub-Himalayan tract upto 4,500 ft. from the Indus eastwards, Salt Range, Bihar, Orissa.

*17 GREWIA MICROCOS Linn (M -Kottei), E. Bengal, Assam, W. Peninsula, Mysore.

18. KALANCHOE SPATHULATA DC.

*19 LODOICEA MALDIVICA Pers. Syn -*L seychellarum* Labill. (H & Bo -Darya-kanariyal), cultivated in India

*20 MARRUBIUM VULGARE Linn , (H -Pahari gandana), Kashmir from 5,000 to 8,000 ft Baluchistan etc.

*21. MOMORDICA CHARANTIA Linn. (H -Karela, B -Karala, Bo -Karla), cultivated throughout India

*22. MUSA PARADISIACA Linn var *sapientum* Kuntze Syn -*M sapientum* Linn. (S -Ramoha, H & Bo -Kela), commonly cultivated.

*23. OLDENLANDIA AURICULARIA K Schum , (B -Muttia-lata, Bo.-Dapoli), almost all over India

*24 POINCIANA PULCHERIMA Linn Syn -*Caesalpinia pulcherima* Swartz (B -Krishnachura, S & Tam -Ratna-gandhi), Gardens throughout India.

*25. PSIDIUM GUAJAVA Linn. (H -Amrud, Bo -Perala), cultivated & naturalized throughout India

26. SAPINDUS TRIFOLIATUS Linn

27 SCHLEICHERA OLEOSA (Lour) Merr Syn -*S trijuga* Willd

*28 SCHWEINFURTHIA SPHEROCARPA Br , (S H & Bo -Sanipat), Rajasthan Desert, Sind, Baluchistan

*29 SOLANUM MELONGENA Linn , (S.-Bartaku, H -Baigun), widely cultivated in India

* 30. SOPHORA TOMENTOSA Linn , (Burm -Thimbawmagyi), Andaman & Nicobar Islands, very occasionally in Indian gardens.

31. STYRCHNOS NUX-VOMICA Linn

32 TERMINALIA CHEBULA Retz.

*33 TERMINALIA CITRINA Fleming , (H -Harira, B -Haritaki), Assam, Dacca, Mymensingh

*34 TERMINALIA PANICULATA Roth , (Bo -Kindal, M -Pekarakai; Western regions of the Peninsula from Bombay through Kanara & Malabar to Travancore, upto 2,000 ft., Coorg, Nilgiris, Anamalais Cuddapah, Ballary

35 TRACHYSPERMUM AMMI (Linn) Sprague, Syn.-*Carum copticum* Benth.

*36. ZANTHOXYLUM BUDRUNGA Wall., (Bazinali, Bo -Tessul, H -Badrang), Konkan, Deccan, S. M. Country, N Kanara, W Ghats in S Kanara, Mysore, Malabar, Annamalais and Travancore at low elevations, Orissa, Sylhet, Khasia Hills, Chittagong.

Indian Plants considered to have Emmenagogue and Abortifacient Properties

1. ABRUS PRECATORIUS Linn.
2. ALOE BARBADENSIS Mill.
3 ANANAS COMOSUS Linn
4. ANNONA SQUAMOSA Linn.
5. APIUM GRAVEOLENS Linn.
6. ARECA CATECHU Linn
7 ARISTOLOCHIA BRACTEATA Retz.
8 ARISTOLOCHIA INDICA Linn
9. ARTEMISIA VULGARIS Linn.
10 CALOTROPIS GIGANTEA (Linn.) Dryand (*C gigantea* R Br)
11. CALOTROPIS PROCERA (Linn) Dryand (*C. procera* R. Br.)
12. CARICA PAPAYA Linn.
13 CELASTRUS PANICULATUS Willd
14 CINCHONA CALISAYA Wedd
15 CINNAMOMUM CAMPHORA Nees & Eberm
16 CITRULLUS COLOCYNTHIS Schrid
17 CROCUS SATIVUS Linn
18 CUCUMIS TRIGONUS Roxb.
19 CUSCUTA REFLEXA Roxb.
20 DAUCUS CAROTA Linn.
21. DOLICHANDRONE FALCATA Seem.
22. EUPHORBIA TIRUCALLI Linn.
23 EXCOECARIA AGALLOCHA Linn.
24 GARCINIA MORELLA Desr.
25 GLORIOSA SUPERBA Linn.
26. GOSSYPIUM HERBACEUM Linn.
*27 LEPIDIUM SATIVUM Linn , (S -Chandrasuru, H -Chansaur, B -Halim), cultivated throughout India.
28. MOMORDICA CHARANTIA Linn.
*29. MOMORDICA TUBEROSA Cogn. (*M cymbalaria* Fenzl ex Naud), Bo.- Kadavanchi), Western parts of India
30. MORINGA OLEIFERA Lam. (*M. pterygosperma* Geratn.)
31. NERIUM INDICUM Mill. (*N. odorum* Soland.)

32 NIGELLA SATIVA Linn.

33 PEGANUM HARMALA Linn.

34. PLUMBAGO INDICA Linn (*P rosea* Linn)

35. PLUMBAGO ZEYLANICA Linn

*36. PLUMERIA RUBRA Linn var *acutifolia* Bailey (*P acutifolia* Poir, *P. acuminata* Ait), (S -Kahira champa, H & Bo -Khair champa), cultivated throughout India, naturalized in many parts

37 RANDIA DUMETORUM Lam

38. RUBUS MOLUCCANUS Linn

39 RUTA GRAVEOLENS Linn

40 SALICORNIA BRACHIATA Roxb

41 SAPINDUS TRIFOLIATUS Linn

42 SEMECARPUS ANACARDIUM Linn f.

43 SESAMUM ORIENTALE Linn (*S. indicum* Linn)

44 STACHYTARPHETA JAMAICENSIS (Linn), Vahl var *indica* H J. Lam. (*S. indica* Vahl)

45. TAXUS BACCATA Linn

46. THEVETIA PERUVIANA (Pers) Merr (*T. neriifolia* Juss ex Steud).

47. TRIANTHEMA PENTANDRA Linn

48 TRIANTHEMA PORTULACASTRUM Linn (*T monogyna* Linn)

*49 URENA LOBATA Linn (H -Bachata, B -Bonokra), hotter parts of India, a weed of waste places, forest clearing and roadsides, common in bambo and mango clumps of Bengal

50 WITHANIA somnifera Dun

Indian Plants liable to produce Dermatitis

1 ABROMA AUGUSTA Linn

2 AILANTHUS ALTISSIMA (Mill) Swingle (Syn *A glandulosa* Desf)

3 ANACARDIUM OCCIDENTALE Linn.

4 ANAGALLIS ARVENSIS Linn.

5 ANTHEMIS COTULA Linn.

6 APIUM GRAVEOLENS Linn

7 ARISAEMA SPECIOSUM (Wall.) Mart.

*8 ARISAEMA TORTUOSUM (Wall.) Schott , (P.-Don, Nep -Birbanka), from Simla to Sikkim & Bhutan at 8,000 ft , Khasia hills, Manipur, Ranchi, W Ghats

*9. ASPARAGUS OFFICINALIS Linn., (P , H. & Kash.-Allipalli), from Kashmir to Bhutan, Khasia Hills, Assam.

10. CALOTROPIS GIGANTEA (Linn) Dryand.

11. CALOTROPIS PROCERA (Linn) Dryand.

12 CANNABIS SATIVA Linn

*13. CISSUS SETOSA Roxb , (H -Karmal, Bo -Khaj-goli-cha-vel), Deccan, W. Ghats of Madras State

14 DATURA STRAMONIUM Linn

15. DAUCUS CAROTA Linn

*16. DELPHINIUM AJACIS Linn., commonly cultivated in gardens.

*17. DICTAMNUS ALBUS Linn , from Kashmir to Kunawar.

18 ERIGERON CANADENSIS Linn

19 EUPHORBIA ACAULIS Roxb

20. EUPHORBIA ANTIQUORUM Linn.

21 EUPHORBIA CATTIMANDO W Elliot (*E trigona* Fl. Brit. Ind. in part)

22. EUPHORBIA HELIOSCOPIA Linn

23. EUPHORBIA NERIIFOLIA Linn.

24 EUPHORBIA NIVULEA Buch Ham.

25. EUPHORBIA PEPLUS Linn

26 EUPHORBIA ROTHIANA Spreng.

27 EUPHORBIA ROYLEANA Boiss

*28 EUPHORBIA THOMSONIANA Boiss , (Kash -Hirtiz), Kashmir 10,000 to 12,000 ft.

29. EUPHORBIA TIRUCALLI Linn

30 EUPHORBIA TRIGONA Haw

31 EXCOECARIA AGALLOCHA Linn.

32 FAGOPYRUM ESCULENTUM Moench.

33 FLEURYA INTERRUPTA Gaudich.

*34. GINKGO BILOBA Linn., (English-Maiden-hair-tree), rarely cultivated in gardens

*35 GIRARDINIA HETEROPHYLLA Decne., Mount Abu, Chota Nagpur, W Peninsula.

36. HEDERA HELIX Linn.

*37. HIPPOMANE MANCINELLA Linn , (English-Manchineal tree), introduced into Indian gardens.

38. HOLIGARNA ARNOTTIANA Hook. f.

39. HOLIGARNA GRAHAMII (Wight) Hook. f.

40. HOLIGARNA LONGIFOLIA Buch. Ham ex. Roxb.

41. HUMULUS LUPULUS Linn

42. HYPERICUM PERFORATUM Linn

43. LAPORTEA CRENULATA Gaudich

*44. LAPORTEA TERMINALIS Wight, sub-tropical Himalayas from Kumaon to Mishmi at 4,000 to 8,000 ft., Madhya Pradesh, W Ghats of Madras State.

*45. LASIOSIPHON ERIOCEPHALUS Decne, (Bo -Rametha) W Ghats of Bombay and Madras States also Nilgiris

*46 LEONURUS CARDIACA Linn, Kumaon, Kashmir, Hazara, Kurrum valley.

*47. LOBELIA EXCELSA Lesch, W Ghats of S India, Nilgiris, Pulney, Travancore above 6,000 ft

*48 LOBELIA NICOTIANIFOLIA Heyne (B & H -Nala, Bo -Dhavala), W. Ghats from Bombay to Travancore at 3,000 to 7,000 ft., Konkan, Deccan, Nilgiris, Malabar

*49. MUCUNA ATROPURPUREA DC, plains of Western India

*50. MUCUNA GIGANTEA DC, (Malay -Kaku-vali), a littoral species found on the Indian Coast

*51. MUCUNA HIRSUTA Wight & Arn, plains of Western India.

*52. MUCUNA MONOSPERMA DC, (Bo, Sonogaravi, M -Thelu-Kodi), E Himalaya, Khasia Hills, Assam, Chittagong, Konkan, S M Country.

53. MUCUNA PRURITA Hook. (*M pruriens* Fl. Brit Ind., non DC.)

54. NERIUM OLEANDER Linn.

55. PODOPHYLLUM HEXANDRUM Royle (Syn — *P. emodi* Wall. ex. Hook. f & Thoms)

56. POLYGONUM HYDROPIPER Linn.

57. RANUNCULUS SCELERATUS Linn.

58 RHUS INSIGNIS Hook. f.

59 RHUS PUNJABENSIS J. L. Stew ex Brand.

60. RHUS SUCCEDANEA Linn.

61. RHUS WALLICHII Hook. f.

62. RUMEX ACETOSA Linn.

63. RUMEX ACETOSELLA Linn

*64. RUTA GRAVEOLENS Linn,, var *angustifolia* Hook. f, (S -Somalata, H.-Gadab, B.-Ermul, Bo.-Satap), cultivated in Indian gardens.

65. SAPIUM INSIGNE Trimen.

66. SCHIMA WALLICHII Choisy
67. SEMECARPUS ANACARDIUM Linn f
68. SEMECARPUS TRAVANCORICUS Bedd
69. TRAGIA BICOLOR Miq
70 TRAGIA INVOLUCRATA Linn
71 URTICA DIOICA Linn
72. URTICA HYPERBOREA Jacq.
73 URTICA PARVIFLORA Roxb
74 URTICA PILULIFERA Linn.
75 WALLICHIA DISTICHA T. Anders
76. XANTHIUM STRUMARIUM Linn

Indian Plants considered to have Insecticidal & Piscicidal Properties.

1 ACACIA PENNATA (Linn) Willd
2. ACORUS CALAMUS Linn
*3 ACORUS GRAMINEUS Soland, Sikkim Himalaya upto 6,000 ft , Khasia hills 4,000-5,000 ft
4 ACRONYCHIA PEDUNCULATA (Linn) Miq (Syn —*A laurifolia* Blume).
5 ADINA CORDIFOLIA (Roxb) Benth & Hook f.
6 AGAVA AMERICANA Linn.
*7 ALBIZZIA CHINENSIS (Osbeck) Merr (Syn -*A stipulata* Boiv), (B.-Amluki, Bo.-Udala), throughout India, ascending to 4,000 ft. in the Himalaya
8 ALBIZZIA PROCERA (Roxb.) Benth.
9 ANACARDIUM OCCIDENTALE Linn
10 ANAGALLIS ARVENSIS Linn
11 ANAMIRTA COCCULUS (Linn.) Wight & Arn.
12 ANDRACHNE CORDIFOLIA Muell Arg.
13 ANNONA RETICULATA Linn.
14. ANNONA SQUAMOSA Linn
*15. APAMA TOMENTOSA Engl (Syn -*Bragantia tomentosa* Blume); Assam & Manipur.
16 ARENGA OBTUSIFOLIA Mart
17. ARISAEMA SPECIOSUM (Wall.) Mart.
18. ARISAEMA TORTUOSUM (Wall.) Schott.

19. ARISTOLOCHIA BRACTEATA Retz.

20. ARTEMISA ABSINTHIUM Linn.

21 ARTEMISIA VULGARIS Linn

22 ASCLEPIAS CURASSAVICA Linn

23 AZADIRACHTA INDICA A. Juss (Syn -*Melia azadirachta* Linn)

24 BALANITES AEGYPTIACA Delile (Syn -*B roxburghii* Planch)

25 BAMBUSA BAMBOS Druce (Syn -*B. arundinacea* Willd)

26 BARRINGTONIA ACUTANGULA (Linn) Gaertn

27 BARRINGTONIA ASIATICA (Linn) Kurz Syn *B speciosa* Forst)

*28. BARRINGTONIA RACEMOSA (Linn) Roxb , (S -Nipi, H -Ijjul), Western sea coast of India, from Konkan to Travancore, Assam, Sunderbans.

29. BERBERIS ARISTATA DC (possibly some other species of Berberis also)

30 BUTEA MONOSPERMA (Lam) Kuntze (Syn -*B frondosa* Koen ex Roxb.)

31 CAESALPINIA NUGA (Linn) Ait.

32 CALLICARPA LONGIFOLIA Lam. var. *lanceolaria* C B Clarke

33. CALONYCTION MURICATUM (Linn.) G Don (Syn -*Ipomoea muricata* Jacq)

34. CALOPHYLLUM INOPHYLLUM Linn

35 CANNABIS SATIVA Linn

*36. CARDARIA DRABA Desv (Syn -*Lepidium draba* Linn.), (Afgh -Bijindak) a weed of cultivation in the Punjab & N W. Frontier Pakistan

37 CAREYA ARBOREA Roxb

38. CASEARIA GRAVEOLENS Dalz

*39 CASEARIA TOMENTOSA Roxb., (H & Bo -Chillara), throughout India, ascending to 3,000 ft in Himalayas

40 CASSYTHA FILIFORMIS Linn

41 CENTRATHERUM ANTHELMINTICUM (Willd) Kuntze (Syn.-*Vernonia anthelmintica* Willd)

42 CERBERA MANGHAS Linn. (Syn -*C. odollam* Gaertn.)

43. CHRYSANTHEMUM CINERARIAEFOLIUM Vis.

44. CHRYSANTHEMUM COCCINEUM Willd.

45 CIMICIFUGA FOETIDA Linn.

46. CINCHONA CALISAYA Wedd.

47. CINCHONA OFFICINALIS Linn.

48. CINCHONA SUCCIRUBRA Pav. ex Klotzsch.

49 CINNAMOMUM CAMPHORA Nees & Eberm.

50 CLEISTANTHUS COLLINUS Benth. & Hook f

51. CORYPHA UMBRACULIFERA Linn.

52 CROTON OBLONGIFOLIUS Roxb.

53 CROTON TIGLIUM Linn

54 CUCUMIS SATIVUS Linn

55 CURCUMA LONGA Roxb

*56 CYMBOPOGON NARDUS (Linn) Rendle (Syn -*Andropogon nardus* Linn) (Bo -Kamkher, H -Ganjni), throughout the hotter parts, wild or cultivated

57 CYNANCHUM ARNOTTIANUM Wight

58 DALBERGIA STIPULACEA Roxb.

59 DELPHINIUM BRUNONIANUM Royle

60. DELPHINIUM CAERULEUM Jacquem ex Cambess

61 DELPHINIUM ELATUM Linn.

62 DERRIS ELLIPTICA (Roxb) Benth

63 DERRIS FERRUGINEA (Roxb.) Benth

64 DERRIS SCANDENS (Roxb.) Benth

*65 DERRIS TRIFOLIATA Lour , var *uliginosa* (Roxb ex Willd) (Syn -*D uliginosa* Benth , *Robinia uliginosa* Roxb ex Willd)

66 DIOSCOREA HISPIDA Dennst (Syn -*D. daemona* Roxb)

67 DIOSCOREA PRAZERI Prain, Burkill (Syn -*D deltoidea* Wall , var sikkimensis Prain)

68 DIOSPYROS EBENUM Koen

*69 DIOSPYROS MONTANA Roxb , (S -Tumala, H -Lohori, B -Bangab), throughout India, from Kanara eastwards in the sub-Himalayan tract

*70 DIOSPYROS PANICULATA Dalz , (S -Thinduka, M -Karinthuvari), S M Country, Travancore, Kanara, Malabar

71 DODONAEA VISCOSA (Linn) Jacq

72 DOLICHANDRONE FALCATA Seem

73 DURANTA REPENS Linn. (Syn.-*D plumieri* Jacq)

74. EDGEWORTHIA GARDNERI Meissn

75. ENTADA PURSAETHA DC (Syn -*E. scandens* Benth)

*76 EREMOSTACHYS SUPERBA Royle ex Benth., W. Himalaya, Punjab, Garhwal.

77. EREMOSTACHYS VICARYI Benth

78. EUCALYPTUS GLOBULUS Labill

79 EUPATORIUM ODORATUM Linn

80 EUPHORBIA ANTIQUORUM Linn

81 EUPHORBIA NERIIFOLIA Linn

82. EUPHORBIA ROYLEANA Boiss

83. EUPHORBIA THYMIFOLIA Linn.

84. EUPHORBIA TIRUCALLI Linn

85. EXCOECARIA AGALLOCHA Linn

86. FLUGGEA LEUCOPHYRUS (Koen) Willd

87 FLUGGEA VIROSA (Roxb. ex Willd) Baill. (Syn -*F. microcarpa* Blume)

*88. GARDENIA CAMPANULATA Roxb , (Burm -Hsathanpaya), foot of Sikkim Himalaya, Assam, Sylhet, Chittagong, Bihar.

89 GAULTHERIA FRAGRANTISSIMA Wall

90 GLORIOSA SUPERBA Linn

91. GNETUM SCANDENS Roxb.

92 GYNANDROPSIS GYNANDRA (Linn) Merr. (Syn.-*G pentaphylla* DC.)

93 GYNOCARDIA ODORATA R Br

94 HARPULLIA CUPANIOIDES Roxb.

95 HEDERA HELIX Linn

96. HYDNOCARPUS KURZII (King) Warb (Syn -*Taraktongenos kurzii* King)

97. HYDNOCARPUS LAURIFOLIA (Dennst) Sleumer (Syn -*H wightiana* Blume)

98 HYDROCOTYLE JAVANICA Thunb

99 JATROPHA CURCAS Linn

100 JUGLANS REGIA Linn

101 KALANCHOE SPATHULATA (Poir) DC.

*102 LAGENANDRA TOXICARIA Dalz (Many authors view this species as synonymous with *L ovata* (Linn) Thw of Ceylon), (Bo Rakhalu, M -Maravara, Tsjmul), Konkan to N. Kanara, Travancore, Mysore, Coorg, Cochin

103 LASIOSIPHON ERIOCEPHALUS Decne

*104. LINOSTOMA DECANDRUM Wall , Sylhet, Chittagong

105. MADHUCA LATIFOLIA (Roxb) Macbride (Syn -*Bassia latifolia* Roxb)

106 MADHUCA LONGIFOLIA (Linn.) Macbride (Syn.—*Bassia longifolia* Linn

107. MAESA INDICA Wall

108 MELALEUCA LEUCADENDRON Linn.

109 MELODINUM MONOGYNUS Roxb

110. MILLETTIA AURICULATA Baker ex Brand

*111 MILLETTIA PACHYCARPA Benth , forests of Garo & Khasia Hills, Sikkim & Assam upto 4,000 ft

*112 MILLETTIA PISCIDIA Wight & Arn , Sikkim and Assam

113 MUNDULEA SERICEA (Willd) Greenway (Syn —*M suberosa* Benth.)

114 MYRICA NAGI Thunb.

115 NICANDRA PHYSALOIDES Gaertn

116 NICOTIANA RUSTICA Linn.

117 NICOTIANA TABACUM Linn

118. NIGELLA SATIVA Linn

*119 OCIMUM GRATISSIMUM Linn , (S -Vantulshi, H & B -Ramtulshi); common wild plant in W India cultivated in gardens in Bengal, Nepal, Deccan Peninsula

120 OUGEINIA DALBERGIOIDES Benth

121 PACHYGONE OVATA (Poir) Miers ex Hook f & Thoms.

122. PEGANUM HARMALA Linn

123 PHYLLANTHUS URINARIA Linn

124 PICRASMA JAVANICA Blume, var *nepalensis* Badhwar nov comb. (Syn -*P nepalensis* Benn).

125. PIERIS OVALIFOLIA D Don

126. PITHECELLOBIUM BIGEMINUM Mart (Syn —*Pithecolobium bigeminum* Benth).

127. POGOSTEMON HEYNEANUS Benth (Syn -*P patchouli* Hook. f. in Fl. Brit -Ind , non. Pellet)

128 POLYGONUM FLACCIDUM Meissn

129 POLYGONUM HYDROPIPER Linn

130 PONGAMIA PINNATA Linn Merr (Syn.-*P glabra* Vent)

131 PYGEUM GARDNERI Hook. f., (Bo -Daka), W Ghats of Bombay & Madras States, hills of Travancore, Malabar, Nilgiris, Pulneys the Deccan.

132 RANDIA DUMETORUM Lam. (Split up into three species by Gamble)

133. RANDIA ULIGINOSA DC.

134. RAUWOLFIA SERPENTINA Benth ex Kurz

135 RHODODENDRON BARBATUM Wall ex G Don, (Nepl.-Guras), Kumaon to Bhutan at 8,000 to 12,000 ft., Sikkim

136 RHODODENDRON FALCONERI Hook f, (Nepl -Kurling), East Nepal to Bhutan at 9,000 13,0000 ft

137 RICINUS COMMUNIS Linn.

138 RUTA GRAVEOLENS Linn. var *angustifolia* Hook f

139 SANTALUM ALBUM Linn

140 SAPINDUS MUKOROSSI Gaertn, (S -Phenila, H, B & Bo -Ritha) cultivated throughout N W India, Bengal & Assam, wild on the Himalayas upto 4,000 ft

141 SAPINDUS TRIFOLIATUS Linn

142 SAPIUM INDICUM Willd

143. SARCOSTEMMA ACIDUM (Roxb) Voigt (Syn -*S brevistigma* Wight & Arn)

144 SAUSSUREA LAPPA C. B Clarke

145 SCHLEICHERA OLEOSA (Lour) Merr (Syn -*S trijuga* Willd)

146 SCLERIA PERGRACILIS (Nees) Kuntn widely scattered from Garhwal at 5,000 ft to Assam, Bihar, Chota Nagpur & Deccan

147 SOPHORA MOLLIS R Grah, (P -Buna, Arab -Arghavan), plains and low hills of the North-West, Hazara & the Salt Range to Kumaon & Nepal, upto 7,000 ft., & Bushahr & near Dehra Dun

148 SPHAERANTHUS INDICUS Linn

149 STEPHANIA HERNANDIIFOLIA (Willd) Walp, (S -Vanatik-tika, B -Agnad), W & E. Coast, Dehra Dun, Biharam (Sikkim, Bengal, Assam, Cachar)

150. STRYCHNOS COLUBRINA Linn

151 STRYCHNOS NUX-VOMICA Linn

152. TAXUS BACCATA Linn.

153 TEPHROSIA CANDIDA (Roxb) DC

154 TEPHROSIA PURPUREA (Linn) Pers

*155 TEPHROSIA VOGELII Hook f, cultivated by tea planters in Assam

156 TERMINALIA BELLERICA (Gaertn) Roxb.

157. THEVETIA PERUVIANA (Pers) Merr (Syn -*T nereifolia* Juss ex. Steud)

*158 TRIGONELLA FOENUM-GRAECUM Linn, (H B & Bo -Methi), Punjab & Kashmir, cultivated in many parts of India

159. VERBASCUM THAPSUS Linn

*160 VITEX NEGUNDO Linn , (S. & H -Nirgundhi, B -Nishinda), throughout India in the warmer zone

161 WALSURA PISCIDIA Roxb

162 WIKSTROEMIA INDICA (Linn) C. A Mey var. *viridiflora* (Meissn) Hook f.

163. ZANTHOXYLUM ALATUM Roxb

*164 ZANTHOXYLUM HAMILEONIANUM Wall , (Nep -Purpuray timur), throughout Assam, in low level forests, except in the Khasia and Jaintia hills

CHAPTER X

Bibliography

In the first edition of The Review, a list of the papers published on the subject under the Indian Council of Medical Research (formerly Indian Research Fund Association) was given As the scope of the present review has been widened to include all the work done on Indian Indigenous Drugs anywhere, the question of including a comprehensive and if possible a complete bibliography for the new edition was discussed and agreed to by the Secretary Indian Council of Medical Research. In view of the fact that no such bibliography has so far been published and also because this subject has been assuming increased importance both in India and abroad, it was decided that a comprehensive bibliography should form a particular feature of the present edition, inspite of the fact that size of volume would be considerably increased

During the last 3 or 4 decades the subject of Indian Indigenous Drugs has received considerable attention of many of the scientific workers in this country and their papers have been published in various journals in India and elsewhere. The subject has interested many foreign workers also who have made very important contributions in foreign scientific journals An attempt has been made to get all the references together, particularly of the work done on the plants of Indian origin. This is a special feature of the bibliography which is given in this Chapter

To facilitate consultation by workers references have been arranged plant-wise (which are given in alphabetical order,) and in sequence of years in which the papers have been published In order to attract the immediate attention of the workers the un-orthodox method of giving the title of the paper first and names of authors afterwards has been adopted

Abbreviations of the journals used are also given in a separate list

The inclusion of all the available bibliography in the present volume has involved considerable labour and careful sifting It is hoped that it will be useful to those who wish to consult the previous work done when taking up a plant for investigation It is by no means such an exhaustive bibliography as we would have wished but it is hoped that it may serve as a first compilation of its kind which can be gradually added to.

List of Journals and Books with the Abbreviations.

Agri Jr Ind	.	Agricultural Journal of India
Agri Ledger	. .	Agricultural Ledger (Govt of India)
Amer Chem Jr	..	American Chemical Journal
Amer Jr. Pharm	...	American Journal of Pharmacy.
Analyst	...	Analyst.
Ann Bot.	...	Annals of Botany.

Ann. Chem. ...	Annalen der Chemie.
Ann Chim ...	Annales de Chemie
Ann Chim Phys ...	Annales de Chimie et de Physique
Ann Chim Appl ...	Annales de Chimie analytique appliquee
Ann. Inst Past ..	Annales de l' Institut Pasteur
Arch de Pharm. ..	Archiv de Pharmacie
Arch Exper Path Pharm	Archiv fur experimentelle Pathologie and Pharmakologie
Arch Hyg. ...	Archiv fur Hygiene
Arch Pharm .	Archiv der Pharmazie
Biochem d Pflzen ...	Biochemie der Pflanzen
Biochem Jr .	Biochemical Journal
Biochem Ztsche	Biochemische Zeitschrift.
B M J ...	British Medical Journal
Buchn Repert Pharm ...	Repertorium der Pharmacie (Buchner)
Bull Acad Sci Allahabad	Bulletin of the Academy of Science of Uttar Pradesh of Agra & Oudh, Allahabad
Bull Bot Gard Kew. ...	Bulletin Botanical Garden Kew
Bull Sc Pharm. ...	Bulletin des Sciences Pharmacologiques
Bull. Soc Chim ...	Bulletin de la Societe Chimique de France
Bull Soc Chim Biol ...	Bulletin de la Societe de Chimie Biologique
Bull Tokyo Inst Technical	Bulletin, Tokyo Institute of Technology
C C ...	Chemisches Centralblatt
Chem Abst ..	Chemical Abstract (American)
Chem Weekbl. ...	Chemisches Weekblad
Chem Ztg ...	Chemiker-Zeitung
Chin Jr Physiol ..	Chinese Journal of Physiology
Compt Rend ...	Comptes rendus hebd des Seances de l' Academie des Sciences
Comp Soc Biol ...	Comptes rendus hebd des Seances de la Societe de Biologie.
Curr Sci ...	Current Science
Helv. Chim Act. ...	Helvetica Chimica Acta
Ind Farming ...	Indian Farming
Ind. For Rec. ...	Indian Forest Records (and Bulletin) Dehra Dun
Ind. For. ...	Indian Forester
Ind. Jr. Agric Sci ...	Indian Journal of Agricultural Science

Ind. Jr. Ent. ...	Indian Journal of Entomology.
Ind. Jr. Med. & Phys Sc.	Indian Journal of Medical and Physical Science
I. J. M. R. ...	Indian Journal of Medical Research.
Ind. Jr. Pharm. ...	Indian Journal of Pharmacy
Ind Jr. Vetr Sci Animal Husbandry ...	Indian Journal of Veterinary Science & Animal Husbandry
I. M. G ...	Indian Medical Gazette.
I M J. ...	Indian Medical Journal
J C. S. ...	Journal of the Chemical Society, London.
Jr. Agri Res ...	Journal of Agricultural Research
Jahrb. Prakt Pharm ...	Jahrbuch fur Praktische Pharmazie
Jr. Amer Pharm. Assoc ...	Journal of the American Pharmaceutical Association
Jr Amer Chem Soc	Journal of the American Chemical Society
Jr Anamalai Univ	Journal of Anamalai University
Jr Assoc Agri Chem	Journal of the Association of Official Agricultural Chemists
Jr. Biol Chem ...	Journal of Biological Chemistry
Jr Bombay Nat Hist Soc	Journal of Bombay Natural History Society
Jr Pharm .	Journal der Pharmacie fur Aerzte
Jr Ind Chem Soc ..	Journal of the Indian Chemical Society
Jr Ind Inst Sci ..	Journal of the Indian Institute of Science, Bangalore
Jr Malaria Inst India .	Journal of the Malaria Institute of India
Jr Mysore Univ	Journal of the Mysore University
Jr Pharm Chim	Journal de Pharmacie et de Chimie
Jr Pharm Soc Japan	Journal of the Pharmaceutical Society of Japan.
Jr Pharm Exp Therap .	Journal of Pharmacology and Experimental Therapeutics.
Jr Physiol. ...	Journal of Physiology.
Jr Pract Chem ...	Journal fur Praktisch Chemie.
Jr Russ Phys Chem Ges.	Journal der Russischen Physikalisch-chemischen Gesellschaft.
Jr Sci Indul Res ...	Journal of Scientific & Industrial Research
Jr Soc Chem Ind ..	Journal of the Society of Chemical Industry.

Jr. Univ. Bombay.	..	Journal of the University of Bombay.
Kew Bull.	...	Kew Bulletin.
Lancet	...	Lancet
Nat. Acad. Sci Ind	...	National Academy of Science, India.
Nature.	...	Nature, London.
Pharm Jr.	...	Pharmaceutical Journal and Pharmacist.
Pharm Jr Trans	...	Pharmaceutical Journal and Transactions.
Pharm. Post	.	Pharmaceutische Post
Pharm Rev	..	Pharmaceutical Review
Pharm Weekbl	..	Pharmazeutische Weekblad
Pharm Ztg	..	Pharmazeutische Zeitung
Pharm Ztschrft f. Russl	.	Pharmazetische zitschrift fur Russland.
Philip Aggric	...	Philippine Agriculturist, Las Banos
Phil Jr Sci	...	Philippine Journal of Science
Physiol Abst.	...	Physiological Abstracts
Pharmacogn	...	Pharmacognosie
Proc Acad Sci U P	...	Proceedings of the Academy of Sciences of Uttar Pradesh of Agra & Oudh, Allahabad
Proc Chem Soc	...	Proceedings of the Chemical Society, London
Proc Ind Acad Sci	...	Proceedings of the Indian Academy of Sciences, Bangalore
Proc Roy Soc Lond	..	Proceedings of the Royal Society, London
Quart Jr Pharm	..	Quarterly Journal of Pharmacy and Allied Sciences (Pharmacology), London
Sci Cult.	...	Science and Culture.
Z Physiol Chem	..	Zeitschrift fur Physiologische Chemie

ABIES PINDROW Spach

The Essential oil from the leaves of Abies pindrow Spach, Simonsen J L, *Indian Forest Records*, 1922, *Vol VIII, Part V.*

ABROMA AUGUSTA Linn.

A note on the development of the female gametophyte in Abroma augusta L. and Pentapetes phoenicea L., Banerji, I, *Curr. Sci.*, 1941, 10, 30.

Pharmacognosy of the root bark of Abroma augusta Linn, Mitra, G. C. & Bal, S N, *Ind Jour Pharm*, 1947, 9, 120.

ABRUS PRECATORIUS Linn

The colouring matter of the seed-coat of Abrus precatorius Linn. (Scarlet variety), Ghatak N. N., *Curr. Sci.*, 1933-34, 2, 380.

Chemical examination of the seeds of Abrus precatorius L.

Constitution of abrine, N Ghatak, *Bull. Acad. Sci United Provinces Agra, Oudh, Allahabad*, 295, 1934

Chemistry and toxicology of Abrus precatorius or seed or Jequirity,. Carlos L Carbone Schi. *Semana Med (Buenos Aires)* 1947, 275.

Some components of the seeds Abrus precatorius Akira Yokoo, (Tokyo Inst. Technol) *Bull. Tokyo Inst Technol.*, 13 43 (1948)

Chemical examination of the fixed oil from the seeds of Abrus precatorius, Moolraj Mandiratta and S Dutta; *Ind Soap Jour* , 1949, 195

ABUTILON AVICENNAE Gaertn

The oil from the seeds Abutilon avicennae Gaertn G R. Stepanov., Trudin Inst Novogo, Lulyanogo Surya, 123 1933, *Chem Abst* 1933, 4945

ABUTILON INDICUM G Don

Cambir, Chem. Exam of seeds of Abutilon indicum, G. Don Indr Raj & Joshi, Sham Sunder , *J Ind Chem. Soc.* 29 (1952), 451

ACACIA Willd

Nierenstein, M -The catechins of the Cutch-producing Acacias., *J. Ind Chem Soc* 7 (1930), 279

Indian Wattle Bark *Jr. Sci Indl Res* , 1947, 6A, 447.

ACACIA ARABICA Willd

The tannin content of Acacia arabica pods, *Anon Bull Imp Inst.* 28, 1 1930

Gum arabic, influence of heat on the chemical composition and physical properties Moorjani & Narwani.-*Jr. Ind Chem Soc* 25 (1948), 502

ACACIA CATECHU Willd

Constitution of Catechol M. Nierenstein, *J Am Chem. Soc* 52, 1672, 1930

L Epi-Calechin from Acacia catechu Rao, P R and Sheshadri, T R , *J Sci Indl Res* , 1948, 7B, 59

ACACIA CHUNDRA (Willd) D. C

Acacia chundra or Acacia Sundra Chatterjee, D., *Sci & Cult.*, 1949, 14, 290

ACACIA LEUCOPHLOEA Willd

Chemical examination of flowers of Acacia leucophloea Willd. Mukherjee, S. K. and Murty, V. V. S , *Jr Sci. Indl. Res.*, 1952, 11B, 125.

ACALYPHA Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency, XVIII, Catherlics, Oleum ricini, Oleum tiglii, Aloe, Succies acalyphae, Caines, J. F. and Mhaskar, K. S., *Ind. Jour. Med. Res.*, 1923, 11, 103.

A INDICA Linn.

Chemical investigation of the plant Acalypha indica Isolation of triacetonamine, a cyanogenetic glucoside and quebrachitol. Clande Remington and G C S Roct. Onderstepoort, *J Vet. Sci. Animal Ind.* 9, 193 1937 *Chem Abst* 4629, 1938

The presence of Hydrocyanic acid in stock feeds and other plants. Acalypha indica L Donw O Steyn, *J S African Vet Med. Assoc* 9, 60, 1938 *Chem Abst* 8617, 1938

ACHILLEA MILLEFOLIUM Linn

Oil of Achillea millefolium, L R E Kremens, *J Am Pharm Assoc* 14, 399 1925

The ethereal oil content of fragrant plants of the southern territory. Achilea millefolium, O Sobolevskii, *Chem Abst* 3017, 1928.

The non pre-existence of azulene in milfoil Katherine Graham, *J Am Pharm Assoc* 22, 819 1933

Achillea millefolium Linn constituents of the petroleum ether extract of the blossoms R L Macmurray, *Am. J Pharm* 105, 573, 1933

Phytochemical notes The steroles of Achillea millefolium. Ole. Gisvold, *J Am Pharm Assoc* 24, 107, 1953

Volatile oil from Western yarrow Achillea millefolium, lanulosa R L, Macmurray, *J. Am Pharm Assoc* 25, 304, 1936

Investigation of the antipyretic action of native medicinal plants. Achilleamillefolium Maksym Naksyn Nikonorrow, *Acta Polon, Pharm* 3, 23, 1939 *Chem Abst.* 3766, 1941

ACHRAS SAPOTA Linn

Achras sapota, fatty acids & glycerides of the oil from sapota seeds Vidyarthi, N L & Mallya, M Venkatesh; *J Ind Chem Soc.* 16 (1939), 443.

ACHYRANTHES ASPERA Linn

Chemical examination of the seeds of Achyranthes aspera Linn Gopalachari, R & Dhar M L , *J. Sci Indl Res* , 1952, **11B**, 209

ACONITUM Linn

The Indian varieties of Aconite Their chemical composition and biological assay Chopra, R N , Gupta, J C & Ghosh, N. N ; *Ind. Jour Med Res* , 1928, 15, 873.

Aconite alkaloids Action of methyl alcoholic sodium hydroxide on atisine, iso-atisine and dihydroatisine-Walter, A Jacobs and Hayman E Craig, *J. Biol. Chem.* 147, 567, 1943

A. ANTHORA Linn

Atisine from Aconitum heterophyllum Wall, and authorine from Aconitum anthora, L. Andoregoris, *Comp Rend* 205, 1007, 1937

BALFOURII Stapf

The alkaloids of some Indian Aconites (A. balfourii, A. deinorrhizum and Chumbi Aconite). Thomas Anderson Henry and Thomas Marval Sharp, *J Chem Soc* 1105, 1928

CHASMANTHUM Stapf , ex Holmes.

Aconitine from Cashmerian Aconitum chasmanthum K H Baver and Tara Chand Razdan, *Pharm Zentralhalle*, 72, 146, 1931

. DEINORRHIZUM Stapf

The alkaloids of some Indian Aconities (A balfourii, A deinorrhizum and Chumbi Aconite) Thomas Anderson Henry and Thomas Marvel Sharp, *J Chem. Soc* 1105, 1928

HETEROPHYLLUM Wall.

Atisine from Aconitum heterophyllum Wall and anthorine from Aconitum anthora L Andoregoris, *Comp Rend* 205, 1007, 1937

. NAPELLUS Linn

New alkaloid from Aconitum napellus Henrich Schulz and Gottfried Berger, *Arch. Pharm* 262, 556, 1924

Aconite alkaloids New alkaloids in Aconitum napellus Werner Fredenberg and E F Rogers, *J. Am Chem Soc* 59, 2572, 1937

SOONGARICUM

Alkaloids of Aconitum soongaricum Alkaloids of Ranunculaceae. S Yunusov, *J Gen Chem* (*U S S. R*) 18, 515, 1948, *Chem Abst* 7940, 1948

CORUS CALAMUS Linn

Notes on some Indian Essential Oils Acorus calamus, Linn, (Calamus Root) Rao B Sanjiva, Sudborough J J , and Watson H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol 8A, Part X, pp 149-151

Essential oil from the rhizomes of Acorus calamus, N C Kelkar & B Sanjiva Rao, *J. Ind Inst Sci* 17, 24 1933

Studies in Indian Essential Oils VI – Essential Oil from the rhizomes of Acorus calamus Linn Kelker N C and Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1933, *Vol* **17A,** *Part II, pp.* 25-31

A. GRAMINEUS Soland.

Constituents of the essential oil Acorus gramineus, Soland, Y. Kimura, *J. Pharm Soc Japan* No 431, 380, (1926)

ACTAEA SPICATA Linn

Tinns, Actaea spicata Halmut Thaler Mikrokosmos; 22, 94, 1929, *Chem Abst* 2596, 1929

ACTINODAPHNE Nees

Actinodaphne and Litsea fats as raw material for a valuable new detergent, S. V. Puntambekar: *Ind Forester* 60, 707, 1934.

A HOOKERI Meissn.

Actinodaphne hookeri Meissn; Actionodaphine, an alkaloid from. Krishna, S. & Ghosh T. P , *J Ind Chem Soc.* 9 (1932), 429.

Actinodaphne hookeri Meissn. The fat and oil from the seeds of An indigenous source of Lauric acid Puntambekar, S. V & S Krishna , *J Ind Chem Soc* 10 (1933), 395.

ADANSONIA Linn

Some new data on the pharmacology of Adansonia leaves. Daniel Incent, *Chem Abst* 1698, 1951

ADENANTHERA PAVONINA Linn

Oil from the seeds of Adenanthera pavonina A source of lignoceric acid, S M Mubidri, P. Ramaswami Ayyar and H E Waston, *J Ind. Inst Sci* 11 A, 173, 1928

Chemical examination of the leaves of Adenanthera pavonina Linn Patel, C S , Shah C C , & Parikh, H P , *Curr Sci* , 1947, 16, 344

ADHATODA VASICA Nees

A preliminary note on the pharmacology and therapeutics of Adhatoda vasica (Basak) Chopra, R N & Ghosh, Sudhamoy, *Ind. Med. Gaz* , 1925, 60, 354.

Some observations on the Pharmacological Action and Therapeutic properties of Adhatoda vasica (Basak) Chopra, R N , & Ghosh, S , *Ind Jour. Med Res* , 1925, 13, 205

Vasicine-an alkaloid present in Adhatoda vasica Nees Part II Ghish, Tarak Prasad, *J Ind Chem Soc* 4 (1927), 1

Report of the Imperial Agriculture Chemist, Ash of the Adhatoda vasica Plant, B Viswanath, *Sci. Reports Imp Inst Agr. Research, Pusa*, 1934-35, 103, 1936

A pharmacological study of stem and leaf of Adhatoda vasica Nees. Prasad, S & Prabhu, P N , *Ind Jour Pharm* , 1950, 12, 200

ADINA CORDIFOLIA Hk f

Adina cordifolia Hook A yellow colouring matter from the wood of Jagraj Behari Lal & Dutt, Sikhibhushan, *J Ind Chem Soc* 12 (1935) 257

Fish poison from the wood of Adina cordifolia Rao, N P., *J Sci. Indl Res* , 1949, **8B**, 95

AEGLE MARMELOS Corr

Preliminary chemical examination of Aegle marmelos. Dikshit, Brij Behari Lal, & Dutt, Sikhibhushan *J Ind Chem Soc* 7(1930), 756

On the active principles of the bark of Aegle marmelos Correa Mookerji, Asima (Miss), *Curr. Sci* , 1943, 12, 209

On the alkaloidal isolation from the matured bark of Aegle marmelos. Chakravarty, K. K., *J Ind Chem Soc* 21 (1944), 401

Constituents of the matured bark of Aegle marmelos Correa. Asima Chatterjee and Sudhangsu Sekhar Mitra; *J. Am. Chem. Soc* , 71, 606, 1949.

Aegle marmelos, essential oil from leaves of Baslas & Deshpande, *J Ind Chem Soc*, 26 (1949), 23

L-d-Phellandrene from the essential oil from leaves of. Baslas & Deshpande; *J Ind Chem. Soc.* 26 (1949) 231

Aegle marmelos, Essential oils from the leaves, twigs & fruits of Baslas & Deshpande. *J. Ind Chem Soc* 23 (1951), 19.

Studies on the active principles isolated from the leaves of Aegle marmelos Chatterjee Asima (Nee Mukerjee) & Bose Sukumar, *J Ind Chem. Soc* 29 (1952), 425

AEROTHECIUM LANATUM Wakker

A preliminary note on the isolation and pathogenicity of Aerothecium lanatum Wakker and Helminthosporium tetramera Mckinney on the leaves of Oryza sativa Linn Ganguly, A K., Ganguly, D C, *Sci & Cult*, 1941, 6, 424,

AESCULUS Linn

The physiological significance of the glucosides in Aesculus and Salix, Gerhard Kerston, *Chem Abst* 6772, 1934

A HIPPOCASTANUM Linn

The hydrolysis of phytic compounds derived from seeds of hemp horse bean horse chestnut and flax, wheat and embryos of rye W, Jarosza, *Chem Abst* 5501, 1934.

Production of alcohol from horse chestnuts, Zabrodskii, golde berg Fuks and Kogan *Chem Abst* 7415, 1934

Plant pigments, the occurrence of L and B carotene in different natural products, horse chestnut P Karyer and W Schlientz Helv, *Chem Acta* 17, 7, 1934

Inhibiting effect of Pilocarpine on intestine treated with extract of horse chestnut. H Busquet Comp, Redn Soc. Biol 118, 234, 1935 *Chem Abst* 3035, 1935.

Chemistry of Aesculus saponin and its non sugar components E Bures and K Babor, *Chem Abst* 5454, 1935

Use of Aesculus hippocastanum (horse chestnut) as a raw material for distilleries, B Lampe and R deplanque, *Chem Abst.* 4616, 1936.

Electric treatment of horse chestnuts and acorns. Hayao Muraoka; *Chem Abst* 3268, 1936

Saponin content and varying saponin content of certain plants and seeds Aesculus hippocastanum L M Roberg and E Marchal Jahrb, Wiss Botan. 84, 710, 1937 *Chem. Abst.* 177, 1938.

Saponins from domestic (German) plants Aesculus hippocastanum E Wagner, *Chem Abst* 3032, 1941.

AGAVE AMERICANA Linn

Saponins A new method for preparing the foam producing principle of Agave americana, L. J. Balansard and P. Flandrin; *Bull. Soc. Chem Biol.* 27. 618, 1945.

AGERATUM CONYZOIDES Linn.

Travancore Essential oils Essential oil from Ageratum conyzoides (Appa grass). Moudgill K L, *Journal of the Indian Chemical Society, Calcutta*, 1925, Vol I

Perfumed oils of French upper Ubangi, Ageratum conyzoides L L Jolly, *Chem Abst* 2750, 1637

AILANTHUS GLANDULOSA Desf

Book of Ailanthus glandulosa Desf and its bitter principle ailanthin R Wasichy and S Oerik, *Chem Abst* 255, 1934

Ailanthus wood as paper making material L Vidal and M Aribert, *Paper Trade Jour* 85, 7, 49, 1927

Tests carried out at the French School of paper making, qualities of Ailanthus wood (Ailanthus glandulosa) L Vidal and M Aribert *Papier*, 30, 183 1927 *Chem Abst* 2062, 1927

Essential oils of the flora of Tadshikistan, Ailanthus glandulosa, V Isaev; *Chem Abst* 3180, 1934

Ailanthus glandulosa, Anna M Bernasconi, *Chem Abst* 8473, 1938

A. MALABARICA D. C

Chemical Examination of Ailanthus malabarica DC, Rastogi, R P Sharma, V N & Dhar, M L., *J Sci Indl Res* 1952, 11B, 124

ALANGIUM BEGONIAEFOLIA Roxb

A note on the development on the Embryo sac and Endosperm in Alangium begoniaefolia, Roxb (Marlea begoniaefolia, Roxb) Mitra, J N & Datta, R M., *Sci & Cult.*, 1947, 12, 451

A. LAMARCKII Thw.

Alangium lamarckii Its chemistry and pharmacological action Chopra, R N & Chowhan J S, *Ind Jour Med Res*, 1934, 21,507.

Chemical examination of the seeds of Alangium lamarckii, A Lakshminarasimhaiah, B L Manjunath and B. S. Nagaraj, *J. Mysore Uni* B 3, 113, 1942

Chemical examination of the seeds of Alangium lamarckii, isolation of alangol Prithvi Nath Bhargava and Shakhibhushan, Dutt *Proc. Ind. Acad. Sci.* 16A, 328, 1942

Chemical investigation of Alangium lamarckii Thw. Basu, N K., Nair, N S, & Bhattacharya, N N *Ind. Jour Pharm*, 1950, 12, 98

ALLIUM CEPA Linn.

Examination of some Hangarian onions A modified method for crude fiber. Zoltan sandor., *Chem Abst* 1788, 1934.

Carbohydrates of the bulbs of Allium odorum and Allium cepa, Yoshijiro Kihara. *J Agr. Chem. Soc Japan*, 11, 548, 1935

Food values and contents of Vitamin C of Hangarian, Holland and Spanish onions. Zoltan Sandor, *Chem. Abst.* 1896, 1935.

Essential oil of onion plants Allium cepa establishments; *Antonic Chirs, Perfumes France* 15, 219, 1937.

Pharmacological trials of some domestic plants Allium cepa H. Kreitmair, *Chem Abst* 3149, 1937

The antimycolic effect of ethereal oils of the leek and onion family of plants on pathogenic fungi P W Schmidt and V Marquardt, *Chem Abst* 2642, 1937.

Vitamin C content of chillies, onion & garlic, both inraw-state and when boiled with water Biswas, H G, & Das, K L., *Ind Jour Med Res*, 1939, 27, 139.

Onion juice and bacterial growth. James E Fuller and Ernest R Higgins, *Food Research* 5, 503, 1940.

The significance of nucleoelastic enzymes in the phenomenon of bacteriology, the working hypothesis and its basis Bacteriolytic agents of onion juice Ph L Spainer, E I Chartkova, Sh Gering, *Chem Abst.* 2552, 1941.

Antibacterial properties of yeasts, Fusarium species, onion and garlic C W Carpenter, *Chem Abst*, 1558, 1946

A study of the oils from the seeds of Luffa aegyptica, Benincasa cerifera and Allium cepa, K D Phadnis, A V. Rege, D. C Pishawikar and S V. Shah., *J. Univ Bombay*, **17A**, 62, 1948

Carbohydrates of Garlic (Allium sativum) and Onion (A. Cepa L Srinivasan, M, Bhatia, I S, Satya Narayana, M N, *Curr Sci* 1953, 7, 203

A MINUS

The inactivation of antibacterial agents and their mechanism of action Allium sativum and A minus C. J Cavallito, John Haya Bailey, T H. Haskell, J. R Maccromick and W F Warner, *J Bact* 50, 61, 1945

A ODORUM LINN

Carbohydrates of the bulbs of Allium odorum and Allium cepa Yoshijiro Kihara, *J Agr Chem Soc, Japan*, 11,548,1935

A PORRUM LINN.

The antimycolic effect of ethereal oils of the leek and onion family of plants on pathogenic fungi. P W Schmidt and V. marquardt *Chem. Abst.* 2642, 1937

A. SATIVUM LINN.

Bactericidal properties of acrolein (present in garlic) Richard E. Vollrath, Lucile Walton and Carl C Lindegrem, *Proc. Soc Acad Expt Biol Med* 36, 55, 1937

Vitamin C content of chillies, onion & garlic, both in raw state and when boiled with water, Biswas, H G., & Das, K. L, *Ind. Jour Med. Res.* 1939, 27, 139

Antibacterial principle of Allium sativum. its precursor and essential oil of garlic Chester J. Cavallito, John H Bailley and Johannes S. Buck, *J Am Chem Soc* 67, 1032, 1945

Investigation on Plant antibiotics Pt I Studies on Allicin, the antibacterial principle of Allium sativum (Garbi), Raghunandana Rao, R Srinevasa Rao, S & Venkataraman, P R , *J Sci Indl Res* , 1946, 5, 31

Oxidase of garlic. J Sugihara & W V Cruess, *Chem Abst* , 6508 1946

On the effect of garlic on bacteria, Omer Ozek, *Chem Abst* , 6547, 1946

Pharmacological investigation of Allium sativum general action Action on arterial pressure and respiration Giuseppe santeilppo and Giuliano OHaviono, *Chem Absta* 7384, 1946

Allin the pure mother substance of garlic oil A Stoll and E Scebeck, *Experimentia*, 3, 114, 1947

Experimental studies of the Pharmacology of the active principles of Allium sativum (garlic), Enrique Umbert de Torres Casana, *Chem Abst* 2172, 1947

Antibiotic principles of Allium sativum (Lahsan), Datta, N L, Krishnamurti A & Siddiqui, *S J Sci Indl Res* , 1948, **7B**, 42

Allicin the active principle of Allium sativum, isolation, physical properties and antibacterial action, Chaster J , Cavallito and John, H. Bailley, *J Am Chem. Soc* 66, 1950

A preliminary report on 'Loop Top' of Garlic Mahmud, K A , *Sci & Cult* , 1952, 71, 296

ALOE Tourn ex Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency, XVIII Catherlics Oleum ricini, Oleum tigli, Aloe, Succies acalyphae, Caines, J F , Mhaskar, K S , *Ind Jour Med Res* , 1923, 11, 103

Areca nuts and aloes H Octtel Apoth, *Ztg* 48, 1457, 1933

Chemical examination of Indian Aloes R N Chopra and N N Ghosh, *Arch Pharm* , 276, 348, 1938

Further studies on Cathartic action in mice, Senna, Aloe Casara and bile salts, Lloyd W Hazleton and Kathleen D Talbert, *J Am Pharm Assoc* 33, 170, 1944

Indian Aloes , Mukerji, B , & Karkum, J. N , *Ind Jour Pharm* 1950, 12, 236

A ABYSSINICA Lam

A chemistry of Ethopian Aloe and possibility of use of A abyssinica Lamarck,E Baccari, *Chem Abst* 4547 194

A VERA Linn

Indian Medicinal Plants Studies in the specifications of Aloe vera Linn. Sen Gupta, S S , & Gupta, H N , *Ind Jour Pharm* , 1949, 11, 27.

ALSOPHILA GLABRA Hook

Spore germination in Alsophila glabra Hook Kachroo, P , *Sci & Cult* , 1952, 17, 342

Alsophila glabra (B_1) Hook from Sylhet Islam, A S , *Sci & Cult* , 1952, 17, 473

ALSTONIA Br

Pharmacology and toxicity of Alstonia alkaloids, Patricia Koegh and F. H Shaw, *Australian J. Expt Biol Med Sci* 21, 183, 1943

A SCHOLARIS Br

Total alkaloids in Alstonia scholaris The Popular and well known indigenous antimalarial drug Rakshit, J N , *Sci & Cult* 1944, 9, 303. Quinine and Alstonia scholaris (CHHATM) in Malaria Das Gupta, B M , Siddons, L B , & Chakaravarti, H *Ind Med Gaz* , 1943 79, 408

Total alkaloids in Alstonia scholaris Dutt, A T , *Sci & Cult* 1944 9, 556

ALTHAEA OFFICINALIS Linn

Oils from Althaea officinalis and Malva arborea, H Ya Tropp, *Chem Abst* 1597, 1936

A. ROSEA Linn

Damping off in Hollyhocks Srivastava, H C , *Sci & Cult* , 1951, 17, 91

AMARANTHUS GANGETICUS Linn.

Availability of Calcium in Lady's finger (Hibiscus esculentus) cabbage (Brassica oleracea, capitata), Drumstick (Moringa oleifera) and Amaranth Tender (Amaranthus gangeticus) Pt I Availability of Calcium in vegetables determined by experiments on growing rats Basu, K P , & Ghosh, D , *Ind Jour Med Res* , 1943, 31, 37

Examination of the fatty oil from the seeds of Amaranthus gangeticus Chidambaram & Iyer, *J Ind Chem, Soc* 22 (1945), 117.

AMARANTUS BLITUM Linn

Colchicine induced polyploidy in Amarantus blitum Tandon, S L , & Chinoy, J J , *Sci & Cult* , 1950, 15, 398

AMOMUM XANTHIOIDES Wall

Constituents of Amomum xanthioides Wallich T. Kariyone and Y. Yoshida, *J. Pharm Soc. Japan* 50, 54, 1930.

ANACARDIUM OCCIDENTALE Linn

The food value of the nut of Anacardium occidentale (Hijli Badam). Chatterjee, N. K , *Ind Med. Gaz* , 1930, 65, 12.

Reaction products, Cashew nut-shell oil The Harvil corp. *Chem. Abst* 2584, 1931.

Composition of Rubber and Cashew nut-shell liquid, Mortimer T. Harvey, *Chem Abst* 5720, 1930

The industrial utilization of Cashewnut, Feloz Malberti, *Chem Abst* 1815, 1932.

A new globulin from Cashew nut (Anacardium occidentale) Damodaran and Tarakad G Sivaswamy, *Biochem J.* 30, 604, 1936

Cashew Nut-Shell oil as a mosquito larvicide Wats R C, & Bharucha, K, H., *Curr Sci*, 1937-38, 6, 216.

Cashew Apple a rich source of Vitamin C K Mitra, *Science & Culture*, 6, 186, 1940

Phenols from Cashew nut-shell liquid. Mortimer T. Harvey, *Chem. Abst* 1210, 1941

Study of the nuts of acajou (Anacarduim accidentale) Francica Rosa Bonchristiano, *Chem Abst*, 339, 1941)

Ethers of cashew nut-shell liquid Solomon Caplan, *Chem. Abst* 4779, 1941

Mildew in Cotton, Synthetical experiments in antiseptics for textiles Derivatives of Cashew nut oil R C Gandhi and K. Vankataraman, *Ind Jour Chem Soc & Ind*, 89, 1942

Biological value of the protein of the Cashew nut, Cyro Camargo Nogueira, *Chem Abst*, 3263, 1942

Components of the latex of Anacarduim occidentale Linn. (Cashew nut) H J Backer and N H Haack, *Chem Abst*, 1394, 1943.

Subsidiary constituents of Cashew Nut shells Pt I, Bahl, O P., Rastogi, Vishwa Nath Sharma & Salimuzzaman Siddiqui; *J Sci Indl Res*, 1946, **8B**, 222

Hypoglycemic action of Anacardium occidentale (Cashew) in normal individual, Francisco Arduino and Maria de leuides N G. Soares., *Chem Abst*, 10296, 1951

ANAGALLIS ARVENSIS Linn

Fine suspended poisonous plants, Anagallis arvensis. E. Murray Pullar, *Australian, Vet J* 15, 19, 1939

ANAMIRTA COCCULUS W & A

Preparation of purotoxin, Anamirta cocculus. E. P Clark, *J. Am. Chem Soc*, 57, 1111, 1935.

ANANAS TOURN ex Linn

A preliminary study of the anthelmintic activity in vivo of fresh pine apple juice Conarado F., Assenjo, *J Am. Pharm Assoc.* 29, 8, 1940

ANCHUSA TINCTORIA Lam.

The constituents of Alkanet root (Anchusa tinctoria) Part II Anchusin and its derivatives. Majundar & Chakravary, *J. Ind. Chem Soc* 17 (1940), 272

ANDROGRAPHIS ECHIOIDES Nees

A note on Andrographis echioides Nees, Rangaswami, S., Subbarao, V., *Ind Jour Pharm*, 1951, 13, 63

A PANICULATA Nees

Andrographis paniculata Nees, on the bitter principle from. Part I Moktader, A & Sircar, S S Guha, *J Ind Chem Soc* 16 (1939), 333

Studies in the specification of Indian medicinal plants Pt II Andrographis paniculata Sen Gupta, S B, Dutta, P P., & Banerjee, S, *Ind Jour Pharm* 1948, 10, 72

Studies in the specification of Medicinal Plants Andrographis paniculata Nees Sen Gupta, S B, Banerjee, S, & Chakravarty, D, *Ind Jour Pharm*, 1949, 11, 77

Andrographolide, the active constituent of Andrographis paniculata Nees A preliminary communication Chakravarti R N, & Mrs Chakravarti, D, *Ind Med Gaz*, 1951, 86, 96

ANDROPOGON Linn.

The constituents of some Indian Essential Oils Part VIII The Essential Oil from a new species of Andropogon occuring in the Etawah District, U P Simonsen J L, *Indian Forest Records*, 1924, *Vol X Part VIII*

Toxicity of Sorghums, Em Miege, *Chem Abst* 3013, 1933

Hydrocyanic acid content of Sorghum varieties James F Couch Reinhild R Bruise and J W Martin, *Chem Abst* 4629, 1939

A JWARANCUSA Jones

The Essential oil from Andropogon jwarancusa. Jones and the Constitution of Piperitone—J L Simonsen, (*Trans Chem Soc* 1921, *Vol.* 119 *p* 1644)

The Essential oil from Andropogon jwarancusa and the Constitution of piperetone John Lional Simonsen, *J Chem Soc* 1644, 1921.

The constituents of some Indian Essential Oils Part IV-The Essential Oil from Andropogon jwarancusa, Jones Simonsen J. L. and Rau M Gopal, *Indian Forest Records*, 1922, *Vol IX Part IV, pp* 18-22

The constitution of the terpenes present in the essential oil from Andropogan jwarancusa, Jones. John Lionel Simonsen, *Jour. Chem Soc* 2292, 1922

The constitution of the Terpene present in the Essential oil from Andropogon jwarancusa, Jones J. L Simonsen, (*Trans Chem Soc* 1922, *Vol* 121, *p* 2292)

A. KUNTZEANUS Hack

Studies in Indian Essential Oils Essential oil from the flower heads and stalks of Andropogon kuntzeanus Hack Var. Foveolata, Hack, Rao B. Sanjiva *Journal of the Indian Institute of Science, Bangalore*, 1932, *Vol* **15A**, *Part VII, pp.* 75-77

A SQUARROSUS Linn.

The Cus Cus Oil in India-Puran Singh; (*Chemist & Druggist 1914, Vol. LXXXV*) *p* 51, *Aug*, 8.

ANETHUM SOA Roxb

Notes on some Indian Essential oils Anethum soa, Roxb (Dill oil). Rao B Sanjiva, Sodborough J J and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1925 *Vol* **8A**, *Part X*, *pp* 183-184

Anethum soa, green herb and seeds—*Ind Acad Sc Vol XII, Sept*, 1939, *p* 251)

Chemical investigation of the essential oils derived from Anethum soa Roxb oil from the green and the seeds Braj Kishore Malvia and Sikhibhushan Dutt, *Proc Ind Acad Sci* **12A**, 251, 1940

ANGELICA ARCHANGELICA Linn

The essential oil in the roots of Angelica archangelica L grown in Dutch soil S P, Dijkstra, *Pharm Weekblad*, 81, 387, 1946

ANISOMELES MALABARICA Br

Chemical constitutents of Anisomeles malabarica (N O. Labiatae) Essential oil Pt I., Rao, S B, & Majumdar, D N., *Ind Jour Pharm*, 1945, 7, 123

ANNONA MURICATA Linn.

The alkaloids of Annona muricata Linn Th -M Meyor, *Chem. Abst.* 8206, 1941

A RETICULATA Linn

Alkaloids from Annona reticulata Linnacus Alfred C, Santos Philippine, *J Sci* 43, 561, 1930

A SQUAMOSA Linn

Alkaloids from Annona squamosa leaves N Trimurti, *J. Ind Inst Sci* 7, 232, 1924

Oils and fats from the seeds of Indian Forest, plants Annona squamosa Ram Chandra, R V Ghanckar and P R Ayyer, *J Ind Inst Sci* **10A**, 1927.

Isolation of Anonine from annono sauamosa Linn, Feliciana R Reyes and Alfredo C Santos Philippine, *J Sci* 44, 409, 1931.

Anonine Alfredo C Santos Phillippine, *J Sci* 47, 357, 1932

Seed oil of Formosan Plants Constituents of seed oils leguminosae, Annona squamosa Kinzo Kafulice and Chintu Hata, *J Chem Soc Japan*, 55, 369, 1934

ANOGEISSUS LATIFOLIA Wall

The tanning value of Anogeissus latfolia leaves *Anon Bull Imp Inst* 29, 137, 1931, 425, 1929

ANTHEMIS NOBILIS Linn.

Anthemis nobilis flowers, E Berthlome, *J. Pharm Belg* 15, 595. 1933

Belglan essential oils Anthemis nobilis, Ernst S. Giemtner, *Am. Perfumer*, 33, 71, 1936

ANTIARIS TOXICARIA Leschen

A preliminary note on the pharmacological action of Antiaris toxicaria Chopra, R N & De, Premankur, *Ind Jour Med Res.*, 1934, 21, 513

Glycosides and aglucons I, Antiarin (Antiaris toxicaria) K Dobel, E Schittlar and T Reichstein, *Helv Chem Acta* 31, 688, 1948

APIUM GRAVEOLENS Linn

Oil of Celery (Apium graveolens) Ernest G Guenther, *Amer. Perfumer*, 31, 75, 1935

AQUILARIA AGALLOCHA Roxb

The Nature of "Agaru" formation Bose, S R., *Sci & Cult*, 1938, 4, 89

ARACHIS HYPOGEA Linn

The properties of arachin and conarochin and proportionate occurrence of these proteins in the peanut D Breise Jones and Mullard J Horn, *J Agr Res* 40, 673, 1930, *Chem News*, 141, 38, 1930

Utilization of ground nut meal Basu, U P & Sen Gupta, S K, *J Ind Chem Soc* 21 (1944), 389

ARCHANGELICA OFFICINALIS Hoffm

Addition of nitrosyl chloride to B Phellandrene and the occurrence of the Phellandrene in some essential oils Archangelica officinalis Trustham F West, *J Soc Chem Ind 58*, 122, 1939

ARCTOSTAPHYLOS UVA-URSI Spreng

The constituents of diuretic drugs, the flavoxol glucoside of Folium Uva-Ursi and Folium vaccinii, Harnkitu Nakamura, Tatuo Ota and Genitiro Hukuti, *J Pharm Soc Japan*, 55, 800, 1935

Arbutin content of infusion, decoctions and maceration of bearbery leaves (Uva Ursi) L Zechner, *Pharm Monatsh* 17, 47, 1936

Effect of synthetic growth substances on various types of cuttings of Arctostaphylos uva-ursi J A De France, *Proc Am Soc Hort Sci* 36, 800, 1939

Standardisation of the leaves of Arctostaphylos unva-ursi, D S Blinits-ka, *Chem Abst* 7488, 1939

Conditions under which some substances of Uva Ursi can exert an antibacterial action in urine, G Madaus and Fr E Koch, *Chem Abst* 7489, 1939

Composition of the ashes of the drugs Ash of Spanish Uva Ursi leaves, I, Rosenthaler and G Beck, *Pharm Acta Helv U*, 186, 136

ARECA CATECHU Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XIV Pulvis arecae. Caines, J F & Mhasker, K S, *Ind Jour Med Res*, 1921, 9, 206

Analysis of areca nuts, P Bourect; *Bull. Sci Pharmacol*, 98, 1933

The catechol in the fruit of Areca catechu Linn Ryo Yamamoto and Tosiro Muraoka, *Chem Abst* 727, 1933

Areca nuts and aloes H Octtel, *Apoth Ztg* 48, 1457, 1933

Betel nut as a useful temofuge Hsio lian Lik, *Chinese Med Jour* 50, 1273, 1936.

Steroles, unsaponifiable constituent of Areca catechu, L Sho Kuwada and S. Yoshiki, *J Pharm Soc , Japan* 266, 1937

The Pharmacological action of areca alkalcids, Arecoline hydrochloride Chukadonago Folia, *Pharmacol Japan*, 26, 73, 1938

Koleroga disease of Areca nut Uppal, B N , & Desai, M K *Curr Sci* 1939, 8, 122

Principal physical and chemical characteristics of dum palm nut fat as orginally present and after hydrolysis, Ivo Ubaldini, *Chem Abst* 419, 1939

Anthelmintics containing areca nut, H Schlegel Arch *Expt Path Pharmacol*, 192, 389, 1939

Improved assay for Areca, *Rept Chem Abst* 2134, 1940

The pharmacological action of areca alkaloid, Arecaidene hydrochloride and arecaidenemethyl iodide, Ches Kadonaga Folia, *Pharmacol Japan*, 28, 57, 1940

Possible use of tanin from the kernels of green betel nuts Luz Baens, *Chem Abst* 6143, 1941

Pharmacology of the alkaloids and alkaloid drugs (areca nuts) Hans Braun, *Chem Abst* 4490, 1941

Nicotine like action of arecolina, U S V Euller and B Dome, *Acta Pharmacol, Toxicol* , 263, 1945

The areca and the betel Andre Mericier, *Chem Abst* 956, 1946

ARGEMONE ALBA Lestib

The alkaloids of Argemone alba, Lestib, P A Foots, *J Am Pharm. Assoc* 21, 246, 1932

A. MEXICANA Linn

Alkaloids of Argemone mexicana Alfredo C Santos and Pacifica Adkilen, *J Am Chem Soc* 54, 2923, 1932

Maxican argemone, Ode Almeda costa, *Chem Abst* 1811, 1934

(Maxican Poppy) Argemone mexicana L. Oswaldo de Almeida Costa, *Chem Abst.* 4901, 1935.

Investigation into the epidemiology of Epidemic Dropsy Pt XIII Application of the biological test to modified Argemone oil and its derivatives Lal, R. B., Das Gupta , Agarwal, S. P.. *Ind. Jour. Med. Res* , 1941, 29, 813.

Investigation into the epidemiology of Epidemic Dropsy Pt XIV Feeding experiments on human subjects to test the toxicity of some of the derivatives and modifications of Argemone oil Lal, R B , Das Gupta, Agarwal, A. C , *Ind Jour Med Res* , 1941, 29, 839

A sensitive test for the detection of Argemone oil Sarkar, S N *Curr Sci* , 1941, 10, 405

Chemical test for detection of Argemone oil Mukherji, S P , *Curr Sci* , 1942, 11, 279

A sensitive chemical test for the detection of argemone oil, the specificity of the test S N Sarkar, *Ann. Biochem* and *Exptl Med* 2, 101, 1942

A chemical method for the estimation of alkaloids present in Argemone oil and its application to a mixture of argemone and mustard oils, Sarkar, S N Rahman, Md Bazlur , *Curr Sci* , 1945, 14, 196

Detection of Argemone oil in Mustard oil, A K Sen , *J. Proc Inst Chem (India)*, 18, 102, 1946

Argemone oil, Sen, A K *Ind Med Gaz* , 1946, 81, 126

Argemone and Mustard seeds Sanyal, P K , *Ind Med. Gaz.*, 1950, 85, 498

Epidemic Dropsy A new test for Argemone oil Chakravarti, R N , Chaudhuri, R N., & Chakravarty, N K , *Ind Med Gaz* 1950, 85, 344

ARGYREIA SPECIOSA Sweet

Argyreia speciosa, fatty oil from the seeds of Kelkar, Phalnikar & Bhide, *J Ind Chem Soc* , 24 (1947), 83, 6

ARISTOLOCHIA INDICA Linn

Essential oil from the root of Aristolochia indica, Part I and II Krishnaswami, Rao Manjunath and Menon, *Journal of the Indian Chemical Society, Calcutta*, 1935, Vol XII

Aristolochia indica, Chemical examination of the roots Part I Krishnaswami, P R , Manjunath, B L and Venkta Rao, S , *J Ind Chem Soc* 12 (1935), 476

Aristolochia indica, Chemical examination of the roots of Part II The essential oil Krishna Rao, U S , Manjunath, B L & Menon, K N , *J Ind Chem Soc* 12 (1935), 494

Aristolochia indica Linn Chemical Examination of the root Part III Isolation of the alkaloid Aristolochine Krishna Swami, P R Manjunath, B L *J Ind. Chem Soc.* 14 (1937), 39

ARISTOLOCHIACEAE

Studies on Indian Medicinal Plants, Pt. I Aristolochiaceae. Chakravarty, H L., *Ind. Jour Pharm.*, 1944, 6, 96

ARNICA RUPP. ex Linn

Constituents of Arnica flowers Chemical & Pharmacological study of arnica preparations, H Thies, *Chem Abst.* 3070, 1945.

A. MONTANA Linn.

Arnica montana (Physiological action of). H S. Simpson, *J Am Inst Homeopathy*, 19, 213, 1926

Flower constituents of Arnica montana, L H Dieterle and K Fay., *Arch Pharm*, 277, 65, 1939

Constituents of the flowers of Arnica montana L.H. Dieterle and K Engelhard, *Arch Pharm.* 278, 225, 1940

Action of Arnica montana on the circulation, A W Forst, *Arch Exptl Pharm*, 201, 242, 60, 1943

Pharmacology of Arnica monatana, Otto Gessner, *Abst.* 2121. 1950.

Action of different constituents of Arnica montana on the isolated frog heart. Elma Barz, *Chem Abst*, 46, 86

ARTABOTRYS B_r.

Alkaloids of Artabotrys Artabotrin and a new alkaloid Suaveolin, Alfredo, E Santos and Feliciana R Ryees, *Chem Abst* 2251, 1933.

A SUAVEOLENS B_1

Alkaloidal content of Artabotrys suaveolens Blume Juaquin M Maranon, *Philippine J Sci.* 38, 259, 1929

ARTEMISIA Linn

The correlation between chemical composition of Anthelmintics and their therapetic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb, Melia azadirachta Linn, Punica granatum Linn, Picrasma excelsa Swartz, Vernonia anthelmintica Willd, Cocos nucifera Linn, (coconut), Caines, J F & Mkhaskar, K S., *Ind Jour Med Res*, 1923, 11, 353.

Indian Santonin Chopra, R N & Chandler, A C, *Ind Med Gaz* 1924, 59, 537

Indian species of Artemisia, Chopra, R N & Mukerjee, B, *Ind Med Gaz*, 1931, 66, 666

Indian Artemisias S Krishna and R. S Verma, *Quart Jour. Pharm Pharmacol*, 6, 23, 1933

The cultivation of Artemisia. Verma, R. S and Krishna, S, *Curr. Sci*, 1935-36, 4, 29.

A short report on the economical value of Artemisia growing in the N. W. F. Province Qazilbash, N A, *Curr Sci* 1935-36, 4, 51.

Mode of action of Santonin and Chenopodium oil, H. A. Oelkers, *Arch Exptl Path Pharmacol*, 196, 161, 1940

Synthesis of Santonin. Paranjape, K (Miss), Phalniker, N. L., Bhide, B. V. & Nargund, K S, *Curr Sci.*, 1943, 12, 150.

A new synthesis of Santonin. (Miss) Mukerji, D., *Sci. & Cultr.* 1947. 13, 258.

Commercial utilization of Indian Santonica for manufacturing santonin. Qazilbash, N A, *Ind Jour. Pharm*, 1950, 12, 175.

A. ABSINTHIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency III, Oleum absenthii. Caines, J F. & Mhaskar, K S, *Ind Jour. Med Res*, 1920, 7, 690

The reason for the curative action of Artemisia absinthium (Worm Wood) on rat Scabies G Madaus and Fr F. Koch, *Chem. Abst* 5541, 1944

Formation of bitter glucosides in the glandular hairs of certain plants (Artemisia absinthium) Jean Politis, *Comp Red* 222, 910, 1946

A KURRAMENSIS

The Kurram santonica Qazilbash, N A, *Curr Sci*, 1943, 12, 233

A MARITIMA Linn

Camphor from Erven Worm Wood, A L Mudighoyan, Artemisia maritima *Chem Abst*, 5820, 1943

A SACRORUM Ledb var minor

Oil of Artemisia sacrorum Ledb Var minor Ledh Z. G. Chestova, *Chem Abst* 3940, 1936

A SCOPARIA Waldst. & Kit.

Chemical examination of Artemisia scoparia, isolation of an essential oil and lactone Dharmpal Parihar and Shikibhushan Dutt, *Proc Ind Acad Sci*, 25 A, 153, 1947

A VULGARIS Linn

Nutritive value of the leaves of Artemisia vulgaris, L Var Indica Maxim R Sakai and T Masaki, *J. Pharm Soc Japan*, 53, 251, 1933

Blumea densiflora and Artemisia vulgaris, their insecticidal and larvicidal properties R N Chopra and S M. Ghosh, *J Malaria Inst India*, 3, 495, 1940

ARTOCARPUS HIRSUTA Lamk

Artocarpus hirsuta Lamk Fixed oil from the seeds of Varier *J. Ind Chem Soc* 22 (1945), 275.

A. INTEGRIFOLIA Linn

Artocarpus integrifolia Crystallographic investigation of Artoste none, the stenone isolated from the Indian summer fruit by means of Goniometer & X-rays Nath, M C & Mukherji; D. L.; *J Ind Chem. Soc* 16 (1939), 229

On the occurrence of a mixed inflorescence in Artocarpus integrifolia Linn Guha, M. P., *Sci & Cultr*, 1945, 11, 99

Vivipary in Artocarpus integrifolia Linn Venkate, V., *Sci. & Cult.;* 1950, 15, 361

ARUNDO DONAX Linn

Arundo Tourn. *J Sci Indl Res.*, 1946, 6A, 276.

ASARUM EUROPEUM Linn.

Composition of Asarum oil (Asarum europeum L) Viktor Bruckner and Tibor Szeki, *J. Prakt. Chem* , 134, 107, 1932

Comparative results of toxicological investigations of the Romanian plant Asarum europeum and the Japanese Sieboldii Julium Orient., *Chem Abst* , 6247, 1934.

Active principles of Asarum europeum L. Leaves M A. Abdulmenev, *Chem Abst* , 7411, 1946.

ASCLEPIADACEAE

The existence of Resinole in Asclepiadaceae Rama Murti & Sheshadri, T R , *Ind Jour Pharm* , 1944, 6, 86

ASPARAGUS Linn

Sciponins (Sapogenins) from Indian Medicinal Plants I Sapogenins from Asparagus Rao, S B , *Ind Journ Pharm*, 1952, 14, 131.

A ADSCENDENS Roxb

Studies on tuber mucilages Pt V Mucilage from the tuber of Asparagus adscendens Roxb Rao, P A , Beri, R. M , & Budhiraja, R. P , *J Sci. Indl Res* , 1952, **11B**, 127

A RACEMOSUS Willd.

Studies of Tuber Mucilages Pt IV Mucilage from the tubers of Asparagus racemosus Rao, P S , Beri, R M & Budhiraja, R P , *J Sci Indl Res* , 1951, **10B**, 261

Studies on tuber mucilages Pt VI. Partial hydrolysis of the mucilage from the tubers of Asparagus racemosus Willd Rao, P.S & Budhiraja, *J Sci Indl. Res*, 1952, **11B**, 209

ASTERACANTHA LONGIFOLIA Nees

Investigation of Indian Medicinal plants, Asteracantha longifolia N K Basu, S B Lal and S N Sharma, *Quat Jour Pharm. Pharmacol*, 20, 38, 1947

Investigation of Indian Medicinal plants, Asteracantha longifolia, Trianthema monogyna, Boerhaavia diffusa N K. Basu, S B. Lal and S N Sharma, *Quart Jour Pharm Pharmacol* , 20, 38, 1947

Chemical investigation of Asteracantha longifolia Nees Basu, N K & Gode, K , *Ind Jour Pharm* , 1952, 14, 212

ASYSTASIA COROMANDELIANA Nees

A Cercospora leaf spot of Asystasia coromandeliana Nees. Mahmud, K. A *Sci & Cult* , 1952, 17, 474.

ATALANTIA MONOPHYLLA Corr.

Essential oil from Atalantia monophylla M T. Chobe and B. Sanjiv Rao, *Proc. Soc. Biol Chemist. India*, 2, 16, 1937.

Oil of Atalantia monophylla M T Chobe and B. S. Rao; (*Proc. Soc. Biol Chem India*, V 2, 1937) Part I

Essential oil from Atalantia monophylla Nayak & Guha, *J. Ind Chem Soc* 28 (1951), 713.

ATHYRIUM FILLIX-FOEMINA Roth

A study of Athyrium filix-foemina S Trupp and Forres, J. Goodrich, *J. Am Pharm Assoc* , 29, 286, 1940

ATRIPLEX HORTENSIS Linn

Salt bushes (Atriplex hortensis) and their allies in the United States G L Bidevall and N S Golding, *Chem Abst* , 1120, 1926

ATROPA Linn

Distribution of alkaloids in two species of belladonna. E Est , F. Cabelleron, Josefina Tomas and Y Royo, *Chem Abst* 7600, 1937

Manufacture of Atropine from Belladonna Roots Srivastava, G. P , & Basu, N K , *Ind Jour Pharm* , 1944, 6, 3

Increased alkaloidal contents of induced polyploids of Datura, Atropa and Hyoscyamus Datura species Hack Marris Rowson, *Quart Jour Pharm Pharmacol* 18, 175, 1945

Note on the quality of Tincture Belladonna of Indian manufacture Bhattacharya, A., *Ind Jour Pharm* , 1947, 9, 78

A comparative study of Atropa species grown in Kashmir Kapoor, L D , Handa, K L & Kartar Singh, *Ind Jour Pharm* , 1951, 13, 249

A ACUMINATA Royle.

Assay of Indian belladonna root W A N Markwell, *Pharm J.* 146, 259, 1941

Seasonal variation in the alkaloidal contents of the leaves of Atropa acuminata , I C Chopra, K L Handa and L D Kapoor, *Ind Jour Agri Sci* , 16, 144, 1946

Pharmacognostic and chemical studies of Indian Belladonna, Atropa acuminata Herber W Youngken and William E Hassan, *J Am. Pharm Assoc Sci.* Ed , 37, 450, 1948

Indian belladonna, R Chatterjee and J K Lahiri; *J Am Pharm Assoc* , 38, 11, 1949

Cultivation of Atropa acuminata Royle ex Lendly Kapoor, I, D , Handa K L & Chopra, I C , *J Sci Indl Res* , 1952, **11A**, 534

A BELLADONNA Linn

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects, Clyde C., Hamilton and Lousie G. Gemelle, *J Econ Entmol* 27, 446, 1934

Detection of Scopoletin in radix gelsemi and radix belladonna Robert Fischer and Herbert Ehrlich, *Arch. Pharm* , 274, 268, 1936.

The Bulgarian belladonna root. P Kuiper and P. Van der Weiler, *Pharm Week Bland*, 74, 1546, 1937

Analysis of the alkaloidal mixture in Atropa belladonna A Kuhn and G Schafer, *Chem Abst* , 4277, 1938

Drug extraction The effect of pressure and vaccum on efficiency of extraction belladonna root Em J Husa and Geo R. Jones , *J Am Pharm, Assoc* , 27, 852, 1938

Belladonna root and Bulgarian treatment Angelo Firrari, *Chem Abst* , 6397, 1938

Variation in the alkaloidal content of Atropa belladonna during a vegetative period A Kuhn and G Schafer , *Pharm Exnt Zentralhalk*, 80, 151, 1939

Intubation studies of the human small intestine The effect of atropine and belladonna on the motor activity of the small intestine and colon, Kendall A Elson and J L Drossener, *Am J Digestive Diseases Nutrition*, 6, 589, 1939

Bulgarian treatment of Parkinsons disease, Pharmacological aspects and chemical effects of alkaloids of belladonna root Hermann Vollmer, *Chem Abst* 5539, 1940

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves Bernard S Feinstein and Frank J Slama, *J Am Pharm Assoc* , 29, 370, 1940

Dynamics of accumulation of alkaloids extractives and inorganic compounds in different growth stages and diferent parts of the Belladonna plant N A Valyashko and A Sova; *Farmatsiza*, 4, 20, 1940

The growth effects of thiamine chloride, ascarbic acid and Phytohormones on Belladonna and Ricinus Loxis C Zoft, *J. Am Pharm Assoc* 29, 487, 1940

The determination of the quality of Belladonna N V Kuznestov, *Chem Abst* 856, 1940

The effect of varying temp, upon the total alkaloid content of folia Stromoni and folia Belladonna H Fluck, *Chem Abst* 3878, 1940

Suggestions for the collection, preparing and keeping Belladonna root for Pharmaceutical purposes Arturo Nanniz, *Chem Abst* 2672, 1941

Alkaloids of Bulgarian Belladonna root Harold King and Hancelot L Wave, *J Chem Soc* 331, 1941

Alkaloids of Belladonna root H Kreitmair, *Chem Abst* 9541, 1941 Germination of Belladonna seeds R Melville, *Pharm J.* 146, 116, 1941.

Study of Portugese Belladonna. Manuel Riberiro Cabral, *Chem Abst* 1811, 1942.

Amino neuteric determination of alkaloids in Belladonna, lobelia herb and harmel roots G. Ya Kait, *Chem. Abst* 2996, 1942.

Experimental investigation of the influence of Belladonna, atropine and other substances on the strychnine poisoning of white mice Fr E. Koch and Fr Naumann, *Chem Abst* 6623, 1942.

Some diseases of Belladonna in California and their control Johan T. Middleton, Plant diseases Rept 25, 513, 1941 *Chem Abst* 2985, 1942

Analysis of alkaloidal mixture in Atropa belladonna L. W. Marki, *Pharm Acta Helv* , 18, 215, 1943

Belladonna cultivation in Eastern Washington, Morris Wolfred; *Am. J. Pharm* , 115. 100, 1943

The influence of the application of artificial fertilizer on the concentration of active ingredients (total alkaloids) in the roots of Atropa belladonna E De Conno, *Chem Abst* 3218, 1943

The differentiation of Scopolia and Belladonna leaves. D. Markovic, *Chem. Abst.* 10, 1, 9, 1944

Tincture of Belladonna Timoteo. A. Estevz, *Chem Abst* , 2790, 1944

Tetraploidy in Atropa belladonna, Gyula Szomolany, *Chem. Abst* 3318, 1944

Synthesis of Hyoscyamine in Atropa belladonna and Datura stramonium, B. T. Cromwell, *Biochem. Jour* , 37, 717, 1944

Culture study of the drug plant Atropa belladonna, W R. Brewer and Alex Laurie, *Chem Abst* 3419, 1944.

A belladonna and Hyoscyamus extract which is stable in the air and contains the alkaloids of the drug in mutually unchanged relative proportion, J Gjesing Andersen, *Chem Abst.* 6497, 1944

Biological assay of galenical preparations of Belladonna Jeanne Lévy, *Bull Soc Chem Biol* 27, 431, 1945

Biosynthesis of the Belladonna alkaloids W O. James, *Nature*, 158, 654, 1946

Culture studies on the drug plant Atropa belladonna, W R Brewer and Alex Laurie, *Chem Abst* 4480, 1946

Experimental work in breeding medicinal plants. Brief report concerning the examination of properties of Atropa belladonna in 1948-49 Alenka Mstnak, *Chem. Abst* 1650, 1950

Influence of Belladonna tincture on the gastric activity of children. E. Pere Delgado and Garcia Morato Castano, *Chem Abst* 4576, 1950.

Belladonna roots and leaves cultivated in this country (as compared with those existing in the market). Felix A. Fares Rail; *Chem. Abst.* 6580, 1950.

Cultivation studies of the Solanaceous drugs Alkaloidal content of mature and unmature belladonna leaves W R Brewer and L. David Hiner, *J Am Pharm Assoc* 39, 638, 1950

The inhibition phenomena in Atropa belladonna Juan Bautista Abad Manvique; *Chem Abst* 3098, 1950

ADIRACHTA INDICA A. Juss. (MELIA)AZADIRACHTA Linn)

Chemical composition of Nim or Margosa oil *Ind. Jour. Med Res* 1920, 8, 356

Odorous and bitter constituents of neem oil E R Watson, N G Chatterjee and K C Mukerjee, *J Soc Chem Ind* 42, 387, 1923

Odoriferous principle of Neem Oil—*J S C I*, Sept, 1923, Page **387 T**

The correlation between chemical composition of Anthelmintics and their therapuetic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum Rutae (Ruta graveolens Linn), Butea monosperma Roxb , Melia azadirachta Linn , Punica granatum Linn , Picrasma excelsa Swartz , Vernonia anthelmintica Willd , Cocos nucifera Linn (coconut) Caines, J F & Mhaskar, K S., *Ind Jour Med Res.*, 1923, 11, 353.

A note on the efficacy of neem batties in the destruction of rat and rat fleas in rat burrows Tiwari, C D & Lal R B , *Ind Med Gaz* 1925, 60, 310

Neem oil, The bitter principle of Neem oil Part I Sen, Rajindra Nath & Banerjee, Ganapari, *J Ind Chem Soc* 8 (1931), 773

Melia azadirachta, The Chemistry of Oleo margosa from Neem oil Part I Insolation of the constituents of the oil—Mohmmad Qudrat-i-Khuda, Ghosh, Subbash Kumar and Mukerjee Austosh, *J Ind Chem Soc*, 17, (1940) 189

A Margosa tree without the bitter principle Thirumalachar, M J , *Curr. Sci* , 1941, 10, 413.

A Margosa tree without the bitter principle Jacob, Cherian, K , *Curr. Sci.*, 1941, 10, 335 & 414

Fatty acids of neem oil, C J Dasa Rao and T R Seshadri, *Proc. Ind. Acad. Sci* , **15A,** 161, 1942

Bitter principles of the Neem oil Rangaswami, S., *Curr. Sci* , 1942, 11, 367

Utilization of Nim oil and its bitter constituents (Nimbidin series) in pharmaceutical industry. Salimuzzaman Siddiqui and Chittaranjan Mitra; *Jour Sci Indl Res* , (1945-46), 4, 5.

Chemical examination of the flowers of Melia azadirachta. Subramanian, S. Sankara & Rangaswamy, S ; *Curr. Sci* , 1947, 16, 182

Chemical examination of Nim blossoms (Melia azadirachta, Flora), Chittaranjan Mitra, Narasimha Rao, P., Satyendra Bhatacharji & Salimuzzaman Siddiqui, *J Sci Indl Res* , 1947, **6B**, 19

Denaturation of alcohol by Neem cake distillate Godbole, N N & Pande, G D , *J Sci Indl Res* , 1947, **6B**, 121

A note on the chemical examination of Nim blossoms (Melia azadirachta) Mitra, Chittaranjan & Siddiqui, S , *Curr Sci* , 1948, 17, 51

A note on the chemical examination of Nim (Melia azadirachta) exudate Satyendra Bhattacharji, Chittaranjan Mitra & Siddiqui, S , *J Sci Indl Res* , 1949, **8B**, 187.

A note of the chemical examination of Nim (Melia azadirachta) Trunk bark Satyendra Bhattacharji, Mitra C and Siddiqui S , *J Sci Indl Res* 1949, 813, 188

Structure of the Neem (Azadirachta indica) Gum, Mukerjee, S , & Srivastava, H C , *Curr Sci* , 1951, 20, 127

The active constituents of Nim seeds (Azadirachta indica Linn) Nimbidin and its composition. Sen, A. C , Sriram Singh & D N Majumdar *Ind Jour Pharm* , 1951, 13, 265

Preliminary study of antibacterial substances from Melia azadirachta, Chopra, I C , Gupta, K C , Nazir, B N , *Ind J Med Res* , 1952, 40, 511.

Chemical examination of the root bark of Nim (Azadirachta indica Syn Melia azadirachta) Siddiqui S , Mitra C , Rao P N , *J S I R* , 1953, **4B**, 152

Chemical examination of the Trunk of Nim (Azadirachta indica Syn Melia azadirachta) Siddiqui S , Bhattacharji S., Mitra C , *J S I R* , 1953, **4B**, 154

BALANITES AEGYPTIACA Wall

Sapogenins of Balanites aegyptiaca George A R , Kon and W T Weller , *J Am Chem Soc* 794, 1939

Some tropical seeds and their contents. Balanites aegyptiaca, Franz Berger , *Chem Abst* , 1528, 1939

Steroles from Balanites aegyptiaca Russel E Marker, Wagner T.P , Paul R , Ulshafer Dale, P J Goldsmith and Clarence H Ruof , *J Am Chem Soc.*, 65, 1247, 1943

Oil from fruit of Balanites roxburghii C B Patel, *Curr. Sci* 12, 58, 1943.

A note on the oil from the fruit of Balanites roxburghii Patel, C B *Curr. Sci* , 1943, 12, 58.

Balanites aegyptiaca and its various uses in Tehad territory P Creach , *Chem Abst.* 3777, 1944

Oil from the kernels of Lalob fruit, Balanites aegyptiaca, S A. Hussain, F G Gollear and R.T O. Connor; *J. Am Oil Chemists Soc* 26, 730, 1949

A chemical study of Balanites aegyptiaca Jean S. Goldfiem , *C hem Abst* 8572, 1949.

Balanites roxburghii, saponin from the seeds of Dhekni & Bhide J *Ind. Chem Soc.* 28 (1951), 588

BALANOPHORA POLYANDRA Griff

On the occurrence of monoecious Balanophora polyandra Griff in the Visakapatam District, Madras Presidency Dutt, B.S.M., *Sci. & Cult* , 1951, 17, 131

BALSAMODENDRON MUKUL Hook.

The chemical investigation of gum resin from Balsamodendron mukul Hook Ghosh, S & Chopra, R N , *Ind Jour. Med Res* 1942, 309, 331.

BAMBUSA Schreb

Constituents of Bamboo K Azami and S Sengoku , *Chem. Abst* 2248, 1926

Japanese plants, the occurrence of free pentose in Bamboo shoots, Shigeru Komatsu and Yasuo Sasaoka, *Bull Chem Soc. Japan*, 2, 57, 1927

The composition of Japanese Bamboo. Y. Veda, K. Kasama and K. Kimura , *Chem Abst* 3977, 1928

Biochemical study of Bamboo Shigeru Komatsu , *Chem Abst* 4807 1930

Biochemical studies in Bamboo Chemical development in the growth of shoots Choji Tanaka , *Chem Abst* 983, 1931

Chemical investigation of Bamboo The pentosans of Bamboo Sulezo Oguri , *J Soc Chem Ind Japan*, 34, 233, 1931

Biochemical studies on Bamboo , Wie Sun Tac , *J Chem Soc Japan*, 52, 882, 1931

Chemical study of Bamboo Study of Bamboo cellulose. Sutez Oguri , *J. Soc Chem Ind Japan*, 35, 347, 1932

Lethal properties of aqueous extract of young Bamboo shoots Stewart, A.C , Moorthy, U N , *Ind Med. Gaz.*, 1935, 68, 320.

B ARUNDINACEA Willd

Chemical and pharmacological examination of young sprouts of Bambusa arundinacea. N N. Ghosh & R N. Chopra ; *Arch. Pharm* 2760, 351, 1938.

BARRINGTONIA ACUTANGULA Gaertin.

Chemical examination of the seeds of Barringtonia acutangula Gaertin J K. Lahiri and S Ghosh , *J Am Pharm Assoc.* 31, 193, 1942

BASSIA LATIFOLIA Roxb

The uses of Mowrah (Bassia latifolia) , A S Carlos , *Chem Abst* 1719, 1927

Production of alcohol from spent Mohwra (Bassia latifolia) flowers D N Sahasraduthe and V G Patwardhan , *Poona Agr Coll Mog* 22, 45, 1930

Indian seed fats, Mowha (Bassia latifolia) and Tamal (Garcinia morella fats D R Dhingra, G L Seth and P C Speers , *J Soc Chem Ind* 52, 116, 1933

Some Indian vegetable oils (Bassia latifolia), G N Bhattacharya , *Ind J Phys* 10, 403, 1936

The unsaponifiable of Mahua oil from the United Provinces of India, (Bassia latifolia) N N Godbole and P D Srivastava, *Chem Abst* 5193, 1937.

Kenetics of saponification of Mohawa flowers (Bassia latifolia , R N Gobhill and R N Gobhill and N G Chatterjee , *Ind Soap J* 5, 101, 1938

Fatty acids and glycerides of solid seed fats, seed fat of Madhuca (Bassia latifolia), Mowarahfat T P Hilditch and M B Ichaporia , *J Soc Chem Ind* , 57, 44, 1938

Mohua oil soap and rancidity (Bassia latifolia) A N Ghose , *Ind Soap J* ; 4, 265, 1938

Sapogenins, Bassic acid (Bassia latifolia) B Jason Heywood George, A R Kon and Lancelot L Wane , *J Chem Soc* 1124, 1939

A preliminary note on the preparation of syrup from Mahua flowers (Bassia latifolia) V S Abhyankar and N Narayana , *Poona Agr Coll* 33, 168, 1942

Industrial Utilization of Mahua Chatterji, N G , *J Sci Indl Res* , 1944-45, 3, 265

Causation of rancidity in Mahwa oil (Bassia latifolia) B P Srivastava and S N. Sethumadhya Rao , *Soap Perfumer & Cosmestics*, 24, 673, 1951.

B. LONGIFOLIA Linn

Chemical examination of the essential oil derived from the fruit of Bassia longifolia Salgursarn Nigam and Shikhibhushan Dutt , *Ind Soap J* , 11, 131, 1945

BAUHINIA ACUMINATA Linn

Some abnormal flowers of Bauhinia acuminata Linn Trivedi, B S , & Nigam, P N , *Sci & Cult* , 1952, 17, 438

B VARIEGATA Linn

Bauhinia variegata Linn. The fatty oil from the seeds of. Punlambekar, S V & Krishna, S , *J Ind Chem Soc* 17,1940), 96

Bauhinia variegata leaves as cattle feed Kehar, N D & Goswami, M N., *Sci. & Cult* , 1951, 16, 476.

BELAMCANDA CHINENSIS Leman

Components of commercial Herbageranı (Belamcanda chinensis) Toshio Tominaga , *J Pharm Soc Japan* 62, 189, 1924

Glucoside from Belamcanda chinensis. L.C Mannich P. Schumann and Wan Ho Lin., *Arch Pharm* 275, 317, 1937

BENINCASA CERIFERA Savi

A study of the oils from the seeds of Luffa aegyptica, Benincasa cerifera and Allium cepa K D Phadnis, A V Rege, D G Pishawikar and S.V Shah , *J Univ Bombay*, **17A** 62, 1948

BERBERIS Linn

Berberine in the treatment of oriental sore Das Gupta, M B & Dikshit, B B., *Ind Med Gaz* , 1928, 64, 67.

Pharmacological action of Berberins Chopra R N , Dikshit B B & Chowhan, J S , *Ind Jour Med Res.*, 1932, 19, 1193

Berberine and Berberine-containing plants in pharmacology and therapeutics Chopra, R N , Dikshit, Chowhan, J S , *Ind Med Gaz* , 1932, 67, 194

Importance of Berberis in German therapy Walther Awe and Gg Schweizer , *Chem Abst* 5997, 1938

Chemical assay of Rasaunt and Hing from the Punjab market. Grewal, K S , Kochhar, B D , *Ind Jour Med Res* , 1940, 28, 463

Alkaloids in the Himalayan Berberis, Histological distribution of. Chatterjee, R., *Sci & Cult* , 1942, 7, 571

Himalaya species of Berberies, chemical studies on Chatterjee, R., *Sci. & Cultr* , 1942, 7, 619

The histological distribution of Alkaloids in the Himalayan Berberis. R Chatterjee , *J Am Pharm Assoc* 32, 1, 1943.

Berberine in Malaria A preliminary note Brahmachari, U. N , *Ind. Med. Gaz.*, 1944, 79, 259.

B AQUIFOLIUM Pursh.

Estimation of alkaloids in preparations of Berberis vulgaris and B. aquifolium. H. Neugebauer and K. Brunwer, *Chem. Abst.* 3669, 1939.

B. ARISTATA DC

Berberine salts from "Rasot" roots, Berberis aristata Ray, J N, Roy, B S., *Sci & Cultr*, 1941, 6, 613

Alkaloidal constituents of the Bark of B aristata (Rassaunt), Chakravarti, K K, Dhar, D C & Siddiqui, S, *J Sci Indl Res*, (1950) **9B,** 161

Alkaloidal constituents of the bark of Berberis aristata (Rassaunt). Chakaravarti, K K, *J Sci. Indl Res*, 1950, **9B,** 306

B FLORIBUNDA Wall

Berberis floribunda, isolation of Oxyacanthine, barbamine, berberine, epeberberine, Palmatine, dehydrocorvadaline, Jatrorrhizine & columabmine Chatterjee, *J. Ind Chem Soc*, 28 (1951), 225

B. HIMALAICA Arndt

Plant alkaloids, Part IV Berberis himalaica & B. tinctoria Chatterjee, R. Guha, M P. & Das Gupta, A K. *J Am Ind Chem Soc*, 29 (1952), 921.

B. INSIGNIS Hook

Umbellatine from Berberis insignis Hook R Chatterjee, *J Am. Pharm Assoc* 30, 247, 1941.

B. INSIGNIS Hook. f & B. UMBELLATA Wall

Berberis umbellata. The alkaloid of Part I Isolation and examination of Umbellatine Chatterjee, R *J Ind Chem Soc*, 17 (1940), 289

The alkaloid of Berberis umbellata Part II Chatterjee, R. *J, Ind Chem Soc* 19 (1942), 233

The alkaloid of Berberis umbellata, Part III. Chatterjee, G R, *J Ind Chem Soc* 19 (1942), 385

General pharmacology of umbellatine, a new alkaloid isolated from Berberis umbellata Wall and Berberis insignis Hook f and its use in the treatment of oriental sore. Gupta; J C & Kahali, B S, *Ind. Jour. Med Res*, 1944, 32, 53

B VULGARIS Linn

Berberis vulgaris (pharmacological action). H. S Stimpson, *J. Am Inst Homeopathy*, 19, 235, 1925

Berberine in the "Common Berberry" (Berberis vulgaris). E R Schultz, *J Am. Pharm Assoc* 15, 33, 1926

Pharmacology of Berberis vulgaris, L Henri Leclerc, *Chem Abst* 3731, 1935.

Estimation of alkaloids in preparations of Berberis vulgaris and B aquifolium. H. Neugebauer and K. Brunwer, *Chem. Abst* 3669, 1939.

Pharmacological and chemical investigation of Berbery (Berrberis vulgaris) Z Supek and D Tonnie, *Chem Abst* 9246, 1949

BETA VULGARIS Linn

Secondary products in the extraction of glutamine from beets (Beta vulgaris), C Ravenna and R Nuccorni, *Chem. Abst* 5486, 1929

Study of the activity of enzyme in the live leaves of Beta vulgaris A S Okanenko, *Chem Abst* 5688, 1931

A hypoglucemic substance in Beets W Rychlik, *Chem Abst* 6842, 1934

A chemical study of Sugar beet during the first growth year Frank Knowles J E Watkin and F W F Hendry, *J Agr Sci* 24, 368, 1934

The estimation of sugar in the leaf of the manglod (Beta vulgaris). James E. Van der Plank, *Biochem J* 30, 475, 1936

BETULA Tourn

Changes of some of the physical constants of Birch tar on prolonged storage with access to air T G Kovalev and V V Illanionc, *Chem Abst* 7061, 1935

B UTILIS D Don

Sesquiterpenes in Birch (Betula utilis D Don) K A Vesterberg and F. Nydahl, *Chem Abst* 4059, 1927

Chemical composition of tree bark Birch bark (Betula utilis D Don) M. Sharkov and Belyaevskii, *Chem Abst* 5368, 1933

Tannin extract from Birch bark (Betula utilis D Don) A A. Mukhamedov, *Chem Abst.* 5573, 1933

The pigment from Birch bred K H Bauer and Dierich Ber **66B,** 1053, 1933

Birch Sap and its use (Betula utilis D Don) C Voss, *Chem. Abst* 3260, 1934

The paraffin hydrocarbon in the oil from Birch trees (Betula utilis D Don) F Petru and J Hadaeek, *Chem Abst* 3770, 1935

BIXA ORELLANA Linn

Pharmacolgically utilizable constituents Bixa orellana L Friedrich W Friese, *Pharm. Zentrachale* 76, 4, 1935

Philippine Annato dye as colouring agent (Bixa orellana) Santiago Tachico and Augustus P. West, *Philippine J Sc* 61, 429, 1936.

South American drugs. Chemical composition of Boldus, bixol a new alcohol from the oil of Bixa orellana, M. Bachstez and G. Cavellini, *Chem Abst.* 8107, 1937.

Alleged presence of carotene in annato seeds (Bixa orellana). Gilberto Guimaraes Villela, *Chem. Abst.* 224, 1943

BLEPHARIS EDULIS Pers.

Blepharis edulis Pers Constituents of the seed of Part I Jagraj Behari Lal, *J Ind Chem Soc* 13 (1936), 109

Blepharis edulis Pers. Constituents of the seeds Part II The comp of the oil. Pendse, G P & Jagraj, Behari Lal, *J Ind. Chem Soc.* 14 (1937), 362

BLUMEA BALSAMIFERA D C

Sombaong Blumea balsamifera D C (Chemical investigation), Henri Leclerc, *Chem Abst* 2891, 1941

B DENSIFLORA D C

Blumea densiflora and Artemisia vulgaris, their insecticidal and larvicidal properties R N Chopra and S. M. Ghosh, *J Malaria Inst India*, 3, 495, 1940

B. ERIANTHA DC

Blumea eriantha DC Natural flavones Part IV On the constitution of Evianthin, the yellow colouring matter of. By Prafulla Kumar Bose & Dutt, Phanibhushan, *J Ind Chem Soc* 17, (1940), 45.

B. LACERA D. C

Blumea lacera, essential oil from the leaves Baslas & Deshapande, *J Ind Chem Soc* 27 (1950), 25

B. MALCOMII Hook f

The Essential Oil from Blumea malcomii—J L Simonsen and M Gopal Rau, (*Trans Chem Soc* 1922, Vol 121, p 876)

BOERHAAVIA DIFFUSA Linn

The pharmacology and therapeutics of Boerhaavia diffusa (Punarnava) Chopra, R N , *Ind Med Gaz* , 1923, 58, 203

Chemical examination of Purer-Nava or Boerhaavia diffusa Linn. Radha Raman Agarwal and Shikibhushan Dutt, *Proc Acad Sci* 4, 73, 1934

Chemical examination of Punarnava or Boerhaavia diffusa Linn Isolation of an alkaloid punarnavine Radha Raman Agarwal, Shikhibhushan Dutt, *Proc Acad Sci* 5, 240, 1935

A comparative study of Boerhaavia diffusa Linn and the white and red flowered varieties of Trianthema portulacastrun Linn *Ind Jour. Med Res.*, 1940, 28, 475

BOMBAX MALABARICUM D C.

Bombax malabaricum T P Ghosh; *Ind Forester*, 61, 93, 1935.

Bombax malabaricum Oil from the seeds of Rao, Rao & Venkateswarlu, *J Ind Chem. Soc.* 20 (1943), 403

BORASSUS FLABELLIFER Linn

Investigation on the Palm juice Analysis of the fresh and fermented juice (Borassus flabellifer) V S Basrur and M Qureshi, *J Osmania Univ* 7, 19, 1939

Edibles from Borassus flabellifer (Palmyra palm) with special reference to Nira or sweet toddy K Mitra and H C Mitra, *Ind Jr Agr. Sci* 10, 824, 1940

Biological value of proteins of Palmyra fruits—Borassus flabellifer Linn S N Sarkar and A C Bose, *Ann Biochem Exptl Med.* 5, 59, 1945.

BOSWELLIA SERRATA Roxb

The Constituents of some Essential oils Part VII—The Essential oil from the gum-oleo-resin of Boswellia serrata, Roxb Simonsen J L, *Indian Forest Records*, 1923, Vol IX, Part XI

Constituents of the essential oil from the gum-oleo-resin of Boswellia serrata Roxb O Roberts, *Chem Abst* 566, 1924

Chemical constitution of gum from Boswellia serrata, M A Malandkar, *J. Indl Inst Sci* **8A**, 240, 1925

Preparation of turpentine from Boswellia serrata (Roxb), gumoleoresin. R S Pearson and Puran Singh, (*Indian Forest Records* Vol VI, Part VI)

BOTRYCHIUM VIRGINIANUM Sew.

Phyllody in Botrychium virginianum Sew Govindu, H C. & Narayana, H S., *Sci & Cult*, 1949, 15, 199.

BRAGANTIA WALLICHII Br

Chemical examination Manjunath, B L. & Shankara Rao, M S. *J Ind Chem Soc* 15 (1938) 646.

BRASSICA CAMPESTRIS Linn

The nitrogenous compound in oil cake of Brassica campestris Var. Chinensis Kiyohisa Yoshimura and Shiro Fujise, *J. Chem. Soc. Japan*, 45, 427, 1924

B JUNCEA Hook f & T

A simple method for the removal of the toxic alkaloids from mustard oil adultrated with Argemone Roy, A C., *Curr Sci*, 1950, 19, 91

Argemone and Mustard seeds Sanyal, P K., *Ind Med Gaz.* 1950, 85, 498.

Purification of mustard oil contaminated with Argemone Oil. Om Parkash, Atma Ram & Brahma Prakash, *Curr. Sci.* 1951, 20, 16.

B. OLERACEA Linn

Availability of Calcium in Lady's finger (Hibiscus esculentus), Cabbage (Brassica oleracea Var capitata), Drumstick (Moringa oleifera) and Amarnath Tender (Amaranthus gangeticus) Pt I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K P & Ghosh D., *Ind Jour Med Res.*, 1943, 31, 29.

Availability of calcium in Lady's finger (Hibiscus esculentus), Cabbage (Brassica oleracea Var capitata), Drumstick (Moringa oleifera) & Amarnath Tender (Amaranthus gangeticus) Pt. II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P Ghosh, D., *Ind. Jour. Med Res*, 1943, 31, 37.

Colchicine induced polyploidy in Brassica oleracea Var Botrytis L Tandon, S L, *Sci & Cult* 483, 16, 1951

BROSIMUM GALACTODENDRON D Don ex Sweet

Cow tree Chatterjee, D, *Sci & Cult* 1950, 16, 116

Introduction of 'Cow Tree' in India, Biswas, K, *Sci & Cult.*, 1950, 16, 195.

BRUCEA SUMATRANA Roxb

Composition of the seeds of Brucea sumatrana. Heikei Uno, *J. Pharm. Soc. Japan* 63, 579, 1943

BRYONIA ALBA Linn.

Bryonia alba, a typical animal proving T G Mitchell, *J. Am Inst. Homeopathy*, 19, 105, 1926

BRYOPHYLLUM CALYCINUM Salisb

Some common indigenous remedies, Picrohiza kurroa, Erythrina indica, Sansiviera zeylanica, Pongamia glabra, Hygrophylla spinosa, Bryophyllum, colycisa Rheum emodi, Solanum indicum R N Chopra, & S Ghosh, *Ind Med Record* 55, 77, 1935

Organic acids of the leaves of Bryophyllum calycinum, Geo W Pucher, *J Biol Chem* 145, 511, 1942.

BURSERA Linn

Mysore Linaloe oil *J Sci Indl Res*, 1944-45, 3, 516

BUTEA Roxb,

The treatment of intestinal worms with the Indigenous drugs. Butea, Embelia and Kamala Mukerji, A.K., Bhaduri, N V, *Ind Med. Gaz*, 1947, 82, 66.

B. MONOSPERMA (Lam.) Kuntze. (B FRONDOSA Roxb)

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Santonin, Oleum rutae (Ruta graveo-

lens Linn), Butea monosperma Roxb , Melia azadirachta Linn , Punica granatum Linn , Picrasma excelsa Swartz , Vernonia anthelmintica Wild , Cocos nucifera Linn. (coconut), Caines, J. F. & Mhaskar, K S , *Ind Jour Med Res* , 1923, 11, 353.

The chemical examination of the oil from the seeds Butea frondosa Roxb Tummin Katti, M C & Manjunath, B L , *J Ind Chem Soc* 6 (1929), 839

Butea frondosa flowers, isolation of a crystalline glucoside of Butin Jagaraj, Behari Lal & Dutt, Sikhibhushan, *J Ind Chem Soc* 12 (1935), 262

Butea frondosa Roxb On supposed occurence of acids with uneven number of C atoms in vegetable oils and fats Part I. The acid fraction of Mean M W 354 from the seeds Krishna Rao, U. S & Manju Nath, B L , *J Ind Chem Soc* 12 (1935), 611

Constitution of Burtin Isolated from Butea frondosa flowers. Behari, L J , *Curr Sci* , 1936-37, 5, 426

Butea frondosa Studies on the enzymes of the seeds. Part I Proteolytic enzymes Chatterjee, N. R , Ghosh, S & Chopra, R N , *J Ind Chem Soc.* 15 (1938), 101

Butea frondosa Studies on the enzymes Part II Lipolytic enzymes. Chatterjee, N R , Ghosh, S & Chopra, R N , *J Ind Chem Soc* 15 (1938), 107

Occurence of free buteen and butin in the flowers of Butea frondosa. P Bhaskara Ramamurti and T R Sheshadri, *Proc Ind Acad. Sci.* **12A**, 477, 1940

Preliminary report on the presence of an estrogenic substance in the storage root of Butea frondosa Supvanta, *Chem Abst* 2929, 1940

The Phlobatannins of Kino and buteagums G V Krishnamurti and T R Sheshadri, *Proc Ind Acad Sci* **22A**, 134, 1945.

Chemical investigation of the seeds of Butea frondosa, Examination of the fixed oil Dharmpal Parihar and Sikhibhushan Dutt, *Ind Soap Jour* 12, 26, 1946.

B SUPERBA Roxb

Butea superba Roxb Khasem Pangsri Vongse *Chem Abst* 3088, 1938.

Chemical composition of flowers of Butea superba Subba Rao, V., Sheshadri, T R , *J Sci. Indl. Res* , 1949, **8B**, 178.

CAESALPINIA BONDUCELLA Fleming

Some constituents of Caesalpinia bonducella nut. Part II Bonducella Nut oil Godbole, S. N Paranjpe, D. R. & Shirkhande, J. G ; *J. Ind Chem Soc* 6 (1929), 295.

Caesalpinia bonducella Flem. Chemical examination of seeds. Part I. Tummin Katti, M. C.; *J. Ind. Chem. Soc.* 7 (1930), 207.

Search for an antimalarial drug in the indigenous materia medica Pt. II. Caesalpinia bonducella Fleming Mukherji, B , Ghosh B K & Siddons, L B , *Ind Med Gaz.* 1943, 78, 285

C DIGYNA Rottl

Investigation on the isolation and constitution of the tannin from Indian Teripods. Biswas, H G , *Sci & Cult* , 1943, 9, 90

Gallic acid from Teripods. Biswas, H G , *J. Sci Indl Res* , 1948, **7B,** 27

CALENDULA OFFICINALIS Linn

Saponin Series, Guaic saponin and saponin from Calendula officinalis Alfred Winterstein and Gertraud Stein, *Chem Abst* 5170, 1951

CALLITRIS RHOMBOIDEA R Br

Notes on some Indian Essential Oils Callitris rhomboidea R Br (Syn Frenela rhombidea. Endl) Rao B Sanjiva, Sudborough J J and Watson H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol **8A**, Part X, pp 144-145

CALOPHYLLUM INOPHYLLUM Linn

The proximate composition of Palomaria seed, oil and resin (Calophyllum inophyllum) F. A. Soliven, *Chem Abst* 3118, 192, 1924

CALOPLACA ELEGANS

Indian Lichens Caloplaca elegans Neelakantan, S & Seshadri, T R *J Soc Indl Res* , 1952, **11B,** 126

CALOTROPIS Br

African arrow poisons Heart poisons in Calotropis latex, Gerhard Hesse Franz Reicheneder and Hans Eysenboch, *Chem Abst* 1742, 1939

African arrow poisons Crystalline Calotropis resin Gerhard Hesse, Hans Eilbracht and Franz Reicheneder, *Chem Abst* 4387, 1941

The Digitalis like principles of Calotropis compound with other cardioactive substances K K , Chem C I Bliss and E Brown Robbins *J Pharmacol.* 74, 223, 1942.

C GIGANTEA Br.

Calosterol, a sterol present in the milky juice of Calotropis gigantea Kalipoda Basu and Madhab Chander Nath; *Biochem. J* 28, 1561, 1934.

Calotropis gigantea. On the proteinasa in the milky juice, its purification and activation by ascorbic acid and gluthathione. Basu,

Kalipada & Madhab Chandra Nath, *J Ind Chem Soc.* 13 (1936), 34 Chemical composition of Calotropis gigantea wax and resin, components of the latex P Bhaskara Ramamurti and T R Seshadri, *Proc Ind Acad Sci.* **18A**, 145, 1943

Chemical composition of Calotropis gigantea The resinols of the root bark K J Balkrishna, P Bhaskara, Ramamurti and T R Sheshadri, *Pro Ind Acad Sci* **22A**, 138, 1945

Chemical composition of Calotropis gigantea flowers A comparison of the various parts of the plant P Bhaskara, Ramamurti and T R. Sheshadri, *Proc Acad Sci* **22A**, 304, 1945

Madar Juice, N Pitchandi (Calotropis gigantea) *J Proc Inst of Chemists*, 20, 34, 1948

A preliminary study of the toxicity of Calotropis gigantea Rathnasabapathy, V , Rao K & Krishnaswamy, A , *Ind Jour Med. Res* , 1949, 37, 483.

C PROCERA Br

African arrow poison calotropin (Calotropis procera) Gerhard Hesse and Franz Reicheneder, *Chem Abst* 1031, 1937

CALYCOPTERIS FLORIBUNDA Lamk

The anthelmintic constituent of the leaves of Calycopteris floribunda Araya Puram, Ratnagiriswaran, Kumar B Sehra and Krishanna Swami Vankataraman, *Bioch J* 28, 1964, 1934

The constituents of calycopterin The yellow colouring matter of the leaves of Calycopteris floribunda Shah, R C Virkar, V V & Venkataraman, K , *J Ind Chem Soc* 19 (1942), 135

Anthelmintics—Anthelmintic activity of Calycopteris floribunda M L Khorrana, D K Motiwala and K Venkataraman, *Proc Ind Acad Sci* **27A**, 121, 1948

CANANGA ODORATA Hk f & T

Cultivation of Canaga odorata and experimental production of oil of Ylang-Ylang in Java. W Bobloff, *Chem Abst* 985 1927.

C PATCHOULI

Analytical characteristics of Dutch East Indian oils of Cananga patchouli and Vetiver D R Koolhas and P A Rowan, *Perfum France*, 15, 245, 1937

Cananga patchouli and Vetiver oil, D R Koolhas and P A Rowan, *Perfumery Essential Oil Record* 29, 53, 1938

CANARIUM COMMUNE Linn

The fatty oil of the seeds from Canarium commune. Alph. Steger and J Van Loon; *Chem Abst* 3937, 1940

C STRICTUM Roxb

Travancore Essential Oils Black Dammar Moudgil K L and Aiyar K Sitarama, *Department of Industries, Trivandrum*, 1923, *Bulletin* No XVII, pp 11-17

CANAVALIA ENSIFORMIS D C

A toxic principle in the seeds of Canavalia ensiformis A Orru and A Fractoni, *Chem Abst* 6123, 1946

CANNABIS Tournef

The hydrolysis of phytic compounds derived from seeds of hemp, horse bean, horse chestnut, flax, wheat and embryos of rye W Jarosza, *Chem Abst* 5501, 1934

Resin content and Physiological activity of hemp grown in the region of Paris Rene Paris and Marce Lousie Merac, *Chem Abst* 9377, 1949

C SATIVA Linn (C INDICA Lamk)

Chemical examination of India hemp and its preparations R Weitz and Dardanne, *Bull Sci Pharmacol* 31, 321, 1924

The valuation of Charas Chatterjee, D N & Roy, N B, *Ind Med Gaz*, 1929, 64, 373

Cannabis indica, resin, Part II, Robert Sidney Cahn *J Chem Soc* 630, 1931

Cannabidiol and Cannabol constituents of Cannabis indica resin. A Jacob and A R Todd, *Nature*, 145, 350, 1940

The use of Hemp Drugs in India for Euphoric purposes Chopra, R N, *Sci & Cultr*, 1940, 5, 761

Cannabis sativa in relation to mental diseases and crimes in India Chopra, R N, Chopra, G S, & Chopra, I C, *Ind Jour Med Res*, 1942, 30, 155

Observations on the physiologically active fraction of Indian Hemp, Cannabis sativa Linn, Bose, B C & Mukerji, B, *Ind Jour. Med Res*, 1945, 33, 265

CAPPARIS SEPIARIA Linn

Vivipary in Capparis sepiaria Linn Mahobale, T S., *Sci & Cultr*, 1939, 5, 372

C SPINOSA Lam

Analysis of Capparis spinosa seed oil D Zabramnyl, A Ochakovskil and N. Petrova, *Chem. Abst* 6822, 1941.

CAPSELLA BURSA-PASTORIS Moench

Medicinal action of Capsella bursa-pastoreis, and also of its parasites, Cystopus candidus and Peranospora parasitica W Harste, *Arch Pharm*, 266, 133, 1928

Alcoholic extract of Capsella bursa-pastoris Attilio Bartola, *Chem Abst* 6063, 1932

C ANNUUM L

Pharmacodynamic action of the active principles of chillies (Capsicum annuum) Jose Lille and Eliseco Ramerez, *Chem Abst* 4836, 1935

Pungency in chillies (Capsicum annuum) a Mendelian character Deshpande, R B, *Curr Sci*, 1935-36, 4, 418

The nitrogen complex of Indian foodstuffs Condiments Pt II Chillies (Capsicum annuum) Coriander seeds (Coriandrum sativum) Narasimhamurthy, G, *Ind Jour Med Res*, 1968, 25, 863

(Capsicum) chillies, Vit C contents of Indian foodstuff Rothenheim, C A, Shaik Mohammad, H S & Cowlagi, S S, *J Ind Chem Soc* 15 (1938), 15

A case of variegation in Capsicum annuum L, Deshpande, R B, *Curr Sci* 1939, 8, 313

Vit C content of chillies, onion & garlic, both in raw state and when boiled with water Biswas, H G & Das, K L, *Ind Jour Med Res*, 1939, 27, 139

CARICA PAPAYA Linn

Protolytic enzymes of Carica papaya, N C Nag and H N Banerjee, *Trans Bose Research Inst Calcutta*, 1, 1930-31

The colouring matter of Carica papaya Ryo Yamamato and Scityu, *Chem Abst* 3480, 1933

Carica xanthin, Kryptoxanthin monopalmitate (Carica papaya), P Karrer, W Schlientz, *Helv Chem Acta*, 55, 1934

Carica xanthin of the Papaya and Citrus poonensis Hort Ryo Yamamoto and Akimasa Kato, *J Agr Chem Soc Japan*, 10, 754, 1934.

Carica xanthin a colouring matter in the fruit pulp of Carica papaya and Citrus poonenis, Ryo Yamamoto and Yosiakin Kato, *Chem Abst* 806, 1935

An alkaloidal principle isolated from Carica papaya seeds Tryambak B. Panse and Anand S Pranjpe, *Rascyanam* 1, 215, 1941

Some chemical constituents of papayas and their relation to flavour. S J. Lynch and W.M Fifield; *Chem Abst* 1135, 1941,

Carica papaya Linn , Intro-carpellary Fructification in Salam, M A , *Sci. & Cultr* 1942, 8, 143

The biological value of the proteins of papaya (Carica papaya) and Lady's fingers *Esptl Med* 2, 71, 1942

Carpasemine isolated from Carica papaya seeds T B Panse and A S Pranype, *Proc Ind Acad Sci* **18A,** 140, 1943

Puetro Rican fatty oils Characteristic composition of expressed papaya (Carica papaya) seed oil C F Asenjo and J A Hoyco, *Oil and Soap*, 20, 217, 1943

Internal proliferation in Carica papaya L Sattar Khan, *Sci & Cultr* , 1946, 12, 194.

A case of vivipary in Carica papaya Linn Sen, P , *Sci & Cultr.* 1946, 12, 153

Papain Activation of Ray, *J Ind Chem Soc* 23 (1946), 313.

Carica papaya L , study of the fruit, C A Lolano, Salcedo, *Chem Abst* 2507, 1947

Damping off of Papaya (Carica papaya L) seedlings Babu Singh & Paharia, K. D., *Sci. & Cult* 1952, 17, 477

Safflower oil The component glycerides of Vidyarthi, *J Ind Chem Soc* 20 (1943), 45.

CARTHAMUS TINCTORIUS Linn

Australian research on the use of Safflower oil C, Macmilin, *Chem Abst* 6546, 1948.

A rapid method for determining the oil content of Safflower and sunflower seeds W Keith Kennedy and John Urau, *Chem Abst* 5209, 1949

Utilization of the Tobacco seed and Safflower seed oil in varnish and paints. Pt I Sharma, P G , Budhiraja. N C & Aggarwal, J S , *J Sci Indl Res* , 1951, **10B,** 33

CARUM CARVI Linn

Shiyah Zirah (black caraway), H Stanley, *Redgrove Pharm J* 127, 496, 1931.

The essential oil of Japanese orandahakka (Crapemint), Carvone Toice Nagasawa, *J Chem Soc Idn. Japan*, 41, 252, 1938.

C COPTICUM Benth

The manufacture of Thymol from Ajowan Lahhani I V , Sudborough J J and Watson H E., *Journal of the Indian Institute of Science, Bangalore*, 1921, Vol 4, Part V pp 59-84

Ajowan Oil—R R Sobte and P Singh; (*Perfum. Record*, 14, 1923 399)

The Manufacture of Thymol Inugauti N N, Bhate S R and Habib Uasan, *Department of Industries and Commerce, Hyderabad*, 1924, *Bulletin No.* 8

Production of Thymol from Ajowan Iseeds Verghese James Gulati K C & Joshi, M L, *Curr Sci*, 1949, 18, 17

CARYOPHYLLUS AROMATICUS Linn

Clove industry in India (Article), *Sci & Cultr* 1938, 4, 221

CASSIA ABSUS Linn

Oil of Cassia absus Zafaruddin Ahmed, *Chem Abst* 8376, 1935

Cassia absus A new formula for Chaksine The alkaloid of C. absus and some expts On its constituents Kapur, Hansraj, Gaind, Kidar Nath, Narang, Kartar Singh, Raj, Jiranendra Nath, *J Ind Chem Soc* 17 (1940), 28

Studies in the alkaloids of Cassia absus Linn Pt I Vishwa Nath Puri, Vishwa Nath Sharma & Salimuzzaman Siddiqui, *J Sci Indl Res*, 1945-46, 4, 701

Chaksine Aggarwal, M L, Ray, J N, Sen, D C, *Sci & Cultr*, 1946, 12 201,

Pharmacological investigation of Chaksine, an alkaloid from Cassia absus Linn Pradhan, S N, Varadan, K S, Rov C, De N N., *J S I R* 1953, **8B,** 358

A note on the new formula for Chaksine, Siddiqui, S & Ahmad, Z, *J Ind Chem Soc* 18 (1941), 589

C ANGUSTIFOLIA Vahl

The effective constituent of Senna leaves. W Straub and H Gebhardt, *Arch Exptl Path Pharmacol* 181, 399, 1936.

A new crystalline component of Senna leaves (Cassia angustifolia), P Bhaskara, Ram Murti and S. Rangaswami, *J. Pharm* 2, 203, 1940

Further studies on Cathartic action in mice, Senna, Aloe, Cascara and bile salts, Lloyd W Hazleton and Kathleen D Talbert, *J Am. Pharm Assoc* 33, 170, 1944

Factors effecting the cathartic activity of Senna in mice Lloyd W Hazleton and Kathleen D. Talbert, *J Am Pharm Assoc* 34, 260, 1945

Crystalline active Senna glucosides, Sandoz Ltd, *Chem Abst* 587, 1945

Evaluation of the purgative activity of Senna extract, composition of bioassay and chemical assay of Senna H. O J Collier, E C. Fillier, S. K. Paris and D. M. Bellis Wuart, *J. Pharm. Pharmacol* 21, 252, 1948,

Crystallised glucosides from Senna leaves and pods Sandoz Ltd, *Chem Abst* 2376, 1949

Biological assay of vegetable purgatives, Senna leaf, fruit and their preparations, T C Lou, *J Pharm Pharmacol* 1, 673, 1949.

The repeated administration of Tinnevelly and Alexandria senna to mice Maribelle Woods and I W Crote, *J Am Pharm Assoc* 40, 198, 1951

C. FISTULA Linn

Abnormal flowers of Cassia fistula Linn Raghavan, T S , & Venktasubban, K R , *Curr Sci* 1934, 35, 3, 256

The use of Cassia fistula in the treatment of Black-water fever Venkatachalam K & Ratnagiriswarn, *Ind Med Gaz.* 1941, 76, 211

Pharmacological investigation of Cassia fistula L and some Brazillian species of Cassia Richard Wasiky, *Chem Abst* 1562, 1943

The fruit-pulp of Cassia fistula Linn Karim, M A & Guha-Sircar, S S *Sci & Cultr* 1946, 12, 194

A study of Cassia fistula Pulp Miss Modi, F K. & Khorana M L , *Ind Jour Pharm* , 1952, 14, 61

C OCCIDENTALIS Linn

Poisoning with Cassia occidentalis is due to a toxic albumin Raymond Moussa, *Chem Abst* 3320, 1925

The fatty oil of Cassia occidentalis Linn seed A. Steger J Van Loon, *Chem Abst* 2208, 1934

C TOMENTOSA Linn

Anthracene derivatives from the genus Cassia Cassia tomentosa and C corymbosa E Maurin; *Chem Abst* 1289, 1927

C. TORA Linn

Constituents of Folium Ketsumii (Cassia tora) Genichire Fukuchi and Kanco Imai, *J. Pharm Soc. Japan*, 70, 568.

Cassia tora Linn. Chemical examination of the seeds of. Part I Subba Jois, H & Manjunath. B L., *J Ind. Chem Soc.* 7 (1930) 521.

CASSYTHA Linn.

Cassytha L on Viscum Dutt. B.M.S., *Sci. & Cultr.*, 1951, 16, 373.

C. FILLIFORMIS Linn.

On Cassytha filiformis Linn. Srivastava, J , *Curr. Sci* 1945, 14, 242

On the range of Hosts of Cassytha filiformis Linn. Nayar, B.K & Nayar, P N , *Sci & Cultr.*, 1952, 17, 383.

CASUARINA EQUISETIFOLIA Forst.

Sex in Casuarina equisetifolia Forst. Chandrasekharan, S N., Parachasarathy, S V & Sundararaj, D D., *Sci & Cult*, 1949, 15, 158.

CATHA EDULIS Forsk.

Central stimulating action of the alkaloid Cathine (Catha edulis) *Arch. Exptl Path and Pharmacol* 198, 100, 1941.

CEDRELA TOONA Roxb

Essential oil from the wood of Cedrela toona, Roxb. P. Parameswara Pillai and B Sanjiva Rao, *J Soc. Chem Indl* 50, 220, 1931.

Cedrela toona Roxb Isolation of a lactone, an essential oil and a colouring matter from the wood of. Parihar & Dutt, *J Ind. Chem Soc* 27 (1950), 77.

CEDRUS DEODARA Loud.

The constituents of some Indian Essential Oils Part III. The Essential Oil from Cedrus deodara, London Simonsen J L and Rau M Gopal, *Indian Forest Records* 1922, Vol IX, Part IV, pp. 13-17

CELASTRUS PANICULATUS Willd.

Seed and fruit coat fats of Celastrus paniculatus, B.G Gunde and T P Hilditch, *J Chem. Soc.* 1980, 1938

A note on the chemical examination of Celastrus paniculatus, Warsi. Sharifuddin Ahmad *Curr Sci*, 1940, 9,,134

Chemical investigation of Celastrus paniculatus Willd. N.K Basu, P R Pabrai, *J Am Pharm Assoc* 35, 272, 1946

A note on the chemical investigation of the fruit of Calastrus paniculatus Willd (N O Celastraceae)

Shah, M M, Phalinkar, N L, Bhid, B V, *Curr Sci* 1947, 16, 57

CENCHRUS PAUCIFLORUS Benth

New Plant immigrant, Cenchrus Pauciflorus Benth (Gramineae, A). Sen, S K, *Sci & Cultr*, 1942, 459, 7.

CENTIPEDA ORBICULARIS Lour.

Centipeda orbicularis. Vyas, B N & Sinha, H K, *Ind Med. Gaz.*, 1930, 65, 75

Chemical examination of Centipeda orbicularis Lour Rastogi, R P Sharma, V N. & Dhar, M L, *J S I R*, 1953, **7B**, 332.

CEPHAELIS IPECACUANHA Rich

Review of the Pathogenesis of Ipecac and its alkaloids, Megavack, T H, *J. Am Inst of Homeopathy*, 21, 300, 1928

Indian Ipecacuanha. Chopra, R.N. & Mukerjee, B., *Ind. Med. Gaz*, 1932, 67, 88.

Treatment of chronic intestinal amoebiasis with Gavano, a derivative of Ipecacuanha. Chopra, R N , Sen, S. & Sen, B, *Ind. Med Gaz* , 1934, 69, 130,

Action of emetine on the activity of the adrenal & thyroid glands. Chopra, R.N , Gupta, J.C & Roy, A C , *Ind Jour Med. Res.*, 1935, 22, 771.

Radix Ipecacuanha and its alkaloids as expectorant and emetics Pfisers A , *Pharm Ztg* 81, 503, 1936.
The emetic action of Ipecacuanha, Mukhnacheva A I , *Chem. Abst* 6449, 1936.

Biological studies on tincture Ipecac Astrue A , Girovex J and S Barrau, *J Pharm Chem* 1, 185, 1940
Assay of Indian Ipecacuanha Guha, R C , & Mukerji, B , *Sci. & Cultr.*, 1945, 11, 204.

Pharmacognostic studies on 'Ipecacunha'. Bal, S N., & Datta, S C *Sci & Cult.*, 1945, 10, 448

Pharmacognostic studies on Indian Ipecacuanha Bal, S N , & Datta S C , *Sci & Cult* , 1946, 12, 201.

Pharmacognostic studies on Indian Ipecacuanha Bal, S N & Datta, S C *Ind Jour. Pharm* , 1946, 8, 76.

Ipecacuanha (Cephaelis ipecacuanha) of Brazil Arnaldo Augusto Addor, *Chem Abst* , 1812, 1947

Observations on the growth of Cephaelis ipecacuanha (Bort) A Rich in Calcutta. Mitra, G C & Chakravarti, D , *Sci & Cult* 1948, 13, 504.

On the cultivation of Ipecacuanha in India Biswas, K P & Sampathkumaran, M A , *Sci & Cult* 1948, 14, 160.

A process for the preparation of emetine HCl from Indian Ipecacuanha Shah, S K , *J Sci Ind Res* , 1951, **10B**, 76

CEPHALANDRA INDICA Naud

Cephalandra indica (Telakucha) in Diabetes Chopra, R.N , & Bose, J P , *Ind Med Gaz.*, 1925,13, 11

Observations on the antidiabetic preparations of Cephalandra indica (Telakucha) Chopra, R N , & Bose, J P , *Ind Med. Gaz* 1925, 60, 201.

CERBERA ODOLLAM Gaertn

Oil and fats from the seeds of Indian forest plants, oil from the seeds of Cerbera odollam. Ram Chandra, Ghanekar R V. and Ayyar, P.R. *J Ind Inst Sci* **10A**, 20, 1927

A preliminary study of pharmacological action of the glucoside of Cerbera odollam Chopra, R N., Bose, B.C., Gupta, J.C. & Chopra, I.C., *Ind. Jour Med Res.*, 1942, 30, 107.

CHAMAENERIUM ANGUSTIFOLIUM Schur.

Comparative plant chemistry Chamaenerium angustifolium Konstantia Puringer, *Montash* 44, 247, 1924

CHEIRANTHUS CHEIRI Linn.

Heart active glucosides of Cheiranthus cheiri Jaretzky R and Wilke M , *Arch Pharm* 270, 81, 1932.

CHENOPODIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Oleum Chenopodii Caines, J.F & Mhaskar, K S , *Ind Jour Med Res* , 1920, 7, 570.

Indian Chenopodium Chopra, R N., & Mukerjee, B , *Ind Med Gaz* 1932, 67, 5

A comparative record of anthelmintic treatment with tetrachloroethylene and oil of Chenopodium Manson, D , *Ind Med Gaz* , 1934, 69, 500

Pharmacological study of Chenopodium oil M Aissi Manicini and Leonardo Donatelli, *Chem Abst* 4451, 1935.

Pharmacological investigation on Chenopodium oil. Leonardo Donatelli, *Chem Abst* 4217, 1936.

The comparative value of oil of Chenopodium and tetrachloroethylene as anthelmintic for use in mass treatment. Hare, K P., Dutta, S C , *Ind Med Gaz* , 1939, 74, 198

A short note on the mass anthelmintic treatment by oil of Chenopodium and tetrachloroethylene Majumdar, D C , *Ind Med Gaz* , 1940, 75, 652

Pharmacology of chenopodium oil Oelkers H A , *Arch Exptl Path. Pharmakol* 195, 315, 1940

Effect of heat and light on the Ascaridol content in oil of Chenopodium Mukerji, A R , & Sen Gupta, K K , *Ind Med Gaz.*, 1945, 80, 347

C ALBUM Linn.

A phytochemical study of the fruits of Chenopodium album Tonuta J and Kaufmann K.L , *Pharm Arch* 12, 27, 1941.

C AMBROSIOIDES Hance

Saponin and sapogenin of Chenopodium ambrosiodes L, O. Dafert, Bauer, F Baur, M Capesius V and Grifinger S , *Chem Abst* 6937, 1934.

Studies on saponins of Chenopodium ambrosioides, Grifinger L, S , *Chem. Abst* 4839, 1937.

Investigation of the oil from Chenopodium ambrosioides L. Var anthelminticum Gray. cultivated in the gardens of medicinal plants of the Stepan Botany Univ. in Wilno. Tadensz Burchacinshi; *Chem. Abst* 8916, 1939.

A preliminary investigation of the yield and composition of the oil distilled from Chenopodium ambrosioides L. Var anthelminticum Gray. R.E Shapter, *J. Coun Sci Ind Res.* (Australia) 14, 201, 1941.

Culture tests on Chenopodium ambrosioides L. Var anthelminicum. The value and determination of the volatile oil. Schmonthuin, A & Schmied, E, *Chem. Abst.* 216, 1944

CHLOROXYLON SWIETENIA DC

Alkaloid isolated from Chloroxylon swietenia, constitution of. Mookerjee & Basu, *J. Ind Chem Soc.* 23 (1946), 1.

Chloroxylon swietenia. Alkaloid from the bark of. Mookerjee & Bose, *J. Ind Chem Soc* 23 (1946)

CHRYSANTHEMUM CINERARIIFOLIUM VIS.

Insecticidal principle in Chrysanthemum cinerariifolium. Insecticidal principle produced on dry distiliation of Chrysanthemum cinerariifolium Ryo Yamamoto and Mizuho Sumi H.; *Chem Soc Japan*, 44, 1070, 1923

Pyrethrum and Derris insecticides as arsenical substitute Roy Huston, *Chem Abst* 6234, 1934

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects Clyde C. Hamilton and Lousie G Gemelle, *J Econ Entmol.* 27, 446, 1934.

Determination of Pyrethrum I in commercial insecticides containing Pyrethrum and pyrethrum extract D A. Holaday, *Chem Abst* 3078, 1938

Pyrethrum cultivation in Kashmir Fotidar, M R, *Curr Sci*, 1940, 9, 360

Treatment of Scabies and Pediculosis with Pyrethrum Roy, D N, Ghosh, S.N & Chopra, R N, *Ind Med Gaz*, 1941, 76, 333.

Pyrethrum cultivation in Kashmir (*Med & Public Health Sec*), *Sci Cultr.*, 1941, 7, 217

Further work on Pyrethrum in the treatment of Pediculosis. Roy, D N. & Ghosh, S M, *Ind Med Gaz.*, 1942, 77, 480.

A water emulsion of pyrethrum extract for spray killing adult mosquitoes Russel, P P, Knipe, F W, *Ind. Med. Gaz.*, 1942, 77, 477.

Pyrethrin content of Indian Pyrethrum Puntambekar, S.V., *Curr. Sci.*, 1943, 12, 232

Pyrethrum in Kashmir Chopra, I.C., Dhar M.L, Handa, K.L. Habibullah, M. & Asa Nand, *Curr. Sci*, 1945, 14, 104.

Carotenoid pigments of Pyrethrum flowers Subbaratnan, A.V., & Parameshwaram Pillay, P, *J Sci Indl Res.*, 1947, **6B**, 100.

Pharmacognostic studies on homegrown Pyrethrum flowers Mitra, G C.; *Ind Jour. Pharm*, 1950, 12, 100

Cultivation of Pyrethum in Kashmir, Chopra R.N., Kapoor L.D., Handa K.L. and Chopra I C *Ind. Farming* Vol. VIII 78, 1947

CICER ARIETINUM Linn.

γ Galactan contained in the seeds of Cicer arietinum and its molecular constitution N Castoro, *Chem Abst.* 2642, 1925.

The food value of Cicer arietinum. V Zagami, *Chem Abst.* 3764, 1934

CICHORIUM INTYBUS Linn

Cichorium Intybus Linn Chemical examination of the seeds of. Constituents of the oil from the seeds Misra, Ramnath & Dutt Sikhibhushan, *J Ind Chem Soc* 14 (1937), 141

CICUTA VIROSA Linn

The toxin in Cicuta virosa E. Svagr. *Chem Abst* 407, 1924.

A case of the poisoning of six boys by the root nodules of Cicuta virosa V M. Karasek, *Chem. Abst* 7418, 1940.

CIMICIFUGA RACEMOSA Nutt

Constitution of the rhizomes of Cimicifuga racemosa Fern Mercier and J Balansaid, *Comp. Rend Soc Biol* 118, 166, 1935

CINCHONA Linn.

The relative value in malaria of the Cinchona alkaloids Quinine, Cinchonine, Quinidine, Cinchonidine & Quindoidine and the two derivatives Hydro-quinine and Ethyl Hydro-Cuprine MacGilchrist, A C , *Ind Jour. Med Res* , 1915, 3, 1.

On the behaviour of Paramecium caudatum towards the Cinchona alkaloids Acton, H W , *Ind Jour. Med Res* , 1921, 9, 339

Comparative action of the principle alkaloids on the dog heart (quinine, quinidine, Cinchonine, (Cinchonidine) A Elerc, C Pezzi and G Perro Gahaud, *Chem Abst* 293, 1924

Pharmacological notes on the principal alkaloids of Cinchona. L. Barthe, *Chem Abst* 700, 1925

A comparative study of the action of Cinchona alkaloids on the uterus Chopra, R N , David, J C & Dikshit, B B., *Ind Jour. Med. Res* , 1928, 15, 571

The cultivation of Cinchona in India A K Y , *Curr Sci.*, 1939, 8, 435

Cinchona plant. Bal, S.N , *Sci & Cultr* , 1939, 4, 617

Bengal Cinchona Plantation (*Med & Public Health Sec*), *Sci. & Cultr.*, 1941, 7, 300

Cinchona cultivation in India—Sen, S.C., *Curr Sci* , 1941, 10, 223

Quinine and Alstonia scholaris (CHHATM) in Malaria. Das Gupta, B.M., Siddons, L.B & Chakaravarti, H., *Ind Med. Gaz.*, 1943, 79, 408.

Cinchona, pharmacognostic studies on Datta S C, & Bal S N, *Ind. Jour Pharm*, 1946, 8, 85

Pharmacognostic studies on Indian Cinchona Datta, S.C., & Bal, S N, *Sci & Cult*, 1946, 12 246

Influence of temperature on growth and alkaloidal content of Cinchona seedlings Harold F Winters, Arnaud, J Loustalalot and Norman F Childens, *Chem Abst* 2126, 1947

Waste Cinchona materials for production of Quinine Mukerjee, S, *Sci & Cultr* 1949, 13, 372

Cinchona alkaloids of newly cultivated plants of Assam. Mukerjee, S, *Ind Jour Pharm*, 1949, 10, 107

Quinine from Cinchona bark extractions Pt I & II Bhunvara, N.B, & Khorana, M L, *Ind Jour Pharm*, 1949 11 Pt I, 148 and Pt II 152

Structure of Quinamine and Cinchonamine A note on the Str of Quinamine and Cinchonamine, two minor alkds of Cinchona Tyabji, A, *Ind Jour Pharm*, 1950, 12, 243

Cinchona finds a new home Chatterjee, R, *Sci & Cultr*, 1950, 15, 339.

C FEBRIFUGA

Cinchona febrifuga in the treatment of malaria Gore, R N, *Ind Med Gaz.*, 1938, 73, 608

Soluble salts of "Cinchona febrifuga" as a cheap and effective antimalarial drug Chatterjee, R P, *Sci. & Cultr* 1952, 17, 524

C LEDGERIANA Moens ex Trimen

Vegetative propogation of Cinchona ledgeriana from gootes (Marcotte) and cuttings by treatment with auxins Chakutra, A G, *Sci & Cult*, 1944, 9, 409

CINNAMOMUM Bl.

Cinnamon leaf oil Krishna, S, Kamath, H R., Kudva, K T & Kudva, K G, *J Sci Indl Res*, 1945-46, 4, 464

C CAMPHORA Nees.

Note on the possibilities of camphor cultivation from Cinnamomum camphora in Northern India. Howard S H, Robertson W.A. and Simonsen J L, *Indian Forest Records*, 1923 Vol IX, Part VII, pp 34.

Notes on some Indian Essential Oils Cinnamomum camphora, Nees & Eberm (Syn Camphora officinarum, Baugh, Laurus Camphora, Linn) Rao B Sanjiva, Sudborough J J and Watson H E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. **8A**, Part X, pp 160-174

Camphor seed oil. Hao Chen, *Ind Res. (China)* 6, 92, 1937.

Constituents of Camphor oil, Acids and Phenols in the brown camphor oil, Kashichi Ono and Monoru Imoto, *J. Chem Soc Japan*, 58, 531, 1937

C JAPONICUM Siebold.

Fat of the seeds Cinnamomum japonicum, as a substitute for Cacoa butter. Talsoo Karyone and H Iwao, *J Pharm Sco Japan*, 58, 238, 1938.

C ZEYLANICUM Nees.

Essential oil from the leaves of Cinnamomum zeylanicum V P Shintue and B. Sanjiv Rao, *J. Ind Inst Sci* **15A**, 84, 1932

Studies in India Essential Oils X-Essential Oil from the leaves of Cinnamomum zeylanicum, Breyn Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1932, Vol **15A**, Part VII, pp 84-87.

Cinamon-bark oil of the Scychelles (Cinnamomum zeylanicum) W Holdsworth Haines, *Perfumery Essential Oil Records*, 27. 6, 1936.

CISSAMPELOS PAREIRA Linn

Chemical Examination of the roots of Cissampelos pareira Linn Bhattacharji, S , Sharma, V N & Dhar, M L , *J Sci Indl Res* , 1952, **11B**, 81

A preliminary note on the pharmacological action of the total alkaloids isolated from Cissampelos pareira Linn Roy, P K., Dutta, A T , Roy, G K , Mukerjee, B , *Ind J Med Res* , 1952, 40, 95.

CITRULLUS COLOCYNTHIS Schrad

A preliminary note on the chemical examination of the roots of Citrullus colocynthis Scharader. Agarwal & Dutta, S , *Curr Sci* , 1934-35, 3, 250

Citrulus colocynthis Examination of the oil from the seeds of Alimchandani, Badami & Katli, *J Ind Chem. Soc* 26 (1949) 515 & 519

C VULGARIS Schrad

Citrullin a new amino acid in press juice of the Watermelon, Citrullus vulgaris Schard. Mitsunori Wada, *Chem Abst* 5335,1930

Proteins and other nitrogeneous constituents of Watermelon seeds (Citrullus vulgaris) P S Krishnan and T K Krishnaswamy *Bioch J* 33, 1284, 1939

Characteristics and composition of Watermelon seed oil Arthur J. Nolte and Harry W. Von Loeseke; *J Am Chem Soc* 61, 869, 1939

Extracting the therapeutic principle from various parts of Watermelon fruit or plant. Mathew Evertz; *Chem Abst.* 5257, 1941.

The diuretic action of Watermelons I T Isawa, Y Takahasi and T. Togo, *Chem Abst* 521, 1941

Biological value of proteins in Watermelon and Pumpkin seeds Tsoo E King, *Chinese J. Physio* 16, 31, 1941

Cucurbit-seed glabulins, use as substitute for edestin in experimental diets, (Citrullus vulgaris) Hubull R B . Vockery H B and Nolan, L S *J Nutrition.* 25, 99, 1943

Citrullus vulgaris Component fatty acids of the oil from the seeds of Dhingra & Biswas, *J Ind Chem Soc* 22 (1945), 119

Melon seed oil Safet Riz Alpar and Suat Esin; *Chem Abst* 9700, 1950

CITRUS Linn

Preservation of citrus fruit juices Jiwan Singh Pruthi & Girdhari Lal, *J Sci Indl Res* , 1951, **10B**, 36

Nagpur Oranges—(*Proc Nat Acad Sc* Vol I, 1939, pp 175-180)

C ACIDA Roxb

Xanthyletin Isolation of. from Citrus acida bark Mookerjee, *J Ind Chem Soc* 23 (1946), 41

Citrus acida Roxb Natural coumarins isolated from the leaves of Khastagir, *J Ind Chem Soc* 24 (1947), 421-34.

C AURANTIUM Linn

Preparation of Citric Acid from Citrus aurantium or Karana Khatta Dhingra, D R., Gupta, G.N., Nigam, V.N , & Raghvendra Singh, *Ind Jour Pharm.* 1950, 12, 351

C BIGARADIA Loisel.

Notes on some Indian Essential Oils Citrus bigaradia, Risso Rao, B Sanjiva, Sudborough, J J and Watson, H E , *Journal of the Indian Institute of Science, Bangalore,* 1925 Vol **8A**, Part X, pp 176-179

C. DECUMANA Linn.

Citrus decumana On the bitter principles of Miss Asima Mukerjee, *J Ind Chem Soc* 17 (1940), 593

Ascorbic acid of Shaddock (Citrus decumana) availability of by human subjects. De & Barai, *J. Ind. Chem Soc.* 25 (1948), 389

C. LIMON Burm f

Survey of the possibilities of manufacturing Citric Acid from Lemons in Bihar and Bengal. Ghatak, N. & Karimullah, *J., Sci Indl Res* , 1944-45, 3, 110.

A study of species of Citrus limonis. Thomas Oomen, P , *Ind. Jour Pharm* , 1951, 13, 120.

C POONENSIS Hort ex Tanaka

Carica xanthin of the papaya and Citrus poonensis Hort Ryo, Yamamoto and Akimasa Kato, *J Agr Chem Soc , Japan*, 10, 754, 1934

Carica xanthin, a colouring matter in the fruit pulp of Carica papaya and Citrus poonenis Ryo Yamamoto and Yosiakin Kato *Chem Abst* 806, 1935

CLAVICEPS PURPUREA (Fries) Tulasne

The therapeutic activity of liquid preparations of Ergot of the Calcutta market Chopra, R N & De, Premenkur, *Ind Med Gaz*, 1928, 63, 519

Ergot in India Padwick, G Watts *Curr Sci* , 1941, 10, 488

Can the deterioration of Ergot extracts be prevented in the tropics Bose, I B & Dey, N K , *Ind Med Gaz* , 1942, 77, 286

Germination of Ergot Bose, A B , *Curr Sci* , 1942, 11, 439

Ergot in Indian Kulkarni G S , *Curr Sci* 1942, 11, 246

Indian medicinal importance of Ergot and need for its cultivation Mukerji, B , Bose, A B , *Sci & Cultr* , 1942-43, 8, 267, 348

Claviceps purpurea (Fr) Tul and a new species from India Watts Padwick, G & Azmatullah, *Curr Sci* , 1943, 12, 257

Ergot on Sugarcane in Mysore Thirumalachar, M J , *Curr Sci* 1943, 12, 330

Assay of Indian Ergot Mukerji, B. & Dey, N K., *Curr Sci* 1943, 12, 58

A method for the assay of individual Ergot sclerotium Mukerji, B & De, N K , *Curr Sci* 1944, 13, 128

Ergot on Cynodon dactylon Pers Thirumalachar, M J , *Curr Sci* , 1944, 13, 288

Ergot and sphacelial stages on some wild grasses in Mysore Thirumalachar, M J , *Curr Sci* , 1945, 14, 22.

Ergot sclerotia on Sorghum vulgare Pers Ramakrishnan, T S , *Curr Sci* , 1948, 17, 218

Ergot on bamboo Ramakrishnan, T S & Ramakrishnan, K , *Curr. Sci* , 1949, 18, 344.

Ergot on two grasses from South India Ramakrishnan, T S & Sundaram, N V , *Sci & Cultr* , 1950, 16, 214

CLAUSENA WILLDENOWII Wight & Arn

Oil of Clausena willdenowii—B S Rao and K S Sobramanian, *Proc. Indian Acad Sci.* **1A**, 1934, 189.

Oil of Clausena willdenowii—B S. Rao and K S. Subramanian, *Proc. Indian Acad Sci. Sect.* A, 3, 1936.

Chemical investigation on some Minor Forest Products. The Essential Oil from Karibe, Clausena willdenowii, leaves and twigs Rao K Narain, Varadhan C and Janniah S L, *Department of Industries, Bangalore*

Constitution of L Clausenan (Clausena willdenowii W & A.). Narsimha Rao, P L, *J. Sci Indl Res*, 1949, **7B**, 11

CLEOME VISCOSA Linn

Cleome viscosa Linn Chemical examination of the seeds of Part I. The constituents Gupta, Mahadeo Prasad & Dutt, Sikhibhushan, *J Ind Chem Soc* 15 (1938), 532

CLERODENDRON INFORTUNATUM Gaertn

Constitution of clerodin, the active bitter principle of Clerodendron infortunatum H N Banerjee, *Trans Bose Research Inst, Calcutta*, 12, 75, 1936-37

Clerodendron infortunatum, chemical examination of. Part I. M B Rane & Kondaiah K, *J Ind Chem Soc* 14 (1937), 46.

Clerodendron infortunatum Active principle Clerodin of Chaudhry & Dutt, *J Ind Chem Soc* 28 (1951), 295

COCCULUS D.C

Sinomenium and Cocculus alkaloids Constitution of Cepharanthine Heisaburo Kondo and Ichiro Keimatu, *Ber* **71B**, 2553, 1938

C. LAURIFOLIUS D C

Alkaloid of Sinomenium and Cocculus Alkaloid of Cocculus laurifolius H Mondo and T Kondo, *J Pharm Soc. Japan* 524, 876, 1925

C SARMENTOSUS Diels

The alkaloids from Sinomenium and Cocculus The alkaloids of Cocculus sarmentosus Diels Constitution of menisavine Heisaburo Kondo and Masao Tomita, *J Pharm Soc Japan*, 55, 637, 1935

COCHLEARIA ARMORACIA Linn

The content of Lepidium latifolium and Cochlearia armoracia, in essential oils S Rivos Goday and M. Gomez, serranillos, *Chem Abst.* 4432, 1945

COCOS NUCIFERA Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency. Santonion, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb , Melia azadirachta Linn., Punica granatum Linn., Picrasma excelsa Swartz , Vernonia anthelmintica Willd , Cocos nucifera Linn. (coconut). Caines, J F. & Mhaskar, K.S., *Ind Jour Med Res*, 1923, 11, 353.

Chemistry of the products of Cocos nucifera. Juan P.C. Chandersena, *Bioch J* 24, 1493, 1930

Antirachitic effect of Coconut oil and its mode of action on bone Calcification Nimal C Dutt, *Ann Biochem Exptl Med.* 8, 69, 1948.

COFFEA Linn

The chemistry of Coffee bean The unsaponifiable matter of Coffee bean oil Preparation and properties Kahweol, R O. Bengis and R J Anderson, *J Bioch Chem* 97, 99, 1932

Coffee black bean Venkalarayan, S V , *Curr Sci* , 1938, 7, 236.

Chemistry of Coffee Elucidation of the constitution of Cafestrol, Carl H Slotta and Klas Neissar, *Ber* **71B**, 2342, 1938

Chemistry of Coffee A new method for the determination of trignolline Karl Heinrich Slotta and Klaus Neisser, *Ber* **17B**, 1987, 1938

Chemistry of Coffee, recent analytical findings Karl H. Slotta and Klaus Neisser, *Chem Abst* 2604, 1939

A coffee substitute Cassia occidentalis that is toxic before roasting. P Bruni, *J Pharm Chem* 20, 321, 1942

Coffee oil K H Bauer and R Beu, *Chem Abst* 5607, 1943.

Determination of Coffee in coffee mixtures and substitute by the Tillmanns—Hollatz method Jean Deshuses *Chem Abst* 3033, 1944.

C ROBUSTA Linn

Importance of reserve food material in successful establishment of shoot cutting in Coffea robusta L Rallabhiraman, T V & Gopalkrishnan, K S , *Curr Sci* 1947, 16, 385

COLCHICUM Linn

Colchicine Bose, A., *Sci & Cultr* 1940, 5, 511

COMBRETUM PILOSUM Roxb

Combretum pilosum Roxb as an anthelmintic for Ascaris lumbricoides Ramsay, G C , *Ind Med Gaz* , 1922, 57, 374.

COMMELINA ASIATICA Linn

Chemical Examination of the oil from the seeds of Commelina asiatica Linn Aggarwal, J S & Miss Padmini Soni, *J Sci. Indl. Res* 1949, **8B**, 49

CONVOLVULUS HAMADAE (Vved) Petrov.

Alkaloids from Convolvulus hamadae G V Lazurevskii; *Chem Abst.* 4029, 1941.

C PLURICANTES Chois.

On an abnormal flower of Convolvulus pluricantes Chois. Rao, A R ; *Curr Sci.*, 1935-36, 4, 441.

COPTIS TEETA Wall

The constituents of Coptis teeta Wall Chatterjee, A , *Sci & Cultr* , 1950, 15, 330

Alkaloid from Coptis teeta Wall Saw, P L Seshadri, T R , *J Sci Indl. Res* , 1952, **11B**, 308

Plant alkaloids Part II Coptis teeta Wall Chatterji, R Guha, M.P. & Chatterjee, A , *J Ind Chem Soc* 29 (1952), 97.

CORCHORUS Linn

Corchortoxin a cardiac agent from jute seeds P Karrer and P Banerjee, *Helv Chem Acta* 32, 2385, 1949

C OLITORIUS Linn

Corchorus olitorius Composition of the oil from Chakravarti & Sen, *J Ind Chem Soc* 20 (1951), 790

CORIANDRUM SATIVUM Linn

Notes on some Indian Essential oils Coriandrum sativum Linn Rao B Sanjiva, Sudborough J J and Watson H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol **8A**, Part X, pp 179-182

Report on the examination of Indian and foreign coriander (Coriandrum sativum) B. Vishwanath and C V Ramaswamy Ayyar, *Ag Livestock India*, 4, 583, 1934

The fatty and the essential oils from Coriandrum sativum grown in the Northern Caucasus I I Vanin and A A Chernoyarova, *Chem Abst* 3833, 1934

The nitrogen complex of Indian foodstuffs Condiments Pt II Chillies (Capsicum annum) Coriander seeds (Coriandrum sativum) Narasimhamurthy, G , *Ind Jour Med Res* , 1938, 25, 863

Production of faboil from the Coriander seed waste at the Aleksevskii plant F Tansue Sienko, *Chem Abst* 5649, 1938

Production of decaldelhyde from Coriander P P Shorygin and V P Osipova, *Chem Abst* 3781, 1942

Decaldehyde from the oil of blooming Coriander Ya Bryusova Ryu Shagolova and N Novikova, *Chem Abst* 3781, 1942

Production of citral from Coriander oil V I Isagulyants and E K Smolyani *Chem Abst* 3782 1942

COSCINIUM FENESTRATUM Colebr

A note on the alkaloids of Coscinium fenestratum (Colebr) Varier, N S & Pillai, P P , *Curr Sci* 1943, 12, 228

The berberine content of Coscinium fenestratum (Colebr.) Child, R., Narhanael, W.R.N , *Curr Sci* , 1943, 12, 255

CRATAEGUS OXYCANTHA Linn.

Constituents of Crataegus oxycantha H Dieterle and O Dorner; *Arch Pharm.* 275, 428, 1937

Investigation of Crataegus oxycantha Thomas E Pugh, *Chem. Abst* 3462, 1939

A pharmacological study of Crataegus tincture I I Sivertsev, *Chem Abst* 6463, 1939

Pharmacological action of Crataegus oxycantha J D P Graham, *Quart Jour Pharm Pharmacol* 13, 49, 1940

CRATAEVA NURVALA Ham

Chemical constituents of the root bark of Crataeva nurvala Ham. Bhandari, P R , Dhar, M L , Shafma, V N , *J Sci Indl Res* , 2951, **10B**, 105

CRINUM ASIATICUM Linn

Alkaloids of Crinum asiaticum L Var Japonicum, Bak Kinzi Tanaka, *J Pharm Soc , Japan* 57, 652, 1937

CROCUS SATIVUS

Saffron, colouring matter Paul Karner and Harry Salomon, *Chem. Acta* 11, 711, 1928

Picrocrocin the bitter principle of Safran H E W Lutz, *Chem Abst.* 110, 1931

On the possibility of cultivation of Safron (Crocus sativus) in the Hyderabad state and its importance Inam-Ul-Haq & Sayeeduddin, M , *Cur Sci.*, I, (1932-33), 394

Picrocrocin, the terpene glucoside of Safron and the biogenesis of the Carotenoid Carboxylic acids Richard Kuhn and Alfred Winterstein, *Chem Abst* 161, 1934

Kashmir Saffaron with methods of testing its purity K L Budhiraj, *J Ind Chemical Soc Ind and News Ed* 5, 135 1942

CROTALARIA JUNCEA Linn

A note on the occurrence of Tri-cotyledonary seedlings in Crotalaria juncea Linn Purkayastha, K.K *Curr Sci* , 1941, 10, 30

C MEDICAGINEA Lam

A note on breaking dormancy in Crotalaria medicaginea, Lamb. Satyanarayana Rao, N *Sci & Cultr* , 1947, 12, 503

C SPECTABILIS Roth

Isolation and some properties of an alkaloid from Crotalaria spectabilis Roth. W M Neal, L L Rusoff and C F Ahmann, *J Am. Chem Soc* 57, 2560, 1935

Crotalaria spectabilis poisoning in Lorisiana livestock P O. Piercy and L L Rusoff, *J Am Vet Med. Assoc* 108, 69, 1946

Livestock poisoning by Crotalaria spectabilis. P.L Piercy and L.L. Rusoff, *Chem Abst* 6681, 1946.

CROTON TIGLIUM Linn

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XVIII Catharatics, Oleum Ricini, Oleum Tiglii, Aloe, Succies acalyphae Caines, J F., Mhaskar, K S , *Ind Jour Med Res.*, 1923, 11, 103

Poison of Croton oil Isolation of Croton resin, thin oil and Phorbol from Croton oil by alcoholysis Bonifaz Flachentrager and George Wigner, *Helv Chem. Acta* 25, 569, 1942

CRYPTOSTEGIA GRANDIFLORA R Br

Extraction of rubber from Cryptostegia grandiflora Bhatnagar, S S , Karimullah & Uma Shankar, *J Sci Indl Res* , 1944-45, 3, 441

Extraction of rubber from Cryptostegia grandiflora Bhatnagar S S , Karimullah & Uma Shankar, *J Sci. Indl Res* , 1944-45, 3, 263

Cryptostegia grandiflora R Br A war time source of vegetable rubber, Pt VIII Examination of seeds and the fatty oil. Siddiqui, R H & Warsi, S A , *Ind Jour Pharm* 1945, 7, 75

Cryptostegia grandiflora R Br Siddiqui, R H & Warsi, S A.; *Ind Jour Pharm* , 1945, 7, 117

Rubber from Cryptostegia grandiflora Bhatnagar, S S , Karimullah & Uma Shankar, *J Sci Indl Res* , 1945-46, 4, 654

CUBEBA OFFICINALIS Rafin

Notes on some Indian Essential Oils Cubeba officinalis Misc (Syn Piper cubeba, Linn) Rao B Sanjiva, Sudborough J J and Watson H E *Journal of the Indian Institute of Science, Bangalore,* 1925, Vol **8A**, Part X, pp 159-160

CUCUMIS STATIVUS Linn

A proteolytic enzyme in cucumber (Cucumis sativus) Chopra, R N & Roy, A C , *Ind Jour. Med Res* , 1933, 21, 17

Chemical examination of the seeds of Cucumis sativus Linn Miss Padmini Soni, Gupta, S C & Aggarwal J S , *J Sci Indl Res* 1949, **8B**, 210

C UTILISSIMUS Roxb

Chemical examination of the seeds of Cucumis utilissimus Roxb Bhasin, M M , Gupta, A S & Aggarwal, J S , *J Sci Indl Res* , 1950 **9B**, 230

CUCURBITA PEPO D.C

Examination of the pumpkin oil from Pumpkin seeds The constitution of obec acid. J.L Rubsomer and G A. Nesty, *J Am. Chem. Soc* 56, 1784, 1934.

Biological value of proteins in Watermelon and Pumpkin seeds. Tsoo E King, *Chinese J Physio.* 16, 31, 1941.

The anthelmintic power of Pumpkin seeds Giuseppe Sanifilippo, *Chem Abst* 1346, 193, 32.

CUCURBITACEAE

Nomenclature changes of some common plants of the Cucurbitaceae Chakravarty, H L, *Sci & Cult*, 1949, 15, 31

CUMINUM CYMINUM Linn

Notes on some Indian Essential oils Cuminum cyminum, Linn Rao B Sanjiva, Sudborough J J and Watson H E, *Journal of the Indian Institute of Science, Bangalore*, 1923, Vol **8A**, Part X, pp. 182-183

CUPRESSUS TORULOSA Don

The constituents of some Indian Essential Oils Part XI—The Essential Oil from the leaves of Cupressus torulosa Don Simonsen J L, *Indian Forest Records*, 1922, Vol X, Part I

CURCUMA Linn

Active constituents of Curcuma E Franquela, *Chem Abst* 3833, 1934

C AMADA Roxb

Curcuma amada—S Dutt and J N Tayal, *Soap Journ.* Feb 1941, p 200

Chemical examination of the essential oil derived from the rhizomes of Curcuma amada Roxb S Dutta and Jagat Narayan Tayal, *Ind Soap J* 7, 200, 1941

C AROMATICA Salisb

Constituents of some Indian essential oils Curcuma aromatica, Puthan Madhathel, Bhaskara Panicker, B Sanjiva Rao and J L Simonson, *J Ind Inst Sci* **9A**, 133, 1926

Constituents of some Indian Essential Oils Part XX - Essential Oil from the Rhizomes of Curcuma aromatica Salisb Rao B Sanjiva, Shintre V P and Simonsen, J L, *Journal of the Indian Institute of Science, Bangalore*, 1926, Vol 9A, Part VI, pp 140-144

C CAESIA Roxb

Curcuma caesia—*Proc Nat Acad Sc* Vol V, 1940, pp 64-68

C DOMESTICA Valeton

Constituents of the rhizomes of Curcuma domestica H Dieterle and Ph Kaiser, *Arch Pharm* 217, 337

C LONGA Linn

Studies in Indian Essential oils V-Essential oil from the Rhizomes of Curcuma longa, Linn Kelker N C and Rao B Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1933, Vol **17A**, Part II, pp 7-24.

Essential oil from the rhizomes of Curcuna longa L N C Kelkar & B Sanjiva Rao, *J Ind Inst Sci* 17, 24, 1933

Indian essential oils Essential oil from the rhizomes of Curcuma longa N C Kelkar and B Sanjiva Rao, *J Ind Inst. Sci* **17A**, 7 1933

Volatile plant constituents—Tumerone, the aromatic principle of Turmeric oil H Hupe Georges Clar Alexander Sl Pfau and Pl Plattner, *Helv Chim Acta* 17, 372 1934

Turmeric and vegetable oil as repellents against Anopheline mosquitoes Philip, M I , Ramkrishna, V & Rao, V V , *Ind Med Gaz* 1945, 80, 343

Pharmacological action of essential oil of Curcuma longa, Chopra, R N , Gupta, J C & Chopra, G S , *Ind Jour Med Res* 1941, 29, 769

C ZEDOARIA Rose

Note on some Indian Essential Oils Curcuma Zedoaria, Roscoe Rao B Sanjiva, Sudborough J J and Watson H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol **8A**, Part X, pp 153-155

The constituents of some Indian Essential Oils Part XXIV—The Essential Oil from the rhizomes of Curcuma Zedoaria, Roscoe Rao B Sanjiva, Shintre V P and Simonsen J L , *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol **IIA**, Part XV—I, pp 195-199

CUSCUTA REFLEXA Roxb

Chemical examination of Cuscuta reflexa Roxb Radha Raman Agarwal and Sikhibhushan Dutt, *Ind J Chem Soc* 1935, 12, 384

Cuscuta reflexa Roxb Chemical examination of Part II The constitution of cuscutalin Agarwal, Radha Raman & Dutt, Sikhibhushan *J Ind Chem Soc* 12, (1935), 586

Cuscuta reflexa Chemical examination of Part III The constitution of the oil from the seeds Agarwal Radha Raman & Dutt, Sikhibhushan, *J Ind Chem Soc* 13 (1936), 264

Cuscuta reflexa Roxb Chemical examination of Part IV Isolation of a new yellow flavone, colouring matter from the seeds. Agarwal, Radha Raman, *J Ind Chem Soc* 13 (1936), 531

Parasitism of Cuscuta reflexa Roxb & Loranthus longiflorus Desr Sheriar, K C , *Sci & Cult* , 1951, 17, 218

On Cuscuta reflexa Roxb in the Khasi Hills Kachroo, P , *Sci & Cult* , 1951, 16, 431

CYCAS REVOLUTA Thunb

Constituent of seed oil of Cycas revoluta Seuchi Ueno, Sumio Matsuda and Taka Kimura, *Chem Abst* 5618, 1950.

CYDONIA VULGARIS Pers.

The fatty oil of Quince seed, Cydonia vulgaris A Steger and J Van Loon, *Chem Abst* 2207, 1934

CYMBOPOGON Hack

Cymbopogon Grasses—B S Rao and J J Sudborough, *Journal of the Indian Institute of Science, Bangalore*, Vol **8A**, Part II, pp 9 to 27, 1925

Oil of a new Cymbopogon species *Bull Imp Inst* 27, 1929, 459

Studies in Indian Essential oils Essential oil from the flower heads of Cymbopogon, Kotnis M S and Rao B. Sanjiva, *Journal of the Indian Institute of Science, Bangalore*, 1935, Vol. **18A**, Part XVIII, pp 129-133

Cultivation of Cymbopogon grass in Trans-Gangetic tracts, J N Rakshit, *Rakshit Gardens Bulletin No 1, Ghazipur, U.P.* 1936.

Development of Essential Oil Industry Ocimum lemon, O basilicum, O canum, O sanctum, Lemon grass, Motia, Palmarosa grass Jitandra Nath Rikshit, *Sci & Cult*, 1939, 5, 108

C CAESIUS Stapf

Travancore Essential Oils Inchi grass Cymbopogon caesius Stapf. Moudgill K L and Aiyar K R Krishna, *Department of Industries, Trivandrum*, 1923, *Bulletin* No XVII, pp 18-25

Travancore Essential Oils Part VI—The Essential oil from Cymbopogon caesius, Stapf Inchi grass Moudgill K L., *Journal of the Indian Chemical Society, Calcutta*, 1925, Vol II

Kachi grass oil (Cymbopogon caesius). R B Rao and J J Sudborough, *J Ind Inst Sci* **8A**, 9, 1925

Essential oil from Cymbopogon caesius Stapf Moudgill, *J Ind Chem Soc* 2 (1925), 23

C CITRATUS Stapf

Distillation tests on lemon grass, Cymbopogon citratus R Wilbaux; *Chem Abst* 3636, 1937

Essential Oils Lemon Grass Oil. *Department of Industries, Trivandrum, Bulletin* No IV.

C CLANDESTINUS Stapf

Oil of Cymbopogon clandestinus *Bulletin Imp. Inst.* 27, 1929 438.

C COLORATUS Stapf

The constituents of some Indian Essential Oils Part XVII—Essential Oil from the flower heads of Cymbopogon coloratus, Stapf Pillai P Parameswaran, Rao B Sanjiva and Simonsen J L.; *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol **11A**, Part XV—1, pp 181-186

C. MARTINI Stapf

Note on the economic uses of Rosha Grass (Cymbopogon martini, Stapf)—R.S Pearson, *Indian Forest Records*, 1916, Vol. V Part VII.

Some observations on the Essential oil content of Rosha Grass (Cymbopogon martini var. Motia) Girdhari Lall, *The Indian Journal of Agricultural Science*, Vol V, Part III, 1935

CYNOMARATHRUM NUTTALLII A Gray

Volatile oil and resin of Cynomarathrum nuttallii A Gray (or Pencedanum graveolens) E K Nelson, *J Am Chem Soc* 55, 3400, 1933.

CYPERUS ESCULENTUS Linn

Oil value of Cyperus esculentus tubers J Pieraerts, *Chem Abst* 1917, 1924

C ROTUNDUS Linn

Essential oil of Cyperus rotundus B S Rao, P.B Panicker and J J. Sudborough; *J Ind Inst Sci* **8A**, 39, 1925

Essential oil of root of Cyperus rotundus L of Japan Y Kumura and M Ohtani; *J Pharm Soc Japan*, 48, 971, 1928

Oil of Cyperus rotundus—B J Hedge and B S. Rao, (*Journ Soc Chem Ind* 54, 1935, **T 388**)

Some of the constituents of the tuber of Coqui (Cyprus rotundus L) Preliminary examination of the tuber and composition of the fatty oil. Conrado F Asenjo, *J Am Pharm Assoc* 30, 216, 1941

C ROTUNDUS & C SCARIOSUS Br

Chemical investigation of Cyperus rotundus and Cyperus scariosus Basu N K, *Sci & Cultr*, 1944, 10, 131

Oil of Cyperiol (Cyprus scariosus) Gupta, G N & Ganesh Chandra, *Curr. Sci*, 1951, 20, 273

CYSTOPUS CANDIDUS

Medicinal action of Capsella bursa-pastoris and also of its parasites, Cystopus candidus and Peranospora parasitica. W Harste, *Arch Pharm* 206, 133, 1938

DAEMIA EXTENSA (Linn) R Br

Pharmacological action of an active constituent isolated from Daemia extensa Linn. (Syn Pergularia extensa) Pt I. Gupta, J.C., Roy, P K & Dutta, A, *Ind. Jour Med Res*, 1946, 34, 181

Daemia extensa R Br. Datta, A. & Ghosh, S., *Ind Jour. Pharm* 1947, 9, 58.

Chemical examination of Daemia extensa R. Br. Ashutosh Dutta and Sudhamoy Ghosh, *J Am. Pharm. Assoc.* 36, 250, 1947.

Pharmacological action of an active constituent isolated from Daemia extensa Linn (Syn. Pergularia extensa). Pt. II. *Ind. Jour Med Res*, 1950, 38, 75.

DATURA Linn

The alkaloid content of some Datura species L. S Dyson and J. A. Coetzee, *S African J. Sci* 40, 162, 1943

Cultivation studies of the Solanaceous drugs Post harvest alkaloidal movement in Belladonna and Datura species. W R. Brewer and L. David Hiner, *J Am Pharm Assoc* 37, 541, 1949

D ALBA Nees

Alkaloids of Datura alba Nees S Osada, *J Pharm. Soc. Japan* 504, 89, 1924

Isolation and separation of main constituents of Datura alba Nees Cheoa Sakul Pradisth and A C Santos, *Chem Abst* 5593, 1939

Alkaloids of Datura alba Nees Cheoa Sakul Pradisth and Alfrendo E Santos, *Chem Abst* 767, 1940.

Isolation and separation of the main constituents of Datura alba Nees Cheoa Sakul Pradisth and Alfrendo E Santos, *Chem. Abst.* 3441, 1940

Fertilizer effects on the growth and alkaloidal content of Datura alba Nees Prasad, S, *Ind Jour Pharm.* 1944, 6, 13.

A mosaic disease of Datura alba Nees, Capoor, S P. A & Varma, P M, *Curr Sci*, 1949, 17, 151

D FASTUOSA Linn

Datura metel L & D fastuosa L Narayanswami, V *Sci & Cult*, 1948, 14, 38

Datura metel L & Datura fastuosa L Chatterjee, D Narayanswami, V, *Sci & Cult*, 1948, 14, 206

D INOXIA Mill.

Datura inoxia a potential commercial source of Scopolamine Goe H Garlach, *Econ Botany*, 2, 436, 1948

D METEL Linn

Datura metel L. & D fastuosa L. Narayanswami, V, *Sci & Cult.* 1948, 14, 38

Datura metel L & Datura fastuosa L Chatterjee, D., Narayanswami V., *Sci & Cultr*, 1948, 14, 206

Datura metal Chatterjee, D., *Sci & Cult*, 1949, 15, 77.

D STRAMONIUM Linn (D. TATULA Linn.)

A study of Datura stramonium. Frank H. Eby, Frederick. M. Scholl and David H. Phillips, *J. Am. Pharm. Assoc.* 27, 474, 1938.

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves. Bernard S Feinstein and Frank J Slama, *J Am Pharm Assoc.* 29, 370, 1940

The effect of varying temp upon the total alkaloid content of folia stromonii and folia belladonna H Fluck, *Chem Abst.* 3878, 1940.

Synthesis of Hyoscyamine in Atropa belladonna and Datura stramonium B T. Cromwell, *Biochem Jour* 37, 717, 1944

Comparative analysis of normal and tetraploid Datura stramonium and Datura tatula. Ovrville H. Miller and Louis Fischer; *J Am Pharm Assoc.* 35, 23, 1946

DAUCUS CAROTA Linn

Essential oils of Carrot (Daucus carota) G Igolen, *Chem. Abst* 7281, 1936.

Oil of Carrot. Ernest S Guenther, *Am. Perfumer*, 5, 71, 1936.

The carotenoid pigments and the Vitamin 'A' activity of Indian carrot. Sadana, J C & Bashir Ahmad, *Ind Jour. Med. Res.* 1947, 35, 81.

DECALEPIS HAMILTONII W & A

Oil of Decalepis hamiltonii M. G Srinivasa Rao and M. Sesha Iyenger; *Perfumery Records* 14, 1923, 300

Chemical components of roots of Decalepis hamiltonii (Makali veru) Comparison with Hemidesmus indicus (Indian Sarsaparilla). P. Bhaskara Ramamurti and T R. Sheshadri, *Proc. Ind. Acad* Sci **13A**, 399, 1941.

DELPHINIUM AJACIS Linn.

Alkaloids of Delphinium ajacis. Mathew V Hunter, *Chemist & Druggist*, 139, 304, 1943

D. ELATUM Linn

Alkaloids of the seeds of Delphinium elatum L Jhon. A. Goodson; *J. Chem. Soc* 139, 1943.

DERRIS Lour.

Rotenone the active constituent of Derris root. Shankichi Takiee; *Ber.* **61B**, 1003, 1928

Constituents of Derris root Tubaic acid T. Karizone, K Kondo and Makdie; *J. Pharm Soc. Japan.* 48, 674, 1928.

Recent progress in the chemistry of Derris. R.C Roak, J C Con, *Entomol.* 22, 378, 1929

Rotenone the active principle of Derris root Sankichi Takel, Shakivo Mihajima and Minoru Ono, *Chem Abst* 5565, 1932

Pyrethrum and Derris insecticides as arsenical substitute. Roy. Huston; *Chem Abst.* 6234, 1934.

Some field tests showing the comparative efficiencies of Derris, Pyrethrum and Belladonna powders on different insects. Clyde C. Hamilton and Lousie G Gemelle, *J Econ. Entmol.* 27, 446, 1934.

Rotenone the non-crystalline constituent of Derris root. H L Hallerna, F B. Laforge, *J. Am Chem Soc* 56, 2415, 1934

D. ELLIPTICA Benth

Rotenone the physiologically active constituent of Derris elliptica. A. Buten, *Chem. Abst* 3660, 1928

Toxicological studies of Derris elliptica and its constituent Rotenone. H B Haag, *J. Pharmacol* 43, 193, 1931

Occurrence of Derris elliptica in India Krishna, S. & Ghosh, T.P., *Curr Sci*, 1935-36, 4, 857

D FERRUGINEA Benth.

Denis ferruginea from Assam Krishnan, S & Ghose, T P., *Curr. Sci*, 1938, 7, 22.

D SCANDENS Benth

Chemical Examination of the seeds of Derris scandens. Rao, C. J. Dassa, Subramanian, S. S *Curr Sci.*, 1947, 16, 346.

DICHANTHIUM ANNULATUM Stapf

Bulk emasculation technique as applied to Dichanthium annulatum Stapf Oke, J G., *Sci & Cult*, 1950, 16, 30

DICHROA FEBRIFUGA Lour

Chang Shan—a Chinese antimalarial herb, Dichroa febrifuga C. S. Jang, F. Y Fu, C Y Wang, K C. Haung, G Lu and T C Chou, *Science*, 103, 50, 1946.

Alkaloids with high antimalarial activity from Dichroa febrifuga J B Koepfli, J. F Mead and John A Brockman, *J. Am. Chem. Soc* 69, 1837, 1947

Alkaloids of Dichroa febrifuga. Frederick, A , Kuehl. Jr Claude F. Spencer and Karl Folkers, *J Am Chem. Soc* 70, 2091, 1948

Pharmacology of Chan Shang (Dichro febrifuga), a Chinese antimalarial herb. C. S Jang and F. Y. Fu and K C Haang and C. Y Wang, *Nature*, 161, 400, 1948

Alkaloids of Dichroa febrifuga. Isolation and degradation studies. J B Koepfli J F Mead and John A Brockman, *J Am Chem Soc* 71, 1048, 1949

DICTAMUS FRAXINELLA Pers

Oil from the leaves and flowers of Dictamus fraxinella Pers B N. Rutovskii and I V Vingradova, *Chem Abst* 3774, 1926

DIDYMOCARPUS PEDICELLATA Br

Didymocarpus pedicellata. The constituents of Part I Isolation of a new series of colouring matters. Siddiqui, Salimuzzaman; *J Ind Chem. Soc* 14 (1937), 703

Didymocarpus pedicellata. The constituents of. Part II. Comparative studies in the constitution of Pedicine, isopedicin, pedicinin & Pedicellin Sharma, Vishwanath & Siddiqi, Salimuzzaman, *J Ind Chem Soc* 16 (1939)

Constitution of Didymocarpus pedicellata Pt V. Studies in the constitution of Pedicinin Kailash Chander Saloaja, Vishwa Nath Sharma & Salimuzzaman, Siddiqui, *J Sci Ind Res*, 1947, **6B**, 57.

DIGITALIS (Tourn.) Linn

Physiological standardization of Digitalis with special reference to the method suitable for use in India Tate, G, *Ind Med Gaz*, 1921, 56, 505

Variation in the potency of Digitalis preparations Chopra R N, Bose, S C & De P., *Ind. Med. Gaz*, 1925, 60, 93

Clinical studies on Digitalis in Bengal, Bose, S C., *Ind Med Gaz*, 1925, 60, 147

Variations in the potency of Digitalis preparations in tropics Chopra, R. N, De, Premankur, *Ind. Med. Gaz*, 1926, 61, 57

Observations on the potency of Indian Digitalis Chopra, R N, De, Premankur, *Ind Med Gaz*, 1929, 64, 312

Biological assay of Digitalis preparations in the tropics, Pt III Chopra, R N & Chowhan, J S., *Ind Jour. Med. Res.*, 1933, 20, 1189

The value of palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves. Bernard S Eeinstein & Frank J Slame, *J Am Pharm Assoc* 29, 370, 1940

Combined Digitalis and Rauwolfia poisoning in a human subject De, M M & Tarapaido Chatterjee, *Ind Med. Gaz*, 1941, 76, 724

Digitalis, their chemical evaluation Notes & News, *Ind Jour. Pharm*, 1946, 8, 50

D LANATA Ehrh

A study on Digitalis lanata Ehrh S C Grown in India Bose, A N & Bose, S, *Ind Jour Pharm*, 1952, 14, 189

DIOSCOREA HISPIDA Dennst

Starch from the tubers of Dioscorea hispida Dennst Rao, P S & Beri, R M, *Sci & Cult*, 1952, 17, 482

DIOSPYROS EMBRYOPTERIS Pers.

Pectin in fruits of the Indian Persimmon (D embryopteris Pers) Biswas, H G., *Sci & Cultr*, 1944, 9, 501.

DIPTEROCARPUS INDICUS Bedd

On the essential oil from the Oleo-resin of Dipterocarpus indicus Krishna G S, Rao, Sukh Dev & Guha, P C, *Sci & Cult.*, 1950, 16, 74

Studies in sesquiterpenes Part VIII Sesquiterpene of the essential oil from Dipterocarpus indicus, Bedd Krishna Rao, G S , Sukh Dev & Guha, P C , *J Ind Chem Soc* 29 (1952) 589

DODONAEA VISCOSA Linn

Chemical examination of Dodonaea viscosa Linn T P Ghosh, *Indian Forester*, 59, 78, 1933

Chemical examination of the seeds Dodonaea viscosa, isolation of dodonin and a fixed oil Dharmpal Parihar and Shikhibhushan Dutt, *Pro Ind Acad. Sci* **26A**, 56, 1947

DOLICHOS LABLAB Linn

Oil of Dolichos lablab B S Rao and Co-workers , *Perfum Record*, 28, 1937, 413

DROSERA INDICA Linn

A new variety of Drosera indica Linn from Kolhapur (S M C) Parandekar, S A , Diwan M G , *Curr Sci* , 1943, 12, 304

DRYOPTERIS

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Enquiry in the Madras Presidency XI Extractum Filicis Liquidum Cains, J F & Mhaskar, K S , *Ind Jour Med Res* , 1921 9, 191

Constituents of Filexmas, Part I, Aspidinol. Alexander Robertson and William F Sandrock , *J Chem Soc* 819, 1933

Male Ferns of Kashmir Handa, K L , Kapoor, L D & Chopra I C , *Curr Sci* 1947, 16, 56

A short note on Male-ferns Handa, K L , Kapoor L D & Het Singh, *Ind. Jour. Pharm* 1952, 14, 109

EICHHORNIA Kunth.

Studies in the lignacellulose group Part I An investigation into the constituents of Water Hyacinth (Eichhornia crassipes). Hemendra Kumar Sen, Pal, Patitpavan Ghosh, and Sindhu Bhushan., *J. Ind Chem Soc* 6 (1929), 673

A Pamphlet on Water-Hyacinth (Science & Industry Sec) *Sci & Cult* , 1942, 7, 602

The problem of Water-Hyacinth in Bengal Bose, P K , *Sci & Cultr* , 1945, 11, 167

Eradication of Water-Hyacinth by Methoyone on a field scale trial. Mitra, G.P., *Sci & Cult.*, 1948, 14, 143

ELAEIS GUINEENIS Jacq

Red Palm oil in the treatment of Keratomalacia. Aykroyd. W.R. & Wright, R.E , *Ind. Jour Med Res* , 1937, 25, 7.

ELETTARIA CARDAMOMUM Maton.

Notes on some Indian Essential Oils "Elettaria cardamomum", Maton. (Var a-minor) Rao, B Sanjiva, Sudborough, J J and Watson, H E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. **8A**, Part X, pp 155-158

Mysore cardamon oil Krishnan, P P & Guha, P C, *Curr Sci*, 1950, 19, 158

ELEUCINE CORACANA Gaertn

The nutritive value of Indian vegetable foodstuffs, Pt V. Nutritive value of Ragi (Eleucine coracana) Niyogi S P, Narayana, N. & Desai, B G, *Ind Med Res*, 1934, 22, 373

The relative value of the proteins of certain foodstuffs in nutrition Pt. V Supplementary values of the proteins of Eleucine coracana (Ragi) and of certain pulses and stem milk powder studied by the nitrogen balance and certain growth methods Swaminathan, M, *Ind Jour Med Res*, 1938, 26, 107

Eleucine coracana Gaertn A new raw material for the malting Industry Sastri, B N Ragi, *Curr Sci*, 1939, 8, 34

EMBELIA Burm.

The treatment of intestinal worms with the indigenous drugs Butea, Embelia and Kamala Mukerji, A K, Bhaduri, N V, *Ind Med Gaz*, 1947, 82, 66

E RIBES Burm

Constitution of Embelia ribes Part I Kaul, Banerjee, Amarish Chandra Roy, Dutt, & Sikhibhushan, *J Ind Chem Soc.* 6 (1928), 577

Embelia ribes Constitution of the active principles of. Part II Kaul, Ramjee, Amarish Chandra Roy, Dutt, & Sikhibhushan, *J Ind. Chem Soc* 8 (1931) 231

EMBLICA OFFICINALIS Gaertn

Chemical Examination of the seeds of Emblica officinalis Pt I, The fatty oil and its component fatty acids Dhar, D.C., Dhar, M L. & Srivastava, D L., *J. Sci Indl Res.*, 1951, **10B**, 88.

ENHYDRA FLUCTUANS Lour.

Chemical Investigation of Enhydra fluctuans Lour Part I. Chakravarty, R.N. & Dutt, A., *J. Ind Chem Soc* 29 (1952) 374.

ENICOSTEMMA LITTORALE Blume.

Enicostemma littorale Blume in malaria. Rai, B.B., *Ind. Med. Gaz.*, 1946, 81, 506

ENTADA PURSAETHA D.C

A note on the chemistry and pharmacological action of Entada pursaetha DC (E. scandens Benth) Chopra, R.N, Gupta, J.C, Chopra, G.S & Ghosh, B K., *Ind. Jour Med Res.*, 1940, 28, 469

EPHEDRA Linn

The comparative action of Ephedrine and Pseudoephedrine from Indian varieties of Ephedra on heart Chopra, R N, Dikshit, B B & Pillai, K V., *Ind. Jour. Med. Res.*, 1928, 16, 780.

Ephedrine from Indian varieties of Ephedra. Chopra, R.N, Ghosh, S., & Dutt, A. T, *Ind Jour. Med. Res*, 1928, 15, 889

A preliminary note on the action and uses of Pseudoephedrine Chopra, R N, Dikshit, B. B. & Pillai, K V, *Ind Med. Gaz*, 1929, 64, 1

Seasonal variations in the alkaloidal contents of Indian Ephedra Chopra, R N & Dutta, A T, *Ind Jour. Med Res.*, 1930, 17, 647

Indian Ephedra, their Chemistry and Pharmacology Chopra, R N., Krishna, S & Ghosh, T. P., *Ind Jour Med Res*, 1931, 19, 177.

Ephedra and ephedrine in U S S R. *Chem Abst* 6765, 1940.

Indian Ephedra research creates demand (Medicine & Public Health), *Sci & Cultr*, 1940, 6, 174.

Ephedra concentrate and the extraction of ephedrine (Science & Industry Sec.), *Sci & Cultr*, 1943, 9, 195.

Indian Ephedras, N A Qazilbash, *Quart J. Pharm. Pharmacol* 21, 475, 1948.

Ephedra, Pt. I Estimation of total alkd solvent extraction. Datal, V D & Khorana, M L, *Ind Jour. Pharm.*, 1950, 12, 165.

Pharmacognosy of Indian Ephedra. Gupta, B. & Bal, S. N, *J Sci. Indl Res.*, 1952, **11B**, 253.

E NEBRODENSIS Tin

Medicinal flora of Sardinia. Ephedra vulgaris Rich. and Ephedra nebrodensis Tin. Aldo La Floresta, *Arch Pharmacol. Sper.* 69, 67, 1940.

Pharmacological activity of ephedrine hydrochloride prepared from Ephedra nebrodensis from the Oliena mountains. Sante Gahatto, *Chem. Abst* 6209, 1946.

E. VULGARIS RICH.

Medicinal flora of Sardinia, Ephedra vulgaris Rich. and Ephedra nebrodenisis Tin. Aldo La Florista, *Arch. Pharmacol. Sper.* 69, 67, 1940.

The Ephedras of Sardinia Chemical investigation of Ephedra vulgaris Rich. *Chem. Abst.* 8204, 1941

ERIGERIN CANADENSIS Linn

The essential oil of Erigerin canadensis L T K. Geponekov; *J. Applied Chem.* (U.S.S.R.) 6, 111, 1933.

ERYTHRINA INDICA Lam.

Chemical examination of Erythrina indica (White variety) Rao, J Virabhadra, *Curr. Sci.*, 1945, 14, 198

Some common indigenous remedies Picrorhiza kurrooa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum R. N. Chopra & S Ghosh; *Ind Med Record* 55, 77, 1935

ERYTHROXYLUM MONOGYNUM Roxb

The Essential Oil from the wood of Erythroxylum monogynum Roxb Rao B Sanjiva, Shintre V P and Simonsen J L, *Journal of the Indian Institute of Science, Bangalore*, 1926, Vol. **9A**, Part VI, pp. 145-148.

Devadari - root oil B. S Rao and Co-workers, *Perfum. Record*, 28, 1937, 412.

Devadari wood oil S G. Sastry, *Journal of the Mysore Forest Association*, Vol 5, No 4.

EUCALYPTUS L.her

Note on the Eucalyptus Oil industry in the Nilgiris Puran Singh, *Indian Forest Records;* 1917, Vol. V Part VIII

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency VII Oleum Eucalypti Cains, J F & Mhaskar, K S, *Ind Jour Med. Res*, 1920, 7, 372

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency IX The Eucalyptus $CHCl_3$ mixture S C Cains, J F. & Mhaskar, K. S., *Ind Jour Med Res.*, 1920, 8, 384.

E. CITRIODARA Hook.

Leaves of Eucalyptus citriodara yielded 1.3 per cent oil. Noemi Carcia Arrillaga *Chem. Abst* 5615, 1942.

E. GLOBULUS Labill.

The essence of Eucalyptus globulus Martias Gonzales Atilio Lambardo and M A Mesa De Romero, *Chem. Abst* 4666, 1942.

Essential oil of leaves of Eucalyptus globulus from groves in La Plata. Leonor Pelanda Ponce, *Chem. Abst* 5955, 1942.

Essential oil of Eucalyptus globulus. B. V. Ramaswami, P. L. Narasimha Rao and P. C Guha; *J Ind. Inst. Sci.* **28A**, 57, 1946.

Microscopical studies of the leaves of Eucalyptus globulus Labil. and three other species grown in India Amiya Datta & Datta, D, *J. Sci Indl. Res.*, 1952, **11B**, 181

EUGENIA JAMBOLANA Lam.

Eugenia jambolana Subba Rao, N. V & Seshadri, T R, *Curr Sci* 1939, 8, 120.

Jaman (Black Berry) seed as a cattle feed Kehar, N D. & Sahai, K, *Sci & Cult*, 1948, 14, 205.

EULOPHIA R Br

On the occurrence of velamen in Eulophia R Br Lakshminarayana, S & Venkateswarlu, V, *Sci & Cult*, 1950, 15, 327.

EUPHORBIA Linn.

Triterpenic compounds isolated from Euphorbia latices. Gopalachari, R. & Siddiqui, S, *J. Sci Indl Res*, 1949, **8B**, 234

E. GENICULATA Ort.

On the introduction of Euphorbia geniculata Ort. in India. Van Steenis, C G G J, *Curr Sci.*, 1937-38, 6, 119.

Fresh locality record for Euphorbia geniculata Ort in South India Mayuranathan, P V, *Curr. Sci*, 1937-38, 6, 119.

E. HELIOSCOPIA Linn

Seeds of Euphorbia helioscopia L Paul Gillot, *Bull Sci Pharmacol.* 33, 193, 1926

E PEPLIS Linn

Chemical investigation of Egyptian plant Euphorbia peplis, Abd El Aziz Sharaf, *British Vet J* 105, 128, 1949

E PILULIFERA Linn

Pharmacognostic study of Euphorbia pilulifera Linn. Bose, A B., *Sci. & Cultr.*, 1944, 9, 301.

Pharmacology of Euphorbia pilulifera Lloyd W. Hazelton and Rebecca C Hellerman; *J Am. Pharm. Assoc* 37, 491, 1948.

The isolation of L-inositol from Euphorbia pilulifera. Floyd P. Hallet and Lloyd M. Parks *J. Am. Pharm. Assoc.* 40, 474, 1951.

E THYMIFOLIA Linn.

Flavonol glucosides from Euphorbia thymifolia L. Makoto Nagase; *Chem. Abst.* 3625, 1942.

E. TIRUCALLI Linn.

Chemical examination of dried latex from Euphorbia tirucalli. Dutta, N. L. & Karimullah, *J. Sci Indl Res.*, 3, 213, 1944-45.

Chemical Examination of the dried latex from Euphorbia tirucalli Karimullah & Gopalachari, R , *J Sci Indl Res.*, 1949, **8B**, 89.

Chemical Examination of the latex from Euphorbia tirucalli Gopalachari, R & Siddiqui, S , *J Sci. Indl Res.*, 1949, **8B**, 129.

Studies in the const. of Euphoron Pt I (Euphorbia tirucalli). Gopalachari, R & Siddiqui S , *J. Sci Indl. Res.*, 1949, **8B**, 140.

FAGOPYRUM CYMOSUM Meissn

Phytochemical component of Fagopyrum cymosum. Kaneo Imai and Kiyoshi Furrya, *J. Pharm Soc Japan*, 71, 266, 1951

FERONIA ELEPHANTUM Correa.

Feronia elephantum Essential oil from the leaves of Bhatia & Deshapande, *J. Ind Chem Soc* 26(1949), 342.

Investigation on the structure of Ketha (Feronia elephantum, family Rutaceae) Gum Mathur, G P., Mukerjee, S , *J. Sci Indl Res* , 1952. **11B**, 344

FERULA Linn.

Chemical composition of Ferula spp O M. Efimenko, *J Applied Chem.* (U. S. S R.) 12, 202, 1939.

Chemical assay of Rasaunt and Hing from the Punjab market Grewal, K. S., Kochhar, B. D , *Ind. Jour. Med. Res.* 1940, 28, 463.

F. FOETIDA Regel.

Studies in the specification of Indian Medicinal Plants. Pt I Asafoetida, Ferula foetida. Sen Gupta, S B & Das, B. K , *Ind Jour Pharm.*, 1948, 10, 36

F. JAESCHKEANA Vatke.

Chemical investigation of the fruits of Ferula jaeschkeana Vatke V. P Bersutskii, *Chem Abst* 4522, 1940

FICUS ANTHELMINTICA Martius.

Note on Ficus indica Linn. and closely allied American species. Ficus laurifolia Hort. et Lam. and Ficus anthelmintica Martius. *Curr. Sci.*, (1933-34), 2, 51

F. BENGALENSIS Linn.

Fasciated aerial roots of Banyan & (Ficus bengalensis Linn.). Nadkarni, M D., Airan, J. W , Shah, S V., *Curr. Sci.*, 1944, 13, 233.

Investigation on "Ficosterol". a sterol occurring in Ficus bengalensis. Nath, M.C. & Debnath, C R.; *Sci & Cultr.*, 1947, 12, 599.

F. CARICA Linn.

The constituents of the leaves of Ficus carica. Kunio Okahara, *Bull. Chem Soc. Japan*, 11, 389, 1936

Some pharmacological and biological effects of the latex of Ficus carica L. S. B Ullman, L. Helberstaedter and J Leibowitz, *Chem. Abst* 3071, 1945, 530

Preparation of a highly active vegetable Rennet from Ficus corica Linn Krishnamurthy, C R , Jaganathan, V & Subrahmanyan V , *J Sci Indl Res* , 1945-46 4, 720

The proteolytic enzyme of the latex of Ficus carica Linn Krishnamurti, C R & Subrahmanyan, V , *Sci. & Cultr* , 1947, 13, 204.

F INDICA Linn & F laurifolia Hort et Lam

Note on Ficus indica Linn and closely allied American species—Ficus laurifolia Hort et Lam and Ficus anthelmintica Martius. *Curr Sci* , (1933-34), 2, 51

F RELIGIOSA Linn

Fascination in an anomalous foliar organ of Ficus religiosa Linn De Sarkar, K & Datta, R M , *Sci & Cult* , 1952, 17, 381

FOENICULUM PANMORIUM D C

Notes on some Indian Essential oils Foeniculum panmorium D C (Fennel) Rao B Sanjiva, Sudborough, J. J and Watson, H. E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. **8A**, Part X, pp 184-186.

FRITILLARIA ROYLEI Hook

Some minor alkaloids of Fritillaria roylei T Q Chou, *J Pharm Assoc* 36, 215, 1947

FUMARIA OFFICINALIS Linn

Chemical examination of some Indian medicinal plants, Tinospora cordifolia, Solanum xanthocarpum and Fumaria officinalis Pense G P & Dutt, S , *Ind Jour Med Res* 1932 20, 663

The alkaloids of Fumariaceae plants Fumaria officinalis Richard H. F Manske, *Can J. Research*, **16B**, 438, 1938

Action of Fumarine (Fumaria officinalis) Wilhelm Bolm. Arch, *Exptl Path Pharmacol* 195, 304, 1940

FUSARIUM.

Antibacterial properties of Yeasts—Fusarium species, onion and garlic. C Carpenter, *Chem Abst.* 1558, 1946

GARCINIA INDICA Chois.

Garcinia indica. Fatty acids and glycerides of the fat from the seeds of Vidyarthi, N. L. & Desa Rao, C J ; *J. Ind Chem Soc.* 16 (1939), 437.

G. MANGOSTANA Linn

Resin of Garcinia mangostana. L. Otto. Dragendorff; *Ann.* 482, 280, 1930.

Resin of Garcinia mangostana. Otto Dragendorff, *Ann* 487, 62, 1931

G. MORELLA Desr

Indian seed fats. Mowha (Bassia Latifolia) and Tamal (Garcinia morella) fats. D R. Dhingra, G.L Seth & P. C Speers, *J Soc Chem Ind* 52, 116, 1933.

On 'Morellin' the antibacterial principle of the seeds of Garcinia morella. Rao. R., Raghunandana & Natarajan, S, *Curr Sci*, 1950, 19, 59

Antibiotic principles of Garcinia morella Pt. I Preparation and antibacterial activity of Morellin, Morellin-t, Morellin M, Morellin I, & Iso morellin Narsimba Rao, P L & Verma, S C. L, *J Sci Indl Res*, 1951, **10B**, 184

Antibiotic principles of Garcinia morella Part II. Chemistry of Morellin Narasimba Rao, P L & Verma, S C L, *J Sci Indl Res* 1952, **11B**, 206

GAULTHERIA FRAGRANTISSIMA Wall

A pharmacological study of Gaultheria fragrantissima S Prasad, *Ind. J. Pharm* 3, 62, 1941.

Detection of Scopoletin in radix gelsemi and radix belladonnanae Robert Fischer and Herbert Ehrlich, *Arch Pharm.* 274, 268

G PROCUMBENS.

Manufacture of Wintergreen oil in India—Puran Singh, *Indian Forest Records*, 1917, Vol V Part VIII

GENTIANA KURROO Royle

Comparative pharmacognosy of Gentiana kurroo Royle. & Picrorrhiza kurrooa Royle ex Benth Bal S. N & Datta, S. C., *Ind Jour. Pharm.* 1945, 7, 109.

GERANIUM MACRORRHIZUM Linn

Oil from Geranium macrorrhizum. Schimel and Co, *Chem Zentr* 1519, 1927.

GLORIOSA SUPERBA Linn.

A note on the distribution of Gloriosa superba Linn. Sayeeduddin, M & Salam, M Abdusa, *Curr Sci.*, 1934-35, 3, 443.

Alkaloidal constituents of Gloriosa superba Linn. Sulbaratnan A V, *J. Sci. Indl Res.*, 1952, **11B**, 446.

GLYCOSMIS PENTAPHYLLA Correa.

Alkaloids from Glycosmis pentaphylla Correa., Chatterjee, A. & Majumdar, S. G., *Sci. & Cult.*, 1952, 17, 306.

Glycosin, the new alkaloid of Glycosmis pentaphylla Correa. Asima Chatterjee & Majumdar, S. G , *Sci & Cult.* 1953, 10, 506.

GLYCYRRHIZA GLABRA Linn.

The flavanone glucoside of Glycyrrhiza glabra Var. glandulifera. Regel et Herder, Junzo Shinada and Seuchi Ueeda, *Ber* **67B**, 434, 1934

A study of vehicles for medicines The Glycyrrhiza vehicles. Bernard Fantus, H A Dyniewicz and J. M Dyniewicz, *J. Am Pharm Assoc* 23,9 15, 1934

Sweet constituents of liquorice root Tatuo Kariyone and O Nonaka, *J Pharm Soc Japan*, 57, 166, 1937.

GOSSYPIUM HERBACEUM Linn.

Constitution of Herbacetrin and Herbacetin Neelkantam, K & Seshadri, T R , *Curr Sci.*, 1936-37, 5, 476

GYMNEMA AURANTIACUM Wall

Chemical Examination of Meda (Gymnema aurantiacum) Tubers Mukherjee, S & Srivastava, H C , *J Sci Indl Res* , 1951, **10B**, 324.

G SYLVESTRA Br

Gymnema sylvestra in Diabetes Mellitus Chopra, R N., Bose, J P , Chatterjee, N K., *Ind Jour Med. Res* , 1928, 16, 115

HAGENIA ABYSSINICA J. F Gmel

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency XII Kousso Cains, J F & Mhaskar, K S , *Ind Jour Med Res* , 1921, 9, 198

HARDWICKIA PINNATA Roxb

The oleo-resin from Hardwickia pinnata K S Iyer and J J Sudborough, *Journal of the Indian Institute of Science, Bangalore,* Vol. 2, Part III, pp. 29 to 35, 1918

Hardwickia pinnata Essential oil from the oleo-resin of Dev & Guha, *J. Ind Chem Soc.* 25 (1948), 495

Humulene, presence of, in oleo resin from Hardwickia pinnata Dev & Guha, *J. Ind Chem Soc.* 25 (1948), 500

HELIANTHUS ANNUUS Linn.

A rapid method for determining the oil content of Safflower and sunflower seeds. W Keith Kennedy and John Urau, *Chem. Abst.* 5209, 1949

HELMINTHOSPORIUM TETRAMERA Mckinney.

A preliminary note on the isolation and pathogenicity of Aerothecium lanatum Wakker and Helminthosporium tetramera Mckinney on the leaves of Oryza sativa Linn. Ganguly, A.K , Ganguly, D.D. *Sci & Cultr* , 1941, 6, 424.

HEMIDESMUS INDICUS Br.

Chemical examination of the roots of Hemidesmus indicus A T Dutt, S Ghosh and R N Chopra, *Arch. Pharm.* 276, 33, 1938

Preparation of the essential oil from the root of Hemidesmus indicus (Sarsaparilla) B. Sanjiva Rao, K.S Subhramanian and N C Kelkar, *Proc Soc Biol Chemists India*, 3, 35, 1938

Ether soluble constituents of Sarsaparella root. James C E Simpson and Norman E Williams, *J Chem Soc* 2040, 1939

HERACLEUM NEPALENSE D. Don

Crystalline components of the seeds of Heracleum nepalense Chandranath Bhar *Sci & Cult*, 1947, 12, 504

Heracleum nepalense Crystalline components of the seeds of. Bhar. *J Ind. Chem Soc* 25(1948), 139

HERPESTIS MONNIERIA H B.K

The Chemical investigation of the leaves of Herpestis monnieria Basu, N K & Walia, J S, *Ind Jour Pharm*, 1944, 6, 84

Correct name of Herpestis monnieria H B K Chatterjee, D., *Sci & Cult*, 1948, 14, 156

HIBISCUS CANABINUS Linn

The colouring matter of Deccan Hemp (Hibiscus cannabinus) flowers. Cannaluscitrin and Cannabiscetin. Neelakantam, K, Seshadri, T R, *Curr Sci*, 1937-38, 6, 504.

H ESCULENTUS Linn

The biological value of protein of papaya (Carica papaya) and Lady's fingers *Exptl Med* 2, 71, 1942

Availability of Calcium in Lady's finger (Hibiscus esculentus), cabbage (Brassica oleracea var. capitata), Drumstick (Moringa oleifera) and Amarnath Tender (Amaranthus gangeticus) Pt. I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K.P & Ghosh, D., *Ind Jour. Med Res.*, 1943, 31, 29

Availability of calcium in Lady's finger (Hibiscus esculentus), cabbage (Brassica oleracea var capitata), Drumstick (Moringa oleifera) & Amaranth Tender (Amaranthus gangeticus). Pt. II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P & Ghosh, D., *Ind Jour Med Res*, 1943, 31, 37.

Colouring matter of the flowers of Hibiscus esculentus Seshadri, T R. & Viswanathan, N., *Curr. Sci.* 1947, 16, 343.

H SABDARIFFA Linn.

Sida sabdariffa (Hibiscus sabdariffa) Henri Leeleve; *Chem. Abst.* 3891, 1939.

The chemical, botanical and pharmacological characters of the Karkade (Resella) Hibiscus sabdariffa. G. Reabourg and R.H. Moneeu, *J Pharm Chem.* 292, 1940.

Isolation of Hibiscitrin from the flowers of Hibiscus sabdariffa. Constitution of Hibscetin P Surya Prakash Rao and T R Seshadri, *Proc. Indian. Acad Sci* **15A**, 148, 1942.

Constitution of Hibiscetin. Suryaprakasa Rao, *Curr Sci* 1942, 11, 360.

H. SYRIACUS Linn

A note on the Pistillody in Hibiscus syriacus Linn Rao, L. Narayana, *Curr. Sci*, 1936-37, 5, 301

H VITIFOLIUS Linn.

Colouring matter of the flowers of Hibiscus vitifolius K. Visweswara Rao and T R. Seshadri, *Proc Ind Acad Sci* **24A**, 352, 1946.

HOLARRHENA ANTIDYSENTERICA Wall

Conessine Darab Dinsha Kanga, Panch Nanda Ramaswamy Iyer & John Lionel Simonsen, *J. Chem. Soc* 2123, 1926

Oil from the seeds of Holarrhena antidysenterica Ramachandra R V, Ghanikar and P R Ayyar, *J. Ind Inst Sci* **10A**, 20, 1927.

Observations on the pharmacological action of conessine, the alkaloid of Holarrhena antidysenterica Chopra, R N, Gupta, J C., David, J C & Ghosh, S, *Ind Med Gaz*, 1927, 62, 132

The use of a standarized preparation of the total alkaloids of Kurchi Bark in Amoebic dysentery Majumdar, A R, *Ind. Med Gaz*, 1930, 65, 80

New alkaloids from Holarrhena antidysenterica seeds Robert D Haworth, *J Chem Soc* 631, 1933

Pharmacological action of Kurchicine (an alkaloid of Holarrhena antidysenterica) Chopra, R N & Chowhan, J S, *Ind Jour Med Res*, 1933, 21, 277

The treatment of chronic intestinal amoebiasis with the alkaloid of Holarrhena antidysenterica (Kurchi) Acton, H W, Chopra R N, *Ind Med Gaz.* 1933, 68, 6.

A study of the preparations of an efficient extract of Kurchi Hollarrhena antidysenterica Ghosh, A B, *Ind Med Gaz*, 1933, 68, 12

Some new bases from Holarrhena antidysenterica Alfred Bertho, Gustav Von Schuckmann and Walter Sitionberger, *Ber.* **66B**, 786, 1933

The nisoles of the latex of Holarrhena antidysenterica. J.C. Choudhary and D H Peacock, *J Chem Soc.* 1129, 1935

Kurchi alkaloids. Preparation of conessine and its secondary alkaloids. Alfred Bertho, *Arch. Pharm.* 277, 237, 1939.

The Alkaloids of Kurchee Bark (Holarrhena antidysenterica) Part I. Ghosh, Sudhamony & Ghosh, Nagendra Nath, *J. Ind Chem Soc.* 5 (1928), 477.

Holarrhena antidysenterica The alkaloids of. Part I Three new alkaloids from the bark of Indian Holarrhena and new method of isolation and further purification of conessine Siddiqui,' S & Pillai , *J Ind. Chem. Soc* 9(1932), 553

Holarrhena antidysenterica. The alkaloids of Part II. Two further new alkaloids from the bark and the seeds of Indian Holarrhena and their constitutional relationship to conessine. Salimuzzman Siddiqui, *J Ind Chem Soc* 11, 1934, 283

The preliminary pharmacological studies of Kurchi bark Madhev Lal Schroff, S.N Bal and Mohan Lal Dhir , *Ind J. Pharm* 2, 195, 1940

Pharmacological tests on extract and alkaloids of Holarrhena antidysenterica. Alfred Bertho , *Arch Exptl Path Pharmakol* 203, 34. 1944

Studies in the Conessine series, Pt IV Action of sulphuric acid on conessine and its conversion to a new isomer Neo-conessine Salimuzzaman Siddiqui & Shyam Kishan Vasisht, *J Sci. Indl Res.*, 1944-45, 3, 559.

Pharmacognostic studies on Kurchi Bark. Datta, S C & Bal, S N., *Ind. Jour Pharm* , 1945, 113, 7

On the assay of alkaloid in Kurchi Bismuth iodide. Mukerjee, S & Dutta, B C., *Sci & Cultr* , 1945, 10, 506

Studies in the conessine series, Pt 7 Salimuzzaman Siddiqui & Vishwa Nath Sharma, *J. Sci Indl Res.* 1945-46, 4, 435.

Studies in the Conessine series, Pt 8 Salimuzzaman Siddiqui & Shyam Kishan Vasisht, *J. Sci Indl Res.*, 1945-46, 4, 440

Chemistry of Kurchi Seeds Pt I Isolation of a crystalline Glycoalkaloid, Irani, (Miss) R.J.,*Curr Sci* , 1946, 15, 106.

Chemistry of Kurchi Seeds Pt. II Isolation of the Bromide of a Linoleo-Lilinolenin from the fatty oil Irani, (Miss) R.J , *Curr Sci.*, 1946, 15, 161.

Chemistry of Kurchi Seeds Pt. III A new & simple method of analysis by Bromoglycerides. Irani, (Miss) R J , *Curr. Sci.* 1946, 15, 191

Chemistry of Kurchi Seeds, Pt, IV Isolation of Galactose from the Picric acid hydrolysis of glyco alkaloid Irani, (Miss) R.J , *Curr. Sci.*, 1946, 15, 229.

Assay of extract of Kurchi Liq Basu, N.K. & Mithal, B.M., *Ind. Jour Pharm.*, 1947, 9, 118.

Assay of Kurchi Bark Basu, N K & Mithal, B M., *Ind. Jour. Pharm.* 1948, 10, 72.

A method for the assay of Kurchi Bismuth Iodide. Rao, G.K , *Ind. Jour. Pharm.*, 1948, 10, 100.

The decomp of 'Kurchi alkaloids. Sen Gupta, S.B., Dutta, H.B.; *Sci. & Cult*, 1949, 15, 32.

Holarrhena antidysenterica Wall., estimation of total alkaloids in. Datta, A, Ghosh, B K., *Ind Jour Pharm* 1949, 11, 74

Kurchi bases, labile nature of Basu, N.K. & Battacharya, N N., *Ind Jour Pharm*, 1949, 11, 157.

Studies in the conessine series. Pt. 10 Action of fuming Nitric acid (HNO_3) on conessine Vishwa Nath Sharma & Siddiqui, S., *J Sci. Indl Res*, 1950, **9B**, 84.

The seasonal variation of alkaloids in different parts of Holarrhena antidysenterica Wall. and the alkaloidal content in the bark at different ages Datta, A T Ghosh, B K & Gupta, J C, *Ind Jour. Med Res.*, 1950, 38, 467.

Thermolabile nature of Kurchi Bases Thomas Oommen, P., *Ind. Jour Pharm*, 1951, 13, 64.

H. FEBRIFUGA Klotsch

Chemical Examination of Holarrhena febrifuga Klotztsch, Pt I. Salimuzzaman Siddiqui, Shambhu Charan Misra & Viswa Nath Sharma, *J Sci Indl. Res*, 1944-45, 3, 555

HOLIGARNA ARNOTTIANA Hook. f.

Chemical Examination of the constituent of Holigarna arnottiana Hook f. Pt. I latex of Semecarpus travancorica Bed. Pt. II Nair, A.V, Poti, D M & Pillay, P P, *J. Sci Indl Res*, 1952, **11B**, 294, 298.

Holigarna lactone—A new aldehydelactone from the seed kernels of Holigarna arnottiana Hook f Nair G V, Poti A N., Pillay P P, *J S I R.* 1953, **3B**, 119

HYDNOCARPUS Gaertn

Report of a chemical investigation of Chaulmoogra oil in connection with leprosy treatment Sudhamoy, Ghosh, *Ind Jour Med Res*, 1917, 4, 691.

Chemotherapeutic experiments with Chaulmoogras and allied preparations O.H Schohl, *Phillipine J Sci* 23, 533, 1923

Chaulmoogra oil in the treatment of Trachoma. Gubbay, B B & Row, T M., *Ind Med Gaz.* 1929, 64, 563.

A preliminary note on the action and uses of 'Alepol'. Dikshit, B B & Row, T.M, *Ind. Med. Gaz* 1931, 66, 317

Alepole in Leprosy Dikshit, B B., *Ind Med Gaz*, 1932, 67, 7.

Alepole in the treatment of Leprosy Bhandari, A. D., *Ind Med Gaz*, 1932, 67, 244.

Chemical composition of Congo Chaulmoogra seeds and fat L. Adrains, *Chem. Abst.* 6754, 1946.

Examination of the purity of Chaulmoogras aethylicus and oleum chalmoogra. J.T. Kruithof; *Chem. Abst.* 2859, 1947.

H. ANTHELMINTHICA Pierre

Hydnocarpus anthelminthica seeds from Ceylon Anon; *Chem. Abst.* 3391, 1930.

Analysis of Chaulmoogra oils Hydnocarpus anthelminthica oil and Taraktogenos kurzii (Chaulmoogra oil). Howard Cole and Humberts to T. Cardoso, *J Am Chem Soc* 61, 3442, 1939

H. WIGHTIANA Blume

Efficacy of ethyl chaulmograte ethyl hydnocarpate and ethylesters of the total fatty acids of Hydnocarpus wightiana oil Bonifacio De Vera and Casimiro B Lara, *Chem Abst* 1424, 1930.

The constituents of Hydnocarpus wightiana oil Howard Cole. *Chem. Abst.* 1466, 1930

Reduction of irritation by iodised ethyl esters of Hydnocarpus wightiana oil, Howard. Cole, *Chem. Abst* 1466, 1930.

Chemistry of antileprosy oils (Hydnocarpus wightiana) Buu Hoo Paul Caghniant and Joseph Janicaud, *Compt. Rend* 212, 577.

Hydnocarpus wightiana A note on the keeping properties of oil from, and its derivatives Basu, U P & Majumdar, A., *J Ind Chem Soc* 17(1940), 280

HYDROCOTYLE ASIATICA Linn

Chemical examination of the constituents of Hydrocotyle asiatica. M A. Wali and M C Tumunkatti, *Proc Ind Acad Sci*, **5A**, 109, 1937

A preliminary note on the fission of Vascular Cylinder in some of the roots of Hydrocotyle asiatica Linn. Chakraverti, D N, *Curr Sci.*, 1940, 9, 230

Identification of Brahmi Nityendra Nath Sircar, *Sci & Cultr.* 1941, 7, 120.

Investigation of Indian medicinal plants, Hydrocotyle asiatica, Vitex negundo and Monniera cunifolia. Basu, N K., Lamsal, P P and Singh, G B., *Quart. J Pharmacol* 20, 135, 1947

HYGROPHILA SPINOSA T Anders

Hygrophila spinosa. Chemical examination of the roots of Ghatak, Narendra Nath & Dutt, Sikhibushan *J Ind Chem Soc* 8 (1931), 23

Some common indigenous remedies, Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum Chopra, R. N. and Ghosh, S, *Ind Med Record* 55, 77, 1935.

An investigation of oil from seed of Hygrophila spinosa Godbole, N. N. Gunde, B D. & Srivastava, P. D, *Oil and Soap*, 18, 206, 1941.

Chemical investigation of Hygrophilla spinosa. Lagawankar, J. D, Phalinkar, N.L. and Bhide, B V, *J Univ Bombay.* 13, 15, 1945.

HYOSCYAMUS Linn.

The value of Palisade ratios in the differentiation of official Belladonna, Digitalis, Hyoscyamus and Stramonium leaves Bernard S Feinstein and Frank J. Slama, *J. Am Pharm Assoc* 29, 370, 1940.

A belladonna and Hyoscyamus extract which is stable in the air and contains the alkaloids of the drug in mutually unchanged relative proportion, J Gjesing Andersen, *Chem Abst* 6497, 1944.

Increased alkaloidal contents of induced polyploids of Datura Atropa and Hyoscyamus Part I Datura species Jack Morris Rowson, *Quart J Pharm Pharmacol* 18, 175, 1945

HYOSCYAMUS NIGER Linn

Indian Henbane Handa, K L., Kapoor, L D & Chopra, I C, *Curr. Sci*, 1947, 16, 315

Cultivation of Hyoscyamus niger Linn Chopra, I C, Kapoor L.D, Handa, K L. *J S I R* 1953, **5A**, 238

HYPERICUM PERFORATUM Linn

Hypernin, a glucoside of Hypericum perforatum Zofia Jerzmanoroska, *Chem Abst* 7299, 1939

HYPTIS SUAVEOLENS Poit

Essential oil from Hyptis suaveolens Nayak, K C & Guha, P C, *J. Ind Chem Soc* 29 (1952), 183

ILEX VOMITORIA (Soland) Hort

The pharmacological action of Cassine Croceum De and Mundulea suberosa Benth N Sapeika, *S African J Med Sci* 10, 51, 1945

ILLICIUM PARVIFLORUM Michx.

Note on the volatile oil of Illicium parviflorum Michx Foote, P A., *J. Am Pharm. Assoc* 27, 573, 1938

I RELIGIOSUM Sieb & Zucc

A study of Illicum religiosum Chen, K K, *J Am Pharm. Assoc* 15, 861, 1926.

INDIGOFERA ENNEAPHYLLA Linn.

Chemical investigation of Indigofera Enneaphylla Linn. and the isolation of its active principle Satyendra Nath Chatterji and Shikhibhushan Dutt, *Proc Natl Inst India* 3, 374, 1937

I. LINIFOLIA Retz

Chemical examination of Indigofera linifolia Retz The isolation of its active principles. Mahado Prasad Gupta and Shikhibhushan Dutt, *Proc Natl Acad Sci. India.* 8, 49, 1938

INULA HELENIUM Hk. f & T.

Examination of plants for insecticidal principles, Elecampane. F. Tutin, *Chem. Abst.* 4337, 1933

Bitter principles of Elecampane roots Karl, Fr., W Hansen; *J. Prakt. Chem.* 136, 176, 1933.

Pharmacological studies of Korean Inula helenium L. Tinsyo Go, *Japan J Med Sci Pharmacol.* 12, 110, 1938

I. ROYLEANA D.C

Inula royleana D C. Its Chemistry, and pharmacological action Chopra, I C., Kohli J D & Handa, K L, *Ind Jour Med Res*, 1945, 33, 139

IPOMOEA Linn.

The seed structure of Ipmoea, a criticism. Maheshwari, P., *Sci. & Cultr*, 1944, 9, 557.

I. HEDERACEA Jacq

Studies in the specification of Indian Medicinal Plants Pt. III Kaladana Sengupta, S B, Gupta, H N, *Ind Jour Pharm*, 1948, 10, 106.

I. MURICATA Jacq.

Ipomoea muricata. Fatty oil from the seeds of Kelkar, Phalnikar & Bhide, *J Ind Chem Soc* 24 (1947), 87-90

Ipomoea muricata Chemical examination of the seeds of Misra & Tewari *J Ind Chem Soc* 28 (1951), 221

Chemical examination of seeds of Ipomoea muricata Part II Misra, A L & Tewari, J. D, *J Ind Chem Soc* 29 (1952), 63

Chemical examination of seeds of Ipomoea muricata Part III. Misra, A L. & Tewari, J D, *J Ind Chem Soc* 29 (1952), 430

Chemical examination of Ipomoea muricata seeds, Part IV. Misra A L., Tewari J D, *J I C S* 1953, 6, 391

I PESCAPRAE Roth

A phytochemical study of Ipomoea pes-caprae Gustav E. Cwalina and Glenn L. Jenkins, *J. Am Pharm Assoc* 27, 585, 1938.

Study of the leaves of Ipomoea pescoprae B V Chrestense and J A. Reese, *J Am Pharm. Assoc.* 27, 195 1938.

I. PULCHELIA Roth.

Damping of Ipomoea pulchelia Roth, due to Pythium aphanidermatum (Eds) Fitz Mohmud. K. A & Jain, A C.

I. REPTANS Poir.

The carotenoids and some lipoids of Ipomoea reptans Poir. M. Ishi, *Chem. Abst* 3376, 1935

JASMINUM Linn.

Jasmine Oil—J N. Rakshit, *Perfum Record*, 26, 1937, 241.

Jasmin Flowers—Sadgopal; *S.P.C* July, 1939, p. 589.

Development of essential oil industry (Ocimum lemon, O. basilicum, O. canum, O. sanctum, lemon grass, Motia, Palmarosa grass). Jitandra Nath Rikshit, *Sci & Cultr* 1939, 5, 108.

Chromosome numbers in some ornamental Jasmines. Mridula Dutt, *Sci & Cult.*, 1952, 17, 527.

JATEORRHIZA PALMATA Miers

Bitter principles of Columba root, Palmarin F Wesseky, K Schonol and W Iseman, *Monatsh.* 68, 21, 1936.

Bitter principles of Columba root, Methylation of Columbin. F. Wessely and K Jentzsch, *Monatsh* 70, 30, 1937

Evaluation of Colombo root and its preparations H Neugebourer and K Brunner, *Arch. Pharm* 276, 199, 1938.

JATROPHA CURCAS Linn.

Seeds of Jatropha curcas from the Belgian Congo Adriaens, *Chem Abst.* 5440, 1936.

Chemical investigation of the seeds of Jatropha curcas Linn Desai, C M & Vyas, M T , *Curr Sci* , 1949, 18, 49

J GLANDULIFERA Roxb

Jatropha glandulifera Roxb Examination of the seeds of. Alimchandani, Badami Katti, *J Ind Chem Soc* 26 (1949), 523.

JUGLANS REGIA Linn

The Vit C contents of Tender Walnut (Juglans regia) Ratnanathan, S., *Ind Jour Med Res* , 1942, 30, 513

JUNIPERUS Linn.

Oil of Juniper berries Ernst S Guenthur, *Am Perfumer* 33, 64, 1936

Analysis of Juniper berries. P Casparis and W Freund, *Pharm. Acta Helv* 13, 307, 1938

New hydrocarbon from juniper oil P Casparis and W Freund, *Pharm Acta Helv* 14, 1, 1939.

Assay of Juniper for volatile oil content *Chem Abst* 4523, 1940

J COMMUNIS Linn

Constituents of some Indian essential oils, essential oil from the seeds of Zanthoxylum ovaltifolium & Juniperus communis J. L. Simonsen, *Ind Forest Records* 11, 1, 1924

The constituents of some Indian Essential Oils Part XV—The Essential Oil from the seeds of Juniperus communis. Simonsen J L , *Indian Forest Records*, 1925, Vol. XI Part I, pp 6-9

KAEMPFERIA GALANGA Linn.

Constituents of some Indian essential oils Essential oil from the rhizomes of Kaempferia galanga. Puthan Madhathil, Bhaskara Panicker, B. Sanjiv Rao and J. L. Simonsen, *J. Ind Inst of Sci* **9A**, 133, 1926.

LACTUCA SCARIOLA Linn.

Kahu (Lactuca scariola) Seed Oil Component fatty acids of. Dhingra & Pershad, *J Ind Chem. Soc* 22 (1945), 127

LAGENARIA VULGARIS Ser

Wilt disease of Lagenaria vulgaris Ser Ashok Kumar Kar, *Sci & Cult*, 1946, 12, 290

An abnormal flower of Lagenaria vulgaris Ser Ahmed Shamsul Islam, *Sci & Cult*, 1951, 16, 427

LANSIUM ANAMALAYANUM Bedd

Chemical Investigation of some minor Forest Products The Essential oil of Chigatmari (Lansium anamalayanum) Rao K A Narain, Varadhan C and Janniah S L., *Department of Industries, Bangalore*

Studies in sesquiterpenes Part X Sesquiterpenes of the essential oil of Lansium annamalayanum Bedd A Somasekar Rao, Dutta, K B , Subdev & Guha, P C , *J Ind Chem Soc* 29(1951), 604

L. DOMESTICUM Correa

Chemical study of the seeds of Lansium domesticum Correa Kassem Pangsrivongse, *Chemical Abst* 3088, 1938

LANTANA CAMARA Linn

Oil from the flowers and leaves of Lantana camara—D. D Kanga, *Journal of the Indian Institute of Science*, Bangalore Vol. I, Pt IX, pp 93 to 95, 1914

Travancore Essential Oils Lantana camara Moudgill, K. L. and Vrindachalam, P N , *Department of Industries, Trivandrum*, 1923, Bulletin No XVII, pp 7-10.

Travancore essential oils from the leaves of Lantana camara. Moudgil, K L , *Perfumery Essent Oil Record* 16, 9, 1925.

Volatile oil of Lantana camara L. Kinzo Kafuku, Tassaku Ikeda and Yasuji Feijita, *J Chem Soc Japan*, 55, 305, 1924.

Lantana camara (The paper was read in the Annual meeting of the National Academy of Science and has been published in the annual issue of the Academy, 1939)

LATHYRUS SATIVUS Linn

The production and pharmacological action of Khesari Amine. Acton, Hugh W. & Chopra R.N., *Ind. Med Gaz* , 1922, 57, 412.

The toxic effects in Guinea-pigs of diets containing large proportion of Lathyrus sativus Bhagwat, K., *Ind. Jour Med Res.*, 1946, 34, 299

The toxic effects of Ticora (Khesari) on man Jacoby, H., *Ind. Med. Gaz.*, 1947 82, 122.

Study on "Lathyrism" a disease produced by Lathyrus sativus Linn. De, H N. & Datta, P.K., *Sci & Cult*, 1948, 14, 159.

LAURUS CAMPHORA Linn

Laurus camphora Nees (Cinnamomum camphora) V E Vorotzov, *Chem Abst* 5474, 1933

LAVENDULA BURMANNI Benth

Oil from the flowers and leaves of Lavendula burmanni—D D Kanga, *Journal of the Indian Institute of Science, Bangalore* Vol I, Pt VIII, pp 181 to 189, 1916

LAWSONIA INERMIS Roxb

Essential oil from the flowers of Camphine or Henna plant Antia M B & Kaushal, R , *Curr Sci* , 1950, 19, 284

LEUCAS ASPERA Spreng.

Studies on Leucas aspera Shirazi, A M J , *Ind Jour Pharm* , 1947, 9, 116.

LEGUMINOSAE

Seed oil of Formosan Plants Constituents of seed oils Leguminosae. Anona squamosa Kinzo Kafulice and Chintu Hata, *J Chem Soc Japan* 55, 369, 1934

LENS ESCULENTA Moench

Studies in nutrition value of Indian vegetable foodstuff Pt III Nutritive values of Lentil--Lens esculenta Moench , Cow pea—Vigna catjang Walp., and Aconite Bean—Phaseolus aconitifolius Jacq Niyogi, S P , Narayana, N. & Desai, B G , *Ind Jour Med Res* , 1932, 19, 859.

LEPIDIUM LATIFOLIUM Linn

The content of Lepidium latifolium and Cochlearia armoracia in essential oils S Rivos Goday and M Gomez Serranillos, *Chem Abst* 4432, 1945

LEUCAENA GIAUCA Benth

Leucenol, a definite principle obtained from the seeds of Leucaena glauca Benth Marcel Maseve, *Compt-Rend* 240, 890. 1937

Presence of Stachyose (Manotetrose) in the seed of Leucaena glauca Benth H Heressey and M Mascre, *J Pharm Chem* 1, 521, 1941

Structure of leucenine (Leucenol) from Leucaena glauca Bentham A F Biehel, *J Am. Chem Soc* 60, 1801, 1947

LINUM USITATISSIMUM Linn.

The hydrolysis of plytic compounds derived from seeds of hemp, horse bean, horse chestnut flax, wheat and embryos of rye W. Jarosza, *Chem Abst.* 5501, 1934.

LITSAEA Lamk.

Actinodaphne and Litsaea fats as raw material for a valuable new detergent S. V. Puntambekar, *Ind Forester*, 60, 707, 1934.

L. ZEYLANICA Nees.

Oil of Litsaea zeylanica, Nees. Rao, B. S.; *Journal of Indian Institute of Science, Bangalore*, **15A**, 1932, 71.

Bellary leaf oil Sharma, M. R. G., Varadhan C., Sastry, S. G *J. S. I. R.*, 1953, **6B**, 243.

LOBELIA Linn.

Amino nueteric determination of alkaloids in belladonna, lobel herb and harmel roots, G. Ya Kait, *Chem. Abst.* 2996, 1942.

The origin of Haustoria in the value of Lobelia. Maheshwari, P *Curr. Sci.*, 1944, 13, 186

L. INFLATA Linn.

Indian substitute for the drug Lobelia inflata *Ind. Jour. Pharm* 1944, 6, 105.

L. NICOTIANAEFOLIA Heyne.

Pharmacognostic studies on Lobelia nicotianaefolia Heyne. Datt S C. & Bal, S. N , *Sci. & Cult* , 1944, 10, 260

Lobelia nicotianaefolia Heyne as substitute for Lobelia inflata Lin Mukerji, B. & Ghosh, S K., *Curr Sci.*, 1945, 14, 198.

L. PYRAMIDALIS Vahl.

Pharmacognostic investigations on Lobelia pyramidalis Vahl. substitute for Lobelia inflata Linn. Amiya Datta & Datta, S. C *J. Sci. Indl. Res.*, 1951, **10B**, 218.

LOLIUM TEMULENTUM Linn.

Piosonous foodgrain wheat mixed with Lolium temulentum. Greva S. D. S. & Bhandari, P. N , *Ind. Med Gaz* , 1946, 81, 294.

LORANTHUS LONGIFLORUS Den.

Parasitism of Cuscuta reflexa Roxb. & Loranthus longiflorus De Sheriar, K. C., *Sci. & Cult.*, 1951, 17, 218.

LUFFA ACUTANGULA Roxb

Luffa acutangula The Chemical and Pharmacological investigatio of Luffa seeds Grewal, K A & Kochhar, B. D , *Ind Jour. Med. Res* 1943, 31, 63

L. AEGYPTIACA Mill.

A study of the oils from the seeds of Luffa aegyptiaca, Benincas cerifera and Allium cepa. Phadnis, K. D., Rege, A V., Pishawika D. G., and Shah, S V., *J Univ. Bombay*, **17A**. 62, 1948.

L. AMARA Roxb.

Chemical constituents of Luffa amara Roxb Pt. I. Isolation a crystalline bitter from the seeds Chaudhry, G. R., Vishwa Nat Sharma & Siddiqui, S.; *J. Sci. Indl. Res.*, 1951, **10B**, 26.

L. CYLINDRICA Linn.

The crystalline principle of Luffa cylindrica Linn. Roemer, Martin, De, La Torre and A. C. Santos, *Chem Abst* 5593, 1939.

Saponins of Luffa cylindrica Linn. Roemer., Amparo Mendoza, Paraluman Cruz and Alfredo C. Santos, *Chem Abst.* 4915, 1941.

A chemical study of Luffa cylindrica with special emphasis on its saponins. Amparo. S. Mendoza and Alfredo C. Santos, *Chem Abst* 7652, 1941.

L. ECHINATA Roxb. & L. GRAVEOLENS Roxb.

Chemical examination of the seeds of Luffa graveolens Roxb. & Luffa echinata Roxb. Nigam, Ram Gulam Singh, Pandya, K. C & Tayal, J N, *Curr. Sci.*, 1949, 18, 451

LUPINUS (Tourn) Linn

Biological experimental study of the Lupine alkaloids A. Guillaume, *J. Pharm. Chem* 12, 335, 1930.

L. ALBUS Linn.

Phyto-Pharmacological experiments with Lupinus albus. Ludwig Johanning, *Chem. Abst* 167, 89, 1930

Hypoglucmic action of the seeds of Lupinus albus Orestin, G., *Chem Abst* 1511, 1941

The alkaloids of Lupinus albus L and its supposed antimalarial action Liberalli, C H, *Chem Abst* 4101, 1944

LUVUNGA SCANDENS Ham

On the constitution of natural coumarins isolated from Luvunga scandens Ham. Bose, P K & Miss Mukerji, A, *J Ind Chem. Soc.* 21 (1944), 181.

LYCOPODIUM.

Toxicity of the alkaloids of Lycopodium P. Oficialski; *Bull Sci. Pharmacol* 33, 470, 1937

Pharmacological action of certain Lycopodium alkaloids Henry M. Lee and Chen, K K, *J. Am Pharm. Assoc* 34, 197, 1945

L. CLAVATUM Linn.

Alkaloids of Lycopodium (Lycopodium clavatum L.). Achmatowie and W Uzieblo, *Chem Abst* 9092, 1938.

The alkaloids of Lycopodium species, Lycopodium clavatum L. Leo. Marion and Richard H. F. Manske, *Can. J. Res.* **22B**, 137, 1944

L. PHLEGMARIA Linn.

Lycopodium phlegmaria Linn., on occurrence of two sporangiate sporophyllus in. Ghosh, A. K., *Sci & Cult.*, 1941, 7, 410.

MABA NIGRESCENS Dalz.

Occurence of marginal strands in the leaf of Maba nigrescens Dalz Rao, T. A. & Kelkar, S. S , *Sci & Cult* , 1950, 16, 209

MAHONIA Nutt.

Structure of Neprotine A new alkaloid of Mahonia species Chatterjee, R & Guha, M P., *Sci. & Cult*, 1950, 16, 119

M. ACANTHIFOLIA Don.

The alkaloids of Mahonia acanthifolia Don. Chatterjee, R , Guha, M P , *Sci & Cult* , 1949, 15, 163.

MALLOTUS PHILIPPINENSIS Muell

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency XIII Kamala Cairns, J F & Mhaskar, K S , *Ind Jour. Med Res* , 1921, 9, 202

The constitution of Indian Kamala (Mallotus philippinensis) Sikhibhushan Dutt, *J Chem Soc* 2044, 1925.

The Constitution of Rottlerin from Indian Kamala Part II Dutt, Sikhibhushan, Puri, & Dhanraj, *J Ind Chem Soc* 5 (1928), 21.

Rottlerin V Bakshi, H S Ravi, S Jalota, Narang, K S. & Ray, J N., *Curr. Sci.*, 1939, 8, 165.

The constitution of Rottlerin Ray, J. N , Narang, K S. & Roy, B S., *Curr Sci* , 1939, 8, 558

The use of Kamala as an antioxidant of Ghee Govindarajan, S V. & Banerjee, B. N , *Curr Sci* , 1939, 8, 559.

The fatty oil from the seeds of Mallotus philippinensis Muell Arg. (N O Euphorbiaceae) Singh, Bawa Kartar, Saran Brij Mohan, *Curr Sci.*, 1942, 11, 360

The fatty oil from the seeds of Mallotus philippinensis Muell Puntambekar, *Curr Sci* , 1942, 11, 464

Kamala as a colouring agent for hydrogenated vegetable oil Its toxicity on white Rats. Gupta, J C & Chatterjee, M L ; *Sci & Cultr* , 1946, 11, 375

Quality of commercial Kamala and its treatment Subba Rao, V , *Ind Jour Pharm.*, 1947, 9, 81

The treatment of intestinal worms with the Indigenous drugs– Butea, Embelia and Kamala Mukerji. A. K , Bhaduri, N V., *Ind Med Gaz* , 1947, 82, 66.

Chemical Examination of the seeds of Mallotus philippinensis Muell Arg. Pt I. Analysis of drying oil and seed cake Aggarwal, J S , Bhatnagar, S. S , Parkash Narain & Karimullah, *J. Sci Indl. Res* , 1948, **7B**, 136.

The Chemical Constitution of the oxy-acid from the seed oil of Mallotus philippinensis Muell. Arg. (Kamala) Gupta, S. C , Vishwa Nath Sharma & Aggarwal, J S , *J. Sci Indl Res* 1951, **10B**, 76.

Chemical Examination of the seeds of Mallotus philippinensis Muell. Arg (Kamala) II Constitution of the unsaturated Hydroxy acid isolated from the oil Gupta, S. C , Sharma, V. N. & Aggarwal J S . *J. Sci Indl Res* , 1952, **11B**, 463.

Chemical Examination of the seeds of Mallotus philippinensis Muell Arg (Kamala) III—Constitution of Mamlolenic Acid isolated from the oil Gupta S. C., Gupta S S , Aggarwal, J S , *J S I R* 1953, **6B**, 240

MALVA ARBOREA Webb.

Oils from Althea officinalis and Malva arborea H Ya. Tropp, *Chem. Abst* 1597, 1936

M SYLVESTRIS Linn.

Chemical and pharmacological researches on Malva sylvestris Luigi Callegari, *Chem Abst* 1589, 1946.

MANGIFERA Linn

A note on the genus Mangifera Linn. Mukerjee, S. K , *Sci & Cultr.* 1942, 8, 92

M. INDICA Linn.

Mangifera indica. Chem. Examination of mango "Chep"—the exudation of the fruit of Vasistha, Shiam Kishore & Siddiqui, Salimuzzaman, *J Ind Chem Soc* 15 (1938), 110

Chemical Examination of Mango blossoms (Mangifera indica L , Flora) Jogendra Lall Ball & Salimuzzaman Siddiqui, *J Sci Indl Res* , 1948, **7B**, 100

Wild mangoes of India. Mukerjee, S K , *Sci & Cult* , 1950, 15, 464.

MANIHOT ESCULENTA Crantz.

Age of Cassava Plant for max. yield Chaterjee, K N , *Sci & Cult* , 1949, 14, 533.

M UTILISSIMA Pohl.

The poisonous constituents of sweet Cassawa (Manihot utilissima) and the occurrence of hydrocyanic acid in foods prepared from Cassawa B J. W Furnock, *J Trop Med Hyg* 43, 65, 1937

Therapeutically utilisable constituents of Manihot utilissima and Mercurialis annua or perennis Freedrich W Friese, *Chem Abst.* 2288, 1938

MANSONIA DIPIKAE C. S. P.

Mansonia dipikae C. S. Purkayastheo sp Nov A new species from Assam Jain, B. C., Iyer, B. H., & Guha, P C , *Sci & Cultr* , 1947, 12, 405

MARRUBIUM VULGARE Linn.

Comparative marrubiin content in Marrubium vulgare from European Vs. American seeds Adelia Mccrea; *J. Am Pharm Assoc* 19, 231, 1930

Maruburr, the bitter principle of hove bound (Marrubium vulgare) Alexander Lawson and E D. Euktice; *J. Chem. Soc.* 587, 1939.

Substance contained in Marrubium vulgare Gualtenio Seppi *Chem Abst.* 4893, 1947

MARTYNIA DIANDRA Glox

Fatty oil from the fruit of Martynia diandra (N O. Pedaliaceae) Shah, S. V., Airan J W. & Rege, A V., *Curr Sci.* 1942, 11, 291.

MATRICARIA Linn.

Biochemical studies of the glucosides of Camomile flowers. Charlev Beguim; *Pharm Acta Helv* 7, 332, 1932

Estimation of essential oil in comomile flowers. F. Gestirner, *Chem. Abst.* 5477, 1933.

Anti-inflammatory action of Camomile oil Wolfgang Henlner and Franz Grabe; *Arch. Expt Path Pharmakol* 171, 329, 1933

Biochemical studies of the glucosides of camomile flowers, comparison of samples of different origin collected in 1932. Ch Beguin, *Pharm. Acta Helv.* 10, 147, 1935

Camomile Orlando Gulminelli, *Chem Abst.* 1517, 1936

M. CHAMOMILLA Linn.

New results for the evaluation of the pure azulene content of Matricaria chamomilla and its preparations Kaiser, H. and Frez, H , *Chem Abst* 1441, 1939.

MELIA AZEDARACH Linn

Fatal intoxication by fruit from Melia azedarach Regelio, L , Cavrrabala, E.; *Chem Abst* 6951, 1939

Chemical Examination of the Bakayan fruit (Melia azedarch Linn.) Amir Chand, Chittaranjan Mitra & Salimuzzaman Siddiqui, *J Sci Indl Res* 1948, **7B**, 69

MELILOTUS PARVIFLORA Desf.

Powdry Mildew on Melilotus parviflora Desf. Mahmud, K. A. & Namen, K. G., *Sci. & Cult.*, 1952, 17, 473.

MENTHA AQUATICA Linn.

Studies in the genus Mentha. The non-volatile constituents of Mentha aquatica Linn. Samuel M. Gordon; *Am. J. Pharm.* 100, 433, 1928.

Mentha aquatica L. Crestoforo Masino, *Chem. Abst.* 2741, 1937.

M. VIRIDIS Linn.

Japanese peppermint oil. Oil of Japanese Mentha viridis L. Var Crispa Benth. Carvone Tetu Nagasawa; *Chem. Abst.* 219, 1940

MERCURIALIS AMNUA Linn.

Therapeutically utilisable constituents of Manihot utilisima and Mercurialis amnua or perennis Frcedrich. W. Friese; *Chem Abst.* 2288, 1938.

MESUA FERREA Linn.

Mesua ferrea Preliminary note on Mesuol. The bitter principle of. Dhanindra Bhushan Dutt, Deb, Narendra Chandra & Bose, Prafulla Kumar, *J. Ind. Chem. Soc* 17 (1940), 277

Refining of Nageshwar oil Gupta, A. C., *J. Sci. Indl Res.*, 1951, **10B**, 24.

MILLETTIA PACHYCARPA Benth.

Occurrence of Rotenone in Millettia pachycarpa Ghose, T.P. & Krishna, S , *Curr. Sci.*, 1937-38, 6, 57.

MIMOSA PUDICA Linn.

The chemical properties of the harmone from Mimosa pudica Hans, Fitting, *Chem Abst.* 8300, 1936.

Chemical Examination of the seeds of Mimosa pudica Linn. Pt I. Joti Sarup Aggarwal & Karimullah, *J Sci Indl Res* , 1945-46, 4, 80.

MIRABILIS JAIAPA Linn.

A note on starch in Mirabilis jalapa soluble in cold water Chaudhri, T. C., *Sci & Cultr.*, 1947, 12, 449

MOMORDICA CHARANTIA Linn.

Balsam pear seed oil (Momordica charantia L) Yoshiyuki Toyama and Tomotaro Tsu Chiya, *J. Soc. Chem. Ind. Japan.* 39, 220, 1936.

Preliminary study of an alkaloid like material obtained from Cundeamor or Momordica charantia. Lius Torres Diaz, *Chem Abst* 6134, 1936.

Chemical Examination of Karla plants Airan, J.W & Ghatge, N D , *Curr. Sci.* 1950, 19, 19.

M. DIOICA Roxb

The fatty oils from the seeds of Momordica dioica (N. O. Cucurbitaceae) Airan, J W & Shah, S.V., *Curr. Sci* , 1942, 11, 246

MONNIERA CUNEIFOLIA Michx.

Investigation of Indian medicinal plants Hydrocotyle asiatica, Vitex negundo and Monniera cuneifolia. Basu, N K , Lamsal, P P., and Singh, G.B., *Quart. J. Phar. Pharmacol* 20, 135, 1947.

MORINDA CITRIFOLIA Linn

Notes on the constituents of Morinda citrifolia. John Lionel Simonsen *J Chem. Soc* 561, 1920

MORINGA CONCANENSIS Nimm

Chemical investigation of seed oil of Moringa concanensis Patel, C B , *Curr Sci* , 1943, 12, 272.

M OLEIFERA Lamk

Availability of calcium in Lady's finger (Hibiscus esculentus), Cabbage (Brassica oleracea var. capitata), Drumstick (Moringa oleifera) and Amaranth Tender (Amaranthus gangeticus). Pt I. Availability of Calcium in vegetables determined by experiments on growing rats. Basu, K P & Ghosh, D , *Ind. Jour Med. Res* , 1943, 31, 29.

Availability of calcium in Lady's finger (Hibiscus esculentus), cabbage (Brassica oleracea var capitata), Drumstick (Moringa oleifera) & Amaranth Tender (Amaranthus gangeticus) Pt II Availability of calcium in vegetables determined by metabolism experiments on a human adult Basu, K P & Ghosh, D , *Ind Jour Med Res* , 1943, 31, 37

M PTERYGOSPERMA Gaertn

A preliminary note on the action of the alkaloids of Moringa pterygosperma (N O Moringae) Chopra, R N , *Ind Med Gaz* , 1932, 67, 128

Moringa pterygosperma (N O Moringae). Chopra, R N., Premankur D C & De, Nripendra Nath, *Ind Jour Med Res* , 1933, 20, 533

A Chemical examination of the bark of Moringa pterygosperma. Ghosh, S , Chopra, R N & Datta, A *Ind Jour Med Res.*, 1935, 22, 785

Antibiotic principles from Moringa pterygosperma Kurup, P A & Rao, P L., *Curr Sci* , 1950, 19, 54

A note of physiological and chemical findings of the active principle (Spirochin) of the Moringa pterygosperma Chatterjee, G. S , Mitra, S R., *Sci & Cult* , 1951, 17, 43

Chemical investigation of the fatty oil from the seeds of Moringa pterygosperma Subba Rao, B C , Rao, K N., Jois, H S , *J.I.C.S* , 1953, 7, 477.

MORINGACEAE

Notes on the affinities of Moringaceae. Chatterjee, D., *Sci. & Cult* , 1948, 14, 253

On the affinities of Moringaceae. Datta, R.M , Mitra, J N. & Chatterjee, D., *Sci. & Cult.*, 1949, 15, 115.

MORUS INDICA Linn

On the Hermaphrodite flowers of Morus indica Linn Datta, R M, *Sci & Cult*, 1938, 4, 301.

On the Hermaphrodite flowers of Morus indica Linn Datta, R M, *Sci. & Cult*, 1939, 5, 373

Fasciation in the Female inflorescence of Morus indica Linn Datta, R M, *Sci & Cultr.*, 1941, 6, 425

M. NIGRA Linn.

The blood sugar lowering action of the leaves of Morus nigra L Henri Leelve, *Chem Abst.* 1507, 1935

MUCUNA PRURIENS D C

Indian Medicinal plants Pt V Mucuna pruriens (N O Papelionaceae), Pt I Mehta, J C & Majumdar, D N, *Ind Jour Pharm*, 1944, 6, 92

MUNDULEA SUBEROSA Benth.

The pharmacological action of Cassine Croceum De and Mundulea suberosa Benth N Sapeika, *S African J Med Sci* 10, 51, 1945

MUNDULEA SUBEROSA Benth

Chemical constituents of Mundulea suberosa Th M Meyer, *Chem Abst* 6607, 1947

MURRAYA EXOTICA Linn & M KOENIGI Spreng

Essential oils from the leaves of Murraya koenigii (Spreng), Murraya exotica Linn and Murraya exotica Var Ovalifoliata (Ergler) A R Penfold and J. L Simonsen, *J Proc Royal Soc* 59, 146, 1925

Murrava exotica Natural glucosides Part I The constitution of the Glucoside present in Murraya exotica Praphela Kumar Bose & Miss Asima Mukerjee, *J Ind Chem Soc* 14 (1937), 489

MYRISTICA FRAGRANS Houlf

Essential oil from the leaves Nutmug (Myristica fragrans) M Meyer, *Chem Abst* 4549, 1941

Chemistry of Nutmeg—Aril (Jay-Patri) Ashok D Pishawikar, Pishawikar, D. G., *Curr Sci* 1953, 3, 81.

MYRSINE AFRICANA Linn.

Myrsine africana Linn. Active principles of M. africana Krishna, S. & Varma, B. S ; *J. Ind Chem Soc* 13 (1936), 115

The active principle of Kurjan seed (Myrsine africana) Annon; *Chem. Abst.* 36, 319, 1938

NARCISSUS TAZETTA LINN.

A new alkaloid in the bulbs of Narcissus tazetta L Yosiziro Kihara; *J. Agr. Chem. Soc. Japan*, 15, 128, 1939

NARDOSTACHYS JATAMANSI D C.

Chemical constituents of Nardo stachys jatamansi. Pt. I. Isolation of a crystalline acid and an essential oil Chowdhry G R, Vishwa Nath Sharma & Siddiqui, S., *J Sci Indl Res.*, 1951, **10B**, 48

NERIUM ODORUM Soland.

Chemical examination of the bark of Nerium odorum (Soland.) Gajanan P Pendse and Sikhibhushan Dutt, *Bull Acad Sci. United Provinces.* 3, 209, 1934

Chemical Examination of the root of "Nerium odorum". Part I Garde, S.T, *Journal of the Indian Institute of Science, Bangalore,* Vol I, Part XVIII, pp 181 to 189, 1910

Pharmacological action of odorin, a glucoside of Nerium odorum. Isao Niimoto, *Chem Abst.* 4734, 1940

N. OLEANDER Linn

A pharmacological study of folinerin, a new glucoside from Oleander leaves. Italo Simon, *Chem Abst* 4799, 1943.

Medicinal use of the glucoside of Nerium oleander L. Gibert Queralto, *Chem. Abst.* 5786, 1943.

Pharmacology of the new Cardiac glucoside folinerin (From Nerium oleander) Effect on the cardiovascular system. V I Simon, *Chem. Abst* 5299, 1944

Glucosides of Nerium oleander of the Argentine, Oscare A Rossi and Jaun A. Izquierdo; *Chem Abst* 7940, 1948

NICOTIANA Linn

The occurrence of crossing over in Nicotiana species Hybrids Kostoff Dontcho, *Curr Sci*, 1933-34, 2, 370

Tobacco seed oil Fatty acid comp of. Rao, Rao & Venkateswarulu, *J. Ind Chem. Soc.* 20 (1943), 374

Glyceride comp of Tobacco seed oil Venkata Rao, C Narsingarao, M & Venkateswarlu, A, *J. Ind Chem. Soc* 21 (1944), 249

Manufacture of 40% Nicotine as a soln. of nicotine sulphate in water Subramanian, T. S & Varia, M. C, *J Sci Indl Res.*, 1944 45, 3, 143

N. RUSTICA L. & N GLAUCA Grab

Nicotine and citric acid content in the progeny of the Allopolyploid Hybrid Nicotiana rustica L X. N glauca Grab Kustoff, Dontcho; *Curr. Sci*, 1939, 8, 59

N. TABACUM Linn

Australian Tobacco Industry, article on Charles Lynch, *J. Sci Indl. Res.*, 1945-46, 4, 760.

Products of Tobacco industry Tobacco seeds and its utilization. Mahant, S. D. & Pandit P. N., *J. Sci. Indl. Res.* 1948, 229, **7A.**

Cytogenetics of Nicotiana tabacum Var. Virii resistant to the common Tobacco Mosaic virus Kostoff, Dontcho, *Curr. Sci.* 1948, 17, 315.

Tobacco Research in India. (Article); *J Sci. Indl. Res.*, 1950, **9A,** 69.

Utilization of Tobacco seed and Safflower seed oil in varnish and paints. Pt I, Sharma, P G, Budhiraja, N C, & Aggarwal, J. S, *J Sci Indl. Res.*, 1951, **10B,** 33.

NUPHAR LUTEUM Sibth & Sm.

The glucoside of Nuphar luteum Marjan Hulajewski and Jerzy, Modrakovski, *Chem. Abst* 5106, 1937

NYMPHAEA (Tourn) Linn

The pharmacological action of nymphacin Zderek Kocher; *Chem Abst* 9282, 1938

N ALBA Linn

Alkaloids from flowers of Nymphaea alba Jerzy, Modrakoswki; *Chem Abst* 9279, 1938

N. LOTUS Linn.

Axial floral Prolification and Metamorphosis of stamens in a Flower of Nymphaea lotus Linn Bose, P K, *Sci & Cultr*, 1940, 5, 636.

OCIMUM Linn.

Development of essential oil industry (O lemon, Ocimum basilicum, Ocimum, canum, O sanctum, Lemon grass, Motia, Palmarosa grass) Jitandra Nath Rikshit, *Sci & Cultr*, 1939, 5, 108.

O BASILICUM Linn

The volatile oil of ocimum of North India—Rakshit J N., *The Perfumery and Essential Oil Record,* March, 1938

Ocimum basilicum Pt I & II A chemical study of the oil and antibacterial properties Khorana, M L, Miss M. B Vangikar, *Ind Jour Pharm* 1950, 12, 132 & 134

O CANUM Sims

Oil of Ocimum canum—B. S. Rao and Co-workers; *Perfum. Record,* 28, 1937, 412.

Ocimum canum oil of North India—Rakshit, J N, *The Perfumery and Essential Oil Record,* October, 1938

Chemical examination of Ocimum canum Sims. Mehta, R. C, Mehta T. P., *Curr Sci*, 1943, 12, 300

O. GRATISSIMUM Roxb.

Essential oil from Ocimum gratissimum Roxb Nayab, U. G. & Guha D. C.; *J. Ind. Chem. Soc.* 29 (1952), 203.

O KILIMANDSCHARICUM Guerke

Camphor production from Ocimum kilimandscharicum Guerke Ribeiro, D J., *J. Sci Indl Res* 1950, **9B**, 230.

O SANCTUM Linn

Ocimum sanctum (*Ind Acad. Sc.* Voc. IX, Jan 1939, pp 72-77 Sikhi Bhusan Dutt,

A seedling blight of Ocimum sanctum Linn. caused by Rhizodonia solani Kuhn Mohmud K A, *Sci & Cult*, 1950, 16, 161.

ODINA WODIER Roxb.

Gum Jeol (Odina woider Roxb) *J Ind Chem Soc* 25 (1948).

(i) Solution & purification of Mukerjee & Bannerjee, 59

(ii) Hydrolysis of & sugar in Mukerjee & Banerjee, 63

(iii) Electrodialysis of. Mukerjee, 323

(iv) Dependence of Viscous electro-chemical prop of aquous solution of, on concentration. Mukerjee & Rohatgi, 339

(v) Influence of neutral salts on some electrochemical & Viscous properties of the solution of Mukerjee & Rohatgi, 531

Aldobionic acid from Gum jeol & its hydrolysis. Mukerjee & Dutta, *J Ind Chem Soc* 25 (1948), 113

OLDENLANDIA BIFLORA Hk. f

Chemical examination of Oldenlandia biflora R N S Clauba & Tewari, J D, *J Ind Chem Soc* 29 (1952) 386.

OLEA Linn

On the occurrence of Sclereide in the Genus Olea Linn Rao, T A, *Sci & Cultr*, 1948, 13, 299

ONONIS SPINOSA Linn

New constituent from the root Ononis spinosa L F Neuwald, *Arch Pharm* 277, 130, 1939

ORIGANUM VULGARE Linn

Volatile oil of Origanum vulgare B N Rutovskii, K A Guseva and L Koroleva, *Chem. Abst* 5476, 1933

Essential oils of Origanum vulgare Linn Var formosanum Yasuji Fugita, *J. Chem Soc Japan*, 57, 574

Essential oil of the flowering tops of Origanum vulgare infected with Eriophyes thomasi R Salgues, *Chem Abst* 4270, 1936

OROXYLUM INDICUM Vent.

Natural flavones Part II On the colouring matter of the bark of Oroxylum indicum Vent. Prafulla Kumar Bose & Sachindra Nath Bhattacharya, *J Ind Chem. Soc* 15 (1938) 311.

Baicalein from the seeds of Oroxylum indicum Vent Mehta, C. R & Mehta, T P , *Curr Sci* , 1943, 12, 274.

Constitution of Oroxylin Sastri, V.D N. & Seshadri, T R., *Curr Sci* , 1946, 15. 235

Tetuin, A Glucoside from the seeds of Oroxylum indicum Vent Mehta C R , Mehta T P , *Curr Sci* 1953, 4, 114

ORTHOSIPHON STAMINEUS Benth.

Preliminary study of Orthosiphon stamineus Benth. Teng Hang Tang and Chao Hsii, *J Chinese Chem Soc* 7, 111, 1940.

OXYTROPIS THOMSONI Benth

A note on Oxytropis thomsoni, Benth. Siddiqui, R.H , Basha, S. K., *Ind Jour Pharm* , 1944, 6, 83

PANDANUS ODORATISSIMUS Roxb.

Kewda Perfumes Vol IV, No 1—Sadgopal,—*Indian Soap Jour.*, July 1936

Kewda—Sadgopal,—*Soap Perfumery and Cosmetics*, May, 1937

Pandanus oil--Sadgopal, *Soap Perfumery and Cosmetics*, 10, 1937, 396 Essential oil from flowers of Kewda, Pandanus odoratissimus. S S Deshpande, *Calcutta University Press*, 1938

Kewda (Pandanus odoratissimus Essential oil from S S Deshpande, *J Ind Chem Soc* 15(1938), 509

PAPAVER SOMNIFERUM Linn

A preliminary note on addiction to "Post" (unlanced capsules of Papaver somniferum) Chopra, R N , *Ind Med Gaz.*, 1930, 65, 361

Addiction to "Post" (unlanced capsules of Papaver somniferum) in India Chopra, R N , Grewal, K S , Chowhan, J S & Chopra, G S , *Ind Jour Med Res* , 1930, 17, 985

Addiction to "Post" Part II Chopra, R. N & Ghosh, N N , *Ind Jour. Med Res* , 1931, 19, 415

Opium and Albuminuria Chopra, R N & Bose, J P , *Ind Med Gaz* , 1931, 66, 299

Administration of Opium to infants in India Chopra, R N , Chopra, G S , *Ind. Med Gaz.*, 1934, 69, 489

Indian Opium Its principal groups of chemical constituent Rakshit, J.N., *Sci & Cult* , 1942, 8, 16

Opium alkaloids Recent development in India Mukhopadhyay, B K , *J Sci. Indl. Res* , **8A**, 118, 1949

Papaverine from Indian opium Mukhopadhyay, B. K & Partha-sarthy, C , *J. Sci Indl. Res.*, 1949, **8B**, 23

PARIS POLYPHYLIA Sns.

A preliminary note on the Glucosides from Paris polyphylla Sns. Chopra, R. N , Gupta, J C , Bose, B. C. & Chopra, I C., *Ind. Jour. Med Res* , 1942, 30. 103.

PEGANUM HARMALA Linn

Oil from Peganum harmala H. Ya Tropp, *Chem Abst.* 1598, 1936.

On Peganine isolated from the flowers of Peganum harmala L. A Rosenfile, D G , Kolismikov, *Chem Abst* 9306, 1939

Amino neuteric determination of alkaloids in belladonna, lobelia herb and harmel roots, G Ya Kait, *Chem Abst* 2996, 1942

The action of Peganum harmala and of its alkaloids on the circulation. Arsenio Fraile Ovejero; *Chem Abst* 3214, 1947

Peganine from the blossoms and stems of Peganum harmala L Remarks on the papers of E Spath on Peganine. A D. Rosenfeld and D G Kolesnfeld and D.G Kolesnikov, *Ber* 28, 1704.

PENICILLIOPSIS CLAVARIAEFORMIS Solms Laub

Penicilliopsis clavariaeformis Solms Laub, A new record in India, Ganguly, D & Ghosh, T , *Sci & Cult.*, 1947, 12, 553.

PENTAPETES PHOENICEA Linn

A note on the development of the female gametophyte in Abroma angusta L and Pentapetes phoenicea L , Banerji, I , *Curr. Sci* 1941, 10, 30

PERANOSPORA PARASITICA

Medicinal action of Capsella bursa-pastoris and also of its parasites, Cystopus candidus and Peranospora parasitica W Harste, *Arch Pharm.* 266, 133, 1928

PEROWSKIA ATRIPLICIFOLIA Benth.

The Essential oil from the flower heads of Perowskia atriplicifolia, Benth, M G Rao, *Journal of the Indian Chemical Society, Calcutta* 1926, Vol III, p. 141.

PHALARIS MINOR Retz

Phalaris minor Retz A new promising winter fodder grass. Gandhi, R T , Wadhwa, N D , *Sci & Cult* , 1951, 17, 113

PHASEOLUS ACONITIFOLIUS Jacq

Studies in nutrition value of Indian vegetable foodstuff Pt III Nutritive values of Lentil-Lens esculenta, Moench., Cow pea - Vigna, catjang, Walp. and Aconite Bean - Phaseolus aconitifolius, Jacq , Niyogi, S P , Narayana, N & Desai, B G , *Ind Jour Med. Res* 1932, 19, 859

P RADIATUS Linn

Constituents of Phaseolus radiatus L Var Aureus Prain E Miyamichi and H Yamada, *J Pharm. Soc Japan*, 50, 1095, 1930.

Saqonin glucoside in adzuki beans (Phaseolus radiatus L Var. Aureus Prain) E. Miyamichi and S Onishi, *J. Pharm Soc. Japan.* 52, 168, 1932.

PHEBALIUM ARGENTUM Sm.

Phebalium argentum, occurence of Psoralen in P. K Bose & Finlayson, *J Ind Chem Soc* 15 (1938), 516

PHYLLANTHUS EMBLICA Linn

Ascarbic acid contents of some Indian plant material M Damodaran and M Srinivasan, *Proc Ind Acad Sci* **2B**, 377, 1935.

Indian Gooseberries (Phyllanthus emblica Linn) as a source of Vit C Giri, K V, *Ind Jour Med Res*, 1939, 27, 429

A new technique for propagating Aonla (Phylanthus emblica) Singh, L B, *Sci & Cult*, 1952, 17, 345

P NIRURI Linn

Bitter principles of Phyllanthus niruri G V Krisana Murti and T R Seshadri, *Proc Ind Acad Sci* **24A**, 357, 1946

Investigation of quinine in Phyllanthus niruri L Richardo Martin Serra, *Chem Abst* 1812, 1947

PHYSOCHLAINA PRAELTA (Don). Miers

Chemical investigation of Physochlaina praelta Miers Handa, K L, Nazir, B N, Chopra, I C & Jamwal, K S, *J Sci Indl. Res*, 1951, **10B**, 182

Chemical investigation of the roots of Physochlaina praelta Miers Handa, K L, *J Sci Indl. Res*, 1951, **10B**, 234

Chemical investigation of Physochlaina praelta Pt II Handa, K L & Onkar Nath Channa, *J Sci Indl. Res*, 1952. **11B**, 505

PICRASMA EXCELSA Swartz

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb, Melia azadirachta Linn, Punica granatum Linn, Picrasma excelsa Swartz Vernonia anthelmintica Willd, Cocos nucifera Linn (coconut) Cairns, J F. & Mhaskar, K S, *Ind Jour Med. Res*, 1923, 11, 353

PICRORHIZA KURROA Royle ex Benth

Some common indigenous remedies, Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum, Chopra R. N. and Ghosh, S., *Ind Med Record* 55, 77, 1935

Comparative pharmacognosy of Gentiana kuroo Royle & Picrorhiza kurroa Royle. ex Benth. Bal, S N. & Datta, S C, *Ind Jour Pharm*, 1945, 7, 109.

Chemical Examination of Picrorhiza kurroa Benth. Part I Rastogi, R P, Vishwa Nath Sharma & Siddiqui, S, *J Sci Indl Res*, 1949, **8B**, 173.

Chemical investigation of Picrorhiza kurroa Benth Attempted synthesis of Kutkin Rastogi, R P, Sen, A B, *J I C S*, 1953, 7, 514

PIMPINELLA SAXIFRAGA Linn

Constituents of the roots of Pimpinella saxifraga Fritz, Wessely and Editha Nadler, *Monatsh.* 60, 141, 1922

Constituents of the roots of Pimpinella saxifraga Fritz, Wessely and Ferdinad Kallal, *Monatsh*, 59, 161, 1932

PINUS Linn

A plea for the distillation of Pine needle oil in India Puran Singh, *Indian Forester*, 1914, Vol XL, No 10

Oleum trebinthinae *Ind Jour Med Res*, 1920, 8, 125

Turpentine oil - H M Hulsny and E. P. Watson, *Quarterly Journal of the Indian Chemical Society*, 3, 1926, 258

The constituents of some Indian Essential Oils Part XVII - Derivatives of Abietic Acid Rau M Gopal and Simonsen J L, *Journal of Indian Institute of Science, Bangalore*, 1926, Vol 9, Part IV, pp 111-116 Continued from Indian Forest Records

The Pinene content of Indian Turpentine, Mulany H M & Watson, E R, *J Ind Chem Soc* 3 (1926), 258.

The preparation of Camphor from Pinene Mulany, H M & Watson, E. R, *J Ind Chem Soc*, 3 (1926), 253

The Pinene content of Indian Turpentine Mulany and Watson, *Journal of the Indian Chemical Society, Calcutta*, 1928, Vol III, p. 258.

Some pharmacological action of Sandal wood oil Vitivert oil and Pine needle oil Tameno Okanishi, *Chem Abst* 658, 1930

Vitamin C in pine needles Iyengar, N. K, Bose, B. C & Mukerji, B, *Ind Jour Med Res*, 1944, 32, 165

On the possibility of using oil of Turpentine for the treatment of Scabies Roy, D N & Ghosh, S M., *Ind Med Gaz* 1944, 79, 589.

Abietic acid from Indian Rosin Ghatak, N & Mitra, S P., *J Sci Indl Res* 1945-46, 4, 653

Studies in Indian Turpentine—production of p cymene (3) Sindhi, H K; *Sci. & Cult*, 1949, 15, 202

P EXCELSA Wal.

The constituents of some Indian Essential Oils Part II The Essential oil from the oleo-resin of Pinus excelsa Simonsen, J L and Rau M Gopal, *Indian Forest Records*, 1922 Vol IX, Part IV, pp. 6 - 12.

The leaf oil from Pinus excelsa. Simonsen, J.L. *Indian Forest Records*, Vol. IX. Part IX, 1922.

•Indian resin oil from Pinus excelsa. Patel, I.S. & Guha, P.C., *Curr Sci*, 1950, 19, 128.

P GERARDIANA Wall.

The constituents of some Indian Essential Oils Part IX—The Essential oil from the oleo-resin of Pinus gerardiana, Wall Simonsen, J L, *Indian Forest Records*; 1923, Vol IX, Part VIII, pp. 5-8

Oil from Pinus gerardiana Chilgoza oil Part I. Hardikar, S.D. *J. Ind. Chem Soc* 5 (1928), 69.

P KHASYA Royle.

Turpentine oil and resin from Pinus khasya (Khasi Pine, Dieng Kesh or Saral). Saikia, B., Das, P.K. & Roy, B.K, *Curr. Sci.*, 1951, 20, 275

P. LONGIFOLIA Roxb.

Turpentine oil from Pinus longifolia (Chir)—Puran Singh; *Indian Forest Records*, 1913, Vol. IV, Part I.

The constituents of Indian Turpentine from Pinus longifolia Roxb. John Lionel Simonsen, *J. Chem Soc.* 570, 1920.

The Constituents of Indian Turpentine from Pinus longifolia Roxb Part I Simonsen, J.L., *Trans Chem Soc.* 1920, Vol. 117, p. 570.

The Constituents of Indian Turpentine from Pinus longifolia Roxb. Part II Simonsen, J L and Gopal Rao, M, *Trans Chem. Soc.* 1923, Vol 123, p. 549.

The Constituents of Indian Turpentine from Pinus longifolia Roxb. Part III. Simonsen, J L, *Trans Chem Soc.* 1923, Vol. 123, p 2642.

The Constituents of Indian Turpentine, from Pinus longifolia Roxb. John Lionel Simonsen, *Jour. Chem Soc.* 2642, 1923

The Constituents of some Indian Essential Oils Part XVII Abietic Acid from the resin of Pinus longifolia, Roxb (Preliminary note) Rau M Gopal and Simonsen, J L, *Indian Forest Records*, 1925, (Chemistry Series) Vol. XI Part VI.

The Constituents of Indian Turpentine from Pinus longifolia, Part IV Pillay, P P and Simonsen, J.L. *Journal of the Indian Institute of Science, Bangalore*, 1928, Vol. **11A**, Part XV pp. 200-205.

The Constituents of Indian Turpentine from Pinus longifolia Roxb. John Owen and John Lionel Simonsen, *J. Chem. Soc.* 3001, 1931.

On the carenes in Indian Turpentine Oil from Pinus longifolia Roxb. Verghese, James, *Curr. Sci*, 1950, 19, 212.

Turpentine oil from Pinus longifolia Roxb. James Verghese & Gulati, K C., *J. Sci Indl. Res.*, 1951, **10A**, 112.

Liquid phase production of *p*-cymene from Indian Turpentine, oil of Pinus longifolia (Roxb.) & 3-carene. James Verghese & Lourdu M. Yeddanapalli; *J. Sci Ind. Res.*, 1951 **10B**, 100.

Studies in oil of Pinus longifolia Roxb. II. Aromatization of 3-carene to *p*-cymene James Verghese & Lourdu M. Yeddanapalli, *J. Sci. Indl Res.*, 1952, **11B**, 36.

Studies in oil of Pinus longifolia Roxb III—Two directional Cyclopropane ring of 3-carene in Sulphur Aromatization. Yeddanapalli, L.M , James Verghese , *J S I R.* 1953. **3B**, 121.

Studies in oil of Pinus longifolia Roxb II—Isomerization of 3-carene & Terpinene. Yeddanapalli L M., Krishnan T S., Verghese J; *J S I R* 1953, 38, 122

PIPER BETLE Linn.

Studies in human nutrition Pt IV Availability of Calcium ingested in the process of Chewing Betel-leaves with lime Basu, K.P.; Basak, M N & De, H.N., *Ind Jour Med Res* , 1942, 30, 309.

Tipburn of Piper betel in the Central Provinces. Asthana, R P. & Mahmud, K A , *Curr. Sci.*, 1944, 13, 234

The areca and the betel. Andre Mericier, *Chem. Abst* 956, 1946.

On the control of *Rhizoctonia*, root-rot of Pan (Piper betel L.) Chowdhury, S , *Sci. & Cult.*, 1948, 13, 507.

Propagation of Pan (Piper betel L.) in the Northern districts of the Central Provinces. Mahmud, K A., *Sci. & Cult* , 1950, 15, 324.

The antibacterial principle of Betel leaf. A preliminary note. Pai, Narayana & (Miss) Irani, R J., *Ind Med Gaz.*, 1950, 85, 302.

P. CUBEBA Linn

Constitution of some Indian essential oils, essential oil from the fruit of Piper cubeba Linn. B S Rao, V P. Shintre and J L. Simonsen, *J Soc. Chem Ind.* 74, 92, 1928.

The constituents of some Indian Essential Oils. Part XXIII—The Essential Oil from the fruits of Piper cubeba, Linn Rao B Sanjiva, Shintre, V P and Simonsen J L , *Journal of the Indian Institute of Science, Bangalore,* 1928, Vol **11A**, Part XV-1, pp 187-194.

P NIGRUM Linn

The Nitrogen complex of Indian foodstuffs condiments. Pt I. Black pepper (Piper nigrum) Narasimhamurthy, G. & Ranganathan, S , *Ind. Jour. Med Res* , 1937, 25, 373.

PISTACIA INTEGERRIMA Stew.

Chemical investigation on Kakra Singi Karimullah, Subba Rao, V. & Uma Shankar, *J. Sci. Indl Res.*, 1944-45, 3, 423.

A note on the crystalline constituents of (Pistacia integerrima Stew. N O. Anacardiaceae) Kakra Singi. Ghose, S.K., *Sci. & Cultr.*, 1945, 11, 46.

Chemical investigations of Kakra Singi (Pistacia integerrima). The essential oil. Karimullah & Uma Shankar ; *J. Sci. Indl. Res.*, 1946, 5, 60.

Essential oil from the galls of Pistacia integerrima. Baslas & Deshpande, *J. Ind Chem. Soc.* 27 (1950), 441.

P TEREBINTHUS Linn.

Ethereal oil from the resin of Pistacia terebinthus. A Tsatsas, *Chem. Abst* 5989, 1939

PITHECOLOBIUM SAMAN Benth

Composition of 'Rain Tree' fruit Rao, V R., Noshir, N, *Curr. Sci.*, 1946, 15, 250.

PLANTAGO MAJOR Linn.

A holloside extracted from the seeds of Plantago major L and Plantago ovata Forsk. (P Isphagulata Roxb), N Wathez and H. Hans, *Chem Abst* 4849, 1945.

Wound healing preparations from plantain (Plantago major) leaf. R.K Ahev, *Chem Abst.* 2210, 1947

P. OVATA Forsk

Plantago ovata, ispaghul in chronic diarrhoeas and dysenteries. Chopra, R N, *Ind Med. Gaz*, 1930, 65, 428

A holloside extracted from the seeds of Plantago major L and Plantago ovata Forsk (P isphagulata Roxb), N Wathez and H Hans, *Chem Abst* 4849, 1945

PLUMBAGO ROSEA Linn

Chemical examination of the root bark of Plumbago rosea Linn. M G Tumin Katti and V N Patwardhan, *J. Ind Inst Sci.* **15A**, 9, 1932

P ZEYLANICA Linn.

The pharmacological action of Plumbago zeylanica and its active principle (Plumbagin). Bhatia, B.B & Lal S., *Ind. Jour Med. Res*, 1933, 20, 777

PLUMERIA ACUTIFOLIA Poir

The aerial roots of Plumeria acutifolia Poir. Banerji, I, *Curr Sci* 1943, 12, 85

PODOPHYLLUM Linn.

Components of Indian Podophyllum. Seshadri, T.R. & Subramanian, S.S., *J. Sci. Indl. Res* 1950, **9B**, 137.

A revision of the structure of Podophyllotoxin and picropodophyllin Chatterjee, R., Chakravarti, S C, *Sci & Cult* 1951, 17, 136.

POLYGALA CHINENSIS Linn

Polygala chinensis Linn as substitute for official senega Hossain, T., Guha, R.C. & Mukerji, B.; *Sci. & Cult.* 1943, 9, 167

The Senegas of Indian Market. Gupta, B. & Bal, S.N., *J. Sci. Indl Res*, 1952, **11B**, 116.

POLYGONUM Linn.

Anthracenic derivatives in the genera Polygonum and Rumex. Maurin, E., *Bull. Sci. Pharmacol.* 33, 138, 1926.

PONGAMIA GLABRA Vent.

The oil of Pongamia glabra. Beal, C D. and Katti, M.C.T.; *J. Am. Pharm Assoc.* 14, 1086, 1925

Pongamia glabra Leaf-Gall-Farmer. Cherian, M C., *Curr. Sci.*, 1934-35, 3, 564.

Some common indigenous remedies, Picrrorhiza kurrooa, Erythrina indica, Sanseviera zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum. Chopra, R.N. and Ghosh, S., *Ind. Med. Record* 55, 77, 1935.

Pongamia glabra. A note on the occurrence of Behenic acid in the oil from the seed of. Manjunath, B L. & Shanker Rao, M.S., *J. Ind. Chem Soc* 15(1938), 653.

Pongamia glabra. A note on the occurrence of free fatty acid in the cake of Subba Rao, N.V. & Veerbhadra Rao, J., *J. Ind. Chem. Soc.* 175(1940), 26.

POTENTILLA ANSERINA Linn. & P. ARGENTEA Linn

The comparative activity of Potentilla anserina L. and Potentilla argentea L. on the isolated guinea pig uterus. Youngken, H and Fischer, E.B , *Am J. Pharm.* 114, 417, 1943.

PRANGOS PABULARIA Lindl.

Essential oil of Prangos pabularia Lindl. L. Grach , *Chem. Abst.* 7529, 1940.

PREMNA INTEGRIFOLIA Linn.

Premna integrifolia Linn , Chemical investigation of. Basu, N K. & Joneja, A.N., *Ind. Jour Pharm* 1949, 11, 191.

PRINSEPIA UTILIS Royle.

The fatty oil from the seeds of Prinsepia utilis. Puntambekar, S V.; *J. Ind Chem. Soc.* 19(1942), 183

PRISTIMERA INDICA Willd.

Pristimerin, the antibacterial principle of Pristimera indica Isolation, Toxicity and antibacterial action. Bhatnagar, S.S., Divekar, P V., *J. Sci. Indl. Res* , 1951, **10B**, 56.

Pristimera indica—A source of Dulcitol. Bhatnagar, S.S , & Divekar, P.V.; *J. Sci. Indl. Res.*, 1951, **10B**, 117.

PROSOPIS JULIFLORA D.C.

Antibiotic activity of ext of Prosopis juliflora. Shankarmurty, P. & Siddiqui, S., *J. Sci. Indl. Res.*, 1948, **7B**, 188.

PRUNUS ACUMINATA Hk. f.

Chemical investigation of the bark of Prunus acuminata. Chakravarti, D., Kundru, N., *Sci & Cult.* 1948, 14, 36.

P. AVIUM Linn.

Pharmacological studies on Prunus avium. Detection and determination of carbohydrates present in an infusion of stems of Prunus avium L Gaetano Di Manggio , *Chem Abst.* 566, 1947

P CERASUS Linn.

Indian Wild Cherry bark Kapoor, L. D. & Handa, K. L., *Cur. Sci*, 1948, 17, 54.

P. NEPALESIS Koch.

Chemical investigation of the bark of Prunus nepalenis Chakarvarti, D. & Momen, S A , *Sci. & Cult.*, 1950, 15, 329.

P. PUDDUM Roxb.

Isolation of a new flavone from the bark of Prunus puddum (N. O. Rosaceae), Chakravarti, D. & Ghosh, R. P., *Sci. & Cultr.*, 1943, 8, 463.

Isolation of a new iso-flavone from the bark of Prunus puddum (N. O Rosaceae), Chakravarti, D. & Bhar, C. N., *Sci. & Cultr.*, 1943, 8, 498.

Identity of Puddumetin and Genkwanin, Venkataraman, K., *Sci & Cultr.*, 1943, 9, 45.

Crystalline components of the bark of Prunus puddum Roxb. Part I. Identity of Puddemtin with Genkwanin. Chakravarti, G D. & Ghosh, R. P', *J. Ind. Chem Soc* 21 (1944) 171

Prunus puddum Roxb Crystalline components of Chakravarti & Behari, *J. Ind. Chem Soc* 22 (1945), 301.

Prunus puddum Roxb. Crystalline components of the bark of. Chakravarti, Kundu & Ghosh, *J. Ind Chem Soc* 25 (1948), 329

Prunus puddum. Isolation of Sakuranin from the bark of. Chakravarti & Sen *J. Ind. Chem. Soc* 27 (1950), 148

PSIDIUM GUAJAVA Linn

Effect of harmones on the rootage of Psidium guajava L Singh, S N. *Sci. & Cult.*, 1950, 16, 198

PSILOTUM TRIQUERUM Sw.

On the occurrence of Psilotum Sw. in the East Godawri Distt. Venkateswarlu, V., *Sci. & Cultr.*, 1943, 9, 166.

PSORALEA CORYLIFOLIA Linn.

Psoralea corylifolia (Babchi). Its constituents, their pharmacological action and therapeutic properties. Chopra R. N., Chatterjee, N. R., *Ind. Jour. Med. Res.*, 1927, 15, 11.

Psoralea corylifolia. Chemical examination of the seeds of Part I Jois, H. S , Manjunath, B L & Venkta Rao, S., *J. Ind Chem Soc.* 10 (1933), 41

Oil of Psoralea corylifolia—T. R. Seshadri and C. Venkat Rao, *Proc. Indian Acad. Sci. Sect* **5A,** 1937, 351

Components of Psoralea corylifolia Linn. T R Seshadri and C. Vekata Rao, *Proc Ind Acad. Sci* **5A,** 1937, 351

Nicotinic acid in Bavchi (Psoralea corylifolia - Bharucha, F. R , *Sci & Cult* , 1941, 7, 169

Histological study of fruit and seed of Psoralea corylifolia Linn. Prasad, G. S., *Ind Jour Pharm* , 1944, 6, 61.

Chemical examination of the seed of Psoralea corylifolia Linn. (Bebchi) Kamler Kinkar Chakravarti, Anil Kumar Bore & Salimuzzaman Siddiqui, *J. Sci. Indl. Res* , 1948. **7B,** 24.

PTERIS ACQUILINA Linn.

Poisonous properties of bracken (Pteris acquilina) G. D. Shearar, *Chem Abst* 6643, 1946

PTEROCARPUS MARSUPIUM Roxb

The Phlobatannins of Kino and buteagums G V Krishnamurti and T R Seshadri, *Proc Ind Acad Sci* **22A,** 134, 1945

A preliminary note on the action of Pterocarpus marsupium Roxb. on blood sugar Ojha, K N , Parba, P R & Venkatachalam K., *Ind Jour Pharm* , 1949, 11, 188

PUERARIA THUNBERGIANA Benth

Kudzu vine (Pueraria thunbergiana) as fodder. Dhar, N D & Goswami, M. N , *Sci. & Cult* , 1950, 16, 77

PUNICA GRANATUM Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm Inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb., Melia azadirachta Linn , Punica granatum Linn , Picrasma excelsa Swartz , Vernonia ahthelmintica Willd., Cocos nucifera Linn (coconut) Cairns. J. F. & Mhaskar, K. S., *Ind Jour. Med Res* , 1923, 11, 353.

PUTRANJIVA ROXBURGHII Wall

Putranjiva roxburghii Wall The oil from the seeds of Krishna, S. & Puntambekar, S. V ; *J Ind Chem Soc* 8 (1931), 301.

Development of the Microspores and the Nuclear behaviour in the Tapetal Cells of Putranjiva roxburghii Wall. Dutt, M. K., *Sci. & Cultr.*, 1943, 8, 309.

QUERCUS Linn.

Electric treatment of horse chestnuts and acorus. Hayao Muraoka; *Chem Abst.* 3268, 1936.

Q. DILATATA Lindl., Q ILEX Linn., Q INCANA Roxb.

Quercus incana Roxb, Q. dilatata Lindl. and Q. ilex Linn. Some Indian acorn oil Puntambekar, S. V. & Krishna, S, *J. Ind Chem. Soc.* 11 (1934), 721.

A note on the variable tannin content of the wood of Quercus incana H. S Chaturvedi & Watson, E. R, *J Ind Chem Soc.* 3 (1926), 211

QUISQUALIS INDICA Linn.

Chemical study of the seeds of Quisqualis indica. Composition of the crude oil Chit Fang Hsii and Pao Him King, *J. Chinese Pharm. Assoc* 2, 132 1940

RANDIA DUMETORUM Lamk.

Chemical and Pharmacological examination of Randia dumetorum N O. Rubiaceae Hardikar, S W & Mohiuddin, M. G, *Ind Jour. Med Res*, 1937, 25, 131.

RAUWOLFIA Linn

Combined Digitalis and Rauwolfia poisoning in a human subject. De, M. M. & Tarapaido Chatterjee, *Ind Med Gaz* 1941, 76, 724.

Rauwolscine & derivatives. Mookerjee, *J. Ind Chem Soc.* 20, (1943), 15

Pharmacognostic studies in the leaves of a few species of Rauwolfia. Determination of Palisade Ratio and Stomatal Index value Datta, S C, *Ind Jour. Pharm*, 1944, 6, 77

Rauwolfia roots, Pharmacognostic studies on commercial varieties Datta, S C.; *Ind Jour Pharm*, 1949, 11, 105

A preliminary note on the excretion of Rauwolfia total alkaloids in urine Gupta, J. C, Roy P K, Ray, G K & Ganguly, S. C., *Ind. Jour. Med Res.*, 1950, 38, 67

Rauwolscine methiodide Chatterjee, *J. Ind Chem Soc* 28 (1951) 31

Iso Rauwolscine & its HCl, perchlorate, picrate and Chloroplatinate Chatterjee, *J. Ind Chem. Soc* 28 (1951) 31, 33

R. CANESCENS Linn.

Alkaloids of Rauwolfia canescens Linn. Part I. (Miss) Mookerjee, A., *J. Ind. Chem Soc.* 18 (1941), 33

The Alkaloid of Rauwolfia canescens Linn. Part I. (Miss) Mookerjee, A, *J. Ind. Chem. Soc* 18 (1941), 485.

Function of Rauwolscine in Rauwolfia canescens Linn. (Miss) Mookerjee, A., *Sci. & Cultr.*, 1942, 8, 40.

The Alkaloids of Rauwolfia canescens. Mookerjee, A; *J. Ind. Chem. Soc.* 20 (1943), 11.

Rauwolfia canescens. Constitution of the alkaloid (Rauwolsina) Mookerjee, *J. Ind. Chem. Soc.* 23 (1946), 6.

Studies on the Pharmacology of Rauwolscine, the alkaloid of Rauwolfia canescens Linn. Mukherjee, J. N.; *Sci. & Cult.*, 1953, 7, 338

Yobyrine, the selenium dehydrogenation product of Rauwolscine, the alkaloid of Rauwolfia canescens. Chatterjee, A., Pakrashi, S. *Sci & Cult.*, 1953, 9, 443

R. SERPENTINA Benth

Chemical Examination of the Roots of R. serpentina. Siddiqui-Siddiqui, *J. Ind. Chem. Soc.*, 8, 667 (1931)

Chemical examination of roots of Rauwolfia serpentina Benth By Salimuzzaman Siddique & Rafat Hussain Siddiqui; *J. Ind. Chem Soc.* 8 (1931), 667.

The alkaloids of. Rauwolfia serpentina Benth. Part I. Salimuzzaman Siddiqui & Rafat Hussain Siddiqui; *J Ind Chem Soc* 9, (1932), 539.

The Alkaloids of R serpentina Benth , Part I Siddiqui-Siddiqui, *J. Ind Chem. Soc* 9, 539 (1932),

R. serpentina. van Itallie-Steenhauer, *Arch. Pharm.* 270, 313 (1932),

The pharmacological action of an alkaloid obtained from Rauwolfia serpentina, Benth A preliminary note. Chopra, R N., Gupta, J C. & Mukerji, B., *Ind Jour Med. Res* , 1933, 21, 261

The Alkaloids of R serpentina Benth , Part II, Studies in the Ajmaline-Series Siddiqui-Siddiqui, *J Ind Chem Soc.*, 12, 37 (1935)

Die Wirkung von Rauwolfin auf das Herz, Hartog *Arch Int. Pharmacodyn Therap* 51, 10 (1935).

The alkaloid of Rauwolfia serpentina Benth. Part II Studies in the Ajmaline Series. Siddiqui & Rafat Hussain Siddiqui, *J Ind. Chem. Soc.* 12 (1935), 37

Sur un nouveau paralysant electif des vasoconstricteurs adrenal-inosensible l'ajmaline, aclaloide cristallise de R. serpentina Raymond-Hamet, *Bull. Acad Med* 115, 452 (1936)

The Action of some Drugs on Fibrillation of the Heart (Rauwolfin). van Dongen, *Arch Int Pharmacodyn Therap.* 53, 80 (1936).

Ajmalinine, Alkaloid from R. serpentina of selective Paralyzing Action on Vaso-constrictors Reactive to Adrenaline Raymond-Hamet, *Bull Sci-pharmacol*, 43, 364 (1936)

The Action of Ajmaline on Nerve Impulses Chopra-Das-Mukerjee, *Ind J Med Res* 24, 1125 (1937).

A note on the Alkaloids of R. serpentina Benth. Siddiqui, *J Ind. Chem Soc* , 16, 421 (1939).

Rauwolfia serpentina Benth. A note on the alkaloids of. Salimuzzaman Siddiqui, *J. Ind. Chem Soc* 16 (1939), 421.

Influrence de la serpentinine sur les effets de l'adrenaline, de l'occulsion carotidiene et de la faradisation du pneumogastrque. Raymond-Hamet, *C. r.* 211, 414 (1940).

Sur quelques proprietes physiologiques de la serpentine, alcaloide cristal lise du R. sepentina Benth Raymond-Hamet, *C. r Soc. biol.* 134, 94 (1940).

Effets intestinaux d'ajmaline pure. Raymond-Hamet, *C. r Soc. biol* 134, 369 (1940).

A method of assay for Rauwolfia serpentina, Benth. Mahadev Lal Schroff & Rattan Lal Bhatia, *Ind Jour Pharm*, 3, 59, 1941.

A preliminary note on the pharmacological action of the alkaloids of Rauwolfia serpentina Chopra, R. N. & Chakravarti. M., *Ind Jour Med Res*, 1941, 29, 763

Alkaloids of Rauwolfia serpentina, A comparative study of pharmacological action and their role in experimental hypertension Chopra, R N., Bose, B C., Gupta, J C. & Chopra, I C., *Ind Jour. Med. Res.*, 1942, 30, 319

Preliminary observations on the use of Rauwolfia serpentina Benth. in the treatment of mental disorders Gupta, J. C, Deb, A K & Mahali B. S., *Ind Med Gaz*, 1943, 78, 547

Comparative pharmacology of the total alkaloids of Rauwolfia serpentina Benth obtained from Bengal, Bihar & Dehra Dun. Gupta, J. C, Kahali, B S, *Ind Jour Med Res*, 1943, 31, 215.

Hypnotic effect of Rauwolfia serpentina The principle underlying this action, its probable nature Chopra, R N, Gupta, J C, Bose, B C & Chopra, I. C, *Ind Jour Med Res.*, 1943, 31, 71

Pharmacological action of alkaloid of R. serpentina Benth. Pt I Neo-ajmaline & iso-ajmaline Bhatia, B. B & Kapur, R D, *Ind Jour Med Res.*, 1944 32, 177

The Hypnotic Effect of a Resin-Fraction from the Root of Rauwolfia serpentina Benth obtained from Dehra Dun Gupta-Kahali-Dutt, *J Ind Med. Res*, 32, 183 (1944)

Effects tenseurs vasculaires d' un des alcaloides colores du R serpentina Serpentinine Raymond-Hamet, *C r* 223 927 (1946).

Rauwolfia serpentina Benth Comparative chemical investigation of some of the constituents of the drug obtained from different sources and isolation of the active resin Dutt, A, Gupta, J. C, Ghosh, S & Kahali, B S, *Ind Jour Pharm*, 1947, 9, 54.

A note on the Hypnotic Principle of Rauwolfia serpentina. Gupta, J C, S Ghosh, A T Dutta & B. S Kahali, *J Amer. Pharmac. Assoc Sci Ed* 36, 416 (1947).

Pharmacological action of alkaloids of R. serpentina Benth Pt. II Total alkaloidal exts of Bihar & Dehra Dun varieties. Kapur, R. D, *Ind. Jour Med. Res.*, 1948, 36, 57.

Chemistry of Ajmaline (Rauwolfine of van Itallie and Steenhauer). Mukherji, Robinson-Schlittler, *Exper.* 5, 215 (1949)

Ajmaline, serpentine and Serpentinine. Raymond-Hamet, *C. R.* 229, 1165 (1949).

A Clinical Trial of R. serpentina in Essential Hypertension Vakil, *Brit Heart J* 11, 350 (1949).

Serpentina alkaloids. Bakshi, V. M , *Ind. Jour Pharm* , 1950, 12, 172.

Ueber das alkaloid Serpentin aus R serpentina Benth Schlittler-Schwarz, *Helv.* 33, 1463 (1950),

Structure of Reserpine. Neuss, Boaz, Forbes, *J A. C S* 75, 4870 (1950)

Ex-cretion of Rauwolfia total alkaloids in Urine. Gupta-Roy-Ray-Ganguly, *Ind. Jour. Med. Res.* 38, 67 (1950)

Rauwolfia serpentina in essential hypertension Chakravarty, N K , Rai Chaudhuri, M N & Chaudhuri, R N , *Ind Med Gaz* , 1951, 86, 348.

A new alkaloid from the root of Rauwolfia serpentina Benth. Chatterjee, A & Sukumar Bose, *Sci & Cult* , 1951, 17, 139

Ueber Rauwolfia serpentina, eine neue Droge in der Behandlung des Hochdrucks. Mayer, Deutsche Apothekerzeitung 1952, 435

Rauwolfinine, the new Alkaloid from the Root of R serpentina Bose, *Science and Culture*, 18, 98 (1952)

Ueber R. serpentina, eine neue Droge in der Behandlung des Hochdrucks Meyer, *Deutsche Apoth Ztg* 1952, 435

Zur Konstitution des Serpentins. Bader-Schwarz, *Helv.* 35, 1594 (1952).

Rauwolfia in Hypertension Mazumdar-Mukherji, *J. Ind. Med. Ass.* 19, 362 (1952).

The Use of R. serpentina in High Blood Pressure. Bhatia, *J. Ind Med. Ass* 21, 262 (1952)

Die Behandlung der chronischen arteriellen Hypertonie (II. Teil) Arnold, *Therapie der Gegenwart* 1952, 167.

Behandlung der Hypertonie mit der Indischen R. serpentina. Vida, *Die Medizinische* 1952, 1157

Raupin, ein neues Alkaloid sus R serpentina Bodendorf-Eder, *Naturwiss* 40, Heft 12, (1953)

Sarpagin, ein neues Alkaloid aus R serpentina Stoll-Hofmann, *Helv.* 36, 1143 (1953)

The Constitution of Ajmaline Chatterjee-Bose, *Experr.* 9, 254 (1953)

The Constitution of Reserpine Furlenmeier, Lucas, MacPhillamy, Mueller Schlittler, *Experr.* 9, 331, (1953)

On the Constitution of Reserpine from R serpentina Dorfman, Huebner, MacPhillamy, Schlittler, St. Andre; *Experr* 9, October 15th. (1953).

Alkaloids of Rauwolfia serpentina, The Characterization of Reserpine and its Hydrolysis Products. Klohs, Draper, Keller, Petracek, *J. A. C. S.* 75, 4867 (1953).

Indian Medicinal Plants in the Treatment of Hypertension Chakravarti, *Brit. Med J.* 1953, 1390.

Zur Pharmakologie des Reserpins, eines neuen Alkaloids aus R. serpentina. Bein, *Exper* 9, 107 (1953)

Sedative and Hypnotic Action of Reserpine in Unasthetized Animals Plummer, Barrett, Wagle, Yonkman *Fed Proc* 12, 357 (1953)

Pharmacological Effects of Reserpine, a new Crystallized pure Alkaloid from R serpentina in the Dog Trapold, Osborne, Yonkman, *Fed. Proc* 12, 213 (1953)

Wirkung von Reserpin (Serpasil) auf isolierte Kreislauforgane. Tripod-Meier, Submitted to *Arch Internat Pharmacodyn*

The Cardiovascular Effects of Reserpine (Serpasil) Trapold, Plummer, Yonkman, Submitted to *J Pharmacol exp. Therap*

Combinations of Drugs in the Treatment of Essential Hypertension (pure Reserpine=Serpasil) Wilkins, *The Mississippi Doctor*, 1953, 359 (April 5th)

The Use of R serpentina in Hypertensive Patients (Rauwolfia extracts). Wilkins-Judson, *New England J. Med* 248, 48 (1953)

Medical Management of Arterial Hypertension (Rauwolfia extracts), Meilman, *New England J. Med.* 248, 936 (1953)

The Action of Rauwolfia Serpentina on Vasomotor Reflexes Ray, Roy, Dasgupta and Werner, *Arch exper Path. u Pharmakol Bd.* 219, S 310-314 (1953)

The Sympathicolytic Activity of Rauwolfia Serpentina (Benth) Ray, Roy, Dasgupta and Warner, *Ind. J Med. Sciences*, 7, No 5, 229-235, (5/53).

REVENELIA SESSILIS

Lysis in the germination of the Urediospores of Revenelia sessilis Mathur, R L, *Sci & Cult*, 1952, 17, 380.

RHAMNUS Linn.

Further studies on Cathartic action in mice, Senna, Aloe Cascara and bile salts Lloyd W Hazleton and Kathleen D Talbert, *J Am. Pharm Assoc* 33, 170, 1944.

R VIRGATUS Roxb

Indian Cascara, Rhamnus virgatus Roxb Chopra, I C, Kohli, J D. & Handa, K L, *Ind. Jour Med. Res.*, 1950, 38, 473.

RHEUM Linn.

Assay of anthracene purgatives by the estimation of the contents of Hydroxymethylanth raquinone Pt I Rhubarb. Ghosh, S., Gupta, J. C, & Mahali, B S., *Ind Jour Med Res.* 1945, 32, 129

R. EMODI Wall.

Rhapoticin and anthraquinone derivatives from Rheum emodi Wall. (Indian or Himalayan Rhubarb). M. Ghouse and M. C. Tumin Katti, *J. Ind. Inst Sci.*, **16A**, 1, 1933.

A note on the essential oil from the Rhizomes of Rheum emodi Wall. M. Ghouse Mohiuddin, *J. Ind. Inst. Sci.* **18A,** 134, 1935.

Some common indigenous remedies Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum Chopra, R.N. & Ghosh, S , *Ind. Med. Record* 55, 77, 1935

A note on the Essential oil from the Rhizomes of "Rheum emodi Wall". Mohiuddin M Ghouse, *Journal of the Indian Institute of Science, Bangalore,* 1938, Vol **18A,** Part XVII, pp 134-135.

Indian Rhubarb as substitute for "official" Rhubarb Mukerji, B., *Curr. Sci* , 1943, 12 275.

Examination of Indian Rhubarb-Rheum emodi Wall Ray, G. K , Guhar, R. C., Bose, A B. & Mukerjee, B , *Ind Jour. Pharm* , 1944, 6, 55.

Indian Rhubarb Hocking, M , *Ind. Jour Pharm* , 1945, 7, 89

Studies on Indian Rhubarb, Rheum emodi Heber W Yongken, *J Am. Pharm. Assoc* 35, 145, 1946.

RICINUS Linn.

The Growth of effects of thiamine Hydro chloride ascarbic acid and phytohormones on beladonna and ricinus Loxis C Zoft, *J Am Pharm Assoc.* 29, 487, 1940.

R. COMMUNIS Linn.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency XVIII Cathartics Oleum Ricini, Oleum Tiglii, Aloe, Succies Acalyphae Chains, J F , Mhaskar, K S , *Ind. Jour Med Res.* 1923, 11, 103

On the description of the infloresence of Ricinus communis Linn Datta, R M , *Sci & Cult* , 1946, 10, 431.

ROSA Linn

Rose Oil - Chatterjee, N G.; *Indian Soap Journ* 3, 1937, 293.

Rose Distillation - P E O R Jan. 21, page 14 *Journal of Indian Industries and Labour,* Vol 2, Part I, page 86.

RUMEX Linn.

Anthracenic derivatives in the genera Polygonum and Rumex E. Maurin, *Bull Sci. Pharmacol* 33, 138, 1926

RUTA GRAVEOLENS Linn.

The correlation between chemical composition of anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency Santonin, Oleum rutae (Ruta graveolens Linn), Butea monosperma Roxb , Melia azadirachta Linn , Punica granatum Linn., Picrasma excelsa Swartz., Vernonia anthelmintica Willd., Cocos nucrifera Linn. (coconut). Cains, J F & Mhaskar, K S., *Ind Jour Med. Res.*, 1923. 11, 353.

SACCHARUM SPONTANEUM Linn.

Wilt Rot in grasses I. Involution, Saccharum spontaneum L. Srivastava, R. K., *Sci. & Cult.*, 1952, 17, 529

SALIX Linn

The physiological significance of the glucosides in Aesculus and Salix Gerhard Kerstan, *Chem Abst* 6772, 1934

SALVADORA OLEOIDES Dcne.

Fat from Salvadora oleoides, Khakanfot Patel, C. K., Iyer, S. N., Sudborough, J. J. & Watson, H. E., *J Ind Inst Sci* **9A**, 117, 1926

SAMBUCUS Linn

Presence of alkaloids in Sambucus species Henri Yardin, *Chem Abst* 5723, 1936

SANSEVIERIA ZEYLANICA Willd

Some common indigenous remedies, Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum Rheum emodi, Solanum indicum. Chopra, R. N. & Ghosh, S., *Ind Med Record* 55, 77, 1935

A Phytochemical study of Sansevieria zeylanica Stanley Scheindlin & Austin A. Dodge, *Am J Pharm* 119, 232, 1947

SANTALUM ALBUM Linn

A further note on the oil value of some sandalwoods from Madras Puran Singh, *Indian Forester*, 1915, Vol XII, No 4

West Australian Sandalwood Oil Rao B. Sanjiva & Sudborough J. J., *Journal of the Indian Institute of Science, Bangalore*, 1923, Vol 5 Part XII, pp 163-176

Some pharmacological action of Sandalwood oil, Vitivert oil and Pineneedle oil Tameno Okanishi, *Chem Abst* 658, 1930

Historial regeneration of Sandal (Santalum album Linn.) and its significance *Curr Sci*, 1934-35, 3, 251

East Indian Sandal Wood Oil A.E. Bradfield, Penfold, A. R. & Simonsen, J.L., *Journ. Chem Soc* 1935, 309

The wax from the leaves of Sandal (Santalum album Linn.) Chibnall, A. C., Piper, S. H., H. A. Cl., Mangouri, Williams, E. F. & Iyenger, A. V.V., *Biochem J.* 31, 1981, 1937.

Santalum album Chemical examination of the oil from the seeds Madhurnath, M.K & Manjunath, B. L., *J. Ind. Chem. Soc.* 15, (1938), 389

The white Sandal (Santalum album). Majumdar, G.P., *Sci. & Cultr.* 1941, 6, 492.

Studies in the santalol series. Part IV. Chemistry of Guerbets Santalic acid . Guha, P.C. & Bhattacharyya, S C , *J. Ind. Chem Soc.* 21(1944), 333

Studies in the Santalol series. Part V. Isolation of β-Santalic acid, a new constituent of Sandal wood oil. Bhattacharya, S.C.; *J. Ind. Chem Soc.* 21(1944), 337

Studies in the Santalol series. Part I. Separation of the Santalol and the Santalenes. Guha, P C. & Bhattacharyya, S.C , *J. Ind Chem. Soc.* 21(1944), 262.

Manufacture of Sandal Wood oil in India S G. Sastry , *Jour Sci. Indl Res.*, 1944-45, 3, 75.

Studies in the Santalol series. Bhattacharyya, S C., *Sci. & Cultr* , 1947, 13, 158-159

Studies in the Santalol series Parts XI, XII, XIII & XIV. Bhattacharyya, S.C., *Sci. & Cultr* , 1947, 13, 206-209.

Memorandum on the oil value of some sandal woods from Madras. Puran Singh , Forest Bulletins New Series No 6

SAPINDUS Plum.

A note on the Sapogenin from Soapnuts Shah, S.V., *Curr. Sci.*, 1935-36, 4, 436

S LAURIFOLIUS Vahl & S MUKOROSSI Gaertn

Sapindus laurifolius Vahl Soponin from Biswas , *J. Ind. Chem Soc.* 25(1948), 151

Saponins from Indian soapnuts Sapindus mukorossi Gaertn. & Sapindus laurifolius Vahl Gedeon, J , *J. Sci. Indl. Res* , 1952, **11B**, 84.

SAPONARIA Linn.

Damping off in Saponaria L Srivastva, H.C , *Sci. & Cult* , 1951, 17, 226.

SARACA INDICA Linn.

Saraca indica. S K Gupta, *Ind Med Record* 59, 112, 1939.

SARCOCOCCA PRUNIFORMIS Lindl.

Chemical Examination of Sarcocca pruniformis Lindl. Chopra, I.C. & Handa, K.L., *Ind Jour Pharm* , 1951, 13, 129.

SAUSSUREA LAPPA Clarke.

A preliminary note on the pharmacological action therapeutic properties of Kuth root (Saussurea lappa) Chopra, R.N & Premankur, De ; *Ind. Med. Gaz.*, 1924, 59, 540.

The treatment of Bronchial Asthma with Saussurea lappa (Kuth root). Chopra, R.N., *Ind. Med. Gaz.*, 1928, 63, 186.

Saussurea lappa (Kuth root) in Pharmacology and Therapeutics Chopra, R N. & De, Premankur., *Ind Jour Med Res.*, 1929, 17, 351.

Chemical examination of roots and leaves of Saussurea lappa Part I, Sudhamoy Ghosh, Nibarranjan, Chatterjee & Ashutusl Dutta , *J. Ind. Chem. Soc.* 6(1929), 517.

Root oil of Saussurea lappa, Clarke Establishment Antoine Chiris , *Chem. Abst.* 1553, 1937

A histological study of the root of Saussurea lappa C.B. Clarke, Prasad, S , *Ind Jour. Pharm.*, 1945, 7, 81.

Chemical Examination of roots of Saussurea lappa, Clarke, Pt. I On the reported isolation of the alkaloid "Saussurine" Salooja, K C , Vishwa Nath Sharma & Salimuzzaman Siddiqui , *J. Sci. Indl. Res* 1950, **9B**, 1.

SCIRPUS GROSSUS Linn

Studies in enzymes. Part II. Purification of amylose from Kaseru (Scirpus grossus Linn) Shukla, J P , *J Ind Chem Soc* 19(1942), 121

SCOPOLIA Jacq.

The differentiation of Scopolia and Belladonna leaves. D Markovic, *Chem Abst* 10, 1, 9, 1944

SECALE CEREALE Linn.

The hydrolysis of plytic compounds derived from seeds of hemp, horse bean, horse chestnut flax, wheat and embryos of rye W Jarosza , *Chem. Abst* 5501, 1934

SECURIGERA SECURIDACA Linn.

The pharmacological action on the circulatory system of a bitter principle isolated from Securigera securidaca Linn Dagen el Dorfler (N.O. Leguminosae) De, P., *Ind Jour Med Res* 1943, 31, 67.

The constituents of the seeds of Securigera securidaca Linn (Syn S. coronilla D C.). Ghosh, B.K & Dutta, A , *Ind. Jour Pharm.*, 1950, 12, 233.

SEMECARPUS ANACARDIUM Linn.

Constituents of the marking nut, Semecarpus anacardium. D. Satyanarayana Naidu ; *J. Ind. Inst. Sci* **8A**, 129, 1925.

Semecarpus anacardium Linn. Chemical examination of the Marking Nut. Parameswaran, P. & Salimuzzaman, Siddiqui , *J. Ind Chem. Soc.* 8 (1931), 517.

Studies in the Thermal degradation Prdts of naturally occuring resinals, Pt. I. Bhilawan shell liquid. Ram Prakash Rastogi, Vishwa Nath Sharma & Salimuzzaman Siddiqui , *J. Sci. Indl. Res.*, 1948, **7B**, 61.

Chemo-therapeutic studies in Bhilawan series, Pt I. Syn of sulphuric acids and arsines from Bhilawanol and its derivations Vishwa Nath Sharma, Ahmed Kamal & Salimizzaman Siddiqui , *J. Sci. Indl. Res* , 1948, **7B**, 67

S. TRAVANCORICA Bed

Chemical Examination of the constituent of Holigarna arnottiana Hook. F Pt I , latex of Semecarpus travancorica Bed Pt II Nair, A V , Poti, D N & Pillay, P P , *J Sci Indl Res.*, 1952, **11B**, 294, 298

SENECIO JACOBAEA Linn. & S RETRORSUS D C

The alkaloids of Senecio species The necines and necic acids from S. retrorsus and S. Jacobaea. Richard, H F Manske , *Can J. Res.* 5, 61, 1931.

S. VULGARIS Linn

Alkaloids of species of Senecio. Alkaloids of Senecio vulgaris. Degradation of Sinecionine R A Konovalova & A P. Onekhov, *Chem. Abst* 7436, 1937

SESELI INDICUM Wight & Arn

Essential oil of Seseli indicum W & A Farooq , M.O., Gupta, G S , *Curr. Sci.* 1953, 2, 46

SETARIA PALMAEFOLIA Stapf

Fodder value of Setaria palmaefolia Stapf Mirchandani, T.J & Dabadghao, P M , *Sci & Cult* , 1949, 14, 432

SIDA CORDIFOLIA Linn

The action of Sympathomimetic alkaloid in Sida cordifolia (Brela). Chopra, R N., Premankur, De , *Ind. Jour. Med Res* , 1930, 18, 467.

Sida cordifolia Linn Chemical examination of Sudhamoy Ghosh & Ashutosh Dutt, *J Ind Chem Soc* 7, (1930), 825.

SISYMBRIUM IRIO Linn

Chemical Examination of the fatty oil from the seeds of Sisymbrium irio Linn. Joti Sarup Aggarwal & Karimulla, *J. Sci Indl Res.*, 1946, 5, 57

SKIMMIA LAUREOIA Sieb & Zucc ex Walp

Chemistry of essential oils Investigation of the oil of Skimmia laureola Heinrich Weinhans & Tara Chand Rajdhan, *J. Parkt. Chem.*, 147, 113, 1936

A preliminary note on the chemistry and pharmacology of the leaves of Skimmia laureola Chopra, R N , Chatterjee, R G., De, N & Ghosh, S.; *Ind Jour. Med. Res* , 1938, 26, 481.

Studies on the active principles of Skimmia laureola, Asima Chatterjee, Bhattacharya, A., *J.I.C.S* 1953, 1, 33.

SOLANUM INDICUM Linn.

Some common indigenous remedies. Picrorhiza kurroa, Erythrina indica, Sansevieria zeylanica, Pongamia glabra, Hygrophila spinosa, Bryophyllum calycinum, Rheum emodi, Solanum indicum Chopra, R N & Ghosh, S , *Ind Med Record* 55. 77, 1935

S NIGRUM Linn.

Chemical examination of the fruits of Part I The comp of the oil from the seeds Pendse, G P , *J Ind Chem Soc* 14 (1937), 366

S PSEUDOCAPSICUM Linn

The action of an alkaloidal product from the leaf of Solanum pseudocapsicum Watt, L J M , Heimann, H L & Mellzer, E ; *J Pharmacol* 39, 387, 1930

Chemical Examination of the fatty oil from the seeds of Solanum pseudocapsicum Linn Pt I Husain, R , Siddiqui, S., Warsi, S A , *Ind Jour Pharm* , 1944, 6, 68.

S TORVUM Sw

A note on the chemistry of Solanum torvum Krishnamurti, G V , Sheshadri, T R., *J Sci Indl Res* 1949, **8B**, 97.

S XANTHOCARPUM Schrad & Wendl

Chemical examination of some Indian medicinal plants. Tinospora cordifolia, Solanum xanthocarpum and Fumaria officinalis Pense. G P. & Dutt, S., *Ind Jour Med. Res* , 1932, 20, 663

Fruits of Solanum xanthocarpum Saiyed, I Z. & Kanga, D D , *Proc Ind. Acad Sci* **4A**, 255, 1936

The chemical examination of Solanum xanthocarpum Schard & Wendel Part I The constitution of the oil from the seeds Gupta, Mahadeo Prasad & Dutt, Sikhibhushan, *J Ind Chem Soc* 13 (1936), 613

Chemical examination of the seeds of. Solanum xanthocarpum. Schard & Wendel Part II The constituents Gupta Mahadeoprasad & Sikhibhushan Dutt, *J. Ind. Chem Soc.* 15 (1938), 95

SONNERATIA ACIDA Linn.

Constitution of the colouring matter from Sonneratia acida Linn Govind Rai Chaudhry, Vishwa Nath Sharma; Siddiqui, S , *J Sci Indl. Res* , **7B**, 292, 1949

Chemical Examination of Archa (Sonneratia acida Linn). Pt I. Isolation of three crystalline products from the wood Chaudhry, G R & Siddiqui S ; *J. Sci. Indl. Res.*, 1950, **9B**, 137.

Chemical Examination of Archa (Sonneratia acida Linn.) Pt II. Studies in the constitution of Archin and Archiniry Chaudhry, G R., Vishwa Nath Sharma & Siddiqui, S , *J. Sci. Indl. Res.*, 1950, **9B**, 142.

SPARTIUM SCOPARIUM Linn.

Lupine alkaloids. Secondary alkaloids of Spartium scoparium Winterfield, K. and Fritz Nitzsche, *Arch. Pharm.* 278, 393, 1940.

SPHAERANTHUS INDICUS Linn.

Chemical investigation of Sphaeranthus indicus Linn. Basu, N K. Lamsal, P.P. *J. Am Pharm Assoc* 35, 274, 1946.

SPILANTHES ACMELA Linn

Chemical investigation of Spilanthes acmela Linn. Gokhale & Bhide *J. Ind Chem Soc.* 22, (1945), 250

SPIRAEA ARUNCUS Linn.

An antibiotic from Spiraea aruncus L E P Abraham, N G. Heatley R. Rolt and E M Osborn, *Nature*, 157, 511, 1946.

STELIARIA MEDIA Linn.

A note on the embryology of Stellaria media L Niranjan Pal *Sci. & Cult*, 1950, 16, 37.

STEPHANIA GLABRA Miers

The alkaloids of Indian Stephanias Pt. I Isolation of three crystallin alkaloids from the tubers of Stephania glabra Miers Chaudhry, G R & Siddiqui S., *J. Sci Indl. Res*, 1950, **9B**, 79

The alkaloid of Stephania glabra Miers. Identity of Aindarine Gindarinine with Tetrahydropalmatine and Palmatine respectively Chaudhry, G.R, Sharma, V.N. & Dhar, M L, *J Sci Indl. Res* 1952 **11B**, 337.

STRYCHNOS NUX-VOMICA Linn

Nux vomica, chromatographic analysis of its tincture and extract Notes and News. *Ind Jour Pharm* 1946, 8, 105

SWERTIA CHIRATA Ham.

A modified method for the assay of Chirata (Swertia chirata) Hande K L & Vidya Sagar Parbhakar; *Ind Jour Pharm*, 1952, 14, 87.

S. DECUSSATA Nimmo

Chemical investigation of Indian Medicinal plants Part I. Investigation of Swertia decussata Isolation of decussatin and Swertinin Shah, R.C., Dalal, S R, (Miss) Sethna, S, *J I C S* 1953, 7, 457.

Chemical investigation of Indian Medicinal plants Part III Structures of decussatin and Swertinin. Shah, R.C, Dalal, S R., (Miss Sethna, S, *J I C S*, 1953, 7, 463.

SWIETENIA MACROPHYLLA Linn.

Chemical investigation of the seeds of Swietenia macrophylla Linn Chakravarty, T. & Guha, S.S., *Sci. & Cultr*, 1947, 12, 450

Swietenia macrophylla. Chemical investigation of the non bitte principle of. Guha, Sircar & Chakravarty, *J · Ind Chem. Soc* 28 (1951), 207.

Swietenin from the seed of Swietenia macrophylla. Sircar, Guha, & Chakravarty, *J. Ind. Chem. Soc.* 28 (1951), 208.

SYMPLOCOS RACEMOSA Roxb.

Pharmacognostic studies of Symplocos racemosa Roxb. Baddomu ,S , Clarke, C B , Chatterjee, K C. & Khorana, M L., *Ind Jour. Pharm.*, 1950, 12, 290.

TABERNAEMONTYNA CORONARIA Br.

Phytochemistry of the bark-of Tabernaemontana coronaria Br Ratangiriswarn, A.N. and Vankatachalum, K , *Quart J. Pharm. Pharmacol* 22, 174, 1939.

TAGETES ERECTA Linn.

Petaloidy of sepal in a ray floret of Tagetes erecta. Farooq, M , *Sci & Cult* , 1951, 17, 92.

T PATULA Linn

Colouring matter of the flowers of Tagetes patula Isolation of a new flavonol, patulatin and its constitution. P Suryaprakasa Rao & Seshadri, T.R., *Proc Ind. Acad Sci.* **14A,** 643, 1941

TAMARINDUS INDICA Linn.

Tartaric acid from Tamarind fruit Iyenger, A V , Varadaraja, *Curr Sci* , 1937-38, 6, 610

Tamarind seed 'Pectin' Hanji, H R , Savur, G R. & Sreenivasan, A., *Curr Sci* 1945, 14, 129.

Tamarind seed 'Pectin' Damodran, M & Rangachari, P N., *Curr. Sci* , 1945, 14, 203.

Tamarind seed 'Pectin'. Ghose, T P. & Krishna, S , *Curr. Sci.*, 1945, 14, 299.

Tamarind seed 'Pectin', Ghose, T P., Krishna, S & Suryaprakasa Rao, P , *J Sci Indl. Res.*, 1945-46, 4, 705.

Tamarind seed 'Pectin', Damodran, M & Rangachari, P N , *Curr. Sci* , 1946, 15, 20 & 133

Tamarind seed 'Pectin', Rao, P S & Krishna, S , *Curr. Sci.*, 1946, 15, 133 & 163.

Tamarind seed 'Pectin',Savur, G R. & Sreenivasan, A., *Curr. Sci.*, 1946, 15, 43, 134 & 168.

Tamarind seed Polysaccharides Rao, P S , Krishna, S , *Curr. Sci.*, 1947, 16, 256.

Tamarind seed Jellose ("Pectin"), its jellying properties. Rao, P S., *J. Sci Indl Res.*, 1948, **7B**, 89.

Tamarind (Temarindus indica) seeds as Protein rich feed for Livestock, Kehar, N.D. & Sahai, B., *Sci. & Cult.*, 1649, 14.534.

Jellies and rerated products from Tamarind seed kernels. Rao, P.S., *J. Sci. Indl. Res*, 1949, **8A**, 354.

Studies on Tamarind seed kernel powder, Pt I. Preparation and utilization as a sizing material in the Jute Industry. Macmillan, W.G. & Chakarvarti, I.B, *J Sci Indl Res*, 1951, **10B**, 13

Studies on Tamarind kernel powder Pt II Physical Prop T K Paste & T K P Sind Yarns Macmillan, W G & Chakarvarti, I B, *J Sci Indl Res*, 1951, **10B**. 270

Studies on Tamarind kernel powder. Pt III Comparative study of antiseptics used in Jute sizing Mixtures Macmillan, W G. Chakarvarti, I B & Pal, P N, *J Sci Indl Res*, 1952, **11B**, 438

TANACETUM. Linn

Oleum Tenaceti, Caius, J.F & Mahaskar, K S, *Ind Jour Med Res.* 1920, 7, 606

TAXUS BACCATA Linn

Physiologically active constituents of the Yew, Taxus baccata, Taxine Robert K Callow, John M Gulland & Cyril J Virden; *J Chem Soc* 2138, 1931.

Detection and estimation of the alkaloid Taxine in preparations of Taxus baccata, Kuhn, A & Schafer, G, *Chem. Abst* 723, 1938

T. BREVIFOLIA Nutt.

Taxus brevifolia Georges Masson, *J Am. Pharm Assoc.* 28, 493, 1939.

TECTONA GRANDIS Linn.

The oil from the seeds of Tectona grandis (Teak) Puntambekar, S V & Krishna, S, *J. Ind Chem Soc.* 10 (1933), 401

TEPHROSIA Pers

Source of Indian Tephrosia sp. as rotenone Krishna, S & Ghose. T P, *Curr Sci*, 1937-38, 6, 454

Retenone containing Tephrosia as a source of insecticides. M G. Timoshenko, *Chem Abst.* 3530, 1943.

Effect of extracting agents & methods of extraction on the toxicity of Tephrosia extracts. Z K. Bogatova, *Chem. Abst.*

T. PURPUREA Pers

Studies on the hypoglycaemic effect of Tephrosia purpurea Var pumila Swift, H B N, Sethi, M S, Sareen, K N, *Ind. Med. Gaz*, 1949, 84, 243

T. VILLOSA Pers.

Tephrosia villosa. A plant useful in diabetes Harbhajan Singh; *Ind. Jour. Pharm.*, 1945, 7, 60

TERMINALIA ARJUNA W. & A.

Terminalia arjuna. Its chemistry, pharmacology and therapeutic action. Chopra, R N & Ghosh, S , *Ind Med Gaz.*, 1929, 64, 70

Chemical Examination of Terminalia arjuna. Isolation of arjunin. Radharaman Aggarwal & Sikhibhushan Dutt , *Proc Acad. Sci. Ind.* 5, 50, 1935

Chemical examination of the bark of Terminalia arjuna Isolation of arjunetin from the alcoholic extract Radharaman Aggarwal & Sikhibhushan Dutt , *Proc Natl Acad. Sci* 6, 305, 1936

T. BELERICA Roxb.

Preliminary examination of the Fruits of Terminalia belerica Roxb Chakravarti, M D & Tayal, J N , *Sci & Cultr* , 1947, 13, 122

T CHEBULA Retz

Distribution of Tannin in different parts of the Myrobalan plant (Terminalia chebula) Biswas, H G , *Sci & Cult* , 1944, 9, 399.

Tannic acid B P from Myrobalan (Terminalia chebula) Biswas, N G , *J Sci Indl. Res* , 1947, **6B**, 122

THALICTRUM FOLIOLOSUM D.C

Chemical examination of Thalictrum foliolosum D.C Isolation & characterisation of a new alkaloid Thalictrin Vashistha, S K & Siddiqui, S., *J Ind Chem Soc* 18 (1941) 641

Plant alkaloids Part III On Thalictrum foliolosum Chatterjee R , Guha, M D & Chatterjee, A , *J Ind Chem Soc* 29 (1952) 371

THEA SINENSIS Linn

The use of tea in the treatment of burns Peiris, M V P , *Ind Med. Gaz* , 1937, 72, 718.

THEOBROMA CACAO Linn

Coca leaves, the determination of ether soluble alkaloids and ecgonineesters Notes and News, *Ind Jour Pharm* 1946, 8, 104

THEVETIA NEREIFOLIA Juss.

The Pharmacological action of Thevetia glucoside occurring in Thevetia nereifolia (Yellow oleander) Chopra, R N & Mukerji, B , *Ind Jour Med Res* , 1933, 20, 903

Chemical examination of the roots of Thevetia nereifolia (Juss) N. Ghatak and G P Pendse , *Bull Acad Sci. Ind* 2, 259, 1933

The constituents of be-still nuts, Thevetia nereifolia K K Chen. & A. Ling Chen , *J Biol. Chem* 105, 231, 1934

The pharmacological action of Thevetoxin. A second glucoside from Thevetia nereifolia. Bhatia, B B. & Lal, S , *Ind. Jour. Med. Res.*, 1934, 21, 605.

Nevefolin, a new digitalic heleroside from Thevetia nereifolia Marcel Frerefacque , *Compt. Rend.* 221, 645, 1945.

THYMUS SERPHYLLUM Linn.

Studies in Indian Essential oils. III—Essential oil from leaves o Thymus serphyllum, Linn. Jagjit Singh and Rao B. Sanjiva *Journal of the Indian Institute of Science*, Bangalore, 1932, Vol **15A** Part VII, pp 78-83.

THYSANOLAENA MAXIMA Kuntze

Thysanolaena maxima as cattle feed Das, B K & Mukherjee, N.C. *Sci & Cult* , 1951, 17, 137

THYMUS VULGARIS Linn

Thymus vulgaris L Christine Rosenthal , *Pharm Ind.* 10, 22, 1943.

TINOSPORA CORDIFOLIA Miers.

Chemical Examination of Tinospora cordifolia Miers Kidwai, A R , Salooja, K C , Vishwa Nath Sharma & Siddiqui, S , *J Sci. Indl. Res.*, 1949, **8B**, 115

Chemical examination of some Indian medicinal plants Tinospora cordifolia, Solanum xanthocarpum and Fumaria officinalis. Pense, G.P & Dutt, S., *Ind Jour Med Res* , 1932, 20, 663.

The nature and germination of seeds of Tinospora cordifolia Miers Ajrekar, S L. & Oza, J D , *Curr Sci* , 1934-35, 3, 379.

Chemical investigation of Tinspora cordifolia Bhide, B V., Phalinkar, N.L. and Pranjpe, K , *J. Univ Bombay.*,3, 89, 1941

Chemical investigation of Tinospora cordifolia (Miers) Bhide, B.V , Phalinkar, N L and Pranjpe, K , *J Univ. Bombay.* 10, 89, 1941

T. CRESPA Miers

The wax of Tinospora crespa Miers. Lucienene Beauquesne ; *Chem. Abst* 868, 1942

TODDALIA ACULEATA Pers

On the failure of Toddalia aculeata in the treatment of malaria. Bhatia, B B , *Ind Med Gaz* , 1932, 67, 192

The chemistry and pharmacological action of Toddalia aculeata. Dey, B B., Pillay, P P , David, J C. & Rajamanikam, N. , *Ind. Jour. Med. Res* , 1935, 22, 765

Botany, Chemistry and Pharmacodynamus of Toddalia aculeata. Lobstien, J.E and Hesse, P.; *Chem Abst.* 2811, 1937.

On the constitution of natural conmarins of Todalia aculeata. Phanibhushan Dutta , *Jour Ind. Chem. Soc.*

TRAPA BISPINOSA Roxb.

The food value of the nut of Trapa bispinosa. Brahmachari, B.B. & Chatterjee, N K., *Ind. Med. Gaz.*, 1927, 62, 365.

TRESPASIA POPULNEA

Colouring matter of Indian Tulip (Trespasia populnea) flowers. Populnin and populnetin. Oeelakatam, K & Seshadri, T R, *Curr. Sci.*, 1938, 7, 16

TRIANTHEMA PORTULACASTRUM Linn.

A comparative study of Boerhaavia diffusa Linn. and the white and red flowered varieties of Trianthema portulacastrum Linn. Chopra, R N, Chatterjee, N R and Ghosh, S, *Ind Jour Med Res*, 1940, 28, 475

TRIBULUS TERRESTRIS Linn

Chemical Examination of the fruits of Tribulus terrestris Linn. Ghatak, N, *Bull Acad Sci* 2, 163, 1933.

TRICHOSANTHES ANGUINA Linn

Chemical examination of drying oil from the seeds of Trichosanthes anguina Linn Miss Padmini Soni, Aggarwal, J S., *J Sci Indl Res.*, 1949, **8B**, 150

T DIOICA Roxb

Chemical examination of the Indian Medicinal plants, Trichosanthes dioica Nag, N C, *Trans Bose Research Inst, Calcutta*, 10, 113, 1934-35,

TRIGONELLA FOENUM-GRAECUM Linn

Composition of Fenugreek seeds Mixture of fenugreek seeds with grain intended for flour E Fleurent, *Compt. Rend* 182, 944, 1926

Proteins of Indian Food Stuffs V Alcohol-soluble protein of fenugreek (Trigonella foenum-graecum) Sreenivasa Rao, Y V Sastri, B N and Narayana, N, *J Ind Inst Sci.* **16A**, 85, 1933

Thesaponin of fenugreek seeds. Gabra Soliman and Zahira Mustafa, *Nature*, 151, 195, 1943

Study and improvement of funugreek (Trigonella foenum graecum L.). Toao Marquis De Almeida, *Chem Abst* 419, 1943

Sapogenins The sapogenins of Trigonella foenumgraecum Russel E Marker, Wagner, R B, Ulshaper and Clarence H Rouf, *J Am. Chem. Soc.* 65, 1247, 1943.

TRITICUM Linn

The hydrolysis of plytic compounds derived from seeds of hemp, horse bean, horse chestnut, flax, wheat and embryos of rye. W. Jarosza, *Chem. Abst* 5501, 1934

Poisonous food grain wheat mixed with Lolium temulentum. Greva, S.D S & Bhandari, P.N, *Ind. Med Gaz*, 1946 81, 294

TRIUMFETTA RHOMBOIDEA Jacq.

The development of the female gametophyte in Triumfotta rhomboidea Jacq Bhattacharyya, S.S. & Mitra, J.N., *Sci. & Cult.*, 1952, 137, 45.

TUSSILAGO FARFARA Linn

Plant pigments Taraxanthin from Tussilago farfara (Colts foot) Karner, P and Morf, R., *Helv. Chem Acta* 15, 863, 1932.

TYLOPHORA ASTHMATICA W & A

The chemical examination of Tylophora asthmatica and isolation of the alkaloid Tylophorine and Tylophorinine Ratnagiriswaran, A N & Venkatachalam, K , *Ind Jour Med Res* , 1934, 22, 433

Chemical and Pharmacological investigation of Tylophora asthmatica Chopra, R N , Ghosh, N N , Bose, J B & Ghosh, S , *Arch Pharm* 275, 236, 1937

URGINEA INDICA Kunth

Chemical Examination of Indian squill Seshadri, T R & Subramanian, S S , *J Sci Indl Res* , 1950, **9B**, 114

Optical activity of preparations of Indian Squill Subramanian, S , *Ind Jour Pharm* , 1952, 14, 25

VACCINIUM Linn.

The constituents of diuretic drugs, the flavonol glucoside of Folium Uvae Ursi and Folium Vaccinii-Harnkitu Nakamura, Tatuo Ota and Gentiro Hukuti, *J. Pharm Soc , Japan* 55, 800, 1935.

VALERIAN Linn

Investigation of the Pharmacological action of fresh and dried Valerian Janina Romanowska Majcherezyk, *Chem Zent* 906, 1939

V OFFICINALIS Linn

An acid ester present in the roots of the Valeriana officinalis, Emil Corinna, *Comp Rend* 201, 1152

Pharmacological investigation of four strains of Valeriana officinalis var Labfolia W Rusieki , *Chem Zent* 1404, 1939

VANDA ROXBURGHII R. Br

Pharmacological action of an active constituent isolated from Vanda roxburghii R Br Gupta, J P , Roy, P. K & Sen Gupta, K K , *Ind Jour Med Res* , 1946, 34, 253.

VANGUERIA SPINOSA Roxb

The fat from the seeds of Vangueria spinosa (N O. Rubiaceae). Airan, S. W & Shah, S V , *Curr Sci* , 1942, 400, 11

Fat from the seeds of Vengueria spinosa. Nadkarni, Airan & Shah , *J Ind Chem. Soc.* 24(1947), 25-30

VERBENA OFFICINALIS Linn.

Constituents of Verbena officinalis L Identity of Verbenalin with verin. Benno Reichert *Arch. Pharm.* 273, 357, 1935.

V. VENOSA.

Presence of Stychose in the stems and roots of Verbena officinalis, and the underground parts of Verbena venosa. J. Cheymol *J. Pharm. Chem.* 25, 110, 1937.

Dessication of Verbena officinalis L. Loss of holosides and Verbenaloside J Cheymol, *J. Pharm. Chem.* 25, 581, 1937.

Constituents of Verbena officinalis L., constitution of Corin. Benno Richert and Walter Hoffmann, *Arch. Pharm.* 275, 474, 1937

Action of Verbenaloside on two isolated animal organs. J. Cheymol; *J. Pharm. Chem.* 27, 386, 1938

Glucide (total holoside and verbenaloside) contents of different parts of Verbena officinalis Jean Cheymol, *Chem Abst* 2977, 1938.

Verbanalin. E. Buris & D. Susterova. Rihova, *Chem Abst.* 6004, 1938.

Chemical and Pharmacological Researches on verbenin, a galactagog-acting glucocide from Verbena officinalis L Kotoku Kawazima, *Chem Abst* 7396, 1939

Verbenalin (Verbena officinalis). P. Karner & H. Salomon, *Helv Chim Acta* 26, 1544, 1946

VERNONIA ANTHELMINTICA Willd.

The correlation between chemical composition of Anthelmintics and their therapeutic values in connection with the Hookworm inquiry in the Madras Presidency. Sentonin, Oleum rutae (Ruta graveolens Linn.) Butea monosperma Roxb., Melia azadirachta Linn, Punica granatum Linn, Picrasma excelsa Swartz, Vernonia anthelmintica Willd, Cocos nucifera Linn (coconut). Caius, J.F & Mhaskar, K.S, *Ind Jour Med Res.*, 1923, 11, 353.

Anthelmintic properties of Vernonia anthelmintica Willd (Syn Serrantula anthelmintica) Chopra, R N., Ghosh, N N & Mukerji, A K., *Ind. Jour Med Res.*, 1934, 22, 183.

VETIVERIA ZIZANIOIDES Stapf

Notes on some Indian Essential Oils "Vetiveria Zizanioides" Stapf Rao B. Sanjiva, Sudborough, J J & Watson, H E *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. 8A, Part X, pp. 147-149

Analytical characteristics of Dutch East Indian oils of Cananga patchouli and Vetiver Koolhas, D R. & Rowan, P.A, *Perfumes France*, 15, 245, 1937.

Cananga patchouli and Vetiver Oil. Koolhas, D.R. & Rowan, P.A, *Perfumery essential oil Record* 29, 53, 1938.

Indian Vetiver, Khus. Menon, A K, Ittyachan, C.T., *J.Sci. Indl. Res*, 1947, 6, 274.

VIGNA CATJANG Walp.

Studies in nutrition value of Indian vegetable foodstuff. Pt. III. Nutritive values of Lentil—Lens esculenta; Moench, Cow pea—Vigna catjang, Walp and Aconite Bean—Phaseolus aconitifolius,

Jacq. Niyogi, S.P., Narayana, N. & Desai, B.G , *Ind. Jour.Med. Res.*, 1932, 19, 859.

VINCA ROSEA Linn.

Abnormalities in the flowers of Vinca rosea Linn. Kapadia, G A , *Sci. & Cult* , 1950, 15, 402

VIOLA ODORATA Linn.

Viola odorata L. the fragrant violet considered from the old and the new point of view Ludwig Kroeber. *Chem. Abst.* 6022, 1947

VISCUM Linn

Cassytha L. on Viscum. Dutt, B M S , *Sci. & Cult* , 1950 16, 258

V. ALBUM Linn.

Active principle of Viscum album A Endirs, O Feuchtinger & S Jannssin , *Arch. Expl Path. Pharmcol* 196, 290, 1940.

Pharmacology of Viscum album. Stamatis C Comninos, *Chem Abst* 5996, 1941

The pharmacologically active substances in Viscum album. Alfred A. Andres, *Chem Abst* 5996, 1941.

VITEX NEGUNDO Linn.

An introductory note on the chemical investigation of the plant Vitex negundo (N O Verbenaceae) Basu, N K , Singh, G B , *Sci & Cultr.*, 1944, 9, 508

A note on the chemical investigation of Vitex negundo Pt I Basu, N K & Singh, G B , *Ind Jour. Pharm* , 1944, 6, 71

Investigation of Indian medicinal plants, Hydrocotyle asiatica Vitex negundo and Monniera cuneifolia, Basu, N.K , Lamsal, P P & Singh, G B , *Quart J. Phar. Pharmcol* 20, 135, 1947

V PEDUNCULARIS Wall.

On the failure of Vitex peduncularis in the treatment of malaria Chopra, R N., Knowles, R & Gupta, J.C , *Ind. Med. Gaz.*, 1924, 59, 133.

Vitex peduncularis—an antihaemolytic agent Gupta, J.C , Kahali, B.S. & Ganguly, S.C. , *Trop. Diseases Bull* 40, 677, 1943.

WRIGHTIA TINCTORIA Ber.

Chemical examination of the fixed oil from the seeds of Wrightia tinctoria. Parihar & Dutt, *J. Ind. Chem Soc* 23 (1946), 307.

XANTHIUM STRUMARIUM Linn.

Oil from the seed of Gokhru (Xanthium strumarium), Shrivastava, R.C., Krishnamurthy, R.S. & Athawale, C R , *J Sci. Indl. Res.*, 1950, **9B**, 282.

ZANTHOXYLUM ACANTHOPODIUM D C. & Z. ALATUM Roxb.

The constituents of some Essential Oils. 42 (a). Part V—The Essential Oil from the seeds of Zanthoxylum alatum, Roxb. 42 (b). Part VI—The Essential oil from the seeds of Zanthoxylum acanthopodium D C 42(c). Part VII —The Essential Oil from the seeds of Zanthoxylum budrunga, Wall. Simonsen, J L. & Rau M. Gopal; *Indian Forest Records*, 1922, Vol IX Part IV. pp. 23-26

Z ALATUM Roxb

Zanthoxylum alatum ; —*Proc. Nat. Acad Sc.* Vol. IX, 1939, pp 187-192

Z BUDRUNGA Wall.

The constituents of some Essential Oils. 42(a) Part V—The Essential Oil from the seeds of Zanthoxylum alatum, Roxb 42 (b). Part VI—The Essential Oil from the seeds of Zanthoxylum acanthopodium, D C. 42(c). Part VII The Essential Oil from the seeds of Zanthoxylum budrunga, Wall Simonsen, J L & Rau M Gopal, *Indian Forest Records*, 1922, Vol. IX Part IV, pp. 23-26.

The alkaloids of Zanthoxylum budrunga Khastagir, Harinarayan, *Curr Sci* 1947, 16, 185.

A study of the colouring matter of Tambul seeds. Seshadri, T.R , *J. Sci Indl., Res.*, 1949, **8A**, 232.

Z OVALTIFOLIUM Wight.

Constituents of some Indian essential oils Essential oil from the seeds of Zanthoxylum ovaltifolium, Simonsen, J L , *Ind Forest Records*, 11, 1, 1924.

Constituents of some Indian essential oils from the seeds of Zanthoxylum ovaltifolium, Juniperus communis Simonsen, J L ; *Ind. Forest Records*, 11, 1, 1924.

The Constituents of some Indian Essential Oils Part XIV—The Essential Oil from the seeds of Zanthoxylum ovaltifolium Simonsen, J L , *Indian Forest Records*, 1925, Vol. XI, Part I, pp. 1-5.

Z RHETSEA D.C.

Notes on some Indian Essential oils. Zanthoxylum rhetsea D C. Rao, B Sanjiva, Sudborough, J.J. & Watson, H E , *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol **8A**, Part X, pp. 174-177

ZINGIBER OFFICINALE Roscoe.

Notes on some Indian Essential Oils Zingiber officinale. Roscoe, Rao, B Sanjiva, Sudborough, J J & Watson, H.E., *Journal of the Indian Institute of Science, Bangalore*, 1925, Vol. **8A**, Part X, pp. 151-153.

Travancore Essential Oils. Part VII—The Essential Oil from the Rhizomes of Ginger, Zingiber officinale Moudgill, K.L ; *Journal of the Indian Chemical Society, Calcutta*, 1928, Vol V.

Essential oils of Travancore, Part VII, from rhizome of Ginger, Zingiber officinale. Kishori Lal Moudgill, *J. Ind. Chem Soc.* 5 (1928), 251.

A note on the essential oil from Ginger scrapings. Varier, N S, *Curr. Sci*, 1945, 14, 321.

ZIZYPHUS VULGARIS Lam.

Anaesthetics from leaves of Zizyphus vulgaris Lam E N. Taran, *Farmatsiya*, 4, 20, 1941.

Z XYLOPYRUS Willd.

Oil from the seeds of Zizyphus xylopyrus Willd Airan, J W., *Curr. Sci* 1948, 17, 150

GENERAL

ALKALOIDS.

Estimation of total alkaloid by Chromatography. Datal, V. D. & Khorana, M. L., *Ind. Jour Pharm.* 1950, 12, 173

The formation of alkaloids in plants Chatterjee, R *Sci & Cult-*, 1950, 16, 58.

ALKALOID CHEMISTRY.

Trends in alkaloid chemistry. Chatterjee, R. G., *Sci. & Cult*, 1951, 17, 237.

ANIMALS.

CONORHINUS RUBROFASCIATUS De Geer

Some notes on Conorhinus rubrofasciatus (De Geer). Awati, P R. *Ind Jour. Med. Res.* 1921, 9, 371.

LABEO ROHITA.

Bile from Labeo rohita Basu, U. P., *Ind. Jour. Pharm.*, 1944, 6, 81

MYLABRIS

Synthesis of Cantharidin. Paranjape, K. (Miss), Phalnikar, N L, Nargund, K. S., *Curr. Sci.*, 1943, 12, 256.

M MACILENTA.

Cantharidin content of Mylabris macilenta Beetles. Mukerji, B, Hassain, T. & Karim, B. A., *Curr. Sci*, 1944, 13, 315.

UROLONCHA MALABARICA Linn.

On a Trypanosome of the white throated Munia—Uroloncha malabarica Linn Das Gupta, D. M. & Siddons, L B., *Ind Med. Gaz*, 1941, 76, 151.

ANTHELMINTICS.

Correlation between the Chemical composition of Anthelmintics and therapeutic values in connection with the Hookworm enquiry in the Madras Presidency. XXII. Summary and conclusions Cairns, J F & Bhaskar, K. S., *Ind. Jour. Med. Res.*, 1923, 11, 371.

ANTIBACTERIAL SUBSTANCES.

Antibacterial substance in plants. Notes & News., *Ind. Jour. Pharm.* 1946, 8, 60.

ANTIBIOTICS.

A search for Antibiotic substances in some Indian Medicinal Plants. Mariam George, Venkataraman, P R & Pandalai, K M , *J Sci. Indl. Res.* 1947, **6B**, 42.

Investigation on plant antibiotics. Pt. IV Further searches for antibiotic substances in Indian medicinal plants George, Mariam & Pandalai, K M , *Ind Jour Med Res.*, 1949, 37, 169

Lichens and Antibiotic Activity Rangaswami, S & Subramanian, S. S., *Ind. Jour. Pharm* , 1952, 14, 214

Antibiotic activity of some Indian Medicinal plants Joshi, C G. & Magar, N. G , *J Sci. Indl Res* , 1952, **11B**, 261

AROMATIC PLANTS.

Supplement. Aromatic Plants of India. Pt IX, X, XI, XII Krishna, S , Badhwar, R. L , **2A, 12A, 10B, 11B.** *J Sci Ind Res.*

Supplement Aromatic Plants of India Pt XV *J Sci Ind. Res*

Aromatic Plants of India XIII Krishna, S & Badhwar, R L , *J. Sci. Ind. Res*

Supplement : Aromatic Plants of India Krishna, S & Badhwar, R. L., *J. Sci. Ind Res.*,

Aromatic supplement, plants of India, I, II, III, IV Krishna, S & Badhwar, R. L., *J. Sci Indl. Res*

CRUDE DRUGS

Report on (the assay of) crude drugs H. W Youngken, *J Assoc Official Agr Chem.* 18, 516, 1935

The present position of the crude drugs used in the Indigenous medicine. Pt. I Handa, K L , Kapoor, L D , Chopra, I C & Som Nath; *Ind. Jour. Pharm.*, 1951, 13, 29

CULTIVATION OF MEDICINAL PLANTS

Cultivation of Medicinal Plants in India Bal, S N , *Ind. Jour. Pharm.*, 1944, 6, 42

Medicinal Plants and their cultivation. Mitra, G C , *J. Sci. Indl. Res.*, 1948, **7A**, 319.

Scope for the cultivation of medicinal plants in India. Chopra, R N , Kapoor, L. D , Handa, K. L., Chopra, I C. & Nayar, S L , *J. Sci, Indl Res.*, 1948, **7A**, 527.

Cultivation of Medicinal plants in Kashmir Chopra, I C., Kapoor, L. D , Handa, K. L., *J. S. I. R* 1953, **7A**, 31.

DEHYDROASCORBIC ACID

Significance of dehydroascorbic acid in plants. Inderjit Babbar, *Ind. Jour. Med. Res* , 1950, 38, 263.

DRUG ADULTERATION

Drug adulteration and spurious drugs in India. Chopra, R. N., *Ind. Med. Gaz.*, 1935, 70, 693.

DRUG RESOURCES

Drug Resources of Himachal Pradesh. Handa, K L., Kapoor, L.D , & Chopra I C , *Ind. Jour Pharm.*, 1951, 13, 118

Drug Resources of Kangra Valley Handa, K L , Kapoor, L.D & Chopra, I C , *J Sci Indl Res* 1951, **10B**, 173.

DRUG SCARCITY.

Scarcity of drugs Mohammad Hamid, *Ind Jour Pharm* 1944, 6, 46.

ESSENTIAL OILS.

Indian Essential oils—Sudborough, J.J , *Journal of the Indian Institute of Science, Bangalore*, Vol. I & II. Part II. pp 13 to 27, 1913.

A further note on the efficacy of essential oils in the prevention and treatment of cholera Tomb, J W , *Ind Med Gaz* 1924, 59, 233.

Notes on Indian essential oils. Rao, B S , Sudborough, J J. & Waston, H E , *J Ind Inst Sci* **8A**, 143, 1925

The constituents of some Indian Essential oils Note on the rate of oxidation of d-3 Carene and other Terpenes in the presence of Catalysts Rao, M. Gopal, *Indian Forest Records*, 1925 (Chemical Series), Vol XI, Part V

Observations on the treatment of Cholera with essential oil, mistura pro-diarrhoea and permanganate of potash. Bharti, S R.; *Ind. Med Gaz* , 1926, 61, 596.

The constituents of Indian essential oils. B. Sanjiva Rao & John Lionel Simonsen, *J. Chem Soc* 2496, 1928

Notes on some Essential oils. *Perf. & Ess Oil Rec.* 1937, p. 28.

Notes on some Essential oils—*Perf. & Ess Oil Rec.* Nov. 1937, p 411.

Scent Factor of Flowers—Rakshit, J.N *The Perfumery & Essential Oil Record.* July 20, 1937, Vol 28, No. 7, page 241.

Essential oils from Flowers, Grasses and Plants—Rikshit, J.N.; *Rakshit Gardens, Bulletin No. 2, Ghazipur*, U.P 1938.

Development of Essential Oil industry—Rakshit, J.N.; *Science & Culture*, Vol V, No 2, August 1939.

Can Indian Natural Essential Oils compete with Synthetic Products of foreign countries—Rakshit, J.N.; *Science & Culture*, Vol. 5, No 6, December 1939, pp. 355-357.

Exploitation of Phytosynthesis of Essential oils—Rakshit, J.N.; Applied Chemistry Students' Reunion, 1940, published from the *University College of Science and Technology, Calcutta* p. 13-18.

Position of essential oil & Perfumery Trade in India. Rao, P A., *Sci. & Cultr*, 1941, 6, 710.

A preliminary study of the rideal walker-coefficient values of certain indigenous essential oils. Gupta, J C, Bose, B.C., Ganguly, S C, *Ind Med. Gaz.*, 1942, 77, 210.

ESTUARIAL FLORA

Estuarial flora of the Godavari, a preliminary Note on Venkateswarlu, V, *Sci. & Cultr.*, 1943, 8, 351.

A note on the Estuarine Flora of Kistna. Venkateswarlu, V, *Sci. & Cultr.*, 1946, 12, 295.

HEART POISONS

Heart poisons from plants Chakravarti, J.K., *Sci & Cultr* 1952, 17, 282.

INDIAN DRUG INDUSTRY.

Indian Drug Industry and the medical profession Mukerji, B, *Ind. Jour Pharm*, 1949, 11, 120.

INDIAN FOODS

Nicotinic acid contents of Indian foods Saharia, G.S, *Univ Bombay*, 13, 5, 1945.

INDIAN MEDICINE.

A note on Indian Medicine. Clifford, A, *Ind Med Gaz*, 1928, 63, 53.

INDIAN PHARMACOPOEIA.

Indian Pharmacopoeia, Pt. A.; *Indian Jour Pharm.* 1950, 12, 85

INDIAN PHARMACOPOEIAL LIST.

Note on the Indian Pharmacopoeial List Subramanian, S.S., *Ind. Jour. Pharm.* 1948, 10, 4

I.P.L, Banerjee, J.N. Tamhene, R.G & Kulkarni, V P, *Ind Jour. Pharm.*, 1950, 12, 12.

INDIGENOUS DRUGS

The field of research in Indian Indigenous Drugs Chopra, R N & Ghosh, B.N., *Ind Med. Gaz.* 1923, 58, 99

Indigenous substitute for imported drugs Bal, S N & Prasad, S, *Ind Jour Pharm.* 5, 30, 1943

Indentification of Indigenous Drugs Sircar, N N, *Ind. Jour Pharm*, 1944, 6, 9.

Symposium on utilization of indigenous drugs of India. Ghosh, S, *Ind. Jour. Pharm.*, 1944, 6, 10

Action of some indigenous drugs on Uterus. A preliminary Note. Kapur, R.D., *Ind. Jour. Med. Res*, 1948, 36, 47

Symposium on utilization of Indigenous drugs. Salimuzzaman Siddiqui; *Ind. Jour Pharm.*, 1950, 12, 221.

Phagocytic response in Guinea pigs to certain indigenous Medicinal extracts. Broker, R., Gaffar, K T., *Curr Sci* 1953, 2, 44.

INDIGENOUS REMEDIES.

Chemistry and pharmacology of some common indigenous remedies. Chopra, I C., Kohli, J.D. & Handa, K.L *Ind Jour. Med. Res.*, 1945, 33, 157

INSECTICIDES OF VEGETABLE ORIGIN.

Insecticidal Plants. Kabir, S A & Ramaswamy, M N., *Curr. Sci*, 1939, 8, 82.

Insecticides of vegetable origin. Siddiqui, S, *J Sci. Indl. Res*, 1950, **9A**, 181.

MEDICINAL PLANTS.

Some medicinal plants growing in Himalayas, II Chopra, R N, Ghosh, N.N., Ratnagriswaran, A N, *Ind Jour Med. Res.* 1928, 16, 777.

The scientific and economic importance of researches on Indian Medicinal Plants Ghosh, S, *Ind. Med Gaz.* 63, 650, 1928.

Observations in certain medicinal plants used in indigenous medicine Chopra, R.N & Ghosh, S., *Ind Jour. Med Res* 1929, 17, 377

Medicinal plants; from Bt Col R N. Chopra's presidential address, National Institute of Science in India at Madras *Sci & Cultr*, 1940, 5, 620.

Research on medicinal and poisonous plants. (Article). *Curr. Sci*, 1940, 9, 94

Medicinal plants in Chitral state. Datta, S C., *Sci. & Cultr.*, 1945, 10, 519

Scope for the cultivation of medicinal plants in India Chopra, R N, Kapoor, L D, Handa, K.L, Chopra, I.C. & Nayar, S.L., *J Sci Indl. Res*, 1949, **8A**, 14.

NOMENCLATURE OF DRUGS.

The nomenclature of Drugs. (An Article). *Ind. Jour. Pharm.* 1952, 14, 153.

PECTIN FROM INDIAN PLANT MATERIAL

Pectin from Indian Plant materials. Damodaran, M. & Rangachari, P.N.; *J Sci Indl Res* 1945-46, 4, 298.

PLANT PIGMENTS.

Plant pigments. Datta, C., *Sci & Cult*, 1951, 16, 458 & 512.

POISONOUS PLANTS

The toxicity of known and unknown poisonous plants in the union of South Africa Douw G. Steyn, Onderstepoort ; *J. Vet. Sci Animal. Ind* 9, 573, 1937.

Poisonous plants of India Chopra, R N., *Sci & Cultr* 1939, 5, 9

Research on medicinal and poisonous plants (Article), *Curr Sci.* 1940, 9, 94

SILAJIT

Chemical composition and antidiabetic properties of Silajit, Chopra, R N , Bose, J P & Ghosh, N N , *Ind Jour Med Res* 1926. 14, 145

STERIOD HORMONES

Isolation of steriod hormones from Indian plants Chakravarti, *Ind Jour Pharm* , 1952, 14, 80.

VEGETABLE DRUGS

Vegetable drugs Editor, *Ind Jour Pharm.*, 1951, 13, 81.

VEGETABLE DRUG RESOURCES

Vegetable drug resources of Jammu & Kashmir Pt I Plants yielding B P drug and their substitutes Chopra, R N , Kapoor, L D , Handa, K L. & Chopra, I.C., *J Sci Ind Res* , 1947, **6A**, 485.

Supplement to the Bibliography.

ACORUS CALAMUS.

Pharmacological action of some common essential oil bearing plants used in indigenous Medicine Part I Pharmacological action of *Acorus calamus*, Curcuma zedoria, Xanthoxylum alatum and Angelica archangelica Chopra, I C , Jamwal, K S. & Khajuria, B N , *Ind Jour Med Res* , 1954, 42, 381

ADHATODA VASICA.

Antitubercular action of *Adhatoda vasica* N O Acanthacea, Part I. Gupta, K.C , & Chopra, I C , *Ind Jour Med Res* , 1954, 42, 355.

ALPINA GALANGA.

Pharmacological action of some common essential oil bearing plants used in Indigenous Medicine Part II Pharmacological action of *Alpinia galanga*, Pistacia integrima, Piper betel, and Nardostychas jatamansi Chopra, I. C , Jamwal, K S & Khajuria, B. N , *Ind Jour Med. Res* , 1954, 42, 385

ARGEMONE MAXICANA.

Removal of toxic alkaloids from Argemone oil and oils adulterated with argemone oil. Roy, A C., *Jour. Sci. Indl Res* , 1954, **13B**, 376.

BERBERIS ASIATICA Roxburgh

Plant alkaloids Pt. VI, Berberis asiatica Roxburgh. Barva, A K & Das Gupta, A K , *Jour Ind Chem Soc* , 1954, 31, 253

CARUM ROXBURGIA

Pharmacological study of an Indigenous Drug 'Ajmod' Carum Roxburgiana Benth and Hook Gujral, M L , Bhargava, K P , Srivastava, R S and Kishore, K A., *Ind Jour Med Res* , 1954, 42, 389

CEASALPINIA LIGYNA Rottles

Chemical examination of the roots of Caesalpinia Ligyna Rottles Chaudhry, G. R , Sharma, V N & Dhar, M L , *Jour Sci. Ind. Res* 1954, **13B**, 150.

CHENOPODIUM AMBERIOSOIDES

A short note on Chenopodium raised in Kashmir. Handa, K. L Kapoor, L. D & Abrol Hira Lal. *Ind. Jour Pharm* , 1954, 16.

COFFEE POWDER.

Chemical standards for coffee powder. Narayaniayer, S & Ram Chandran. *Curr. Sci* , 1954, 192, 23.

CURCURBITA PEPO

Chemical Examination of the oil from the seeds of Curcurbita pepo. Narayanmurthy, N L., Iyer, B H , Gangadharam P R J & Sirsi, M , *Ind Jour Pharm* , 1954, 7, 148

DATURA METEL Linn.

Variation & alkaloid content due to difference in size of the leaves of Datura Metel Linn. Chaudhuri, B. N., *Ind. Jour. Pharm* , 1954, 16, 34

ENTADA SCANDENS BENTH

A note on sulphur containing glucoside in the seeds of Entada scandens Benth Rangaswamy and Subba Rao, *Ind Jour. Pharm.* 1954, 7, 152

FERONIA ELEPHANTUM

A new aldobiouronic acid 3—(Glucuronopyranosyl)—D-Galactopyranose from Kata (Feronia elephantum) Gum. Mathur, G. P. & Mukherjee, S , *Jour Sci Indl Res.*, 1954, **13B**, 452.

HYOSCYAMUS MUTICUS

Chemical investigation of Hyoscyamus muticus , Handa, K. L , Abrol, Hira Lal, *Jour Sci Indl Res* , 1954, **13B**, 376.

JATROPHA GLANDULIFERA Roxb.

Chemical examination of the seed oil of Jatropha glandulifera Roxb , Sheth, M C , Desai, C M., *Jour Ind. Chem Soc* , 1954, 31, 407.

MELODINUS MONOGYNUS Roxb

Chemical examination of the root bark of Melodinus monogynus Roxb Part I. Chatterjee, S K , Sharma, V N & Dhar, M L , *Jour. Sci Indl Res.*, 1954, **13B**, 546

MENTHA ARVENSIS.

A short note on Japanese Mint raised in Kashmir Handa, K. L , Kapoor, L D & Abrol, H L , *Ind. Jour Pharm* , 1954, 16, 32.

MELIA AZEDARACH.

Chemical examination of the heart wood of Melia azedarach. Bhola Nath, *Jour. Sci Indl Res*, 1954, **13B**, 740.

MUSUA FERRAE

Refining of Nahor seed oil (Musua ferrae) Kusturi, T. R., Narayana Murty, N L and Ujer, B. H , *Jour. Sci. Indl Res.*, 1954, **13B**, 453.

OCIMUM KILIMANDSCHARICUM Guerke.

Camphor and camphor oil from Ocimum Kilimandscharicum Guerke. Choudhry, J. K., *Sci. & Cult* , 1954, 19, 354.

POLIANTHESE TUBEROSA Linn.

Glucofructosan from Polianthese tuberosa Linn and Garlic (Allium sativum Linn.)., Srinivasan, M., and Bhatia, I. S., *Curr. Sci.*; 1954, 23, 192.

PARMELIA KAMTSCHADALIS.

Chemical investigation of the Lichens. Parmelia kamtschadalis and Parmelia arnoldii Shah, Latika, G (Mrs). *Jour Ind. Chem. Soc*, 1954, 31, 311.

RAUWOLFIA CANESCENS.

The alkaloids of R. canescens, Part VI Yobyrine, the Se-Dehydrogenation Product of Rauwolscine. Chatterjee, A & Pakrashi, S; *J. Ind. Chem. Soc* 31, 25, 1954

The Alkaloids of R. canescens, Part VII Studies on the I R Spectra of Rauwolscine and its Acetyl derivatives Chatterjee, A. & Pakrashi, S. *J. Indian Chem. Soc* 31, 29, 1954.

The Alkaloids of R. canescens Part VIII. SeO_2-Oxidation of Yobyrine, the Se-Dehydrogenation Products of Rauwolscine. Chatterjee, A. & Pakrashi, S, *J Ind Chem Soc* 31, 31, 1954.

Conformation of Rauwolscine, allo-Yohimbine and their Congeners, Identification of Rauwolscine with a-Yohimbine Chatterjee, A, Bose, A K. & Pakrashi, S. *Chem and Industry* 1954, 491

On the Stereochemistry of Rauwolscine, the Alkaloid of R canescens. A. Chatterjee and S Pakrashi. *Naturwiss.* 41, 215, 1954.

The Isolation of Reserpine from R canescens M W Klohs, Draper, M D. Keller, F. Petracek F J *J A C.S.* 76, 1381, 1654

RAUWOLFIA SERPENTINA.

Reserpin, der sedative Wirkstoff aus R Serpentina. J. M Mueller, E. Schlittler & H.J. Bein. *Exper.* 8, 338, 1952

On the Constitution of Reserpine from R. Serpentina L Dorfman, C F. Huebner, H B. MacPhillemy, E Schlittler & A F. St. Andre. *Exper* 9, 368, 1953.

Chemistry of the Rauwolfia Alkaloids, including Reserpine. E Schlittler, H B MacPhillamy, L. Dorfman, A Furlenmeier, C F Huebner, R Lucas, J M Mueller, R Schwyzer and A.F St. Andre *Ann. N Y Acad. Sciences* 59, 1, 1954

Structure of Reserpine. N. Neuss, H E Boaz, J W. Forbes: *J A C S* 75 4870, 1953

Rauwolfia serpentina Alkaloids; Structure of Reserpine. N Neuss. H E Boaz & J W Forbes: *J A C S* 76, 2463, 1954

Rescinnamine, a new hypertensive and sedative Principle. Klohs, M.W, M.D. Draper & F Keller; *J A C.S.* 76, 2483, 1954,

Reserpinine, ein neues Alkaloid aus Rauwolfia serpentina. E Haack, A. Popelak, H. Spingler & F. Kaiser, *Naturwiss* 41, 214, 1954.

Sarpagin, ein neues Alkaloid aus R. Serpentina. A. Stoll und A. Hofmann *Helv. Chim. Acta* 36, 1143, 1954

Neue Alkaloids aus R. serpentina. A. Popelak, H Spingler und F. Kaiser *Naturwiss* 40, 625, 1953.

Essai de Classification des Alcaloides des Rauwolfias (Apocynaceaes). Raymond-Hamet *C r Sceances hebd Acad Sci.* 237, 1435, 1953

S Bose . *J. Indian Chem. Soc* 31, 47, 1954 Rauwolfinine, a New Alkaloid of R. serpentina, Part I.

Rauhimbin und iso-Rauhimbin, zwei neue Alkaloide aus R. serpentina A Hofmann. *Helv Chim Acta* 37, 314, 1954

Isolation of S -Yohimbine and a New Related Alkaloid from R serpentina F L Weisenborn, M Moore, P A Diassi *Chem. and Industry*, 1954, 375.

A Quel Type d' Acaloide la Raubasine appartient-elle Raymond-Hamet : *C R Sceances hebd Acad. Sci* 238, 1338, 1954.

Rauwolfia Alkaloids IX Isolation of Yohimbine from R serpentina F E. Bader, D Dickel & E. Schlittler, *J A.C S* 76, 1695, 1954.

Alkaloids of R serpentina II The Isolation of Naturally occurring Py-Tetrahydroserpentine (Ajmalicine) and a Contribution towards its structure M W. Klohs, M D Draper, F. Keller, W Malesh & F J Petracek *J A.C S* 76, 1332, 1954

A Preliminary investigation of R.serpentina W L Holt & Costello C H , *J. Amer Pharm. Ass* 43, 144, 1954

Chemistry of Ajmaline, Part I. Comp Chem and Industry, 1952, 442. F A L Anet, D Chakravarti, R Robinson and E Schlittler . *J.C S* (*London*) 1954, 1242

The Constitution of Ajmaline, Part I Chatterjee, A. & Bose, S *J. Indian Chem. Soc* 31, 17, 1954

Reserpinin, ein neues Alkaloid aus R Serpentina E Schlittler, H Saner & J M. Meller · *Exper.* 10, 133, 1954

The Constitution of Sarpagine A F Thomas · *Chem and Industry* 1954, 488.

Rauwolfia Alkaloide III Die Isolierung weiterer Alkaloide aus R. serpentina A Hofmann *Helv Chim Acta* 37, 849, 1954.

Method for Determination of Serpasil H Sheppard, G.L Wagle & A J. Plummer : *Fed Proc.* 13, 404, 1954.

The Determination of Serpasil in Blood Urine and Tissues (to be published). H. Sheppard, N. Smith, L Stannard & G Wagle.

Zur Pharmakologie des Reserpins, eines neuen Alkaloids aus R. serpentina H J Bein . *Exper.* 9, 107, 1953

Pharmacological Effects of Reserpine, a New Crystalline Pure Alkaloid from R serpentina, in the Dog J H Trapold, M.W. Osborne & F F. Yonkman . *Fed. Proc.* 12, 373, 1953.

Sedative and Hypnotic action of Reserpine in Unanesthetized Animals A.J Plummer, W. Barrett, G. Wagle & F F Yonkman: *Fed Proc* 12, 357, 1953.

Pharmakologische Untersuchungen Uber verschiedene Alkaloide (Serpasil, Ajmalin, Serpentin and Serpentinin) aus R serpentina. H J Bein und F Gross *Verh Ber. Dtsche Gas Kreislaufforsch* 19, 277, 1953.

Experimentelle Untersuchungen uber "Serpasil" (Reserpin), ein neues, sehr wirksames Rauwolfia-Alkaloid mit neuartiger zentraler Wirkung H J Bein, F. Gross, J Tripod und R Meier: *Schweiz. Med Wschr* 83, 1007, 1953

Wirkung von Reserpin (Serpasil) auf isolierte Kreislauforgane. J Tripod und R. Meier *Arch internat Pharmacodyn* (In Press)

Cardiovascular and Respiratory Effects of Serpasil, a New Crystalline Alkaloid from R serpentina J H Trapold, A J Plummer & F F Yonkman *J. Pharm. Exper Therap* 110, 205, 1954

Additional Pharmacological studies on Serpasil, a Crystalline Alkaloid from R serpentina, in the Dog J H Trapold, M W Osborne, A J Plummer & F F Yonkman · *J Pharm Exper. Therap.* 110, 49, 1954

Sedative and Hypertensive Actions of a Group of Esters Related to Serpasil A J. Plummer, W Barrett & R Rutledge: *Fed Proc* 13, 395, 1954 :

Effects of Serpasil (Reserpine) on Monkeys (Motion Picture). A E Earl, R C Dibble & R.D Wolf *Fed Proc* 13, 350, 1954

Action of Serpasil on Gastrointestinal Motility and Secretion W E. Barrett, R A Rutledge & B Rogie, *Fed Proc.* 13, 334, 1954,

Behavioral and Electroencephalographic Studies with Serpasil (Reserpine), a New Alkaloid from R serpentina J.A Schneider & A E Earl *Fed Proc* 13, 130, 1954

Effet Hypotenseur chez l 'Animal de la Reserpine, un nouvel Alcaloide de la R serpentina R Meier, H J Bein, F Gross, J. Tripod & H Tuchmann-Duplessis *C r. Sceances hebd Acad Sci.* 238, 527, 1954

Effetss de la Reserpine, un nouvel Alcaloide de la R serpentina, sur le Systeme nerveux central de l' Animal R. Meier, H J Bein, F Gross, J Tripod et H Tuchmann-Duplesses *C r. Sceances hebd. Acad Sci* 238, 961, 1954.

Hypotensive Agents from R serpentina Reserpine and other Alkaloids G Croneheim, C Stipp & W Brown: *J. Pharm. Exper. Therapy* 110, 13, 1954 ·

Characterization of Central Effects of Serpasil and their Antagonistic Reactions. J. Tripod, H J Bein und R. Meier: *Arch. intern. Pharmacodyn* 96, 406, 1954 :

Pharmacology of Rauwolfia Alkaloids including Reserpine. A.J. Plummer, A.E. Earl, J.A Schneider, J. Trapold & W. Barrett: *Ann. N Y Acad. Sciences* 59, 8, 1954.

Endocrine Aspects of the Pharmacology of Reserpine Gaunt, F., Renzi, A.A., Antonchak, N., Miller, G. & Gilman, M *Ann N Y. Acad. Sciences* 59, 22, 1954

Effects of Reserpine (Serpasil) on behaviuor and autonomic regulating mechanisms in the monkey. Tripod, J und Meier: *Arch Internat Pharmacodyn* (In Press)

Cardiovascular and Renal Hemodynamic Response (acute and chronic) to Reserpine. E. Dennis, W Hughes & J Moyer *Fed Proc.* 13, 347, 1954

Effect of Reserpine (Serpasil) on "sham rage" in the cat J A Schneider (to be published)

Mouse Prosis bioassay of Rauwolfia serpentina for reserpine-like activity B Rubin & J C Burke, *Fed Proc* 13, 400, 1954.

Changes in convulsant threshholds after Rauwolfia serpentina, Reserpine and Veriloid E H. Jenney *Fed. Proc* 13, 370, 1954

Sedative and Antihypertensive effect of Intravenous Reserpine in Man A M Sellers and J H Hafkenschiel, *Fed Proc* 13, 404, 1954

Pharmacological investigation with Serpine Das Gupta, S R & G. Werner *Bull Calcutta School Trop Med* 1, No, 4, 1954.

Indian Medicinal Plant in the Treatment of Hypertension. Chakravarti, M D, *Brit Med. J* 1953, 1390

Neue Alkaloide aus R serpentina J D Achelis und G Kroneberg. *Naturwiss* 40, 625, 1953,

Alterations in Cardiovascular Responses of the Dog following Rauwiloid, an Alkaloid Extract from R serpentina J T Gourzis, J R Sonnenschein & R. Barden *Proc. Soc Exptl Biol Med* 85, 463, 1954.

A Preliminary Note on the Pharmacological investigation of the Hypotensive Action of R serpentina J W Nelson & C A Schlogel · *J Amer Pharm Ass* 42, 324, 1953

Hypertensive Action of another R serpentina Alkaloid (Serpasil) R. J Vakil *J Indian Med. Assoc* 23, 97, 1953

Combinations of Drugs in the Treatment of Essential Hypertension R. Wilkins *Mississippi Doctor* 1953, 359

Hochdruck behandlung miteinem Rauwolfia alkaloid (Serpasil) W. Loffler, A F Essellier, F. Prott und A Wegmann *Schweiz Med. Wschr* 83, 1012, 1953

Reserpine and the Alseroxylon Alkaloids of R Serpentina in Hypertension. T. Winsor *Arizona Med.* 10, 419, 1953

Ueber die Behandlung der arteriellen Hypertonie mit Reserpin G Damm und H. Trautner *Dtsche Med Wschr* 79, 39, 1954

Clinical usage of Rauwolfia Alkaloids, including Reserpine (Serpasil). Wilins, R.: *Ann N. Y .Acad Sciences* 59, 36, 1954.

Clinical and Experimental Effects of Reserpine in Patients with Essential Hypertension. E D. Freis & R Ari · *Ann. N. Y. Acad. Sciences* 59, 45, 1954

Intravenous Reserpine in patients with Essential Hypertension. H. H. Hafkenschiel & A M. Sellers *Ann. N Y. Acad. Sciences* 59, 54, 1954

Human Pharmacology of Reserpine. T. Winsor · *Ann. N. Y. Acad. Sciences* 59, 61, 1954.

Clinical use of Reserpine (Serpasil) in Geriatrics. J. H. Moyer : *Ann. N. Y Acad. Sciences* 59, 95, 1954.

Use of Reserpine (Serpasil) in Certain Gynaecologic Disorders. R. B. Greenblatt *Ann. N Y Acad Sciences* 59, 133, 1954.

The Modern Treatment of Hypertension Comp Med. Times 81, 741. S W. Hoobler, *Univ Mich Med Bull* 20, 1, 1954

Recent Developments in the Treatment of Hypertension. E.D. Freis: *Med Clin N.A.* 38, 363, 1954

Reserpine in the Treatment of Hypertension A note on the Relative Dosage and effects R W. Wilkins, W E. Judson, W. Hollander, W E Huckabee & J. H Friedman *New England J. Med.* 250, 477, 1954.

Observations of Cardiovascular and Renal Hemodynamic Responses to Reserpine and Clinical Results in Treatment of Hypertension J H Moyer, W. Hughes & R Huggins *Clin Res Proc* 2, 72, 1954.

Essais Cliniques sur l'action du Serpasil dans l'hypertension arterielle grave O Ritter und J C. Huguenin · (In print)

Treatment of Hypertension with Serpasil, a R serpentina alkaloid. L Hensel · *Schweiz Med Wschr.* 83, 1162, 1953.

Clinical Experience with Reserpine (Serpasil) A controlled study. H P Dustan, R D Taylor, A C Corcoran & I. H. Page. *Ann. N Y Acad Sciences* 59, 136, 1954.

Reserpine alone and in combination with Hydralazine and Hexamethonium in the Treatment of Hypertension W Hughes, R. McConn and E. Dennis *Fed Proc.* 13, 368, 1954.

Studies on "Essential" Hypertension V An endocrine syndrome. H A. Schroeder and D F. Davies : *Ann. Int Med.* 40, 516, 1954.

Uber die perorale medikamentose Dauerbehandlung des chron. Hochdruckleidens. R. Franke & F. Thiele, *Die. Med.* 50, 1626, 1953.

The Medical Management of arterial Hypertension. E. Meilman : *N. Engl. J. Med.* 248, 894, 1953.

Excretion of Rauwolfia Total Alkaloids in Urine. Gupta, J. C., Roy, P. K., Ray, G.K. & Ganguly, S.C. *Indian J. Med Res.* 38, 67, 1950.

Die Behandlung der chronischcen arteriellen Hypertonie. O.H Arnold *Therapie der Gegenwart* 1952, 167.

Behandlung der Hypertonie mit der indischen R serpentina F. Vida *Die Medizinische* 37, 1157, 1952.

A comparison of Results in the Treatment of Hypertension with R. serpentina alone and in combination with Hexamethonium or Hydralazine I H Moyer, R.W Food, W R Livesay & S I Miller *J Lab and Clin Med* 42, 926, 1953

Extracts of R serpentina in Hypertension. R W. Ford & J H. Moyer *General Practitioner* 8, 51, 1953.

Treatment of Hypertension with R serpentina alone and in combination with Hexamethonium or Hydralazine J H. Moyer and W. R. Livesay *Amer J Med.* 16, 605, 1954

The Pharmacodynaics of Postural Hypertension (Rauwolfia) R A Schneider et al. · *Clin Res Proc* 2, 69, 1954.

Effects of Orally Administered "Veriloid" and "Rauwiloid" on the Blood Pressure of Hypertensive Individuals C T Bello & L. W Turner *Clin Res Proc* 2, 70, 1954

Response of Metacorticoid Hypertension to Bistrium, Apresoline Veriloid and Serpentina F M Sturtevant, *Proc. Soc. Expt Biol Med* 84, 101, 1953

Preliminary observation on R. serpentina in hypertensive patients R W Wilkins, W E Judson & I R Stanton *Proc New Engl Cardiovasc Soc* 34, 1951

R Serpentina in the Treatment of Arterial Hypertension F Klausgraber, *Wien, Med Wschr* 83, 995, 1953

New Drug Therapies in arterial Hypertension. R W Wilkins *Ann Internal Med* 37, 1144, 1952

Preliminary observation of Rauwolfia serpentina therapy of Hypertension R W Ford, W R Livesay, S I. Miller & J H Moyer *Med Res and Ann.* 47, 698, 1953

Antihypertensive Therapy. R. P. McCombs: *Bull Neu Engl. Med Center* 15, 129, 1953.

The use of Drugs in the Management of Hypertension Samuel S Riven & J G. Wells *J. Tenn. State Med Ass* 47. 1, 1954

Management of Essential Hypertension. W. C. Gettelfinger *J Kentucky State Med Ass.* 51, 434, 1953

Rauwolfia, die Neuentdeckung einer alten Heilpflanze. E. Steinegger *Bulletin Galenica* 17, 50, 1954.

1 Soprauna Sostanza ad Azione Sedativa Isolate della R. serpentina. A. Lattanzi *Omnia Therapeutica (Italy)* 5, fasc.

Schneider und A.J. Plummer E Schlittler, J A.

Ueber Rauwolfia-Alkaloide Angew. Chemie (In Print).

Zur Wirkung von R. serpentina. G Werner *Arzneimittelforschung* 4, 40, 1954.

Some Modern Hypertensive Agents. R W Richards : *Pharm J.* 172, 269, 1954

Chemical and Medicinal Aspects of R. serpentina R. Kress : *Pharmazie* 8, 726, 1953.

Indian Medicinal, Plant in the Treatment of Hypertension (R. serpentina) Chakravarti, M D.: *Brit. Med. J.* 1953, 1390.

New Results in the Pharmaceutical Treatment of Arterial Hypertension (Part I & II). O.H. Arnold & K D. Bock . *Dtsche. Med. Wschr* 78, 565, 879, 1953

G L. Maison : *Lancet*, 1953, 1308 The Treatment of Hypertension.

Essential Hypertension . Editorial *Med Times*, 81, 226, 1953.

Hypertension Editorial *Canad. Med Ass J* 68, 287, 1953.

Drug Therapy in Management of Hypertension. C. Levy . *Delaware State. Med J.* 26, 36, 1954.

R. Serpentina and other species G.E. Trease & W.C Evans *Pharmaceut. J.* 172, 351, 1954

A Pharmacognostical study of Rauwolfia see also Pharm 125, 186, 1953. H.W Youngken Sr *J. Amer Pharm. Assoc.* 43, 70, 1954.

Malabar Rauwolfia, R micrantha, H.W Youngken Sr. *J. Amer. Pharm. Assoc.* 43, 141, 1954

The pharmacognosy of Rauwolfia T.E. wallis & S. Rohatgi : *J. Pharm Pharmacol.* 69, 334, 1932

Ueber R serpentina, eine neue Droge in der Behandlung des Hochdrucks. G. Meyer *Dtsche. Apoth. Ztg.* 92, 455, 1952.

The use of R. serpentina in Neuropsychiatric Conditions. N.S Kline *Ann N Y. Acad. Sciences* 59, 107, 1954.

Effects of R serpentina on Maniac patients. P.K. Roy . *Indian J. Neurol Psychiat.* 2, 59, 1950.

Two cases of Paranoia treated with Rauwolfia serpentina. C K. Nagendranath : *Ind. J. Neurol. and Psychiatry* 2, 62, 1950.

The use of R. serpentina in Psychiatry. P K. Roy *Ind. J. Neurol. and Psychiatry*, 3, 380, 1952.

Recent development on Rauwolfia serpentina Benth. Rajgopalan, S : *Jour. Sci Indl Res.* **13B**, 17, 1954.

Rauwolfia serpentina. Iswariah, V *Curr. Sci* 23, 253, 1954

Rauwolfinine. A new Alkaloid of Rauwolfia serpentina Benth. Bose, S.: *Jour Ind Chem Soc*, 31, 311, 1954

RICE HUSK

Studies on rice husk, Betel husk and Bamboo lignins Pt II Nitrolignins Part III. Ali, M. Erfan. & M.H Khundkar *Jour. Ind Chem Soc.*, 31, 471 & 474, 1954.

SCILLA INDICA

A note on the isolation of crystalline Glycoside from commercial Indian Squill Rangaswami, S & Subramanian, Sankara, S *Jour Sci Indl Res* **13B**, 150, 1954

STROPHANTHUS SEEDS

Pharmacognosy of some Strophathus seeds from India Gupta, B. & Bal, S N *Indian Jour. Pharm*, 16, 74, 1954

USNEA HIRTA

Chemical components of Usnea hirta Subha Rao, V.. *Indian Jour. Pharm*, 7, 151, 1954.

INDEX

OF

SCIENTIFIC NAMES

PAGE

A

Abies pindrow Spach. .. 125
Abroma augusta Linn 21, 112, 125
Abrus precatorius Linn .. 21,70, 96, 101, 111, 125
Abutilon avicennae Gaertn. . 126
Abutilon indicum G Don .. 101, 126
Abutilon theopharstic Medic 101
Acacia Willd . 126
Acacia catechu Willd 126
Acacia chundra (Willd) DC . 126
Acacia ferruginea DC . 101
Acacia leucophloea Willd .. 126
Acacia pennata Willd 70,115
Acalypha Linn .. 126
Acalypha indica Linn. .. 127
Acanthaceae . 254
Achillea millefolium Linn 127
Achras sapota Linn. 127
Achyranthes aspera Linn. 127
Aconitum Linn . 21, 127
Aconitum anthora Linn 127
Aconitum balfourii Stapf . 21, 62, 128
Aconitum chasmanthum Stapf . 22, 62, 128
Aconitum deinorrhizum Stapf 22, 62, 128
Aconitum elwesii Stapf . 62
Aconitum falconeri Stapf . 62
Aconitum ferox Wall. .. 22, 62, 109
Aconitum heterophyllum Wall. 22, 127, 128
Aconitum laciniatum Stapf. . 22, 62
Aconitum laeve Royle .. 62
Aconitum lethale Griff .. 62
Aconitum luridum Hk. f & T . 62
Aconitum moschatum Stapf. . 62
Aconitum napellus Linn .. 128
Aconitum spicatum Stapf . 22, 62
Aconitum soongaricum . 62, 128
Aconitum violaceum Jacq. .. 22, 62
Acorus calamus Linn. .. 22, 93, 101, 115, 254

	PAGE
Acorus gramineus Soland	93, 115, 128
Acronychia laurifolia Blume	67, 115
Acronychia pedunculata (Linn) Miq	67, 115
Actaea spicata Linn	62, 128
Actinodaphne hookeri Meissn	23, 129
Actinomyces griseus	52
Adansonia Linn	129
Adenanthera pavonina Linn	129
Adenia (modecca) palmata Engl	74
Adensonia digitata Linn	101
Adenia wightiana Engl	74
Adhatoda vasica Nees	19, 23, 96, 101, 129, 254
Adiantum lunulatum Burm	101
Adina cordifolia Benth & Hk f	77, 101, 115, 129
Adonis aestivalis Linn	63
Adonis chrysocyathus H f & T	63
Aegle marmelos Corr	23, 96, 101, 129
Aesculus Linn	130
Aesculus hippocastanum Linn	130
Aerothecium lanatum Wakker	130
Agaricus compestris Linn	55
Agaricus ostreatus (Jacq) Fries	101
Agave americana Linn	91, 115, 130
Ageratum conyzoides Linn	101, 131
Ailanthus altissima (Mill) Swingle	68, 101, 112
Ailanthus excelsa Roxb	101
Ailanthus glandulosa Desf	68, 101, 112
Ailanthus malabarica DC	101
Alangium begoniaefolia Roxb	131
Alangium lamarckii Thw	23, 96, 131
Alangium salvifolium (Linn f) Wang	23, 96
Albizzia chinensis (Osbeck) Merr	115
Albizzia julibrissin Durazz	96
Albizzia lebbeck Benth	96, 101
Albizzia procera Benth.	70, 115
Albizzia stipulata Boiv	115
Algae	52
Allamanda cathartica Linn.	80
Allium cepa Linn.	96, 131
Allium minus	132
Allium odorum Linn.	131, 132
Allium porrum Linn	132
Allium sativum Linn.	23, 91, 96, 132, 133, 256

PAGE

Allium schoenoprasum Linn. . 97
Alocasia indica Schott . 93
Alocasia macrorhiza Schott . 93
Alocasia montana Schott . 93
Alocasia odora (Roxb) C Koch . 93
Aloe abyssinica Lam 133
Aloe barbadensis Mill 24, 111
Aloe species 92
Aloe Tourn ex Linn 133
Aloe vera ex Linn 24, 134
Alpina galanga Willd Swartz 24, 97, 254
Alstonia Br. . 134
Alstonia scholaris R Br 24, 101, 134
Alsophila glabra Hook 134
Althaea officinalis Linn 134
Althaea rosea Linn Cav . 101, 134
Amanita muscaria Pers . 54, 55, 74
Amanita pantheriana 57
Amanita phalloides Secr . 55, 57
Amarantus blitum Linn . 134
Amaranthus gangeticus Linn . . 101, 134
Amaranthus tricolor Linn . 101
Amaryllidaceae 91
Amito pers Kurrum 55
Amito ostreatus Jacq . . 55
Ammannia baccifera Linn 74
Ammannia senegalensis Lamk 74
Amomum subulatum Roxb 24
Amomum xanthioides Wall 101, 134
Amorphonphallus campanulatus (Roxb) Bl. 93
Amorphonphallus lyratus Engl 93
Amorphonphallus sylvaticus (Roxb) Kunth 93
Anaboena sp . 53
Anacardiaceae . 69
Anacardium occidentale Linn. . 69, 112, 115, 134, 135
Anagallis arvensis Linn 79, 112, 115, 135
Ananas comosus Merr. 91
Ananas sativus Schult . 91
Anamirta cocculus (Linn.) W & A . . 64, 115, 136
Anchusa tinctoria Lam. . . 135
Andrachne cordifolia Muell. Arg . . 87, 115
Andrographis echioides Nees. . . 136
Andrographis paniculata Nees. . 24, 101, 136

PAGE

Andropogon Linn. . 136
Andropogon jwarancusa Jones. . . 136
Andropogon kuntzeanus Hack. . . 136
Andropogon nardus Linn . . 117
Andropogon squarrosus Linn. . . 136
Anemone obtusiloba D Don. . . 63
Anethum soa Roxb . 137
Angelica archangelica Linn. 137, 254
Anisomeles malabarica Br. . . 137
Annonaceae . 64
Annona muricata Linn. . . 101, 137
Annona reticulata Linn . 64, 102, 115, 137
Annona squamosa Linn. . 64, 111, 115, 137
Anogeissus latifolia Wall 137
Anthemis cotula linn . 77, 112
Anthemis nobilis Linn . 137
Anthocephalus cadamba Miq S & H 102
Anthocephalus indicus A Rich . 102
Antiaris toxicaria Leschen . . 25, 89, 102, 138
Apama tomentosa Engl . . 115
Aphanizomenon flos-aquae (Linn) Ralfs 53
Apium graveolens Linn . . 76, 111, 112, 138
Apocynaceae . . 80
Aquilaria agallocha Roxb 138
Aquilegia vulgaris Linn 63
Araceae . . 93
Arachis hypogea Linn . 25, 138
Araliaceae . 76
Archangelica officinalis Hoffm . 138
Areca catechu Linn . 25, 92, 111, 138, 139
Arenga obtusifolia Mart. . 92, 115
Argemone alba Lestib. 139
Argemone mexicana Linn. 60, 64, 139, 254
Argyreia speciosa Sweet 140
Arisaema speciosum Mart 93, 112, 115
Arisaema tortuosum Schott. . 93, 112, 115
Aristolochia bracteata Retz. . . 86, 111, 116
Aristolochia indica Linn . 25, 86, 111, 140
Aristolochiaceae . . 86, 140
Arnica montana Linn . 141
Arnica Rupp. ex Linn. . 140
Arundo donax Linn. . 134
Artabotrys Br. . . 141

PAGE

Artabotrys suaveolens Blume. .. 109, 141
Artocarpus hirsuta Lamk .. 142
Artemisia Linn. . 26, 141
Artemisia absinthium Linn. 26, 77, 116, 142
Artemisia dracunculus Linn .. 26
Artemisia kurramensis. .. 142
Artemisia maritima Linn .. 77, 142
Artemisia pallens Wall. ex DC . 26
Artemisia sacrorum Ledele. . 26, 142
Artemisia scoparia Waldst. & Kit. . 142
Artemisia vulgaris Linn 26, 77, 111, 116, 142
Asarum europeum Linn .. 143
Asclepiadaceae .. 81, 143
Asclepias curassavica Linn . 81, 97, 116
Asparagus Linn . 143
Asparagus adscendens Roxb . 97, 102, 143
Asparagus officinalis Linn .. 112
Asparagus racemosus Willd .. 97, 102, 143
Aspergillus clavatus .. 54
Aspergillus fumigattus .. 54
Aspidium filixmas .. 58
Asteracantha longifolia Nees .. 27, 102, 143
Asystasia coromandeliana Nees .. 143
Atalantia monophylla Corr .. 143
Athyrium fillix-foemina Roth . 144
Atriplex hortensis Linn . 144
Atrocarpus integrifolia Linn 142
Atropa Linn . 144
Atropa acuminata Royle 27, 144
Atropa belladonna Linn 83, 144, 145, 146, 147
Averrhoa carambola Linn 102
Avena fatua Linn 94
Avena sativa Linn. .. 94
Azadirachta Linn 147
Azadirachta indica A. Juss. 27, 68, 97, 116, 147, 148

B

Bacitracin subtilis. .. 52
Bacopa monnieri (Linn.) Pennell. .. 27
Balanites aegyptiaca (Linn) Delile. Wall. 68, 116, 148, 149
Balanites roxburghii Planch 68, 102, 116
Balanophora polyandra Griff. .. 149

	PAGE
Baliospermum axillare Blume.	88
Baliospermum montanum muell Arg.	88
Balsamodendron mukul Hook	97, 149
Bambusa Schreb.	149
Bambusa arundinacea Willd.	27, 94, 116, 149
Bambusa bambos Druce	27, 94, 116
Barringtonia acutangula (Linn) Gaertn	73, 102, 116, 150
Barringtonia asiatica (Linn) Kurz	73, 116
Barringtonia racemosa Bl (Linn.) Roxb.	73 116
Barringtonia speciosa Forst	73, 116
Basella rubra Linn.	102
Bassia latifolia Roxb	118, 150
Bassia longifolia Linn	41, 119, 150
Bauhinia acuminata Linn	150
Bauhinia racemosa Lam.	97, 102
Bauhinia tomentosa Linn	102
Bauhinia variegata Linn	97, 102, 151
Begoniaceae	76
Begonia rex Putzeys.	76
Belamcanda chinensis Leman	151
Benincasa cerifera Savi	151
Berberidaceae	64
Berberis Linn	151
Berberis aquifolium Pursh	151
Berberis aristata DC	28, 64, 116, 152
Berberis asiatica Roxburgh	28, 254
Berberis coriaceae Brandis.	28
Berberis floribunda Hort Wall	28, 152
Berberis himalaica Arndt	28, 152
Berberis insignis Hook f. & B	28, 152
Berberis lycium Royle	28
Berberis nepalensis Spreng	41
Berberis umbellata Wall	29, 152
Berberis vulgaris Linn	29, 152
Bergenia ligulata (Wall.) Engl	102
Betula Tourn.	153
Betula utilis D Don.	153
Beta vulgaris Linn.	153
Bidens tripartita Linn	102
Bignoniaceae	84
Bixaceae	65
Bixa orellana Linn.	153
Blepharis edulis Pers.	29, 154

PAGE

Blumea balsamifera DC 154
Blumea densiflora DC 154
Blumea eriantha DC 154
Blumea lacera DC 109, 154
Blumea malcomii Hook f 154
Boerhaavia diffusa Linn 154
Boerhaavia repens Linn 29
Bombax malabaricum DC 154
Boraginaceae 82
Borassus flabellifer Linn 155
Boswellia serrata Roxb 102
Botrychium lunaria sew 102
Botrychium virginianum Sew 155
Bragantia tomentosa Blume 115
Bragantia wallichii R Br 29, 155
Brassica campestris Linn 155
Brassica cernua (Thunb) Forbes et Hemsl 65, 102
Brassica integrifolia (West) O E Schulz 65
Brassica juncea Hook f & T 65, 155
Brassica napus Linn 65
Brassica nigra (Linn) Koch 65
Brassica oleracea 134, 156
Bromeliaceae . 91
Brosimum galactodendron D Don. ex Sweet. 156
Bryonia alba Linn 156
Bryophyta 57
Bryophyllum calycinum Salisb 156
Brucea amarissima (Lour) Merr. 68, 102
Brucea sumatrana Roxb 68, 102, 156
Bursera Linn 156
Butea Roxb 156
Butea frondosa Koen ex Roxb 70, 116, 156, 157
Butea monosperma (Lam) Kuntze . . 29, 70, 97, 102, 116, 156
Buxus sempervirens Linn. 88
Butea superba Roxb. 29, 157

C

Caesalpinia bonducella Fleming . 30, 157
Caesalpinia crista Linn . 30
Caesalpinia digyna Rottl. . 158
Caesalpinia jayabo Maza. . 102

PAGE

Caesalpinia nuga (Linn) Ait. . . 116
Caesalpinia pulcherima Swartz. . 110
Calendula officinalis Linn. 158
Callicarpa longifolia Lam var lanceolaria C B. Clarke . 84, 116
Callitris rhomboidea R Br 158
Calonyction muricatum (Linn) G Don 82, 116
Calophyllum inophyllum Linn . 66, 116, 158
Caloplaca elegans 158
Calotropis Br . 158
Calotropis gigantea Linn.
Dryand R Br 30, 81, 132, 111, 113, 158, 159
Calotropis procera Br. (Linn) Dryand 81, 111, 113, 159
Caltha palustris Linn . 63
Calycopteris floribunda Lam . 30, 159
Camellia sinensis Linn. (O Kunfize). . 30
Campanulaceae 78
Cananga odorata Hk F & T 159
Cananga patchouli 159
Canarium commune Linn . 159
Canarium strictum Roxb . 160
Canavalia ensiformis D C. . 70, 160
Canavalia virosa W & A 70
Cannabis Tournef . 160
Cannabis indica Lamk . 160
Cannabis sativa Linn 30, 89, 113, 116, 160
Canscora decussata Schult 97
Cantharellus cibarius Fr 55
Capparis aphylla Roth . 65
Capparis decidua Edg 65
Capparis sepiaria Linn . . 160
Capparis spinosa Linn . 97, 160
Capparis zeylanica Linn . 109
Capparidaceae . . 65
Caprifoliaceae . 76
Capsella bursa-pastoris Moench . . 161
Capsicum annuum Linn. . 83, 161
Capsicum frutescens Linn . . 83, 109
Capsicum minimum Roxb. . . 83, 109
Carapa moluccensis Lam. . . 103, 109
Cardaria draba Desv. . . 116
Cardenia campanulata Roxb. . . 118
Cardiospermum halicacabum Linn. . . 69
Carex cernua Boott. . 94

	PAGE
Careya arborea Roxb	73, 116
Caricaceae	75
Carica papaya Linn	31, 74, 111, 161, 162
Carica xanthin	161
Carthamus tinctorium Linn	162
Carum bulbocastanum Koch	103
Carum carvi Linn	31, 103, 162
Carum copticum Benth	50, 110, 162
Carum roxburgia . .	254
Caryophyllaceae . .	66
Caryophyllus aromaticus Linn . .	38, 163
Casearia graveolens Dalz	74, 116
Casearia tomentosa Roxb	75, 116, 164
Cassia absus Linn	31, 71, 163
Cassia acutifolia Delile .	71
Cassia alata Linn . .	71
Cassia angustifolia Vahl	31, 71, 109, 163
Cassia auriculata Linn	103
Cassia fistula Linn	71, 97, 100, 164
Cassia obovata Collad .	71
Cassia obtusifolia Linn	97
Cassia occidentalis Linn . .	164, 175
Cassia tora Linn .	97, 103, 164
Casuarina equisetifolia Linn . .	103, 165
Cassytha Linn	164
Cassytha filiformis Linn	87, 103, 116, 164
Catha edulis Forsk. . .	165
Ceasalpinia digyna Rottles .	254
Cedrela toona Roxb .	31, 103, 165
Cedrus deodara Loud .	165
Celastraceae	68
Celastrus paniculatus Willd	32, 111, 165
Celsia coromandeliana Vahl	103
Cenchrus pauciflorus Benth .	165
Centella asiatica (Linn) Urban. .	32, 76, 103
Centipeda orbicularis Lour .	165
Centratherum anthelminticum Kuntze .	32, 78, 116
Cephaelis ipecacuanha A Rich .	32, 77, 165
Cephalandra indica Naud .	34, 166
Cerbera manghas Linn. .	32, 80, 116
Cerbera odollam Gaertn. . .	32, 80, 116, 166
Ceropegia tuberosa Roxb .	103

PAGE

Chamaenerium angustifolium Schur 167
Cheiranthus cheiri Linn 167
Chenopodiaceae . . 85
Chenopodium Linn . 167
Chenopodium album Linn. . 167
Chenopodium ambrosioides Linn 32, 33, 85, 167, 168
Chenopodium botrys Linn . 33, 85
Chionachne koenigii (Spreng) Thw 97
Chloroxylon swietenia Dc . 168
Chrozophora rottleri A. Juss ex Spreng 88
Chrozophora tinctoria Hook f . . 88
Chrysanthemum coccineum Willd 78, 116
Chrysanthemum cinerariaefolium Vis. 78, 116, 168
Chrysanthemum roseum Adam . 78
Cicer arietinum Linn . 16
Cichorium intybus Linn 33, 196
Cicuta virosa Linn . . 76, 169
Cimicifuga foetida Linn. . 63, 97, 116
Cimicifuga racemosa Nutt 169
Cinchona . 33
Cinchona Linn. 169, 170
Cinchona calisaya Wedd. . 77, 111, 116
Cinchona febrifuga 170
Cinchona ledgeriana Moens ex Trimen . 170
Cinchona officinalis Linn 77, 116
Cinchona succirubra Pavon . 77, 117
Cinnamomum Bl 33, 170
Cinnamomum camphora Nees & Eberm 33, 87, 111, 117, 170
Cinnamomum iners Reinw 103
Cinnamomum japonicum Siebold . 171
Cinnamomum zeylanicum Breyn . . 33, 171
Cirnum asiaticum Linn 91
Cirnum lotifolium Linn 91
Cissampelos pareira Linn . . 19, 33, 171
Cissus setosa Roxb . . 113
Citrullus colocynthis Schrad. . 34, 75, 97, 111, 171
Citrullus vulgaris Schrad. . . 75, 171
Citrus Linn. . . 172
Citrus acida Roxb. . . 172
Citrus aurantii-folia (Christm.) Swingle. 34
Citrus aurantium Linn. 173
Citrus bigaradia Loisel. . . 172

PAGE

Citrus decumana Linn 172
Citrus limon (Linn) Burm f 34, 172
Citrus medica var limonis 34
Citrus poonensis Hort ex Tanaka 173
Claviceps purpurea (Fries) Tulasne 173
Clausena willdenowii Wight & Arn 173
Cleistanthus collinus Benth & Hook f 88, 117
Clematis gouriana Roxb 63
Clematis graveolens Lindl . 63
Clematis napaulensis DC 63
Clematis orientalis Linn. 63
Clematis triloba Heyne .. 63
Clematis wightiana Wall 63
Cleome felina Linn 65
Cleome icosandra Linn 34
Cleome viscosa Linn 65, 174
Clerodendron infortunatum Gaertn 34, 174
Clerodendron serratum. (Linn) Moon. 97
Clitoria ternatea Linn 71, 97, 103
Coccinia cordifolia Cogn 34, 97
Coccinia indica W & A 97
Cocculus DC . 174
Cocculus hirsutus Diels . 97
Cocculus laurifolius DC . 174
Cocculus sarmentosus Diels 174
Coffea arabica Linn 77
Cochlearia armoracia Linn . 174
Cocos nucifera Linn . 97, 174, 175
Coffea Linn 175
Coffea robusta Linn. . 175
Colchicum Linn . 175
Colchicum autumnale .. 14, 92
Colchicum luteum Baker . 14, 34, 92
Colocasia virosa Kunth 93
Collybia albuminosa (Berk) Betch 55
Combretaceae . 73
Combretum pilosum Roxb . 175
Commelina asiatica Linn .. 175
Commiphora agallocha Engler .. 97
Commiphora mukul Engl .. 35, 97
Compositae. 77
Coniferae. . 90
Conorhinus rubrofasciatus De Geer. . 248

	PAGE
Convolvulaceae . .	82
Convolvulus arvensis Linn	82
Convolvulus hamadae (Vved) Petrov	175
Convolvulus pluricantes Chois	175
Convolvulus scammonia Linn .	82
Coprinus comatus (Battara) Fr .	55
Coptis teeta Wall . .	35, 176
Corallocarpus epigaeus Benth ex Hook. f	75, 103
Corchorus Linn.	176
Corchorus capsularis Linn	103
Corchorus fascicularis Lam . .	103
Corchorus olitorius Linn	176
Coriandrum sativum Linn	97, 176
Coriariaceae . .	70
Coriaria nepalensis Wall . .	70
Corypha umbraculifera Linn. . .	92, 117
Coscinium fenestratum Colebr .	35, 176
Crassulaceae . .	73
Crataegus oxycantha Linn . .	176, 177
Crataeva nurvala Ham .	98, 177
Cressa cretica Linn .	98
Cretaria sp.	57
Crinum asiaticum Linn . .	177
Crocus sativum Linn	35, 91, 111, 177
Crotalaria juncea Linn	177
Crotalaria medicaginea Lam	177
Crotalaria spectabilis Roth .	177
Croton sp	89
Croton oblongifolius Roxb	88, 117
Croton tiglium Linn.	88, 117, 178
Cryptostegia grandiflora R Br	81, 178
Cubeba officinalis Rafin	178
Cucumis stativus Linn . .	75, 117, 178
Cucumis trigonus .	75, 111
Cucumis utilissimum Roxb .	178
Cucurbitaceae . .	62, 74, 179
Cucurbita pepo D. C . .	178, 255
Cuminum cyminum Linn . .	98, 103, 179
Cupressus torulosa Don. . .	179
Curcuma Linn . .	179
Curcuma amada Roxb. . .	179
Curcuma angustifolia Roxb. . .	98
Curcuma aromatica Salisb. . .	179

PAGE

Curcuma caesia Roxb 179
Curcuma domestica Valeton . . 35, 179
Curcuma longa Linn. . . 35, 117, 179, 180
Curcuma zedoaria Rose . 35, 98, 180
Cuscuta reflexa Roxb . . 36, 82, 111, 180
Cycas revoluta Thunb . . 180
Cyclamen persicum Miller . . 79
Cydonia oblonga Mill. . 103
Cydonia vulgaris Pers. . . 103, 181
Cylista scariosa Roxb. 103
Cymbopogon Hack 181
Cymbopogon caesius Stapf . . 181
Cymbopogon citratus Stapf . 181
Cymbopogon clandistinus Stapf . . 181
Cymbopogon coloratus Stapf 181
Cymbopogon martini Stapf . 182
Cymbopogon nardus (Linn) Rendle 117
Cynanchum arnottianum Wight 81, 117
Cynanchum vincetoxicum Pers . 81
Cynodon dactylon Pers 103
Cynomarathrum nuttallii A Gray . 182
Cyperaceae 94
Cyperus esculentus Linn . 182
Cyperus scarisous R. Br . . 103, 182
Cyperus longus Linn . 94
Cyperus rotundus Linn . 103, 182
Cystopus candidus . 182
Cytisus scoparius Linn . 71

D

Daemia extensa (Linn) R Br . 36, 183, 117
Dalbergia stipulacea Roxb . 71, 117
Daphne cannabina Wall. . . 87
Daphne oleoides Schreb. . . 87
Datura Linn . 187
Datura alba Nees. . . 183
Datura fastuosa Linn. . 83, 183
Datura inoxia Mill. . . 183
Datura metel Linn. . . 83, 183, 255
Datura stramonium Linn . 36, 83, 113, 183
Datura tatula Linn. . . 183, 184

PAGE

Daucus carota Linn . 76, 104, 111, 113, 184
Decalepis hamiltonii W. & A. . . 184
Delphinium ajacis Linn . 113, 184
Delphinium brunonianum Royle . 63, 117
Delphinium caeruleum Jacquem ex Cambess 63, 117
Delphinium elatum Linn 63, 117, 184
Delphinium vestitum Wall. . 63
Dendrocalamus strictus (Roxb) Nees . 94
Derris Lour 184
Derris elliptica (Roxb.) Benth 71, 117, 185
Derris ferruginea (Roxb.) Benth 71, 117, 185
Derris scandens Benth 71, 185
Derris trifoliata Lour var uliginosa (Roxb ex Willd) 117
Derris uliginosa Benth 71, 117
Desmodium gangeticum DC. 104, 109
Descurainia sophia Linn 104
Dichroa febrifuga Lour . 36, 185
Dichanthium annulatum Stapf 185
Dictamnus albus Linn 113
Dictamus fraxinella Pers . 185
Didymocarpus pedicellata Br . 36, 185, 186
Digitalis (Tourn) Linn . 186
Digitalis lanata Ehrh. . 36, 186
Digitalis purpurea Linn. . 84
Dioscorea bulbifera Linn. 91
Dioscoreaceae . . 91
Dioscorea daemona Roxb 91, 117
Dioscorea deltoidea Wall . 91, 117
Dioscorea hispida Dennst. 91, 117, 186
Dioscorea prazeri Prain & Burk . 91, 117
Diospyros ebenum Koenig 80, 117
Diospyros embryopteris Pers 186
Diospyros melanoxylon Roxb. , 98, 104
Diospyros montana Roxb. 80, 117
Diospyros paniculata Dalz. . . 80, 117
Diospyros peregrina Gurkein. 104
Dipterocarpus indicus Bedd . 186, 187
Dodonaea viscosa (Linn.) Jaxq . 69, 117, 187
Dolichandrone falcata Seem . . 84, 111, 117
Dolichos lablab Linn . . 187
Dregea volubilis Benth. ex Hook 37
Droseraceae . 73
Drosera burmanni Vahl. . 73

PAGE

Drosera peltata Sm. var lunata Clarke 73
Drosera spathulata Labill. 73
Drynaria quercifolia J. Sm .. 98, 109
Dryopteris barbigera . 58
Dryopteris blanfordii .. 58
Dryopteris calcarata . 58
Dryopteris marginata . 58
Dryopteris odontoloma . 58
Dryopteris schimperianum .. 58
Duranta plumieri Jacq .. 84, 117
Duranta repens Linn. . 84, 117

E

Ebenaceae 80
Edgeworthia gardneri Meissn 87, 117
Eichornia Kunth .. 187
Elaeodendron glaucum Pers . 68
Elaeocarpus serratus Linn 104
Elaeis guineensis Jacq . 187
Elaeocarpus tuberculatus Roxb 109
Eleucine coracana Gaertn 188
Elettaria cardamomum Maton . 37, 98, 188
Embelia Burm .. 188
Embelia tsjeriam-cottam A DC. . 98
Embelia ribes Burm .. 188
Emblica officinalis Gaertn. . 188
Enhydra fluctuans Lour .. 188
Enicostema littorale Blume . 188
Entada phaseoloides (Linn) Merr 37, 71
Entada pursaetha DC. 37, 117, 189
Entada scandens Benth .. 37, 71, 117, 255
Entoloma microcarpum Berk & Broome. 55
Ephedra Linn . 189
Ephedra gerardiana Wall. .. 37
Ephedra nebrodensis Tin. .. 189
Ephedra vulgaris Rich. .. 37, 189
Equisetum arvense .. 58
Eremostachys acanthocalyx Boiss. 85
Eremostachys superba Royle ex Benth 117
Eremostachys vicaryi Benth. .. 85, 118
Ericaceae .. 79

PAGE

Erigeron candensis Linn. .. 78, 104, 113, 100
Ervatamia dichotoma (Roxb) Blatter .. 80
Erycibe paniculata Roxb 109
Erythrina indica Lam. .. 190
Erythroxylum coca Lam 37, 67
Erythroxylum monogynum Roxb 109, 190
Eucalyptus Lher .. 190
Eucalyptus L W Erit 38
Eucalyptus citriodara Hook . 190
Eucalyptus globulus Labill 73, 118, 190, 191
Eugenia aromatica Kuntze. . 38
Eugenia jambolana Lam . 191
Eugenia operculata Roxb. 108
Eulophia R Br 191
Eulophia nuda Lindl . 98
Euonymus tingens Wall .. 38
Eupatorium odoratum Linn 78, 118
Eupatorium uticifolium L f 78
Euphorbia Linn 191
Euphorbia acaulis Roxb . 88, 113
Euphorbiaceae 87
Euphorbia antiquorum Linn 88, 104, 113, 118
Euphorbia cattimandoo W Elliot 88, 113
Euphorbia helioscopia Linn. .. 88, 109, 113, 191
Euphorbia hirta Linn . 38, 88, 104
Euphorbia hypericifolia Linn 88, 104
Euphorbia neriifolia Linn 88, 113, 118
Euphorbia nivulea Buh Ham 88, 113
Euphorbia peplis Linn 113, 191
Euphorbia pilulifera Linn 38, 291
Euphorbia rothiana Spreng. 113
Euphorbia royleana Boiss 88, 113, 118
Euphorbia thomsoniana Boiss . 88, 113
Euphorbia thymifolia Linn. . 88, 104, 118, 191
Euphorbia tirucalli Linn 88, 111, 113, 118, 192
Euphorbia trigona Haw. .. 88, 113
Evolvulus alsinoides Linn .. 104
Excoecaria agallocha Linn . 88, 111, 113, 118
Exidea polytricha Mont. . 55

F

Fagonia cretica Linn. .. 104, 109
Fagopyrum esculentum Moench . 62, 86, 113
Fagopyrum cymosum Meissn. .. 192

	Page
Fagopyrum tataricum Gaertn . .	86
Feronia elephantum Corr.	104, 192, 225
Feronia limonia (Linn) Swingle	104
Ferula Linn .	192
Ferula foetida Regel.	38, 192
Ferula jaeschkeana Vatke. .	192
Ferula narthex	41
Ficus anthelmintica Martius .	192
Ficus bengalensis Linn	90, 105, 192
Ficus carica Linn .	193
Ficus glomerata Roxb. .	104
Ficus heterophylla Linn .	104
Ficus hispida Linn.	104
Ficus indica Linn	193
Ficus laurifolia Hort et Lam. .	130
Ficus racemosa Linn. .	194
Ficus religiosa Linn. . .	193
Ficus sp .	90
Fimbristylis junciformis Kunth	104
Fistulina hepatica Fr .	55
Flacourtia indica Merr.	109
Flacourtia ramontchi L' Herit .	109
Flemingia tuberosa Dalz	104
Fleurya interrypta Gaud	90, 113
Fluggea leucophyrus (Koen) Willd . .	88, 118
Fluggea microcarpa Blume	88, 118
Fluggea virosa (Roxb ex Willd) Baill	88, 118
Foeniculum capillaceum Gilb	104
Foeniculum panmorium DC.	193
Foeniculum vulgare Mill. .	38, 104
Fritillaria cirrhosa Don Prodr .	98
Fritillaria imperialis Linn. . .	92
Fritillaria roylei Hook	98, 193
Fumaria officinalis Linn.	193

G

Garcinia indica Chois. .	193
Garcinia mangostana Linn. .	104, 194
Garcinia morella Desr.	66, 111, 104
Gardenia campanulata Roxb. . .	118
Gastrochilus pandurata Ridleys. . .	104

	PAGE
Gaultheria fragrantissima Wall.	38, 79, 118, 194
Gaultheria procumbens .	194
Gentiana kurroo Royle. . .	14, 194
Gentiana lutea	14
Geranium macrorrhizum Linn .	194
Geranium robertianum Linn	98
Ginkgo biloba Linn .	113
Girardinia heterophylla Decne	113
Girardinia leschenaultiana Dcne	90
Girardinia zeylanica Dcne .	90
Gloriosa superba Linn	92, 111, 118, 194
Glycosmis pentaphylla Correa	194
Glycyrrhiza glabra Linn	39, 98, 195
Gmelina arborea Linn	98
Gnaphalium luteo-album Linn .	78
Gnetaceae .	90
Gnetum scandens Roxb	118
Gossypium species	66
Gossypium arboreum Linn	98
Gossypium herbaceum Linn	104, 111, 195
Gramineae	94
Graptophyllum pictum (L) Griff	08
Grewia asiatica Linn	98
Grewia hirsuta Vahl. Symb	104, 109
Grewia microcos Linn . .	104, 110
Gymnema aurantiacum Wall. .	195
Gymnema sylvestre R Br	39, 195
Gynandropsis gynandra (Linn) Merr .	65, 118
Gynandropsis pentaphylla DC	65, 118
Gynocardia odorata R. Br.	65, 118

H

Hagenia abyssinica J F Gmel .	195
Haloxylon recurvum Benge ex. Boiss . .	85
Haloxylon salicornicum Bunge. ex Boiss.	85
Hardwickia pinnata Roxb . .	195
Harpullia cupanioides Roxb. . .	69, 118
Hauclea sessilifolia Roxb . .	98
Hedera helix Linn. . .	76, 113, 119
Hedyotis auricularia Linn. . .	42
Helianthus annuus Linn. . .	195

	PAGE
Helicteres isora Linn	39, 105
Heliotropium eichwaldii Steud	82
Heliotropium indicum Linn.	82
Helminthosporium tetramera Mckinney.	195
Helvella crispa Fr	55
Hemidesmus indicus R Br.	39, 105, 196
Heracleum nepalense D Don	196
Herpestis monniera (Linn) H B & K	27, 196
Hibiscus cannabinus Linn	105, 196
Hibiscus esculentus Linn	134, 196
Hibiscus sabdariffa Linn	196, 197
Hibiscus syriacus Linn	197
Hibiscus vitifolius Linn	197
Hippomane mancinella Linn.	113
Hippocratea indica Willd	45
Hirneola polytricha Mont	55
Holarrhena febrifuga Klotzsch	40, 199
Holarrhena mitis R Br	105
Holarrhena antidysenterica Wall	19, 39, 80, 105, 197, 198, 199
Holigarna arnotiana Hook	69, 113, 199
Holigarna ferruginea March	69
Holigarna grahamii (Wight) Hook	69, 113
Holigarna longifolia Buch Ham ex Roxb	69, 114
Homalomena rubescens Kunth	93
Humulus lupulus Linn	114
Hura crepitans Linn	89
Hydnocarpus anthelmintica Pierre	40, 200
Hydnocarpus Gaertn	199
Hydnocarpus kurzii (King) Warb	65, 118
Hydnocarpus laurifolia (Dennst) Sleumer	65, 118
Hydnocarpus wightiana Blume	40, 65, 118, 200
Hydnum caralloides Scop	55
Hydnum repandum Linn	56
Hydrocotyle asiatica Linn.	32, 76, 103, 200
Hydrocotyle javanica Thunb	76, 105, 118
Hygrophila spinosa T Anders	27, 200
Hyoscyamus Linn	201
Hyoscyamus muticus Linn	83, 255
Hyoscyamus niger Linn	40, 83, 201
Hyoscyamus pusillus Linn.	83
Hyoscyamus reticulatus Linn	83
Hypericaceae	66

PAGE

Hypericum japonicum Thunb . . 105
Hypericum perforatum Linn . 66, 114, 201
Hypholoma fasciculare (Huds) Fr 54, 57
Hyptis suaveolens Poit. . . 201

I

Ilex vomitoria (Soland) Hort . 201
Illicium anisatum Linn. . 63
Illicium griffithii Hk f & T . 63
Illicium parviflorum Michx . 201
Illicium religiosum Sieb & Zucc. 63, 201
Illicium verum Hook f . . 63
Indigofera enneaphylla Linn . . 201
Indigofera linifolia Retz . . 201
Indigofera oblongifolia Forsk . . 105
Indigofera tinctoria Linn . 98
Inula graveolens Desf . 78
Inula helenium Hk f & T . 201
Inula royleana DC . 202
Ipomoea Linn . 82, 83, 202
Ipomoea aquatica Forsk 82
Ipomoea hederacea Jacq 40, 82, 202
Ipomoea muricata Jacq 82, 116, 202
Ipomoea pescaprae Roth 202
Ipomoea pulchella Roth 202
Ipomoea purga Heyne . 82
Ipomoea reptans (Linn) Poir 82, 102
Ipomoea turpethum R Br . 40, 83
Iridaceae 90
Ixora nigricans Br. . 105

J

Jasminum Linn. 202
Jasminum auriculatum Vahal 98
Jateorrhiza palmata (Lam) Miers . . 105, 203
Jatropha curcas Linn . 89, 105, 203
Jatropha glandulifera Roxb . 89, 203, 255
Jatropha gossypiifolia Linn. . 89
Jatropha multifida Linn. . . 89
Juglandaceae . . 90
Juglans regia Linn. . . 90, 105, 118, 203
Juncaceae . . 92

PAGE

Juncus effusus Linn. .. 92
Juniperus communis Linn .. 40, 203
Jussiaea suffruticosa Linn .. 105
Justicia gendarussa Burm .. 105

K

Kaempferia galanga Linn 303
Kalanchoe spathulata (Poir.)DC .. 110, 118

L

Labeo rohita .. 248
Labiateae .. 85
Lactarius deliciosus Fr .. 56
Lactuca tatarica C A Meyer var. tibetica C B Clarke. 78
Lactarius vellerens Fr. . 54, 57
Lactuca scariola Linn .. 98, 204
Lagenandra ovata (Linn) Thw. .. 93, 118
Lagenandra toxicaria Dalz .. 93, 118
Lagenaria siceraria Standl . 75
Lagenaria vulgaris Ser .. 75, 204
Lagerstroemia flosregineae Retz .. 74
Lagerstroemia indica Linn .. 74
Lagerstroemia speciosa (Linn) Pers 74
Lamium amplexicaule Linn . 85
Lansium anamalayanum Bedd . 204
Lansium domesticum Correa 204
Lantana aculeata Linn. 85
Lantana camara Linn .. 85, 204
Laportea crenulata Gaud .. 90, 114
Laportea terminalis Wight .. 90, 114
Lasiosiphon eriocephalus Decne . 87, 114, 118
Lathyrus aphaca Linn . 71
Lathyrus sativus Linn. 61, 71, 204
Lauraceae .. 87
Laurus camphora Linn. .. 205
Lavendula burmanii Benth. .. 205
Lawsonia inermis Roxb. .. 205
Leea aequata Linn. .. 99
Leea indica Merrill. .. 105
Leguminosae. .. 70, 205
Lens esculenta Moench. .. 105, 205
Lens culinaris Medic. .. 105
Lentinus subnudus Berk. .. 56

PAGE

Leonurus cardiaca Linn . 114
Lepidium draba Linn . 65, 116
Lepidium latifolium Linn. 205
Lepidium sativum Linn 111
Lepiota sp. 57
Lepiota albuminosa Berk. 55
Lepiota cristta Quel 55, 57
Lepiota mastoidea Fr 56
Lepiota procera (Scop) Sacc 56
Leucaena glauca Benth. 205
Leucas aspera Spreng 205
Leishmania tropica . 29
Liliaceae 91
Linaceae 67
Linostoma decandrum Wall . 118
Linum usitatissimum Linn 61, 67, 99, 205
Litsaea Lamk. . 205
Litsaea chinensis Lam. 99, 205
Litsaea zeylanica Nees . 206
Lobelia Linn 206
Lobelia excelsa Leschen. 78, 114
Lobelia inflata Linn . . 206
Lobelia nicotianifolia Heyne . 78, 114, 206
Lobelia pyramidalis Vahl . 206
Lochnera pusilla K Schum 81
Lochnera rosea (Linn) Reichb . 81
Lodoicea maldivica Pers . . 110
Lodoicea seychellarum Labill 110
Loganiaceae 82
Lolium perenne Linn . 94
Lolium temulentum Linn 54, 94, 95, 906
Loranthaceae 87
Loranthus asper Lour. . . 99
Loranthus longiflorus Den 206
Luffa acutangula Roxb. . 40, 75, 99, 105
Luffa aegyptiaca Mill & Hook f. . 75, 206
Luffa amara Roxb . . 206
Luffa cylindrica Linn . . 207
Luffa echinata Roxb . . 75, 29, 207
Luffa graveolens Roxb. . 207
Lupinus (Tourn.) Linn. . . 207
Lupinus albus Linn. . . 207
Luvunga scandens Ham. . 40, 99, 207

PAGE

Lycium barbarum Linn . 83
Lycoperdon sp 56
Lycopodium clavatum Linn 207
Lycopodium phlegmaria Linn . 207
Lythraceae . . 74

M

Maba nigrescens Dalz . . 208
Machilus macrantha Nees . 99
Madhuca longifolia (Linn.) Macbride 41, 80, 119
Madhuca latifolia (Roxb) Macbride 41, 118
Madhuca indica J F Gmel 41, 80
Maesa indica Wall 80, 119
Magnoliaceae . 63
Mahonia acanthifolia Don 208
Mahonia nepalensis DC . 41
Mahonia Nutt . 208
Malanogaster durissiums Cooke. . . 56
Mallotus philippinensis Muell -Arg 41, 208, 209
Malvaceae . 66
Malva arborea Webb . . 209
Malva parviflora Linn 67
Malva rotundifolia Linn 105
Malva sylvestris Linn . . 209
Mandragora caulescens Clarke . 83
Mangifera Linn . . 209
Mangifera indica Linn . 105, 209
Manihot esculenta Crantz 89, 209
Manihot utilissima Pohl. 89, 209
Mansonia dipikae C S. P 209
Marrubium vulgare Linn . 110, 210
Martynia annua Linn. . 99
Martynia diandra Glox. . . 210
Matricaria chamomilla Linn . 210
Matricaria Linn. . 210
Meconopsis aculeata Royle . 64
Meconopsis napaulensis DC 64
Melaleuca leucadendron Linn. 73, 119
Melanogaster durissiums Cooke . 56
Melastoma malabathricum Linn. . . 105
Meliaceae . 68
Melianthus major Linn. . 69

PAGE

Melia azadirachta Linn .. 97, 116, 148
Melia azedarach Linn. .. 41, 68, 210, 255
Melilotus alba Desr .. 71
Melilotus parviflora Desf. . 210
Melissa parviflora Benth. . 99
Melodinum monogynus Roxb. . 81, 119, 255
Menispermaceae . 64
Mentha arvensis Linn. .. 42, 255
Mentha aquatica Linn. . 210
Mentha longifolia Hunds .. 105
Mentha sylvestris Linn. .. 105
Mentha viridis Linn. . 211
Mercurialis annua Linn. .. 211
Mesua ferra Linn. . 105, 211
Microcystis flos-aquae (Wittr) Kirch . 53
Millettia auriculata Baker ex Brand. . 71, 119
Millettia pachycarpa Benth . 71, 119, 211
Millettia piscidia Wight & Arn .. 71, 119
Mimosa pudica Linn .. 106, 211
Mimusops hexandra Roxb. .. 99
Mimussops elengi Linn . 106
Mirabilis jalapa Linn. . 211
Momordica balsamina Linn .. 75
Momordica charantia Linn . 75, 110, 111, 211
Momordica cymbalaria Fenzl ex Naud . 75, 111
Momordica dioica Roxb .. 99, 211
Momordica tuberosa Cogn .. 75, 111
Monniera cuneifolia Michx. . 211
Morchella esculenta Pers . 56
Morinda citrifolia Linn 106, 212
Morinda umbellata Linn. . 106
Moringaceae. . 70, 212
Moringa concanensis Nimm. . 212
Moringa oleifera Lamk . 42, 70, 99, 111, 134, 212
Moringa pterygosperma Gaertn. . 42, 70, 111, 212
Morus indica Linn. .. 213
Morus nigra Linn. . 213
Mucuna atropurpurea DC. .. 114
Mucuna gigantea DC .. 114
Mucuna hirsuta Wight & Arn. .. 114
Mucuna monosperma DC .. 114
Mucuna pruriens DC. .. 114, 213
Mucuna prurita Hook. .. 106

PAGE

Mucor sp. . 54
Mundulea sericea (Willd) Greenway. . 119
Mundulea suberosa Benth . 71, 119, 213
Murraya exotica Linn 213
Murraya koenigii Spreng 106, 213
Murraya paniculata Jack 106
Musa sapientum O Kuntze 99, 106, 110
Musa paradisiaca Linn var sapientum Kuntze. 110
Musua ferrae 255
Mylabris macilenta. 248
Myricaceae. 90
Myrica nagi Thunb . 90, 106, 119
Myristicaceae. 86
Myristica fragrans Houth. . 42, 87, 213
Myristica malabarica Lamk . . 87
Myrsinaceae . . 79
Myrsine africana Linn . 42, 213
Myrtaceae . 73
Myrtus communis Linn . 106

N

Nannorhops ritchieana H Wendl . 106
Narcissus tazetta Linn . . 91, 213
Nardostachys jatamansi DC 214
Nepeta elliptica Royle ex Benth . . 106
Nerium indicum Mill 81, 111
Nerium odorum Soland . 81, 111, 214
Nerium oleander Linn 114, 214
Nicandra physaloides Gaertn. . . 83, 119
Nicotiana Linn . 214
Nicotiana glauca Grab. . . 214
Nicotiana rustica Linn . . 83, 119, 214
Nicotiana tabacum Linn. . 83, 119, 214, 215
Nigella sativa Linn . 63, 112, 119
Nodularia sp . 53
Nuphar luteum Sibth. & Sm . . 215
Nymphaea (Tourn) Linn . 215
Nymphaea alba Linn . 106, 215
Nymphaea lotus Linn . . 215
Nymphaea pubescens Willd. . 106

O

Ochna pumila Ham ex D Don. . 99
Ocimum Linn . 215

	PAGE
Ocimum americanum Linn	106
Ocimum basilicum Linn.	106, 203, 215
Ocimum canum Sims	106, 203, 215
Ocimum gratissimum Linn	119, 215
Ocimum kilimandscharicum Guerke.	216, 255
Ocimum lemon	203
Ocimum sanctum Linn	203, 216
Odina wodier Roxb	216
Oldenlandia auricularia K. Schum	42, 110, 106
Oldenlandia biflora Hk. f	216
Oldenlandia umbellata Linn.	99
Olea Linn.	216
Ononis spinosa Linn	216
Operculina turpethum (Linn) Manso	83, 99
Origanum vulgare Linn	216
Oroxylum indicum Vent	106, 216, 217
Orthosiphon staminеus Benth	217
Oscillatoria	53
Ougenia dalbergioides Benth	71, 119
Oxalis corniculata Linn	106
Oxytropis thomsoni Benth	217

P

Paeonia emodi Wall.	63
Pachygone ovata (Poir.) Miers	64, 119
Palmaceae.	92
Pandanus odoratissimus Roxb	217
Panicum maximum Jacq.	94
Papaveraceae.	64
Papaver dubium Linn.	64
Papaver nudicaule Linn.	64
Papaver rhoeas Linn.	64
Papaver somniferum Linn.	42, 64, 217
Paris polyphylla Sns.	217
Parmelia sp	57
Parmelia kamtschadalis.	256
Parmelia molliuscula.	57
Paspalum scrobiculatum Linn.	95
Passifloraceae.	74
Pavonia odorata Willd.	107
Pedaliaceae.	84

PAGE

Peganum harmala Linn 43, 67, 112, 119, 218
Penicilliopsis clavariaeformis Solms Laub 218
Penicillium notatum. 54
Pentapetes phoenicea Linn 125, 218
Peranospora parasitica. 218
Perowskia atriplicifolia Benth 218
Peucedanum graveolens Benth. 43, 107
Phalaris minor Retz. 218
Phaseolus aconitifolius Jacq 99, 108
Phaseolus lunatus Linn 72
Phaseolus radiatus Linn. 218
Phaseolus trilobus Art 99, 218
Phebalium argentum Sm. 219
Phyllanthus emblica Linn 107, 219
Phyllanthus niruri Linn 107, 219
Phyllanthus urinaria Linn 89, 107, 119
Physochlaina praelta (Don) Miers 83, 219
Phytolaccaceae 85
Phytolacca acinosa Hook f B I non-Roxb 86
Phytolacca latbenia (Buch Ham) H Walt 86
Picrasma excelsa Swartz 14, 219
Picrasma javanica Blume var nepalensis Badhwar nov comb 119
Picrasma napalensis Benn. 68, 119
Picrasma quassioides. 14
Picroena quassioides Benth 43
Picrorhiza kurroa Royle ex Benth 43, 219, 220
Pieris ovalifolia D Don 78, 119
Pimpinella anisum Linn 43
Pimpinella saxifraga Linn 220
Pinus Linn 220
Pinus excelsa Wall. 220, 221
Pinus gerardiana Wall 221
Pinus khasya Royle 221
Pinus longifolia Roxb 43, 99, 221, 223
Pinus roxburghii Sargent 43, 99
Piper sp 86
Piperaceae. 86
Piper betle Linn. 44, 86, 222, 254
Piper chaba Hunter. 99
Piper cubeba Linn. 44, 222
Piper longum Linn. 99, 107

PAGE

Piper nigrum Linn. . 86, 107, 222
Pistacia integerrima Stew 44, 99, 107, 222, 223, 254
Pistacia terebinthus Linn . 223
Pistia stratiotes Linn 99, 107
Pithecelobium bigeminum Benth 72, 119
Pithecelobium saman Benth . 223
Plantago ciliata Desf 107
Plantago major Linn . 99, 107, 223
Plantago ovata Forsk 19, 44, 107, 323
Plantanus orientalis Linn 197
Plesmonium margaritiferum Schott 93
Pleurotus cretaceus Massee 56
Pleurotus fimbriatus Bolt 56
Plumbaginaceae 79
Plumbago indica Linn 44, 76, 112
Plumbago rosea Linn 44, 79, 112, 223
Plumbago zeylanica Linn . 45, 79, 99, 197, 112, 223
Plumeria acuminata Ait 81, 112
Plumeria acutifolia Poir 81, 112, 223
Plumeria rubra Linn var acutifolia Bailey 112
Podophyllum Linn 223
Podophyllum emodi Wall ex Hook f. & Thoms 64, 114
Podophyllum hexandrum Royle 45, 64, 114
Podophyllum paltatum . 45
Pogostemon heyneanus Benth . . 85, 119
Pogostemon patchouli Hook f . . 85, 119
Poinciana pulcherima Linn . . 110
Polianthese tuberosa Linn . . 256
Polygala chinensis Linn . 66, 223
Polygala crotalarioides Buch Ham . . 66
Polygalaceae. . . 66
Polygala telephioides Willd. . . 66
Polygonaceae . . 86
Polygonum Linn. . . 224
Polygonum aviculare Linn . . 86
Polygonum flaccidum Meissn. . 86, 119
Polygonum hydropiper Linn. . . 86,114, 119
Polygonum orientale Linn. . . 86
Polygonum persicaria Linn . . 86
Polygonum tomentosum Willd . . 86
Polyporus squamosus (Huds) Fr . . 56
Polytoca barbata Stapf. . 97
Pongamia glabra Vent. . . 45, 72, 119, 224

PAGE

Pongamia pinnata (Linn) Merr . . 45, 72, 119
Potentilla anserina Linn . 224
Potentilla argentea Linn . 224
Prangos pabularia Lindl. . . 224
Premna integrifolia Linn. . 45, 224
Primulaceae . 79
Primula reticulata Wall. . . 79
Prinsepia utilis Royle. . 234
Pristimera indica Willd. . 45, 224
Prosopis juliflora DC . 224
Prunus acuminata Hk f . 225
Prunus amygdalus Batsch . 72
Prunus armeniaca Linn . 72
Prunus avium Linn 72, 225
Prunus cerasus Linn. . . 72, 225
Prunus mahaleb Linn. 72
Prunus nepalensis Koch. . 225
Prunus padus Linn . 72
Prunus persica Stokes . 72
Prunus puddum Roxb . 72, 225
Prunus undulata Buch Ham . 72
Psalliota xanthoderma 57
Psidium guajava Linn . 107, 110, 225
Psilotum triquetrum Sw . 225
Psoralea corylifolia Linn. . 19, 45, 225, 226
Psychotria ipecacuanha Stokes . . 32
Pteridophyta (Vascular Cryptogams) 57
Pteris acquilina Linn . 58, 226
Pterocarpus marsupium Roxb . . 226
Pueraria thunbergiana Benth . . 226
Pulicaria dysenterica Gaertn. . . 107
Punica granatum Linn . . 107, 226
Putranjiva roxburghii Wall . . 226
Pygeum gardneri Hook f . 72, 19
Pyrethrum . . 168, 18
Pyrus aucuparia Linn. . 72
Pyrus malus Linn . . 72

Q

Quercus Linn. . . 266
Quercus dilatata Lindl. . . 227
Quercus ilex Linn . . 227

PAGE

Quercus incana Roxb. . 227
Quisqualis indica Linn . 227

R

Randia dumetorum Lamk 46, 77, 112, 119, 227
Randia uliginosa DC. . 77, 119
Ranunculaceae . . 62
Ranunculus arvensis Linn . 63
Ranunculus falcatus Linn. . . 63
Ranunculus laetus Wall . . 63
Ranunculus lingua Linn 63
Ranunculus pensylvanicus Linn f. 63
Ranunculus sceleratus Linn. 63, 114
Rauwolfia Linn. . 227
Rauwolfia canescens Linn. 46, 227, 228, 256
Rauwolfia micrantha 262
Rauwolfia serpentina Benth 19, 46, 120, 228, 229, 230, 231, 255, 257, 258
Revenelia sessilis . 231
Rhamnus Linn. 231
Rhamnus virgatus Roxb 231
Rheum Linn . . 231
Rheum emodi Wall . 46, 86, 107, 231, 232
Rhizodonia solani Kuhn 216
Rhododendron arboreum Sm . 79
Rhododendron anthopogon D Don. 79
Rhododendron barbatum Wall ex G Don 79, 120
Rhododendron campanulatum D Don. . . 79, 100
Rhododendron cinnabarinum Hook f . 79
Rhododendron falconeri Hook f. 120
Rhododendron setosum D. Don . 79
Rhus insignis Hook. f. . 69, 114
Rhus punjabensis J L. Stew. ex Brand. . 69, 114
Rhus succedanea Linn . . 69, 100, 114
Rhus wallichii Hook. f. . . 69, 114
Ricinus Linn. . . 232
Ricinus communis Linn. . . 46, 89, 120, 232
Robinia uliginosa Roxb ex Willd . . 117
Rosa Linn. . . 232
Rosa damascena Mill . . 47
Rourea santaloides W. & A. . . 100
Rubiaceae. . . 77
Rubia cordifolia Linn. . . 107

PAGE

Rubus moluccanus Gaertn . . 73, 112
Rumex Linn . . 232
Rumex acetosa Linn. . 86, 114
Rumex acetosella Linn . . 86, 114
Rumex scutatus Linn. . . 107
Rumex vasicarius Linn . 107
Rutaceae . . 67
Ruta graveolens Linn var angustifolia Hk f . . 67, 112, 114, 120, 232
Ruta tuberculata Forsk. . . 67

S

Saccharum spontaneum Linn . . 233
Salicornia brachiata Roxb . . 85, 112
Salix Linn . . 233
Salix alba Linn 107
Salsola kali Linn. . . 85
Salvadoraceae . . 80
Salvadora oleoides Dene . 80, 233
Salvadora persica Linn. . 80
Sambucus Linn . . 233
Sambucus ebulus Linn . . 77
Sambucus nigra Linn. . 77
Samydaceae . . 74
Sansevieria roxburghiana Schult . . 100
Sansevieria zeylanica Willd. . . 233
Santaloides minus Schellenb. . . 100
Santalum album Linn. . . 47, 120, 233, 134
Sapindaceae . . 69
Sapindus Plum . . 234
Sapindus laurifolius Vahl. . . 234
Sapindus mukorossi Gaertn. . . 69, 120, 234
Sapindus trifoliatus Linn. . . 69, 100, 107, 110, 112, 120
Sapium indicum Willd. . . 89, 120
Sapium insigne Trimen. . . 89, 114
Saponaria Linn. . . 234
Saponaria vaccaria Linn. . . 66
Sapotaceae . . 80
Sphaeranthus indicus Linn. . . 100
Saraca indica Linn. . . 47, 107, 234
Sarcococca pruniformis Lindll. . . 234
Sarcostema acidum (Roxb.) Voigt. . . 82, 120
Sarcostema brevistigma W. & A. . . 82, 120

	PAGE
Saurornatum guttatum Schott. . .	93
Saussurea lappa Clarke .	47, 78, 120, 234, 235
Sachima wallichii Choisy.	115
Schleichera oleosa (Lour) Merr. . .	69, 110, 120
Schleichera trijuga Willd	69, 110, 120
Schweinfurthia spherocarpa Br. .	110
Scilla indica Baker. . .	14, 47, 92, 263
Scilla maritima .	14
Scindapus officinalis Schott. .	107
Scirpus corymbosus Heyne .	9
Scirpus grossus Linn .	235
Scleria pergracilis (Nees) Kunth.	120
Scopolia Jacq .	235
Scopolia anomola (Linn et Otto) Airy-Shaw.	83
Scopolia lurida Dunal .	83
Scrophulariaceae	84
Secale cereale Linn .	235
Secamone emetica R Br.	82
Securigera securidaca Linn. .	235
Semecarpus anacardium Linn. .	447, 70, 108, 112, 115, 235
Semecarpus travancoricus Bedd.	70, 115, 236
Senecio sp .	78
Senecio jacobaea Linn	236
Senecio retrorsus DC .	236
Senecio vulgaris Linn.	78, 236
Sesbania aegyptiaca Pers . .	100
Sesbania sesban (Linn.) Merr. . .	100
Sesamum indicum Linn .	84, 112
Sesamum orientale Linn	84, 112
Seseli indicum Wight & Arn.	236
Setaria palmaefolia Stapf .	236
Shorea robusta Gaertn f. .	108
Sida cordifolia Linn .	48, 100, 236
Sida rhombifolia Linn	67, 100
Sida spinosa Linn. .	108
Simaroubaceae	68
Sinapis alba Linn. .	65
Sisymbrium irio Linn. .	236
Skimmia laureola Sieb. & Zucc. ex Walp .	48, 68, 236
Smilax prolifera Roxb. . .	108
Solanaceae. . .	83
Solanum coagulans Forsk. . .	84
Solanum dulcomara Linn. . .	84

PAGE

Solanum incanum Linn. 84
Solanum indicum Linn. .. 237
Solanum melongena Linn .. 110
Solanum nigrum Linn. 84, 108, 237
Solanum pseudocapsicum Linn . 237
Solanum spirale Roxb. 84
Solanum trilobatum Linn 100
Solanum torvum Sw. 237
Solanum tuberosum Linn 62, 84
Solanum xanthocarpum Schrad & Wendl 237
Sonneratia acida Linn 48, 237
Sophora mollis R Grah 72, 120
Sophora tomentosa Linn 72, 110
Sorghum halepense Pers . 61, 95
Sorghum sacchoratum Pers 95
Sorghum vulgare Pers 61, 95
Soymida febrifuga A Juss .. 108
Spartium scoparium Linn . 238
Sphaeranthus indicus Linn 78, 120, 238
Spilanthes acmella Murr . 108, 238
Spiraea aruncus Linn 238
Spondias pinnata Kurz 100
Stachytarpheta jamaicensis Vahl var. indica Lam. 85, 108, 112
Stachytarpheta indica Vahl. . 85, 108, 112
Stelaria media Linn . 238
Stephania glabra Roxb 48, 100, 238
Stephania hernandifolia (Willd.) Walp. .. 120
Steudnera virosa (Kunth) Prain . 93
Stipa sp 95
Stipa siberica 61
Streblus asper Lour 100
Streptomyces auriofociens . 52
Streptomyces griseus .. 52
Streptomyces rimosus . 52
Streptomyces veneznelae 52
Stropharia semiglobata (Batsch) Quel . 54, 57
Strychnos colubrina Linn. 82, 120
Strychnos nux-vomica Linn .. 48, 82, 87, 108, 110, 120, 238
Suaeda fruticosa Forsk. 85
Sunantherias sylvatica Schott. . 93
Swertia chirata Ham. 48, 238
Swertia decussata Nimmo. .. 238
Swietenia macrophylla Linn. .. 238, 239

PAGE

Symplocos racemosa Roxb. 49, 108, 239
Syzygium cumini (Linn.) Skeels 108
Syzygium jambos (Linn) Alst. 108
Syzygium operculatum Gamble 108

T

Tabernaemontyna coronaria Br . 239
Tabernaemontana dichotoma Roxb. . . 81
Taccaceae . 91
Tacca pinnatifida Forsk. . . 91
Tagetes erecta Linn. . . 239
Tagetes patula Linn. . 239
Tamarindus indica Linn . 239, 240
Tanacetum Linn . 240
Taraktongenos kurzii King . 65, 118
Taxus baccata Linn . 90, 112, 120, 240
Taxus brevifolia Nutt . . 240
Tectona grandis Linn f 108, 240
Tephrosia Pers. . 240
Tephrosia candida (Roxb) DC 72, 120
Tephrosia purpurea (Linn.) Pers. 72, 120, 240
Tephrosia villosa Pers. . 240
Tephrosia vogelii Hook f . 120
Teramnus labialis Spreng . 100
Terminalia arjuna W & A . . 49, 108, 241
Terminalia bellerica Roxb. 73, 120, 241
Terminalia chebula Retz . 73, 108, 110, 241
Terminalia citrina Fleming. . 108, 110
Terminalia paniculata Roth. 110
Ternstroemiaceae . 66
Thalictrum foliolosum DC . 94, 241
Thea sinensis Linn. . . 66, 241
Thevetia nereifolia Juss. . 49, 81, 112, 120, 241, 242
Thevetia peruviana (Pers) Merr. . . 49, 81, 112, 120
Thomsonia nepalensis Wall. . . 94
Thymeliaceae. . . 87
Thymus serphyllum Linn. . . 242
Thymus vulgaris Linn. . . 242
Thysanolaena maxima Kuntze. . . 242
Tinospora crespa Miers. . . 242
Tinospora cordifolia Miers. . . 49, 242

PAGE

Toddalia aculeata Pers 49, 242
Toddalia asiatica Lam. 49
Trachyspermum ammi (Linn) Sprague 50, 110
Tragia bicolor Miq. 89, 115
Trapa bispinosa Roxb. 242
Tragia involucrata Linn. 89, 115
Trespasia populnea 243
Trianthema monogyna Linn. 76, 112
Trianthema pentandra Linn. 76, 112
Trianthema portulacastrum Linn. 76, 112, 243
Tribulus terrestris Linn. 50, 67, 243
Trichosanthes anguina Linn. 243
Trichosanthes bracteata Voigt. 75
Trichosanthes cucumerina Linn 75
Trichosanthes dioica Roxb. 75, 243
Trichosanthes palmata Roxb. 75
Trifolium repens Linn 72
Trigonella foenum-graecum Linn 120, 243
Triticum Linn 242
Triticum aestivum Linn. 95
Triumpheta bartramia Linn 108
Triumfetta rhombodea Jacq 243
Tussilago farfara Linn 244
Tylophora asthmatica W. & A 50, 82, 108, 244
Tylophora fasiculata Buch 82
Tylophora indica (Burm f) Merr. 50, 82
Typha elephantina Roxb 108
Typhonium trilobatum Linn. Schott 94

U

Umbelliferae 76
Uraria lagopoides DC. 108
Urena lobata Linn. 112
Urginia coromandeliana Hook. f. 92
Urginia indica Kunth. 50, 92, 244
Urginia scilla 92
Urticaceae 89
Urtica dioica Linn 90, 100, 115
Urtica hyperborea Jacq. 90, 115
Urtica parviflora Roxb. 90, 115
Urtica pilulifera Linn. 90, 115

PAGE

Uroloncha malabarica Linn . 248
Usnea hirta 263

V

Vaccinium Linn . 244
Valariana Linn. . 244
Valariana wallichii DC. . . 50
Valariana officinalis Linn . 244
Vanda roxburghii R. Br 50, 244
Vanda spathulata Spreng . 100
Vangueria spidosa Roxb . 244
Vateria indica Linn . 100, 109
Verbascum thapsus Linn . 84, 100, 121
Verbenaceae . 84
Verbena officinalis Linn. 85, 245
Verbena venosa . . 245
Vernonia anthelmintica Willd . 32, 78, 116, 245
Vernonia cinerea Less . 100
Vernonia roxburghii Less . . 100
Vetiveria zizanioides Stapf . . 245
Vicia sativa Linn . 72
Vigna catjang Walp. . 245
Vinca pusilla Murr. . . 81
Vinca rosea Linn 81, 246
Viola odorata Linn . . 246
Viscum Linn. . 246
Viscum album Linn . . 87, 246
Vitex negundo Linn . . 121, 246
Vitex peduncularis Wall. . . 20, 51, 246
Volvaria diplasia Berk & Broome. . . 56
Volvaria gloicephala Gill . . 55, 57
Volvaria terrestria Berk & Broome. . . 56

W

Wallichia disticha T Anders . . 93, 115
Walsura piscidia Roxb. . . 68, 121
Wikstroemia indica (Linn) C.A. Mey. var viridiflora (Meissn) Hook f . . 87, 121
Withania somnifera Dunal. . . 51, 84, 100, 112
Woodfordia fruticosa Kurz . . 109
Wrightia tinctoria R. Br. . . 109, 246

PAGE

X

Xanthium strumarium Linn. .. 51, 78, 115, 246
Xanthyoxylum alatum. .. 254

Z

Zanonia indica Linn. . 75
Zanthoxylum acanthopodium DC. .. 247
Zanthoxylum alatum Roxb. .. 68, 121, 247
Zanthoxylum budrunga Wall. .. 110, 247
Zanthoxylum hamiltonianum Wall . 121
Zanthoxylum ovaltifolium Wight. .. 247
Zanthoxyluu rhetsea DC. . 247
Zea mays Linn . 95
Zingiber officinale Rosc. .. 51, 247
Zizyphus jujuba Lam .. 100
Zizyphus vulgaris Lam. . 248
Zizyphus xylopyrus Willd. . 248
Zygophyllaceae .. . 67

INDEX

OF

VERNACULAR NAMES

	PAGE
A	
Aal (Bo.)	106
Aaraar (H)	40
Ach (H & B)	106
Acha (M)	80
Ada (B.)	51
Adhatodai (M)	23
Adrak (H)	51
Adrakam (S)	51
Adulsa (H & B.)	23
Adutina-palai (M)	86
Afim (H & B.)	42, 64
Agave Family (English)	91
Agla (Kumaon.)	70
Agnigarva (S)	74
Agnisikha (M)	92
Agnad (B)	120
Ahiphena (S)	42, 64
Ailanto (English)	101
Ajakarua (S)	100
Ajmod (Vern)	254
Ajmoda (S. & H.)	76
Ajowan (H , Bo & Vern)	50, 163
Ak (H)	30, 81
Akanda (B & Bo.)	30, 81
Akanadi (Vern.)	33
Akar Kanta (B)	23
Akarkara (Bo & P.)	108
Akasbel (H)	36, 82
Akhroot (H. & B.)	90
Ak-Madar (Vern.)	30
Akola (H)	23
Akoria (H.)	69
Akroda (Bo.)	90
Akrottu (M.)	90
Akshota (S.)	90
Aku-jemudu (M.)	88
Al (Bo.)	106
Alangi (M.)	23
Alari (M)	81
Alarka (S)	81, 100
Alasi (Bo)	67
Algusi (B.)	36, 82
Alkaloids (English)	248
Alli (M)	106
Allipalli (P , H & Kash)	112
Alombe (Bo)	55
Alpayushi (S)	92
Alshiviral (M)	67
Amanakham chedi (M)	46, 89
Amaravela (S)	36, 82
Amaryllis Family (English)	91
Amb (H , B & Bo)	105
Ambari (Bo)	105
Amku lang-kalang (M)	51, 84
Amla (Bo)	107
Amlika (S)	106
Amluki (B)	115
Amra (H , B & Bo)	100
Amrataka (S)	100
Amritphala (S)	103
Amrud (H)	110
Amrul (H & B)	106
Amsania (Vern)	37
Amudanda (P)	41
Amumpatchaiarissi (M.)	38
Amva (S)	105
Anantamul (B.)	39
Anannas (H)	91
Ananta (S.)	39
Anar (H)	107
Anaras (B)	91
Anasha pazham (M.)	91
Anashuppu (M)	63
Anasphal (H)	63
Ankra (H.)	72
Ankari (B.)	72
Ankola (Bo.)	23

	PAGE
Ankota (S)	23
Antamul (H. & B)	50, 82
Anthalmul (Bo.)	50, 82
Anthelmintics (English)	248
Antibiotics (English)	249
Aoula (H)	107
Aparajita (S.)	71
Aparājit (H & B.)	71
Aphu (Bo)	42, 64
Arand (H)	89
Archaka (B)	48
Ardana (H)	109
Ardawal (P)	79
Ardisia Family (English)	79
Arfu (Nep)	70
Arghavan (Arab.)	120
Aripu (M.)	85
Arishta (S.)	51, 78
Arjun (H & B)	49
Arjuna (S. & Bo.)	49, 74
Arka (S)	30, 81
Arlu (H)	106
Arni (H & Bo.)	45
Aroid Family (English)	93
Aromatic plants (English)	249
Aru (H)	72
Arusha (Vern)	23
Arvada (M.)	67
Asek (M)	57
Aseru (Nep)	36
Asgandh (H)	51, 84
Ashvakatri (S)	109
Ashwagandha (S., B. & Bo)	51, 84
Asita kutanja (S)	109
Asok (H , B & Bo)	47
Asoka (S)	47
Aspergillus (English)	54
Asundro (Bo)	102
Aswamantaka (S)	102
Ata (B)	64
Ataicha (B.)	22
Atasi (S)	67
Athiballachetu (M)	67
Ati vadayam (M)	22
Atibala (S)	67
Atimaduram (M)	39
Atis (H & Vern.)	22
Ativisha (S)	22
Atmagupta (S.)	106
Atropa (English)	83
Aurhur (Vern)	34
Avaram (Tam.)	103

B

Babachi (H & B)	45
Babla (B)	101
Bach (H & Vern.)	22, 93
Bachata (H.)	112
Bachnag (Vern)	21
Bacteria (English)	52
Badam (H , B & Bo)	72
Badbar (H.)	98
Badari (S.)	100
Badian (Bo)	63
Badrang (H)	110
Bael (H , B & Bo.)	23
Bael fruit (English)	23
Baer (H)	100
Baghnoki (B)	99
Bagua (H)	84
Bahera (H & B)	73
Bahira (S.)	73
Baigun (H.)	110
Bakas (B)	23
Bakra (H)	68
Bakul (S , H & Bo)	106
Bala (S & B)	48, 107
Balraksha (P)	78
Balunt-shep (Bo)	43
Banbalnag (Kash)	22, 62
Banda (P.)	76
Bandurgi (Bo)	69
Bangab (B.)	117
Ban-gal (B.)	80
Banger chhata (B.)	55
Bankhur-i-Mirya (Ind Baz.)	79
Banlaunga (H.)	105

	PAGE
Bannatia (H.)	86
Bans (B & H.)	27, 94
Bans kaban (H.)	94
Bar (B)	104
Bara-elachi (H. & B)	24
Bara-kanur (B.)	91
Bara-keru (B)	38
Barbatti (Bo)	98
Bare kulinjan (H)	24
Barhanta (H)	89
Bari-saunf (H)	38
Bari sopha (Bo)	38
Barnagi (H)	97
Barola (B.)	69
Barphali (Vern.)	38
Bartaku (S)	119
Bartang (Bo)	99
Baru (H)	61, 95
Barun (H)	98
Bas (Bo)	94
Basak (H)	36
Basant (H & P.)	66
Basingh (Bo)	109
Bassu (Vern)	48
Bayabirang (H)	98
Bean-caper and Guaicum Family (English)	67
Beet Family (English)	85
Begonia Family (English)	75
Behaira (Bo)	73
Belatijau (B)	103
Belladonna (English)	168
Bell-flower Family (English)	78
Berberry Family (English)	64
Bet (H., B & Bo)	102
Betel leaf (English)	44
Bhallatamu (S)	47, 70
Bhallava- anga (S.)	105
Bhang (M)	30
Bhangi (M)	30, 89
Bhant (H. & B.)	34
Bharangi (Vern , S. & Bo.)	43, 97
Bharbhand (Vern)	25
Bhat (H & B.)	34
Bhela (H & B.)	47, 70
Bhimb (Bo)	34
Bhui-chane (Bo)	25
Bhuidari (Bo)	82
Bhui-dumur (B.)	104
Bhui-kanda (Bo)	47
Bhumbi-jambu (S)	108
Bhumya-amalaki (S)	107
Bhunimba (S)	24, 101
Bhuta-pala (Bo)	68
Bhutia-Tsallu (Vern)	79
Bhutta (H & B.)	95
Biba (Bo)	47, 70
Bibu (B)	69
Bichu (H & P)	90, 99, 100
Bichuti (B.)	89
Bignonia Family (English)	84
Bihi (H)	103
Bijindak (Afgh)	65, 116
Bikh (Vern)	22, 62
Bilnalita (B)	103
Bililotan (H)	99
Bimba (S)	34
Bincha (B)	109
Bindimuthi (Santh.)	104
Birama-dandu (M.)	25, 64
Birbanka (Nep.)	112
Birmova (Bo.)	104
Birth-wort Family (English)	86
Bis (Vern & P)	62, 107
Bish (H & B)	22, 62
Bishcopra (Kumaon)	79
Bishkapra (P & Bo)	76
Bislambhi (H)	75
Bishlanguli (B)	92
Biswul (H)	70
Bitter-bari Family (English)	68
Black mustard (English)	65
Blue gum tree (English)	38
Bobawachi (Bo)	45
Bonmethi (B)	180

	PAGE
Bon-nil (B.)	72
Bon-okra (B)	51, 78, 112
Bonpatol (B)	75
Bor (H)	104
Borage Family (English)	82
Borasali (Bo)	106
Braehmbhi (H)	27
Brahmi (Vern & S)	27, 32
Brahmo manduki (H.)	32
Brela (B)	48
Brihat-upa-kunchika (S)	24
Brihmisak (B)	27
Brimposh (Kash)	106
Broyphyta (Livervorts & Mosses) (English)	57
Buchanaka (S)	25
Buck-wheat (English)	62
Buck-wheat Family (English)	86
Bun (P & Kash)	107
Bulu (Nep)	79
Buna (P)	120
Bun-akora (B)	108
Burundi (Bo)	82
Buttercup Family (English)	62
C	
Camphor (English)	255
Caper Family (English)	65
Carnation Family (English)	66
Carrot Family (English)	76
Casearia Family (English)	74
Cashew Family (English)	69
Cassarva (Baz)	89
Castor oil Family (English)	87
Castor seeds (English)	46
Cha (Vern.)	30
Chagul-banti (B)	36
Chaksu (H.)	31, 71
Chakramards (S.)	97
Chakrani (Vern)	29
Chakunda (H. & Bo)	97
Chambul (P.)	63
Champa (H)	66
Champabaha (Santh)	99
Champa Family (English)	63
Chandla (Bo)	25, 89
Chandra (B. & Bo)	46, 81
Chandrasura (S)	111
Chansaur (H)	111
Chanduka (Vern)	25
Chanu (B)	76
Chapra (H)	42
Charayotah (H)	48
Chatium (H)	24
Chatri (Nep)	41
Chattrak (S)	55
Chaulmoogra Family (English)	65
Chaulmoogra (H , B & Bo)	65
Cheraelu (H.)	79
Cherailu (H)	100
Chian (H)	71
Chikana (Bo)	48
Chikri (Kash)	88
Chikti (H)	108
Chilla (M)	88
Chillara (H. & Bo)	74, 116
Chilli (H)	74
Chhatim (B)	24
Chhota-gokru (H)	51, 78
Chhoti dudhi (H)	88
Chiner badam (B.)	25
Chir (Vern)	43
Chiraita (Bo.)	48
Chiraputi (Bo.)	75
Chireta (B)	48
Chiretta (Vern.)	48
Chirghas (Kash)	82
Chirribenda (M.)	48
Chiror Anrudana (Vern.)	41
Chir Pine (English)	43
Chirpoti (H.)	75
Chirval (H)	99
Chita (H.)	45, 79

	PAGE
Chitaro (Bo.)	**45, 79**
Chitra (Vern. & P)	28, 73
Chitraka (S)	44, 45, 79
Chitro (Nep)	28
Chittira (M)	45
Chittur-mol (M)	79
Cholam (M)	95
Chota chand (H)	46, 81
Chota gokhru (H)	50, 67
Chota pilu (H & B)	80
Choti-elachi (H & B)	37
Chuka (H., B & Bo)	107
Chukapalam (B)	86
Chukra (S)	107
Chumbi Aconite (English)	128
Chumlani (Nep)	48, 67
Chutrika (S)	86
Cinchona (English)	33
Clathroxystis (English)	53
Coffee (H)	77
Coffee Family (English)	77
Coffee (English)	175
Coffee powder (English)	245
Colombo (Bo)	105
Coriander Family (English)	76
Convannamilpori (M)	46
Coriaria Family (English)	70
Cotton Family (English)	66
Covannamilpori (M)	81
Croton Family (English)	87
Cruciferae (English)	65
Crude Drugs (English)	249
Cryptogamic flora (English)	59
Cubebs (English)	44
Cucumber Family (English)	74
Cultivation of Medicinal Plants (English)	249
Custard apple Family (English)	64

D

Dahar karanja (B.)	45, 72
Daka (Bo)	119
Dalchini (Vern.)	33
Dama (P.)	109
Dalme (H.)	88
Damahan (H)	109
Danti (S , H. & B.)	88
Dantimul (Bo)	88
Dar-hald (H & B)	28, 46
Darimba (S)	107
Darmiya Gobriya (Nep.)	21
Dapoli (Bo)	110
Daru haridra (S)	28, 64
Daru haridraka (S)	35
Darya kanariyal (H & Bo)	110
Datura Family (English)	83
Davana (Vern)	26
Deo-dhan (H & Bo.)	95
Derris (English)	168, 185
Devadarum (M)	109
Devils cotton (English)	21
Dhai (H & B)	109
Dhakangu (P)	63
Dhakur (B)	32, 80
Dhan (B)	95
Dhania (H & B)	97
Dhanyaka (S)	97
Dharakadamba (S)	77
Dhatriphala (S.)	107
Dhatriphal (S)	73
Dhataki (S)	103
Dhavala (Bo.)	78, 141
Dhingri (Vern.)	56
Dhub (H , B. & Bo.)	109
Dhudi (Vern)	39
Dhumrapatra (S)	86
Dhundul (B)	75
Digitalis Family (English)	84
Dikoor (English)	94
Dirghapattra (S.)	75
Dita-bark (English)	24
Diva (Bo)	91
Dochunty (B.)	101
Dodhi (Bo)	37
Dog-bane Family (English)	80
Dogwood (English)	38
Don (P)	112

	PAGE
Dowla (Bo.)	108
Drek (Vern)	41
Drug adulteration (English)	250
Drug resources (English)	250
Drug scarcity (English)	250
Dryopteris (English)	187
Dudhi (H)	38
Dudiya (B.)	88
Dudla (Bo.)	89
Durga chhata (B)	55
E	
Ebony Family (English)	80
Edakula-pala (M)	24
Ela (S)	37
Elachi (Bo)	37
Ellakai (M)	37
Elam (Tam)	101
English-Maiden-hair-tree	113
English-Manchineal tree	113
English-Moonwort	102
Equisetum (Horsetail) (English)	58
Eranda (S)	46, 89
Erendi (Bo)	46, 89
Ergot (English)	54, 173
Ermul (B)	67, 114
Erukku (M.)	30, 81
Essential oils (English)	250
F	
Ficoideae (English)	76
Fauglimehandi (P)	74
Flax Family (English)	67
Foxglove (English)	37
French sorrel (English)	107
Fungi (English)	53
Fusarium (English)	54, 193
G	
Gab (H. & B)	104
Gadab (H)	114
Gadadhar (S)	26, 77
Gaggar (Kash.)	79, 100
Gajapipal (H & B.)	107
Gajar (H., B. & P)	76
Gajega (M.)	30
Ganjara kelangu (M.)	76
Galot (P)	103
Gani (Bo)	95
Gamboge Family (English)	66
Gangwa (B)	88
Ganja (H , B & Bo.)	30, 89
Ganjika (S)	30, 89
Ganjni (H)	117
Ganiari (B)	45
Ganikarika (S)	45
Garbijaur (H.)	99
Gardal (Bo)	71
Garlic (English)	23
Garudu-mukku (Tel.)	99
Gatta colla (Vern)	42
Gaungchi (H)	21, 70
Gelaphal (Bo)	46, 77
Geva (B)	88
Geyapal (Bo)	88
Ghet-kachu (B)	94
Ghetuli (Bo.)	29
Ghiatarui (H)	75
Ghi kanvar (H)	24
Ghirna (Garhwal)	102
Ghosalata (B)	75
Ghosi (Nep)	69
Ghosali (Bo)	75
Ghrita Kumari (S & B.)	24
Gigantic swallow root (English)	30
Gilagach (B.)	71
Giloe (Vern.)	49
Girbuti (Bo.)	83
Gnetum Family (English)	90
Goagarilakei (Bo.)	82
Gobar champa (B.)	81
Gobari (Nep)	21, 62
Goda (B)	51
Godhum (S.)	95
Godumal (M)	95
Gokhuri (B.)	50, 67
Gokhru (Vern.)	50

	PAGE
Gokshura (S)	50, 67
Golapphul (B)	47
Golappu (M)	47
Golmarich (B)	107
Golmirch (H)	107
Gorakh-amli (H. & Bo.)	101
Gorakmundi (H & Bo)	78, 100
Goruma (H)	98
Goshtam (M.)	47, 78
Guttibira (M.)	75
Gowali (Bo)	109
Grass Family (English)	94
Grecian Foxglove (English)	36
Ground Nut (English)	25
Gubak (S)	25, 92
Guchhian (P)	56
Gudatreaka (S)	33
Guduchi (S)	49
Guggul (B , Bo. & H)	35, 97
Gugul (Vern)	42
Guggula-dhup (Bo)	101
Guggulu (S)	35
Gukhulu (M)	35
Gul (Bo)	47
Gulab ka phul (H)	47
Gulabjaman (H. & Bo)	108
Gulancha (H & B)	49
Gulari (H)	104
Gulsakari (H.)	108
Gulwail (Bo)	49
Gumbhari (S)	98
Gum gugul (English)	35
Gandhagatra (S.)	64
Gundumani (M.)	21, 70
Guniyan (Kashm.)	103
Gunj (P)	71
Gunja (S & Bo)	21, 70
Guras (Nep)	79, 120
Gurbiani (P)	49
Gurdlu (P.)	72
Gurguli (P.)	87
Gur mar (Vern.)	39
Gurgunna (P.)	85
Gurgur (B.)	97
Guttiferae (English)	66

H

Haldi (H)	35
Haligach (Vern)	35
Haldi-gach (B.)	35
Halim (B)	111
Halood (B)	35
Hansraj (Bo)	101
Harara (H)	73
Haridra (S)	35
Hardu (H)	77
Harira (H)	110
Haritaki (S & B)	73, 110
Harivera (S)	107
Hastisunda (S)	82
Hatisura (H & B)	82
Hattut-tumatti (M)	75
Hawar (H)	84
Hayer (B)	97
Hazardana (P)	88
Hazarmani (H & B.)	89
Heart poisons (English)	251
Helga (B)	108
Hemp Family (English)	89
Henna Family (English)	74
Hijal (B.)	73, 102
Hijjal (H.)	102
Hijli badam (B.)	69
Hindisana (H)	31
Hing (H & B)	38
Hingan (H & B.)	68
Hinger (Bo)	68
Hingu (S)	38
Hiranpadi (H)	82
Hirankhori (Bo.)	103
Hiranpag (Bo)	82
Hirda (Bo.)	73
Hirtiz (Kash)	113
Hmyaseik (Burm.)	25, 89
Honey-suckle Family (English)	76

	PAGE
Horse-radish Family (English)	70
Horsetail (English)	58
Hsathanpaya (Burm)	118
Hulugiri (Bo)	69
Hurmal (H. & Bo.)	43, 67
Hurua (B)	89
Hurua (Bo.)	89

I

Ichchura-mula (M.)	25
Ijjul (H)	116
Ilaikal (M)	88
Ilayechi (H.)	101
Indian Drug Industry (English)	251
Indian foods (English)	251
Indian or wild Liquorice root (English)	21
Indian medicine (English)	251
Indian Napellus (English)	22
Indian Pharmacopoeia (English)	17
Indian Pharmacopoeia Codex (English)	18
Indian Pharmacopoeia List (English)	17, 251
Indian Rubarb (English)	46
Indian Sarsaparilla (English)	39
Indian winter green (English)	38
Indigenous drugs (English)	251, 252
Indigenous remedies (English)	252
Indra varuni (S.)	34, 75
Indrajau (Bo)	109
Indrayan (H & Bo.)	34, 75
Ingudi (S)	68
Inji (M)	51
Insecticides of vegetable origin (English)	252
Ipecac (English)	165, 166
Ipecacuanha (English)	166
Ipomoea (English)	82
Iris Family (English)	90
Isabghol (Bo.)	44
Isabgul (B. Vern.)	19, 44
Isband (B.)	43, 67
Ishappukol-virai (M)	44
Isharmul (H & B.)	25, 86
Ispaghul (H.)	44
Ivy Family (English)	76

J

Jadikkay (M.)	42
Jafran (H , B & Bo)	35, 91
Jahari sontakka (Bo)	80
Jalqumbhi (H)	99
Jam (H & B.)	108
Jamalgota (H.)	88
Jamana (H.)	72
Jamma (P.)	72
Jamti-ki-bel (H)	97
Jamun Family (English)	73
Janghibadam (Vern)	40
Jangli-chichonda (H.)	75
Jangli darchini (H.)	103
Jangali-kanda (Bo.)	50, 92
Janghimendi (H)	74
Janghimudrika (Bo)	64
Jangli piyaz (H & B.)	50, 92
Janghisaru (H.)	103
Jar Gandapuro (Vern.)	38
Jar-amla (H)	107
Jarul (H & B)	74
Jashti madhu (B & Bo)	39
Jatifalam (S)	42
Javanala (S.)	95
Jaya (S)	101
Jayanti (H & B.)	100
Jayphal (H , B & Bo.)	42
Jethimadh (H.)	39
Jhilla (S.)	105
Jhinge (B)	40, 75
Jira (H. & B.)	31, 98
Jiraka (S.)	98
Jirkivirai (M.)	40, 82
Jiunti (P.)	63, 97
Jivanti (Vern.)	40
Jonkhmari (H.)	79

	PAGE
Jowan (B.)	50
Jowar (H., B. & Bo.)	95
Joypal (B.)	88
Julpai (B)	104
Jungli anarash (B.)	91
Jun-li-kali-mirchi (Bo.)	49
K	
Kabab-chini (H , B & Bo.)	44
Kabiraj (Pers.)	36
Kabra (H. & P.)	97
Kachlora (H & Bo)	72
Kachnar (H)	97, 102
Kachura (H. & Bo)	35
Kadamba (S & H)	102
Kadatathie (M)	84
Kadavanchi (Bo)	75, 111
Kadukkay-pu (M.)	73
Kafi (B)	77
Kahira champa (S.)	112
Kahoo (H)	98
Kaiphal (H , B & Bo.)	90, 106
Kajra (Bo.)	48, 82
Kaju (H. & Bo)	69
Kakadani (S.)	97
Kakadumbura (S)	104
Kakajangha (S , H & B.)	99
Kakaphala (S)	64
Kakatundi (H)	81
Kakilakshya (S.)	27, 102
Kakkanan (M.)	71
Kakkar tamaku (P.)	83
Kakkata shingi (M.)	44
Kakkay-kolli-virai (M.)	64
Kakmachi (S. & B)	84, 108
Kakmari (H. & B.)	64
Kakra singi (Vern.)	69, 222
Kakra shingi (H. & B.)	44
Kakumullu (Vern)	70
Kakurjiwan (H. & B.)	105
Kaku-vali (Malay)	114
Kaladana (H., B & Bo.)	40, 82
Kala Dhatura (H. & B.)	83
Kakra (Vern.)	44
Kalajira (H. & B.)	63, 102
Kalambi (S)	82
Kala mucha (B.)	61, 95
Kalasaka (S)	130
Kala tulshi (H & B.)	106
Kala vala (Bo)	107
Kalenjire (Bo)	63
Kali-Jhant (H. & B.)	101
Kalihari (H)	92
Kalijiri (Bo.)	32
Kalkora (B)	96
Kalikutki (Bo.)	43
Kalmegh (B)	24
Kalmisak (B.)	82
Kalo bikhmo (Vern)	22, 62
Kalo bikhoma donghi (Vern)	22, 62
Kalu-kera (B.)	109
Kamala (H. & B.)	41
Kambari (H)	98
Kambila (Vern.)	41
Kamini (B.)	106
Kamkher (Bo)	117
Kamugu (M)	92
Kanakaphala (S.)	88
Kanaphata (H)	69
Kanchana (S)	49
Kanchkuri (Bo.)	89
Kanchnine (M.)	102
Kanchnal (H)	97
Kanchuri-vayr (M.)	89
Kanda (S & Bo)	96, 97
Kandiari (Vern)	25
Kanduri-ki-Bel (H.)	34
Kanga (Vern.)	45
Kanghi (H.)	101
Kanguni (S.)	32
Kanj (H)	49
Kanhera (Bo)	81
Kanka (M.)	94
Kankhina (Bo.)	80
Kankri (Bo.)	75
Kansa (H.)	97
Kanta (Simla)	64

	PAGE
Kantabans (Vern.)	27
Kantala (H.)	91
Kapas (H., B. & Bo.)	104
Kapasi (Vern.)	39
Kapila (M.)	41
Kapittha (Vern.)	104
Karabi (B)	81
Karachunai (M)	91
Karail (B)	94
Karala (B.)	110
Karanja (S, H. & Bo.)	45, 72
Karavi (S.)	81
Karber (H.)	81
Karchi (H.)	39
Karela (H.)	75, 110
Karer (H)	65
Kari (Bo & Santh)	65, 109
Karia (P)	65
Karianag (Bo.)	92
Karinthuvari (M)	80, 117
Karira (S)	65
Karivana (Bo.)	32
Karkani (Bo)	105
Karki (Bo)	81
Karkata sringi (S.)	44, 69
Karla (Bo.)	75
Karmal (H)	102, 113
Karmora (Kash)	76
Karpas (S.)	104
Karpo-Karishi (M.)	45
Karpura maram (M & Vern.)	38, 73
Karnaspota (S)	69
Karunaikkizhangu (M)	94
Karun Kanam (N)	31
Karun-shiragam (M)	63
Karu umattai (M.)	83
Karval (M.)	65
Kashappuvetpalarishi (M.)	39, 80
Kashira champa (S)	81
Kashmal (H & P.)	28, 29
Kasni (Vern)	33
Katarali (M.)	32, 80
Kat-Karanj (H.)	30
Katki (H & B)	43
Katnim (H.)	106
Katphala (M)	90, 106
Katson (Ku.)	73
Kattalai (M)	24
Kattu-Shiragam (M.)	32
Kattuvalari (Vern.)	70
Katuka (S)	43
Katuka-rogani (M)	43
Kaundal (Bo.)	75
Kava (Bo.)	40
Kavali (Bo.)	39
Kavitha (H.)	140
Kayam (Bo.)	38
Kela (H. & Bo)	110
Keli kadam (B)	77
Kend (B)	98
Kendu (H.)	98
Ker (Bo.)	101
Kesar (Vern.)	35
Khadira (H.)	101
Khair (H)	101
Khair Champa (H. & Bo.)	81, 112
Khaj-goli-cha-vel (Bo)	113
Khappar kadu (Bo.)	103
Khardi (Bo.)	98
Khareti (Vern)	48
Khesari (H & B)	71
Khesari dal (Vern.)	61
Khinna (H.)	89
Khira (H.)	75
Khirni (H)	99
Khorasani ajowan (B.)	40, 83
Khorasani-owa (Bo.)	83
Khorasani-yomam (M.)	40, 83
Khour (Nep.)	101
Khumb (H. & P.)	55
Khubani (H)	72
Khubakallana (H)	104
Khubazi (H. & Bo)	105
Khumbi (H & P.)	55
Khurasani-ajvayan (H)	40, 83
Khushing (P.)	31
Kikar (H.)	101
Killuk-kay-velai (M.)	72

	PAGE
Kilmora (H)	28
Kinai tihiri (Bo)	70
Kindal (Bo.)	110
Kinsuk (S)	70
Kiralu (P)	93
Kiramani owa (Bo)	26, 77
Kiraruga (M)	95
Kirata (S)	48
Kirithi (M)	79
Kiriwolla (Sinhalese)	105
Kirmala (Vern & H.)	26, 77
Kirmar (H)	86
Kirtana (Bo.)	71
Kiwach (H.)	106
Kiryat (H)	24, 101
Knapa chettu (M)	50
Kodali (M)	74
Koda-todali (B)	49
Kodicpalay (M)	37
Kodo (H)	95
Kodoa (B)	95
Kodra (Vern & Bo)	95
Kodrava (S)	95
Koh-tor (Baluchi)	83
Kokoranj (H)	30
Kolamavu (Tam)	99
Kolhal (Bo.)	103
Kolka-phul (B)	49, 81
Kolumbu (Tam)	105
Kobu-Kalli (M)	88
Kombupudalai (M)	75
Konda vaghe (M)	70
Kondapala (M)	82
Konea-dumbar (H.)	104
Korattai (M)	75
Kori (B.)	70
Koshataki (S)	40, 75
Kosum (H)	69
Kottak (M.)	78
Kottei (M)	110
Kovidara (S.)	97
Kowti (Vern.)	40
Krishnachura (B.)	110
Krishna dhatura (S)	83
Krishnajiraka (S)	63
Kua (Tam)	98
Kuberakshi (S)	30
Kuchla (H.)	48, 82
Kuchila (B)	48, 82
Kuchila-lata (H & B)	82
Kukarchita (B)	99
Kukarlata (H.)	75
Kukarnicha (H)	109
Kukarwele (Bo)	75
Kukronda (P)	88
Kukseem (B).	100
Kukursunga (B)	109
Kul (B)	100
Kulahala (S)	103
Kuljud (H)	94
Kuliakhara (B)	27
Kumbhi (H & B)	73
Kumkuma (S)	35, 91
Kunch (B.)	21, 70
Kundu (Bo.)	80
Kungku (H)	38
Kungumapu (M)	35, 91
Kungyi (H)	48
Kurchi (English & B)	80
Kurelo-jangro (Bo)	75
Kurling (Nepl)	120
Kurlinga (Nep)	79
Kushtha (S)	47, 78
Kusturi (Vern)	255
Kut (H.)	47, 78
Kutaja (S.)	39, 80

L

Labangaphal (B.)	40
Lablab (H.)	76
Ladaki-revanda Chini (Bo)	46, 86
Laghukarni (S)	63
Lahana-gokhru (Bo)	50, 67
Lahuriya (H.)	99
Lajja (S)	105
Lajalu (H. & Bo.)	105
Lakh (Bo)	71

	PAGE
Lalbachlu (H)	102
Lal chitra (H , B. & Bo.)	44, 79
Lal-indrayan (H)	75
Lalpost (H)	64
Lalsag (H.)	101
Lal siris (H)	96
Lankasij (B)	88
Lasan (H & Bo)	23, 91
Laskar (P.)	63
Lasuna (S)	23, 91
Lata palas (S)	29
Laural Family (English)	87
Lavangalata (S)	40
Lichens (English)	57
Life-Plant-Family (English)	73
Lil kathi (Santh)	66
Lily Family (English)	91
Lodh (H , B. & Bo)	49
Lodhra (S)	49
Lohari (H)	80
Lohori (H)	117
Lona (H)	64
Long (Vern)	38
Luban (H & B)	102
Ludduga (M)	49
Lycopodium (English)	207

M

Machni (H.)	94
Machutie (B)	86
Madan (S)	45, 77
Madar (H)	81
Madder Family (English)	77
Maddi-palu (Tel)	101
Madhu malati (S)	37
Madhurika (S)	38
Magadhi (S. & Tel.)	98
Magnolia Family (English)	63
Magrabu (H.)	39
Mahakal (S)	75
Mahalimbu (Vern)	31
Mahmira (Bo.)	35
Mahanimb (H)	101
Mahogany Family (English)	68
Mainphal (H)	77
Makal (B.)	75
Makai (Bo)	95
Makhal (B)	34, 75
Makka-scholam (M.)	95
Mako (Bo)	84, 108
Makoi (H)	84
Malabar nut tree (English)	23
Malkanguni (H)	32
Mamekh (P)	63
Mamira (H.)	35
Mamiran (Bo)	49
Manaka (S)	93
Manattak kali (M)	84
Manchingi (Bo)	84
Manda (M)	103
Mandara (Bo)	28, 94
Mandgay (Bo)	27, 94
Mangal (M)	94
Mango Family (English)	69
Mangustan (H , B & Bo)	104
Maniphal (H)	45
Manith (H)	107
Manja kadambe (M)	77
Manjistha (S & B)	107
Manjit (Bo)	107
Mankachu (B)	93
Mankenda (H)	93
Mansa-sij (B)	88
Manskhel (Kash)	55
Manya (S)	98
Mara-manjal (M.)	35
Maravara (M)	118
Maravara Tsjembu (M)	93
Maravuli (M.)	89
Marchula (H)	106
Maricha (S)	107
Marisha (S.)	101
Marlu-mutta (M)	51, 78
Marudam-pattai (M.)	90
Maruk-kallan-kai (Bo. & M.)	46, 77
Marsada boli (M)	30
Mashagani (Tam)	105

	PAGE
Masha-Parui (S.)	100
Mashparui (H)	100
Masur (H & S.)	105
Masuri (B)	105
Mat (H)	99
Matazor (H)	85
Maurabikh (Vern)	22, 62
Mazri (H)	106
Meetha-tellia (Vern)	62
Menphal (B)	46, 77
Meradu (H)	66
Merasingi (H & B)	39
Meshasringi (S)	39
Methi (H , B & Bo)	120
Mexican Poppy (English)	25, 60
Mezereum Family (English)	87
Mandanak-kattai (S)	47
Milkaranai (M)	49
Milk-weed Family (English)	81
Milkwort Family (English)	66
Mimulus Family (English)	84
Minguta (Bo)	88
Mint (English)	255
Mint Family (English)	85
Mirch (H & P)	83
Mistletoe Family (English)	87
Mitha indarjou (H)	109
Modi-murukkan (M.)	29
Mohra (Vern)	22, 62
Mohri (Vern)	22
Mohua (Vern)	41
Mohwa meal (English)	80
Mohwa Family (English)	80
Mooda cotton (M.)	69
Moonseed Family (English)	64
Moravela (H & Bo)	63
Morwa (Bo)	100
Moulds (English)	54
Muchi-tanki (M)	80
Mugani (H. & B)	99
Muhuri (B)	43
Mukajali (H.)	73
Mukuk-rattai (M.)	29
Mulberry Family (English)	89
Mullanjakka (S.)	101
Mullu-chitta (Tam)	101
Mulluvellari (M)	75
Mumugu (M)	25
Mundiri-kai (M)	69
Munditika (S)	78, 100
Mungas kajur (H)	84
Mungphali (H)	25
Munjal (M)	35
Munjariki (S)	106
Munni-vayz (M)	45
Murba (B)	100
Murmuria (B)	78
Murungai (M)	70
Musabbar (Vern)	24
Mushabbar (Vern)	92
Mushkiara (P)	77
Mushrooms (English)	56
Musna (H)	66
Musta (S & Bo)	103
Mustard cake (English)	62
Mustard Family (English)	65
Mutha (B & H)	103
Muttia-lata (B)	42, 110
Myetpye (Burma)	105
Mylabris (English)	248
Myrobolan Family (English)	73
Myrtle Family (English)	73

N

Nachuta (M)	88
Nagadamani (S)	26, 77
Naga-danti (M)	88
Nagamusadi (M.)	82
Nagar Motha (H & B)	103
Nagar mustaka (S)	103
Nagbala (S)	108
Nagbali (H)	51
Nagchampa (Bo.)	105
Nagdona (B.)	26, 77
Nagadouna (H.)	26, 77

	PAGE
Nagdowan (Bo.)	91
Nagkeshar (S , H & B.)	105
Nahani khapat (Bo)	101
Najunda (M)	68
Nokchikni (H)	37
Nala (B. & H)	114
Nala tige (M)	71
Nali (S)	105
Nalichi baji (Bo)	82
Nandibriksha (S & B)	31
Nandru (P.)	83
Nannari (M)	39
Napali dhania (B)	68
Naranji (M)	82
Naraseja (Bo)	88
Narcha (H. & B)	103
Narel (Bo)	97
Nargis (P)	91
Nari-vengayam (M.)	50, 92
Nariyal (H.)	97
Naro (Bo)	74
Nata-karanja (B)	30
Nattu-ireval-chinni (M)	46, 86
Navaladi (M)	51
Nayeti (Bo)	38, 88
Nay-palai	50, 82
Neem Family (English)	68
Negli (Bo)	66
Ner (P)	48, 67
Nettavil (M)	25, 89
Nettle Family (English)	89
Newrang (Bo)	88
Nightshade Family (English)	83
Nil (H. & Bo)	98
Nila kadalai (M)	25
Nilkattei (H. & P)	82
Nila-nirgundi (S)	105
Nila vakai (M)	31
Nila vembu (M)	24,48
Nilika (S)	98
Nili-nargandi (H)	105
Nim (Vern., H , B & Bo)	27, 74
Nimrudi (Bo.)	109
Nirbisi (H & B.)	33
Nirbrami (M)	27
Nirgundhi (S & H)	121
Nirumel-neruppu (M)	74
Nirmuli (Bo)	37, 82
Nirunji (M)	50, 67
Nishinda (B)	111
Nisomali (S)	86
Nipa (S.)	116
Noalata (B)	71
Nomenclature of Drugs (English)	252
Nona (B)	64
Norvishee (H)	73
Nostoc (English)	53
Nurma (H)	98
Nutmeg Family (English)	86
Nux-vomica Family (English)	82
O	
Oandak (P)	63
Olaktambol (Bo)	21
Olang-karai (M)	104
Oleander Family (English)	80
Oman (M)	50
Opium (English)	43
Osari (Bo.)	101
Ot (Sant.)	55
Ouplate (Bo)	47, 78
P	
Paas lata (H & B)	29
Pachak (B)	47, 78
Pochch-ai-alari (M)	49, 81
Pacholi (H.)	85
Packur-mul (B)	86
Paddam (H)	72
Padmaka (S)	72
Padma-kasta (Bo)	72
Pahari gandana (H)	110
Palai (M)	81
Palandu (S)	96
Palla (M)	99
Palm Family (English)	92

	PAGE
Palarvela (Vern)	29
Palas (Vern , H & B)	29, 70
Palupaghel kalung (M)	99
Pamukh (P)	85
Pan (H , B & Bo)	44, 99
Pandharen-kamal (Bo)	106
Pandharphali (P)	88
Pandhrakura (Bo.)	39, 80
Pandorina (English)	53
Panibira (M)	75
Panirak (H)	67
Panlata (B.)	71
Pan-mauri (B)	38
Pannax Family (English)	76
Papai (Bo)	31, 74
Papas (P)	98
Papaw Family (English)	74
Papaya (H)	31, 74
Papey (B)	31, 74
Papita (Vern)	31
Pappayi (M)	31, 74
Papra (H)	45
Papri (P)	88
Parasikaya (S.)	40
Parasi-kaya (S)	83
Parusha (S)	98
Parvar (H)	75
Pashanbheda (Bo.)	102
Passion-flower Family (English)	74
Patalagaruda (S)	103
Patha (S)	33
Patharchuri (B)	102
Patola (S)	75
Patsan (H)	105
Patta karie (S)	88
Pattudu (Tel)	105
Pavakkachedi (M)	75
Pawal-chhatta (B)	56
Pea Family (English)	70
Pectin (English)	252
Pekarakai (M)	110
Penarvalli (M.)	75
Pendari (Bo.)	77
Penicillium (English)	54
Pepper Family (English)	86
Perala (Bo.)	110
Pera-rattai (M)	24
Periya-yelakkay (M)	24
Petthri (P.)	40
Peyttumatti (M)	34, 75
Phala kantak (S)	36
Phalsa (H & B.)	98
Phanasa-alambe (Cutch & Bomb)	55
Phanasa-alambe (Bo)	101
Phanerogams (English)	59
Phemila (S)	100
Phenila (S)	69, 75, 120
Phytolacca Family (English)	85
Pichli (Vern)	23
Pikumkai (M)	40, 75
Pila-kaner (H & Bo)	49, 81
Pilu (H., S & B)	80
Pilvu (Bo.)	80
Pindalu (H.)	77
Pindaluka (S)	91
Pine-apple Family (English)	91
Pine Family (English)	90
Pinjari (H)	49
Pipal (H)	99
Pippali (S)	99
Piralu (B)	77
Pisa (Bo)	23
Pitadaru (S)	106
Pithvan (H)	108
Pitz (Kash)	108
Piyaj (B)	96
Piyaz (H.)	96
Plant pigments (English)	252
Plesmonium (English)	93
Plumbago Family (English)	79
Poal-chhatea (B.)	56
Podina (H.)	42
Poisonous Plants (English)	253
Pomegranata Family (English)	74
Poonam-penmaraiva (Mal.)	100
Ponnan- kottai (M.)	69

	PAGE
Poppy Family (English)	64
Postakatol (M.)	42, 64
Potaki (S)	102
Potal (B)	75
Potari (B)	101
Prashni (Bo.)	99
Prime-rose Family (English)	79
Priyangu (S.)	72
Pudel (M)	75
Pudina (Bo., H , S & M.)	42, 105
Pudinah (B. & Bo.)	42
Puff-balls (English)	56
Pugai-ilai (M.)	83
Pulan-ki-zhanga (M)	35
Pu-maram (M)	69
Punarnaba (B.)	29
Pungammaram (M.)	72
Punnag (B.)	66
Punnaga (S.)	66
Punnagam (M.)	66
Purpuray timur (Nep.)	121
Pusha (Vern.)	48
Pussar (B.)	109
Pyrethrum (English)	168, 185

R

Racta-vinda-chada (S.)	88
Rati-jaman (H.)	108
Rajadani (S)	99
Rajkoshataki (S.)	75
Rakhalu (Bo)	118
Rakra-posta (S.)	64
Rakta kanchan (Bo)	97
Ramban (Bo)	108
Ram dataum (H.)	108
Rametha (Bo.)	114
Ramoha (S.)	110
Ramphal (Bo.)	64, 87
Ramsita (M.)	64
Ramtulshi (H & B)	119
Ranghevada (Bo.)	103
Ranparul (Bo.)	75
Rasna (S., H., B & Bo.)	50
Rasun (B)	23, 91
Ratna-gandhi (S. & Tam.)	110
Rattanjog (P.)	63
Rattu papillay (M)	78
Rauwolfia (English)	80
Rechanaka (S)	41
Revandchini (H & B)	46, 86
Rhododendron Family (English)	79
Rhubarb Family (English)	86
Rice Husk (English)	253
Ritha (H., B. & Bo)	69, 100, 120
Riti (Sing)	89
Rohan (H , B. & Bo.)	108
Rohuna (S)	108
Rosaceae (English)	72
Rose Family (English)	72
Ruba-barik (P)	84
Rudrajata (S)	25, 86
Rudraksha (S)	109
Rudranti (H & B)	98
Rue Family (English)	67
Rukh-alu (Bo)	93
Rush Family (English)	92
Rusts (English)	53
Rutthraksham (M.)	109

S

Sabuni (B.)	66
Sadab (H)	67
Sada dhutura (B)	36, 87
Sadul kou (B.)	81
Safed bikh (Vern.)	22
Safedchandan (H.)	47
Safed damar (H.)	100
Safed musli (H.)	102
Safed siris (H.)	70
Safran (Vern.)	35
Sag-angur (H. & Vern)	27, 87
Sagaragoth (Bo)	30
Sage Family (English)	85
Sahadevi (S. & H.)	100
Sajnah (H.)	70

	PAGE
Saka (S.)	108
Sal (S., H., B. & Bo.)	108
Salad (B.)	98
Salvadora Family (English)	80
Samudra (M)	73
Samudrapad (S.)	73
Samudraphal (B.)	73
Samudraphala (Bo.)	73
Samp-ki-kumb (P)	93
Samutraphullam (M)	73
Sandan (H)	71
Sankhaphuli (H)	97
Sankhapuspi (S.)	97
Sanipat (S, H & Bo)	110
Sanjna (Vern)	42
Sankha pushpi (H)	104
Sant (H)	29
Saonf (H)	43
Saonjna (Vern)	42
Sapodilla Family (English)	80
Sapta parna (S.)	24
Sarkarei-valli (M.)	82
Sarapunkha (S.)	72
Sarivan (H.)	109
Sarphankha (H & Bo.)	72
Sarpagandha (S.)	46, 81
Sarson (Vern)	65
Sasa (B)	75
Satakuppi (M)	43
Satap (B & Bo.)	67, 114
Satapatri (S)	47
Satavari (Bo.)	97
Sati (S)	35
Saya (M)	99
Sebestan Family (English)	82
Sedge Family (English)	94
Segun (H. & B.)	108
Sehud (H.)	88
Sehund (H.)	88
Selupa (M.)	68
Senna (English)	163
Sesamum Family (English)	84
Shada-buri (B)	28
Shadhurakkalli (M)	88
Shakar-al-lighal (P.)	81
Shakar pitan (H. &. P.)	88
Shallaki (S.)	102
Shal Parni (S & Bo)	109
Shankeshvara (Bo.)	51, 78
Shatamuli (S & B.)	97
Shatapuspi (S.)	43
Shatawar (H)	47
Shayrang (M)	47, 70
Shendri (Bo)	41
Sheoia (B)	100
Shera (Bo)	88
Shetapuspa (S.)	43
Shiajira (H)	31
Shialkata (English)	60
Shikha-mulam (S.)	76
Shimai-azho-vanai-virai (M)	43, 67
Shimai-shombu (M)	31
Shiragam (Tam)	98
Shirisha (S.)	96
Shiru kurunja (M.)	38
Shirunari-vengayam (M.)	47
Shivappu-postaka chedi (M.)	64
Shiyappunelli (M)	89
Shombu (M.)	38, 43
Shori (B.)	35
Shothaghni (S.)	29
Sialkanta (H. & B.)	25, 64
Sij (B.)	88
Silajit (Vern.)	253
Simainayuruvi (M.)	108
Simla-Kashmal (Vern.)	28
Sindilkodi (M)	49
Siora (H.)	100
Siris (H., B & Bo.)	96
Sitama-purgonalu (M.)	36, 82
Sitapalam (M.)	64
Sitaphal (H.)	64
Sittrapaladi (M)	88
Small Caltrops (English)	50
Smuts (English)	53
Snuhi (S.)	88

	PAGE
Soap-nut Family (English)	96
Sobhanjana (S)	70
Sojna (B)	70
Soma (S. & Bo.)	82
Somalata (S.)	67, 114
Somlata (H & B)	82
Somaraj (S , H & B)	32
Sona (B)	106
Sonf (Bo)	43
Sonogaravi (Bo)	114
Soolpha (B)	43
Sopari (Bo)	25, 92
Sourabhi-nimba (S)	106
Spinach Family (English)	85
Spindle-tree Family (English)	68
Srigala-kantaka (S)	25, 64
Sriphal (Vern & S)	23
Steroid hormones (English)	253
Strophanthus seeds (English)	263
Subali (H)	88
Sugandha muricha (S)	44
Sugandha-vacha (B)	24
Sujna (Bo.)	70
Sukasa (S)	75
Sukh-darsan (H & B)	91
Sukra puspita (S)	92
Sultana (H)	66
Sumlu (Vern)	28
Sundew Family (English)	73
Sun-flower Family (English)	77
Supari (H & B)	25, 92
Suphadie-khus (H & B.)	47
Surati sonnamukai (Vern)	71
Surinjan (Vern)	34, 92
Sushavi (S)	75
Sutr-sowa (B)	106
Svetakanchan (S)	97
Swarnakshira (S)	65
Sweet-gale Family (English)	90
Swet-berela (H & B.)	67
Swet-chandan (S)	47
Syonaka (S.)	106

T

	PAGE
Tagar (H. & B)	50
Tagara (S)	50
Tagar-ganthoda (Bo)	50
Takadasingi (Bo.)	69
Talim khona (B.)	27
Talipanai (M)	92
Talmakhana (H.)	27, 102
Tamak (B)	83
Tamaku (H)	83
Taman (B)	74
Tamara valli (S.)	89
Tamarta (M)	102
Tambaku (Bo.)	83
Tambula (S)	44
Tanba (M.)	97
Tannins (English)	73
Tarur (H & B)	103
Tatara (H)	73
Tattu dattura (P)	83, 36
Tatwen (P.)	26
Tazak-tsun (Kash)	79
Tea Family (English)	66
Teak Family (English)	84
Tea plant (English)	30
Tekata sij (B)	88
Tejmal (H)	68
Telakucha (B)	34
Tel-kodukki (M)	82
Tellajonna (M)	95
Tella-motuku (M.)	71
Temru (Bo)	98
Tendu (H & Bo)	80, 104
Tessul (Bazinali, Bo)	110
The Anise (English)	43
The Betelnut palm (English)	25
The Asoka tree (English)	47
The Barberry (English)	28
The Caraway seed (English)	31
The common Jowar (English)	61
The Creat (English)	24
The East Indian Screw Tree (English)	39

	PAGE
The Flowering Plants (English)	59
The greater Cardamon (English)	24
The greater Galangal (English)	24
The Indian Millet (English)	61
The Indigenous Drugs of India (English)	19
The Lemon Tree (English)	34
The Lime Tree (English)	34
The Lodh tree (English)	49
Thelu-Kodi (M)	114
The Mahu tree (English)	40
The marking nut tree (English)	47
The marsh mint (English)	42
The Morell (English)	56
The Mushroom (English)	55
The Neem (English)	27
The Opium (English)	42
The Papaya (English)	31
The Potato (English)	62
The Rose (English)	47
The sweet flag (English)	22
The tru barberry (English)	29
The Upas tree (English)	25
The Wealth of India 1949-53 (English)	20
Thikri (Vernl)	29
Thimbawmagyi (Burm)	110
Thinduka (S)	80, 117
Tholkuri (B)	32
Thorapimple (Bo)	107
Tikhi (Bo)	103
Tikhur (H & B.)	98
Til (S , H , B. & Bo)	84
Tilia Kachnag (Vern)	22
Tinduka (S.)	104
Tinis (B)	71
Tinisasegandun (S.)	71
Tisi (H & B)	67
Tita (B)	35
Titakunga (B)	37
Tithwan (Vern.)	26
Tiwas (Bo.)	71
Torai (H. & Bo.)	40, 75
Toria (Vern)	65
Trayamana (S.)	104
Tridhara-sehund (H.)	88
Triputi (S)	71
Truffles (English)	56
Tsjmul (M)	118
Tubah (Malay)	71
Tudovullay (M)	100
Tukhmalanga (P)	106
Tulkapyre (Tam)	99
Tumala (S)	80, 117
Tumburu (S.)	68
Tun (H)	31
Tuni (Bo)	31
Tunumaram (M.)	31

U

Udala (Bo)	115
Udsalap (H)	63
Udumbara (S)	104
Ughaiputtai (M)	80
Ukshi (Bo)	30
Ulatkambal (H & B.)	21
Umatai (M.)	36, 83
Unankodi (Tam)	109
Undi (Bo)	66
Uravu (Mal)	99
Utangan (Bo.)	29
Utaajan (H)	29
Utarni (Bo. & M.)	36
Utran (H)	36

V

Vachnag (Bo)	22, 62
Vada ganneru (M)	81
Vadam-Kottai (M.)	72
Vahisa (S)	99
Vaj (Bo)	22, 93
Vajrakantaka (S)	88
Vakuchi (S)	45
Vallaimurdu (M.)	73

	PAGE
Vallaipundu (M.)	23, 91
Vallarai (M)	27, 32
Val milaku (M.)	44
Valuluwal (M)	32
Vana-palandam (S)	50, 92
Vanatik-tika (S)	120
Vaduka parni (S)	32
Vansa (S.)	27, 94
Vantulshi (S)	119
Vardara (Bo)	100
Varuna (S)	98
Vasaka (S)	23
Vasanvel (Bo)	97
Vashambu (M.)	23, 93
Vashanavi (M)	22, 62
Vata (S)	104
Vegetable Drugs (English)	253
Vegetable drug resources (English)	253
Vellai-maruda-maram (M)	49
Vellerku (M)	81
Vembu (M.)	74
Verbena Family (English)	84
Verenda (B)	46, 89
Vetasa (S)	102
Vettilai (M)	44
Vhaneri (Bo)	85
Vilayati afsantin (H & Dec)	26, 77
Vilayati mehdi (H.)	106
Vilayeti tamaku (H. & B)	83
Vilayati-vakhandi (Bo)	81
Visha (S.)	22, 26
Vishala (S)	75
Vishamandala (S.)	91
Visha-mushti (S.)	48, 82
Vishnugandhi (S.)	104
Vishomangil (M.)	91
Virali (M)	69
Vitaraka (B)	100
Volvox (English)	53
Vrischikali (S.)	89
Vuir (Kash)	107
Vulgarly phansamba (English)	55
W	
Wagata (M)	77
Walnut Family (English)	90
Walsura (Bo. & M)	68
Wee-chhata (Beng.)	55
White mustard (English)	65
Wormwood (English)	26
St John's Wort Family (English)	66
Y	
Yam Family (English)	91
Yamani (S)	50
Yashti-madhu (S)	39
Yavanala (S)	95
Yebruj (B)	83
Yellucheddie (M)	84
Yetti (M)	40, 48, 82
Ysjudemaram (M)	98
Z	
Zira (H)	98
Zira (Vern)	31
Zirishk (Vern)	29
Zufah (Vern.)	98